Advances in Physics Theories and Applications
ISSN 2224-719X (Paper) ISSN 2225-0638 (Online)
Vol 14, 2013
www.iiste.org

Anagrammatic Knowledge Kills Truth: Quantum World Is Also Simulated And Holographic

Quantum Control Of Spin, Decoherence, And Noise, Negative Curved Space Time, Logic Of Sense And Other Topics: Ánthrōpos Métron Ab Uno Disce Omnes: From One, Learn All -Man [Is] The Measure [Of All Things]

[1]Dr. K.N.P.Kumar

Post Doctoral Scholar, Department of Mathematics, Kuvempu University, Janna Sahyadri, Shankaraghatta, Karnataka, India

[2]Prof. B.S .Kiranagi

Distinguished UGC Professor Emeritus, Department of Mathematics Mysore University, Karnataka, India

[3]Prof. J.S.Sadananda

Director, DACEEFO, Professor Department of Political Science, Local Promoter CSLC, Chairman Department of ACEEFO Jnana sahyadri, Kuvempu University, Shankaraghatta, Shimoga District, Karnataka, India

[4]Prof. S.K. Narasimha Murthy

Chairman and Professor Department of Mathematics and Computer Science, Jnana sahyadri, Kuvempu University, Shankaraghatta, Shimoga District, Karnataka, India

@@

Abstract: We give a consubstantiated and consolidated model for the items classified in the paper and discuss various aspectionalities and attributions of the system as stability analysis, Solutional behaviour, and asymptotic analysis. Concomitant theorems are derived. Physical interpretation is provided according to, **agnate, akin** to situational significance wherever possible and explanations in abstract philosophical points of view are expatiated, notwithstanding the appalling task of the expatiation at some points. We hope more erudite scholars would fill in the blanks with redoubtable, illustrious, imminent puissance. Character constitution and predicational anteriorities are also specified at various points where it was deemed necessary. In literature Hyper entanglement-assisted Bell-state analysis (S. P. Walborn S. P´adua and C. H. Monken) propose a simple scheme for complete Bell-state measurement of photons using hyper entangled states - entangled in multiple degrees of freedom. In addition to hyper entanglement, this requires only linear optics and single photon detectors, and is realizable with current technology. At the cost of additional classical communication Bell-state measurement can be implemented nonlocally possible application of these results to quantum dense coding and quantum teleportation are also done. Decoherence Bounds on Quantum Computation with Trapped Ions (Richard J. Hughes et al) used simple physical arguments to investigate the capabilities of a quantum computer based on cold trapped ions. From the limitations imposed on such a device by spontaneous decay, laser phase coherence, ion heating and other sources of error, they derive a bound between the number of laser interactions and the number of ions that may be used. The largest number which may be factored using a variety of species of ion is determined. Our concentration is manifold in consideration to the variety of scheme and doxa of multifarious subject matter involved. . It is to be stated in ineluctable and inexorable and unmistakable terms that all the variables given are different from one section to another and it is presented for

seemingly leviathans typing work involved as also the technicalities abstraction of solidarity such as subscripts and superscripts containing thousand and odd notation. Models are presented in piece meal of nine modules for this reason, raison d'etre and fons et origio. I beg of the readers to pardon me on this count. In the eventuality of modules wherein there is no connection at all or those where is there is apprehensive doubts about the connectivity accentuation and attrition coefficients are taken to be zero. In addition, not wholly or in full measure the dove tailed calculations are given for the first modules and the left out for the remaining one, because of the difficulties in uploading. Dovetailed and accoutrement and apputernential calculations are to be read along with the bastion and stylobate which are quite explicit both in its thematic and discursive form.

Key Words: High-Fidelity Quantum Computation, **qubit register progressive**, Negatively Curved Space-Time, Quantum Formalism, Holographic Renormalization Group, Artificial Intelligence, Casimir Energy, Zeno's Paradox, Nagarjuna's Tetra Lemma, Identity and Contradiction, Qualia, Lisi's E(8) Theory; holographic quantum computation, Neuron DNA, Nirguna Brahman, Heidegger Dasein,

@@@

QUOTES:

It is undeniable that some organisms are subjects of experience. But the question of how it is that these systems are subjects of experience is perplexing. Why is it that when our cognitive systems engage in visual and auditory information-processing, we have visual or auditory experience: the quality of deep blue, the sensation of middle C? How can we explain why there is something it is like to entertain a mental image, or to experience an emotion? It is widely agreed that experience arises from a physical basis, but we have no good explanation of why and how it so arises. Why should physical processing give rise to a rich inner life at all? It seems objectively unreasonable that it should, and yet it does. If any problem qualifies as the problem of consciousness, it is this one.

(Chalmers, 1995, p. 201)

Many times I am asked the obvious yet pertinent question about the title of my longest book: "so who or what is tickling the ticklish subject?" The answer, of course, is: the object - however, which object? This, in a nutshell (or, rather, as a nut within the shell), is the topic of the present book. The difference between subject and object can also be rendered as the difference between the two corresponding verbs, to subject (submit) oneself and to object (protest, oppose, make an obstacle). The subject's elementary, founding, gesture is to subject itself - voluntarily, of course: as both Wagner and Nietzsche, the two great opponents, were well aware of, the highest act of freedom is the display of amor fati, the act of freely assuming what is otherwise necessary. If, then, the subject's activity is, at its most fundamental, the activity of submitting oneself to the inevitable, the fundamental mode of object's passivity, of its passive presence, is that which moves, annoys, disturbs, traumatizes us (subjects): the object is at its most radical that which objects, that which disturbs the smooth run of things. 1 The paradox is thus that the roles are reversed (with regard to the standard notion of the active subject working on the passive object): the subject is defined by a fundamental passivity, and it is the object, from which movement comes, i.e., which does the tickling. But, again, WHICH is this object? The answer is: the **parallax object**.

(**The Parallax View........ The Stellar Parallax: The Traps of Ontological Difference........ The Subject, this "inwardly circumcised Jew" - the von Paulus version - part I ....Slavoj Zizek The Tickling Object)**

An old argument about whether a computer can think: Descartes' argument that no mechanism could use language appropriately or solve problems bears an uncanny resemblance to the test proposed by the mathematician Alan Turing for deciding whether a computer can 'think'. In this test a number of judges are required to distinguish

between a computer and a human using only the replies that they provide to any questions put to them. To eliminate irrelevant cues all questions and answers are typewritten and the judges are placed in a separate room. If the ability of the judges to identify the computer (on the basis of this linguistic exchange) does not differ significantly from chance, then, Turing asserts, the machine may be said to 'think'. The main difference between Descartes and Turing is that Descartes believes machines will always fail this test whereas, 300 years later, Turing thinks they will eventually succeed

(Understanding Consciousness Max Velmans)

If there were machines which have a resemblance to our body and imitated our actions as far as it was morally possible to do so, we should always have two very certain tests by which to recognise that, for all that, they were not real men. The first is that they could never use speech or other signs as we do when placing our thoughts on record for the benefit of others. For we can easily understand a machine's being constituted so that it can utter words, and even emit some responses to action on it of a corporeal kind, which brings about a change in its organs; for instance, if it is touched in a particular part it may ask what we wish to say to it; if in another part it may exclaim that it is being hurt, and so on. But it never happens that it arranges its speech in various ways, in order to reply appropriately to everything that may be said in its presence, as even the lowest type of man can do. And the second difference is, that although machines can perform certain things as well as or perhaps better than any of us can do, they infallibly fall short in others, by which means we may discover that they did not act from knowledge, but only from the disposition of their organs. For while reason is a universal instrument which can serve for all contingencies, these organs have need of some special adaption for every particular action. From this it follows that it is morally impossible that there should be sufficient diversity in any machine to allow it to act in all events of life in the same way as our reason causes us to act**.**

(Descartes quoted in Haldane and Ross, 1931; also cited in Flew, 1978, p. 127 and Max Velmans)

The self-conscious mind is actively engaged in reading out from the multitude of liaison modules that are largely in the dominant cerebral hemisphere. The self-conscious mind selects from these modules according to attention and interest, and from moment to moment integrates its selection to give unity even to the most transient experiences. Furthermore, the self-conscious mind acts upon these modules **modifying their** dynamic spatio-temporal patterns. Thus it is proposed that the self-conscious mind exercises a superior interpretative and controlling role. A key component of this hypothesis is that the unity of conscious experience is provided by the self-conscious mind and not by the neural machinery of the liaison areas of the cerebral hemisphere. Hitherto it has been impossible to develop any neurophysiological theory that explains how a diversity of brain events come to be synthesized so that there is a unified conscious experience of a global or gestalt nature.

(Eccles, 1980, p. 49 also quoted in Max Velmans)

Quantum mechanics deals with the observed behaviors of macroscopic systems whenever those behaviors depend sensitively upon the activities of atomic-level entities. Brains are such systems. Their behaviors depend strongly upon the effects of, for example, the ions that flow into nerve terminals. Computations show that the quantum uncertainties in the ion-induced release of neurotransmitter molecules at the nerve terminals are large (Stapp, 1993, p. 133, 152). These uncertainties propagate in Conscious souls, brains and quantum mechanics 17principle up to the macroscopic level. Thus quantum theory must be used in principle in the treatment of the physical behavior of the brain, in spite of its size.

(Stapp, 2007a, p. 300)

A paradox for quantum dualist interactionism: how could a visual conscious experience actualise its own prior cause? Although it only takes one photon arriving at the retina to trigger a visual experience, the vitreous humour of

the eye scatters light, so that even under optimal conditions, it requires from five to eight photons to trigger that experience (see Chapter 8). Studies of the visual system and other sensory systems also make it clear that conscious experiences do not arise at the instant that input stimuli arrive at the cortex. Instead, a period of preconscious processing occurs that involves neural activation, analysis, synthesis and so on. This takes time. Experiments reviewed by Libet (1996), for example, suggest it takes at least 200 milliseconds for neural states to form in a way that is adequate to support a tactile conscious experience (see Chapter 10). In short, within neuropsychology, consciousness of external events is thought to take place later in time than the occurrence of the events themselves and, perhaps, hundreds of milliseconds later in time than the arrival time of the neural activation they produce at the sensory cortex. The relevance of quantum mechanics to an understanding of how such systems operate is not obvious. According to the Stapp/Von Neumann interpretation a photon is only actualised once it results in a visual experience. But how could a resulting conscious experience 'actualise' its own prior cause? This would seem to require backward causation in time! And are we also to say, following Von Neumann and Stapp, that preconscious processes responsible for creating neural conditions that are adequate to support a conscious experience also only become 'actualised' once the visual experience arises? Note that such paradoxes would remain if there is any delay between the input stimulus and the consequent experience. That is, the problems remain even if Libet's figure of a minimal 200-millisecond delay turns out to be an overestimate.

(Understanding Consciousness Max Velmans)

Acknowledgements: I humbly and sincerely thank Wikipedia, Google search and all the people and sources who have been responsible for this work. In the eventuality of any omission in references, I make a sincere entreat, earnest beseech, fervent appeal to kindly pardon me. Literally thousands of documents were perused and many of them were not taken in to consideration. Kindly pardon me in case of error of non reference.

Introduction:

CLASSIFICATION OF THE PROTAGONIST OR ANTAGONIST LAWS BASED ON THE CHARACTERSTICS AND PENCHANCE, PREDILECTION, PROCLIVITY AND PROPENSITIES OF THE SYSTEMS UNDER INVESTIGATION: I ACKNOWLEDGE IN UNMISTAKABLE AND UNAMBIGOUS TERMS THE HELP OF STANFORD ENCYCOPEDIA, KANTS WRITINGS, AND DELUZE'S LOGIC OF SENSE, GOOGLE SEARCH AND WIKIPEDIA. CONCERTED AND ORCHESTRATED EFFORTS ARE MADE TO UNIFY THE VARIOUS SUBSTANTIVE STATEMENTS ABOUT THE NECESSITY OF PERCEPTION, REALITY, CONSIUSNESS, TOTALISM, STRATIFICATION AND OTHER FACTORS OF CARDINAL IMPORTANCE. (EMPHASIS MINE)

Autopoesis is both a Concept About Self-Maintaining Systems, Biological Especially ones, and an Epistemological Theory About cognition and Self-Reference. Humberto Maturana and FJ Varela Concept of the Proposed autopoesis (as Opposed to allopoetic) as the Organization of Living Things, Which maintains lasting Their Unity and wholeness in the midst of continual turnover of constituents, specifically through self-production. For them, autopoesis is the necessary and sufficient condition for the minimal units of life. **Autopoesis** defines the minimal mechanistic features of the simplest possible living entity. It describes "how life works." All Living Systems are autopoetic, physical and All Systems, if autopoetic, CAN be said to be Living. An autopoetic system is defined as an Entity That can be Clearly Distinguished and is Realized through a closed Organization of Production Processes such That the products of the process interact in the same way as the organization that produces them and a discernable topological boundary emerges as a result of these same processes. (Artificial Life, p.131) An autopoetic unit is a system that makes itself, through a network of interactions that take place within its own well-defined boundaries. (Cf. autocatalysis) Or, as Maturana puts it in Autopoesis and Cognition, "The living organization is a circular organization which secures the production or maintenance of the components that specify it in such a

manner that the product of their functioning is the very same organization that produces them.” (P.48). The goal of autopoetic systems is the maintenance of their internal organization. Thus self-completion or individuation is distinctive to the living state. The capacity of active self-maintenance and self-generation that underlies regeneration, reproduction, and healing, are ways that organisms become distinctive wholes. The autopoetic system is not entirely autonomous, however. It must be structurally coupled to the World, to the Medium in Which IT exists. According to Katherine Hayles, the "Second Wave" of cybernetics Started with an article Entitled "What the Frog's Eye Tells the Frog's Brain," in Which the authors , including Humberto Maturana, showed that a frog's visual system does not so much represent reality as construct it. As the authors noted, "The frog's eye speaks to the brain in a language already highly organized and interpreted instead of transmitting some more or less accurate copy of the distribution of light upon the receptors." (How We Became Posthuman, p.131, p. 135) (See vision) **Humberto Maturana** developed the epistemological implications of this study by stressing the internal organization and circularity of cognition and of living systems in general. It is this recursivity which characterizes "second order" cybernetics, which entails the observation of observation (in a formal and not psychological frame.) Thus, second-order cybernetic systems are both closed and open. Autopoetic Systems are closed at the Level of Operation or Organization, but are at the Open Level of Structure. For Varela, "An autopoetic Machine Continuously Generates and specifies ITS Own Organization through ITS Operation as a system of Own Production of ITS Components." (Cf. Kant's distinctions between mechanism and organism in mechanism / vitalism) Note, however, that the theory of autopoesis defines the relations of the components without reference to the whole. Fylix Guattari argues for a machinistic autopoesis, "the non-discursive and self-enunciating nexus of the machine." (As a critique of any unifying idea of the signifier - esp. The Lacanian one) .This autopoetic nexus "wrests it from structure." He cites Varela's description of a machine as "the ensemble of the interrelation of its components, independent of the components themselves ". Despite the fact that Varela reserves the qualification of "autopoetic" to the biological domain, Guattari extends the notion accross the range of social and technical processes that he includes in "machinism." Guattari is not so interested in autopoesis as unitary individuation but rather in entities that are "evolutive and collective." According to Guattari, "We envisage THUS CAN autopoesis under the heading of an ontogenesis and phylogenesis specific to the mecanosphere .That **superimposes Itself on the** Biosphere." Following the distiction ESTABLISHED by Maturana and Varela in **Autopoesis and Cognition**, Guattari distinguishes allopoetic from autopoetic Machines. The former produce something beside them, and perform a function useful to agents outside themselves, whereas the latter continuously engender and specify their own organization and their own limits. ("Machinic Heterogenesis" in Rethinking Technologies, p. 17) He also connects the idea of the allopoetic machine with hypertext. ("On Machines" in JPVA no.6) But if Varela's emphasis is on autopoesis as unitary individuation, Guattari stresses a more collective machinism, a "diagrammatic rhizome" of ontological Heterogenesis. (See machinic phylum) See Also Network or Assemblage THUS autopoetic the Basic Unit is closely linked to the cell, defined by ITS boundary, with a Metabolic Network Inside. It is self-bounded, self-generating, and self-perpetuating. The potential for self-reproduction and evolution are consequences of the initial definition of autopoesis. According to Pier Luigi Luisi, the minimal autopoetic unit requires a process of self-generation and a process of decomposition. If the rates of the two processes **are equal,** the system will self-perpetuate in a steady (homeostatic) state. If the rate of the generative process is greater, the unit will tend to grow, and if there are limits to its size, it will tend to self-replicate. In the other case, where the rate of decomposition is greater, it will self-implode. These are the Three Basic Kinetic states. Luisi Claims That according to this Definition, viruses are non-Living. They are certainly not free-living. They are parasites that invade cells, co-opt the cell's metabolic machinery to accomplish their own self-reproduction, escape the host cell, and invade another ... NOR is a robot that assembles Other robots out of parts built by Other Machines, Because in both cases reproduction is not occurring within the Domain of the Organizational Unit. Autopoetic Systems have the special ability to discriminate what is Part of the system and What is not, which provides a link to a theory of observation. System is an autopoetic a Self-Describing system? When Brought to Bear on Theory and Criticism, autopoesis establishes a new Epistemological Condition. The Observor is not separate from what is being observed, but is structurally coupled to the phenomenon she sees. Stating that "everything said is said by an Observor," Maturana and Varela develop a theory of the observation of observation, in which the world is not given - as in the traditional representationalist frame - but is rather brought

forth in the dynamic interaction of observer and observed. "Our intention is to bypass entirely this logical geography of inner versus outer by studying cognition not as recovery or projection but as embodied action." (Tree of Knowledge, p. 172) Autopoesis leads to a theory of Observor in which there is no route back from the act of observing to the data that was used to generate the theory in the first place. (Hayles) Niklas Luhmann describes Complex Systems Self-referential as "hypercomplex," in the Ways That They make Distinctions Between Themselves and Their Environments, but Also internalize the Distinction in a paradoxical "Re-entry" of the Distinction into the Distinguished. Autopoesis: (See The Tree of Knowledge, by HR Mataruna and FG Varela. and also "Autopoesis: the Organization of Living Systems, its Characterization and a Model" Biosystems 5. See also Francisco Varela, Autonomie ET Connaissance), Pier Luigi Luisi, " Defining the Transition to Life: Self-replicating Bounded Structures and Chemical Autopoeisis ", in Stein & Varela, eds. Thinking About Biology. Adaptation ---------- In order to create HIS Theory of evolution, Darwin HAD to Take a Step in Revolutionary Thinking About Organism and Environment. Previously there had been no clear demarcation between internal processes and external ones. Darwin created a dramatic rupture between the internal processes that generate the organism and the external processes, the environment, in which the organism must operate. Adaptation became both a process and a measure of "fitting" the organism to the environment. The organism's interaction with the environment - with "nature" - sets up a feedback process whereby nature "selects" one design over another, depending on how well it solves an adaptive problem. If a modification enhances the ability of a design to cause its own reproduction, a positive feedback occurs and an adaptive design will replace all alternative designs in a population. If it results in fewer (or no) offspring, a modification will set up a negative feedback, and a design will become extinct. According to strict usage in evolutionary biology, a character is an "adaptation" for a particular task only if it can be shown to have evolved in specific ways to make it more effective in the performance of that task, and that the change has occurred due to the increased fitness that results. Perhaps only some problems are adaptive problems, that is, problems whose solution can affect reproduction, however distally. Avoiding predation, choosing nutritious foods, Finding a mate, and Communicating with others are Adaptive Problems That Our hominid ancestors would have faced. Johannes von Uexkll, at the beginning of the twentieth century, criticized the Darwinist separation of organism and environment. For Uexkll, "Every animal carries its environment about with it like an impenetrable shell, all the days of its life." (Quoted in Ernst Cassirer, The Problem of Knowledge, p. 202) Every creature has its own private world of perception and activity, prescribed once and for all by its structure, by its type of receptors and effectors. This environment-world is the animal's Umwelt. It is constituted by a more or less broad series of elements that he calls "carriers of significance" (Bedeutungstrdger) or of "marks" (Merkmaltrdger), which are the only things that interest the animal. Every animal is bound to its environment by "the circle of functions." For Uexkll, "The protoplasm that made an earthworm made it for an earthworm's activities, and set it in an earthworm's world." For Uexkll, a unitary world in which all living beings are situated does not exist. - Nor do a space and a time that are equal for all living things. (See Agamben, Open, p.40) We may see an Umgebung, an objective space in which we see a living thing moving, but in reality the Umgebung is our own Umwelt, to which Uexkll does not attribute any particular privilege, and which can vary according to our point of view. Uexkll sought to draw out the consequences of anatomical type defined as a structure of (static) relationships and to resurrect the program of an "idealistic morphology." In his Theoretical Biology and Umwelt und Innenwelt der Tiere, Uexkll describes the "theory of the living being" as a pure natural science with only one goal: "The research into the blueprint (Bauplan or structural plan) of living beings, their origin, and their results." The relations between the functional spheres of the organism are the basic phenomenon of life. The stimuli of the external world, which an animal is able to receive by virtue of its blueprint, constitute the only reality present for it, and by virtue of this physical limit, it closes itself to all other spheres of existence. (Ernst Cassirer, The Logic of the Cultural Sciences, pp.23-24) Whatever ITS context, Adaptation is the Accumulated output of selection and involves modification of Progressive Structures repeated through the Structural Actions of modifiers, or "operators." (Eg. chromosomes are structures which are modified by mutation, recombination, etc.) Adaptation requires a specific performance measure (eg fitness). The defining feature of Darwinian adaptedness is that it results in the survival of more similar offspring. But Darwinian Natural selection is not just Differential reproduction Differential reproduction but is Due to the Adaptive superiority of Those Who Leave More Offspring. In Adaptation in Natural and Artificial Systems, John Holland describes the Problems of

Adaptation as a form of optimization in which information must be exploited as it is acquired, so that performance improves apace. (See genetic algorithms) How can adaptation occur efficiently? Not through an "enumerative" plan, which simply tries out every structure in an order that is unaffected by the outcome of previous tests. It is extremely inefficient and the time required soon exceeds the age of the universe in systems of even modest complexity. Requires Adaptation "Robust" Plans That make efficient use of the History of Their Interactions with an Environment, within a framework of "Enough time." Relative adaptedness Refers to the relation of one Organism Being Adapted Better than Another (of the Same interbreeding Population) to a particular environment, and is the fundamental concept of neo-Darwinian theory. Adaptation works directly on phenotypes, not on genes or genotypes; so adaptedness is an organismic not a genetic quality. For Mathematical Genetics, the Fitness of a phenotype is the number of ITS Offspring Which Survive to Reproduce. Difficulty One with the way Adaptation is defined in Population Genetics is That Increased Reproductive Success does not describe the Correlations Between traits and Their Contributions to organismic well- being on the basis of their design or function. In addition, because of the possibilities of conversions of functions, or "exaptations," adaptations must be described historically, and their histories are rarely simple teleologies directed towards contemporary fitness. Forms CAN just as well be Vestigial or incidental. Richard Lewontin criticizes the Notion That Organisms Adapt to "fit" specific Requirements of the Environment through the use of Terms such as "Ecological niche," Which imply a kind of Ecological That space with holes are filled by organisms whose properties give them the right "shape" to fit the holes. In this framework, the study of the diversity of organisms becomes a map of the space of the environment, just as sprinkled iron filings map a magnetic field. Lewontin argues that it is time to reconsider the relationship between outside and inside, between organism and environment and suggests that the actual process of evolution is better captured by the metaphor of construction. (The Triple Helix, p. 48) While there is a physical world outside of an organism that would continue to exist in the absence of the species, the environment does not exist without species. Organisms determine which elements of the external world are put together to make their environments and the relations between those elements that are relevant to them. The space of the environment is determined by the activities of the organism itself, not the other way around. In addition to this more selective process, organisms actively construct (and destroy), in the literal sense of the word, the world around them. Every species, not only Homo Sapiens, is in the process of destroying its own environment by using resources that are in short supply and transforming them into a form that cannot be used again by individuals of the species. But every such act of consumption is an act of Production for Other Species. Evolutionary biologists describe the **Relationship Between Adaptation and ongoing** Changes in the Environment as the "Red Queen Hypothesis," named After the Queen in Through the Looking Glass Who found she HAD to keep running just to stay in place. According to Lewontin's constructionist view, the world is changing because the organisms are changing. The Red Queen's running only makes the problem worse. (P.58) For Lewontin, IT is a General Principle of Historical development of Biological Systems That the State of Possible Conditions Which make the **Coming into Being of the system are abolished** by IT. http://www.christianhubert.com/hypertext / affect.html Affect ------

In 1892, Charles Darwin categorized human affects into **seven or eight discrete expressions**, each with its own facial display: happiness, sadness, fear, anger, disgust, surprise, interest, perhaps shame, and their combination. (The Expression of the Emotions in Man and Animals) According to Darwin, "The same state of mind is expressed throughout the world with remarkable uniformity." He postulated that these **innate patterns** of feeling and facial display evolved as social signals "understood" by all members to enhance species survival. (Cf ritualization) For Darwin, our expressions of emotion are universal (that is, innate not learned) and they are products of our evolution. They are Also, at Least to some extent, involuntary, and feigned emotions are Rarely Fully convincing. Neither Our expressions NOR Our emotions are Unique to Human Beings; Other Animals have some of the Same emotions, and some of the expressions Shown by Animals ressemble our own. For Darwin, "He who admits on general grounds that the structure and habits of all animals have been gradually evolved will look at the whole subject of expression in a new and interesting light." (Introduction to First Edition) Thus, while the illustrations below are found in the chapter on "Special Expressions of Animals", Darwin points out at the outset that "With mankind some expressions, such as the bristling of the hair under the influence of extreme terror, or the uncovering of the teeth under that of

furious rage, can hardly be understood, except on the belief that man once existed in a much lower and animal-like condition. " (Intro.) In previous Centuries, Facial expressions HAD Been Explained in Terms of rhetoric or Language, as in Charles Le Brun's 1698 treatise on the expression of the passions. Furthermore, Most Studies of Facial Expression Were Part of the Study of physiognomy - the Recognition of Character through the Study of the Forms of the permanent Features. Did Darwin's Treatise Bypass Expression Language and Extend Across Species? Or did his work, as some scholars have suggested, indicate a **split status in the human face**, becoming simultaneously a symptom of an organism's anatomical and physiological functioning and, in its relative impenetrability, the mark of the success or failure of a process of self -mastery and control implicit in the social construction of a normative individual? (Jean-Jacques Courtine and Claudine Haroche, Histoire du visage: Exprimer et taire ses ymotions, Ref. J. Crary in Suspensions of Perception, p.99) Modern Accounts of Cultural Expression Recognise "display rules" That Regulate the expressions of Emotion. The anthropologist Paul Eckman filmed the expressions of American and Japanese students as they watched gruesome footage of a primitive puberty rite. When the white-coated experimenter was in the room, Japanese students smiled politely during scenes that made the Americans recoil in horror. But when the subjects were alone, the Japanese and American faces were equally horrified. Although emotional Linguistic Expression is not, IT is nonetheless communicative, and is expressive of Internal states. Animals Do Distinguish Between Public and Private? Affect Versus Emotion? Sigmund Freud Affect Related to Instinct. He viewed affect as the qualitative expression of the quantity of instinctual energy and of its fluctuations. (Laplance and Pontalis) In The Interpersonal World of the Infant, the Child psychologist Daniel Stern describes infancy Primarily in Terms of Affect. Stern describes infant experience as more unified and global than that of adults. According to Stern, infants "take sensations, perceptions, actions, cognitions, internal states of motivation, and states of consciousness, and experience them directly in terms of intensities, shapes, temporal patterns, vitality affects, categorical affects, and hedonic tones. (p.67) According to Stern, early in life, affects are both the primary medium and the primary subject of communication. Stern distinguishes "vitality affects" from "categorial affects." The former often take the form of a "rush." They are About a way of feeling, not a specific content of feeling. Music and Dance convey vitality Affects. For Stern, Affective Experiences enter the intersubjective Domain through "Affect attunement," Primarily, but not exclusively in the Mother / Infant Dyad. Stern's Descriptions of affect ressemble Deleuze and Guattari's references to "intensities." (See smooth / striated) In "the autonomy of affect," Brian Massumi takes up Deleuze's identification of affect with "intensity," and opposes affect to emotion. For Massumi, emotion is qualified intensity - "the conventional, consensual point of insertion of intensity into semantically and semiotically formed progressions, into narrativizable action-reaction circuits, into function and meaning." Emotion expresses the "capture and closure" of affect, as well as the fact That "something Has Always and again ESCAPED." This Escape of Affect is "the Perception of one's Own vitality, the sense of one's aliveness." ("The Autonomy of Affect, in Observing Complexity, p.285) In Massumi's Account, Emotion and affect have different logics, and cultural theory would benefit from "an asignifying philosophy of affect." (P.277.) Massumi takes issue with Frederic Jameson's claims about the "waning of affect." (See below) Instead, he claims that "our condition is characterized by a surfeit of it." With Reference to Simondon's concepts of "implicit form," (See form / Matter) Massumi identifies the autonomy of Affect as ITS Participation in the virtual, in IT Openness. In the Civilizing Process, Norbert Elias describes a Historical Process of Restraint and moderation of affect that took place with the rise of court society. This new formation is apparent in the transformations of table manners and the new thresholds of shame and embarrassment, in the "invisible wall of affects" which arises between one human body and another. According to Elias, the "civilizing process" creates new distances between adults and children and requires that children rapidly attain the advanced level of shame and revulsion that has occurred in the West over many centuries. Impulses that were once uncensored, such as "holding out the stinking thing for the other to smell" have disappeared from the waking consciousness of adults under the pressure of conditioning. Only psychoanalysis uncovers these unsatifiable desires, or "infantile" residues, in the form of unconsious, or dream content. For Elias, the superego is the Psychological correlate of these transformations of Society. For Freud, the dream work Often Affect Separates from manifest content. This separation was for Freud one of the defining features of repression. The analysis of dreams can reveal how virtually every dream expresses certain infantile wishes and the conflicts precipitated by them. However, the screening of affect can work in both directions. Not Only CAN infantile wishes

be forbidden, but Contemporary ones CAN be screened as well, Especially Those Which Arise in a Transference Relationship. Freud's theories of Neuro-Psychoses (1894) proposed a general theory of **neuroses by classifying them according to three defense mechanisms: Transformation of Affect in Conversion Hysteria, (**Which Psychological Problems in Medical induce symptoms) displacement of Affect in obsessions, exchange of Affect in Anxiety neuroses and Melancholia. HIS In Studies of Hysteria, Freud Explored the Relationship Between Affect and Memory, Especially Instances When a memory excites an affect that it had not excited as an experience. He describes episodes that only became traumatic after the event. For example, in the Project, He describes a Young Girl (Emma Eckstein?) Who.... For theories of Ahnlehnung "anaclitic" or cathexis - the way Affect is "propped up" by Objects - See sexuality Frederic Jameson Refers to the "waning of Affect" in postmodern Culture. Jean Baudrillard proposes to document IT. Also Does IT lead to a reduced Desire for Truth? http://www.christianhubert.com/hypertext/complexity.html Complexity ----- ---------- "Our impression That life evolves Towards Greater Complexity is Probably Only a Bias Inspired by Parochial focus on Ourselves." (Stephen Jay Gould) The Awakening of Modern Interest in Complexity as a science Began in Vienna in 1928, with Ludwig von Bertalanffy's largely descriptive graduate thesis on living organisms as systems. A few years earlier Alfred North Whitehead had described his similar vision of a "philosophy of organism" in Science and the Modern World. Whitehead describes his theory of the organic conception of nature as based on "self-knowledge of our bodily event." This total bodily event is on the same level as all other events, except for an unusual complexity and stability of inherent pattern. (Science and the Modern World, p. 73) Today the Study of Complex, Self-Organizing Systems is the "Related Opposite" of the Study of chaos: In Self-Organizing Systems, orderly Patterns emerge out of Lower-Level randomness, in Chaotic Systems, unpredictable Behavior Emerges out of Lower-Level deterministic rules. In Contemporary Accounts of Complex Adaptive Systems, a Great number of Independent Adaptive agents are interacting with each Other in a kaleidoscopic array of simultaneous Nonlinear Interactions. Chaotic behavior at one level can give rise to distinctive order at the next. Feedback and feedforward Loops permeate Their Decentralized Structures. (Cf Network) The richness of These Interactions Allows the system as a Whole to undergo spontaneous Self-Organization. "What is Being Recognized within the" Sciences of Complexity "is ... That there are characteristic types of order that emerge from the interactions of many different components ... especially in relation to the large-scale aspects of structure or morphology, and the patterns in time that constitute organismic behavior. " (Goodwin, P. IX) Economies, ecologies, Immune Systems, Developing embryos, and the Brain All are examples of Complex Adaptive Systems. The Impact of These Systems Usually Centers on the Aggregate Behaviour, Which Often Feeds back to the Individual parts, and evolves over time. These are Adaptive Systems. They are not chaotic, nor are they merely complex in a static sense. According to some authors, complex systems have somehow aquired the ability to bring chaos and order into a special kind of balance, often called the edge of chaos. (See phase boundary) Stuart Kauffman often characterizes the three dynamical regimes as solid or frozen (the ordered regime, as gas (the chaotic regime), and as liquid (the complex regime). (See phase boundary) These systems face perpetual novelty, Generally and operate far from a global or Optimum Equilibrium. (cf dissipative Systems) In Seeking to Adapt to changing Circumstance, Complex Systems anticipate. For Stuart Kauffman, "Selection achieves and maintains Complex Systems poised on the boundary, or edge, Between order and chaos. These (poised) systems are best able to coordinate complex tasks and evolve in a complex environmnent. "(P.XV) (see evolution) Stephen Pinker echoes Darwin in claiming that" Natural selection is the only scientific explanation of adaptive complexity. .. "(the Characteristic of)" any system composed of many interacting parts where the details of the parts' Structure and Arrangement Suggest Design to Fulfill some function. " Murray Gell-Mann describes as Complex Adaptive Systems "Pattern-Recognition Devices That seek to find regularities in experience and compress them into schemata. "In his book on vision, David Marr proposes that three levels of understanding are required for the analysis of a complex information-processing system: At one extreme, the top level, is the abstract computational theory of the device, in which the performance of the device is characterized as a mapping from one kind of information to another, the abstract properties of this mapping are defined precisely, and its appropriateness and adequacy for the task at hand are demonstrated. At an intermediate level is the choice of representation for the input and output and the algorithm to be used to transform one into the other. At the other extreme are the details of how the algorithm and representation are realized physically - the detailed computer architecture, so to speak. (Pp24-5) For Marr, "These

three levels are coupled. The speculative object and the practical object of philosophy as Naturalism, science and pleasure, coincide on this point: it is always a matter of denouncing the illusion, the false infinite, the infinity of religion and all of the theologico-erotic-oneiric myths in which it is expressed. To the question 'what is the use of philosophy?' the answer must be: what other object would have an interest in holding forth the image of a free man, and in denouncing all of the forces which need myth and troubled spirit in order to establish their power? (LS 278)

Towards the end of **classificational consummation, consolidation, corporation and concatenation** we take the characterstics of the systems, the predicational interiorities, ontological consonance and primordial exactitude, accolytish representations, functional topology, apocryphal aneurism and atrophied asseveration event at contracted points, and other parameters as the bastion and stylobate of the stratification purposes. Such totalistic entities would have easy paradigm of relational content, differentiated system of expressly oriented actions with primary focus and locus of homologues receptiveness and differentially instrumental activity, variable universalism and particularism, imperative compatibilities and structural variabilities, interactional dynamical orientation, institutionalization and internalisation of **pattern variables(Parsons)**common attitudinal orientation of constituionalisation of internalized dispositions, and a qualitative gradient of structural differentiation and ascribed particularistic solidarity abstraction or interactional dynamics, internal differentiation, structural morphology, comparative variability, normative aspect of expectational prediction, projection and prognostication. There are different superfield formalism and Curci-Ferrari type restrictions for different investigating systems. Equational realisation, consummation and concatenation and consubstantiation are of primal and cardinal and seminal importance Accolytish representation and other **parametricization** could also be taken in to consideration. There is no sacrosanct rule for this. It is to be remembered that there are many **investigatable systems** in the world and the equations these theories specify is applicable to them. Another example is about Off shell nil potency; Three Cf type conditions; 3-form Abelian theory; One Cf condition ; 1-form non abelian Theory B + B9bar) =I (CX C9bar) ; absolute anticommutivity are other properties that is found in many systems. Systemic characterstics thus play a primordial role in the development of the theory be it theoretical or empirical. So their characterstics could form the sentinel for the consubstantiation of the dis integrational purposes which is orderly In case of constants we talk about total invariants like the total **Schrõdinger function**, albeit we have reservations about Everett's syndrome, like for example Total gravity in the universe. However it is to be stated in unmistable terms that **Universal function** is not acceptable to the author, while instead one could talk of the various investigating systems with different conditionalities notwithstanding the fact that say a Non Linear Schrõdinger Equation holds good and the equality holds. The same is true in the Newtonian case of **Total gravity** extant in the world. Universal wave function calls for the conditionalities and functionalities of the investigating agencies and investigating systems so that those could also be taken in to consideration in the stratification scheme. Neutrinos or the protons are the same, but the source and the time factor and the space factor are also to be taken in to attribution and ascription and assignment. Two examples that could be cited are: Personality type refers to the psychological classification of different types of individuals. Personality types are sometimes distinguished from personality traits, **with the latter embodying** a smaller grouping of behavioral tendencies. Types are sometimes said **to involve** qualitative differences between people, whereas traits might be construed as quantitative differences According to type theories, for example, introverts and extraverts are two fundamentally different categories of people. According to trait theories, introversion and extraversion are part of a continuous dimension, with many people in the middle. Myers-Briggs Type Indicator (MBTI) assessment is a **psychometric questionnaire** designed to measure psychological preferences in how people perceive the world and make decisions. These preferences were extrapolated from the typological theories proposed by Carl Gustav Jung and first published in his 1921 book Psychological Types (English edition, 1923). Jung theorized that there are four principal psychological functions by which we experience the world: sensation, intuition, feeling, and thinking. One of these four functions is dominant most of the time. Equational satisfaction and consummation is of paramount and cardinal importance. We have studied discrete Schrodinger equations, NLSE, and Einstein Field equations on the same basis which must provide frame work for further analysis **In the case of Super symmetric Yang Mills Theory integrable systems could be taken in to consideration)**The Coulomb branch of N = 2 **supersymmetric gauge theories** in four dimensions is described in general by an integrable

Hamiltonian system in the holomorphic sense. A natural construction of such systems comes from two-dimensional gauge theory and spectral curves. Ground state at the critical value of the trapping potential two entanglement measures is another example. Notwithstanding the constant ground states the characterstics of the investigating systems form the bastion and stylobate of the situation. Starting from this point of view, we propose an integrable system relevant to the N = 2 SU (n) gauge theory with a hypermultiplet in the adjoint representation, and offer much evidence that it is correct. The model has an **S-duality group** (with the central element −1 of acting as charge conjugation); permutes the Higgs, confining, and oblique confining phases in the expected fashion. **Hegel** also argues strongly against the epistemological emphasis of modern philosophy from **Descartes through Kant**, which he describes as having to first establish the nature and criteria of knowledge prior to actually knowing anything, **because this would imply** an infinite regress, a foundationalism that Hegel maintains is self-contradictory and impossible. Rather, he maintains, we must examine actual knowing as it occurs in real knowledge processes. This is why Hegel uses the term "phenomenology". "Phenomenology" comes from the Greek word for "to appear", and the phenomenology of mind is thus the study of how consciousness or mind appears to itself. In Hegel's dynamic system, it is the study of the successive appearances of the mind to itself, because on examination each one dissolves into a later, more comprehensive and integrated form or structure of mind. We also study more exotic phases. **Supersymmetric Yang-Mills theory and integrable systems Ron Donagia, 1, Edward Wittenb, 2** Kind attention is also drawn to the Boltzmann arrow of time and Everett's many worlds different past and future. [Whitehead's philosophy] is provisionally the last great Anglo- American philosophy, just before the disciples of Wittgenstein spread their mists, their sufficiency and their terror. An event is not only 'a man is crushed': the great pyramid is an event, and its duration for an hour,30 minutes, 5 minutes ..., a passage of Nature, a view of God. What are the conditions for an event so that all is event? The event produces itself in a chaos, in a chaotic multiplicity, under the condition that a sort of sieve intervenes. A global methodology is assumed here. There is **Total energy and Total gravity**. Contextually reflexivity and context justification is most alluded coefficient here. As **Geroch puts it**: Theories consist of enormous number of ideas, arguments, hunches, vague feelings, all arranged in a configuration equivocation. It is the entire body of the material that is the "Theory'. (Written from memory: See Geroch 1978). **Dreyfus and Dreyfus** regard ceteris paribus condition as the universal restriction on rule systems. Classificational rules constitute prevent the regression of prevarication of such axiomatic predications and postulation alcovishness, and is a paradigmatic statement in the salient feature which does not disregard the cause effect definition and other scientific heuristic explanations. Such congressional enterprise is necessary for the totalistic view of the problem. Part of jurisprudence could be considered part of the law itself. Constitutive rules could be disregarded when the priority of the used rules are properly understood. Paradigms guide research even in the conspicuous absence of rules. Characterization **is at the heart of** naturalization. Stratification methodologies are much more vintage and montage than any other fait accompli mutatis mutandis desideratum. Manifest world in its entirety is a **holographic projection of information** embodied in its boundary. Here we are talking of the entire manifest world. But the information obtained is only for a piece of the world for which we have access. Nature's general Ledger is to be down loaded from Neuron DNA. Another factor that is taken in to consideration is **Lancan paradox and sense**, expression and event: proposition and the attribute of the state of affairs, individualistic denotation, institutionalized generalisation, personalized manifestation, organisational individuation, and corporate signification of the systems in general. According to Schrodinger actualization means to extend over a series of ordinary points, to be selected according to the rule of convergence, to be incarnated in a body, to become a state of body, and to be renewed locally namely particle coming back to its original position for the sake of limited new actualizations and extensions. It is in this sense that actualization is collective and individual, internal and external. Geometrical interpretation of the theory of differential equations clearly places in evidence two absolutely distinct realities: (a) there is a field of directions and (b**) topological accidents** which may suddenly crop up in it, as for example the existence of the plane of "singular points" to which no direction has been attached and there are integral curves with the form they take on in the vicinity of singularities of the field of directions…… The existence and distribution of singularities are notions relative to the field of vectors defined by the differential equation. The form of the integral curves **is relative to** the solution of this equation. The forms are assuredly complementary, since the nature of the field is defined by the form of integral curves in its vicinity. But it is no less true that the field of vectors on one hand and integral

curves on the other are two essentially distinct mathematical realities. In this connection of classificational process of the universal theory, it is necessary to recall what Kant had proved. Kant proves that sum of all possibilities excludes all but originary predicates and in this way constitute the completely determined concept of an individual being or objectification being. For only in this case, is a concept is equivalent to a thing. A thing completely determined in and through itself and known as the representation of an individual or a system under consideration. Thus the Universal Theory is **but the form of communication in thought between the cosmic consciousnesses, namely the nature's general ledger, and the finite individualities, namely the histories of** individual debits' and credits. It must therefore be entangled with the play of internationalities and pure retentions. Thus the universal theory has a totality. For example total gravity. Individual systems have characterstics for which say a Non Linear Schrodinger's equation holds, and such a consummation and consolidation of the equation **is built up on the corporation and actualization of the equations in simplistic and individual cases**. Compose is to make up; the parts that compose the whole. The notion of compossibility is thus defined as a continuum of the individual systems that are actualized in accord with the reductionist view which is held dearly in all subjects in this measurement world. In the Saussurean system, potential returns in language as the nonlocalizable excess of signifier over signified that accompanies the local realization of a conventional signifier-signified coupling. This excess is ensconced in the "vertical" dimension of the process (understood very differently than in the present context: as a "synchrony" rather than a temporal disparity, the disparity that is the empty form of time). It constitutes the immanent generative surplus from which the "horizontal" relay to the next linguistic realization is drawn ("diachrony": also different from the "horizontal" lines of encounter at issue here, in that diachrony is not essentially differentiating; the system remains the same across its realizations). An analogous surplus figures in Lévi-Strauss's anthropology as "mana" and its structural equivalents, which constitute the specifically cultural presentation of the staying power of processual remainder (the reserve of potential) in many societies. It is in excess-as-remainder that a reconciling of Deleuze and Guattari's thought and **structuralisms of signification might be found ("mana" converts into Deleuze's "dark precursor" or "object = x"; 1979, 315-324; 1990, 113-115; 1994, 119-123). The deleuzo-guattarian conversion of structuralism entails the signifier and signified generating an excess by escaping down a "line of flight" away from the "despotism" (self-sameness across its variations) of the signifying regime--rather than the excess regenerating signifier-signified following the chain of conventional linguistic realizations**. This amounts to an autonomization of signified-signifier coupling as a pure, "postsignifying" form in which the matter of content melds with the manner of expression. It is the melding that takes the lead, as an "abstract machine" of autonomized, depersonalized expression (Deleuze/Guattari 1986, 3-8; Deleuze/Guattari 1987, 129-135, 189-191; Haghighi. The triad thesis, antithesis, synthesis is often used to describe the thought of German philosopher Georg Wilhelm Friedrich Hegel. Hegel never used the term himself, and almost all of his biographers have been eager to discredit it. The triad is usually described in the following way: The **thesis** is an intellectual proposition. The **antithesis is simply** the negation of the thesis, a reaction to the proposition. The synthesis solves the conflict between the thesis and antithesis by reconciling their common truths and forming a new thesis, starting the process over. According to Walter Kaufmann, **although the triad is often thought to form part of an analysis of historical and philosophical progress called the Hegelian dialectic**, the assumption is erroneous. Hegel used this classification only once, and he attributed the terminology **to Immanuel Kant**. The terminology was largely developed earlier by the neo-Kantian Johann Gottlieb Fichte, also an advocate of the philosophy identified as German idealism. The triad thesis, antithesis, synthesis is often used to describe the thought of German philosopher Georg Wilhelm Friedrich Hegel. Hegel stresses the paradoxical nature of consciousness; he knows that the mind **wants to know** the whole truth, but that it cannot think **without drawing a** distinction. Unfortunately, every distinction has two terms, every argument has a counter-argument, and consciousness **can only focus on one of these at a time**. So it fixes first on the one, then under pressure fixes second on the other, until it finally comes to rest on the distinction itself. Hegel refers to this **process of alternation and rest as dialectic**. In other words, the dialectical method **involves the** notion that the form of historical movement, process or progress, **is the result of** conflicting opposites. Thus this area of Hegel's thought **has been broken down in terms** of the categories of thesis, antithesis, and synthesis. Hegel's philosophy of history embraces the concept that a conflict of opposites is a struggle between actual and potential worlds. Kant's first Critique concerns itself with determining the structure of the objects of experience. His

transcendental logic allows **an empirical manifold, or multiplicity, to be grasped as a unified object**, but only insofar as its unity is provided by the subject (via apperception). The transcendental ego synthesizes manifolds that remain distinct in kind from it, and thus its logic can only provide the a priori conditions for possible experiences, not the generative conditions for actual ones. "Kant's main preoccupation is therefore with the validity of propositions given in advance of our enquiry", for it merely seeks the conditions for subjective knowledge of an object, which are presumed to be identical with the conditions of objects themselves. Limited to schematic overviews, brief asides or even dismissive caricatures these philosophies muster enough strength stride by stride with present day theories(Henry Somers-Hall, Hegel, Deleuze, and the Critique of Representation: **Dialectics of Negation and Difference, State University of New York Press, 2012, 289pp., (hbk), ISBN 9781438440095. Reviewed by Jim Vernon, York University)** Hegel shows how the various categories of logic sublate each other, turning contradiction into the motor that drives, rather than the problematic limit that haunts, representational thinking. The extended physical and biological theory is not just a multidimensional consciousness theory, but it is, after and above all, a philosophical theory. But their connection is not hierarchical. The self-created structural and structure less system of the multi-theoretical components is necessarily (in reality and symbolically also, a multi-centred, in- and out- infinitely expanded, multi-layered spherical) logical and logical spacetime theory. But to fully unfold this concept, neither physics, nor any other partial theory is capable in itself. So, the different unified theories together are cannot be other than the wholeness theory, the unified top theory or the philosophical metatheory, but is this not just to give a name, to explain something, would this be the Creation/Non-creation itself? **On Quantum's Universality: Posted on 5/19/2004 by Dave Bacon (The Quantum Pontiff):** Often when I am thinking about the foundations of quantum theory, I am struck by the universality of the theory. Quantum theory (or its related cousin, quantum field theory) applies generically to all physical systems (disregarding the transition to some "classical" theory and of course, difficulties with both QCD and gravity.) Thus we apply quantum theory to our basic theories of physics, electromagnetism, the weak force, the strong force, but we also apply quantum theory to simple atoms and complex molecules, to single electrons and electron gases in metals, etc. Quantum theory is the universal language we use to describe any physical process. If we are thinking about ways to explain quantum theory, then this universality is a bit mysterious: the explanation had better apply to all of these different physical systems and that seems like a lot of work! Of course, this reasoning is flawed: it seems the universality is an illusion. The reason we can describe a complex molecule by quantum theory is that the fundamental constituents of that molecule obey quantum theory. Separation of different energy scales (and other scales, like localibility) allow us to ignore some of the constituents details, and the complex system behaves like a quantum system. So really any explanation of quantum theory need only apply to some basic level of physics (where this level is I refuse to speculate.) While quantum theory appears mysteriously universal, this is an illusion for those persuing understanding the mystery of the quantum. Whitehead was a mathematician as well as philosopher. Metaphysics was his main interest. He said it is the task of philosophy "to frame a coherent, logical, necessary system of general ideas in terms of which every element of our experience can be interpreted. Parson's theory of social action **is based on his** concept of the society. Parsons is known in the field of sociology mostly for his theory of social action. Action **is a** process in the actor-situation system which has motivational significance to the individual actor or in the case of collectively, its component individuals.

Casti's book provides us with formalizations for appreciating the joke. For example, we can see this as another example of the liar paradox or the self-amendment problem in law, understand the humor of the cartoon as a dimensional jump from level N to level N +1, or as a Gtsdel-like statement of the problems of formalization. (Gregory Bateson Makes a Similar use of the ideals of Logical types and levels of abstraction in HIS discussion of Schizophrenia) For Casti, the relation of formal Modelling Systems to Natural Systems Has Come upon the problem of Complexity and not yet Become a science. For Casti, complexity is an inherently subjective concept, not that it is purely subjective, but rather that the complexity of a system is a joint property of the system and its interaction with another system, most often an observor and / or controller. Thus complex systems often generate counterintuitive behaviour. He points out that human societies have evolved to the point where traditional maps no longer match our collective experience for very long. "Thus, by coming up with a workable (ie scientific) theory of complexity, we

can hope to internally represent the experience of change by describing our collective reality as a process. This, in turn, would be a major step toward the development of a framework within which we can begin to understand how to control and manage what our maps tell us are complex processes. " (Complexification, p. 273) Casti describes objective Ways to characterize some Aspects of a system's Complexity. Algorithmically speaking, for example, complexity is generally measured by compressibility, eg. by the shortest program that will cause a Turing machine to print out a number. randomness is considered to be incompressible, that is to say the description of the object is no more compact than the object itself. Thus to explore a complex system, there are no shortcuts to running the system itself. "Logical depth" is another way of describing this computational complexity. Logical depth is defined as the time needed (measured in number of computational steps) for the shortest possible program to generate the structure. A true Deep Structure is THUS CHARACTERIZED by the Mathematical That property IT cannot be generated by fewer steps Via a Computational Simulation on any Other Computer. Casti Believes That the formalization of explicit Complexity Requires the Recognition of the observor. For him the complexity of the system as seen by the observor is directly proportional to the number of inequivalent descriptions of the system that the observor can generate, when one description bifurcates from another. Descriptions of the Choice of Complex Systems depend on contexts of Understanding and Interpretation. Niklas Luhmann calls autopoetic Systems "Complex Systems Self-referential." These systems make and continue to make a difference between the system and its environment. They are operationally closed. Luhmann describes Complex Systems Self-referential as "hypercomplex," in the Ways That They make Distinctions Between Themselves and Their Environments, but Also internalize the Distinction in a paradoxical "Re-entry" of the Distinction into the Distinguished. non-Linearity - ---------- In Dynamical Terms, such as in the Study of chaos, a non-linear situtation is one where the result is not proportional to the Cause. For instance "the straw that broke the camel's back" (eg. the elastic / plastic limit in building structures) introduces non-linearity. Up until that point, deformation had been proportional to load. Suddenly IT Loses All proportionality. "Sensitivity to Initial Conditions" is an Indication of the non-Linearity of Chaotic Dynamical Systems, where the Most Minor Difference in Conditions Starting CAN lead to differences in VAST Subsequent Conditions. In the Study of A non-life -linearity is used to describe systems that do not obey the superposition principle. They are not the sum of their parts. The parts of linear systems can be analysed in isolation and can be combined to understand the system as a whole. The key feature of non-linear systems is that their primary behaviours of interest are properties of the interactions between parts rather than being properties of the parts themselves, and these interactions necessarily disappear when the parts are studied independently. (Cf organism) (see also Holland, who also uses the term "epistatic" see below) There are very important feedback mechanisms between levels in such interactions. The interactions among low-level entities give rise to the global level dynamics, which, in turn, affects the lower levels by setting the local context within which each entity's rules are invoked. For Langton, analytic methods work for linear systems, but non-linear systems must be explored through synthetic methods. (See Langton, "Artificial Life") The Effects of Different Enzymes in the cell are not additive. This phenomenon is known as epistasis. The Effect of each allele of a Gene Strongly depends on What Other alleles are present, and there is no simple way to apportion Credit to Individual alleles for the Performance of the resulting phenotype. When Cultural or Literary critics use the Term, IT relates to More "gardens of forking paths" or networks. Hypertext has been described as nonlinear because various choices are possible at any Occur did, but as a Process of Natural Self-Organization. " Unknown photographer: Photograph of multiple Chico Marx c.1909 Self-Organizing Systems aquire new Structure without specific Interference from the outside. They Exhibit Qualitative macroscopic Changes such as bifurcations or phase transitions . Stuart Kaufman calls this "Order for Free." Self-organization is the capacity of a field to generate patterns spontaneously, without any specific instructions. What exists in the field is a set of relationships among the components of the system such that the dynamically STABLE State into Which IT Goes Naturally - What mathematicians call the Generic (typical)) State of the system Has Spatial and Temporal Patterns. Fields of this type are now Called excitable Media. (See for example Belousov-Zhabotinsky Reaction.) HTTP: / / www.christianhubert.com/hypertext/self_organization2.jpeg "Self-Organizing Systems ... ACCOMPLISH Their Remarkable feats by Chance Incorporating into the Substances of Their Forms, Which then persist, Maintained by Their Own Functions" (Yates, P. 10) ... deterministic in Principle, not exactly knowable in Practice. Self Organizing Systems May have an absolute requirement for Random

Fluctuations at some Point in Their histories. (See clinamen for a Classical Account of fluctuation.) Most Descriptions of Self-Organization stress cooperative behaviour between elements. For example, the difference between ordinary light and lasers can be described as follows: "In the ordinary lamp, electrons of individual atoms make their optical transitions independently of each other, but in the laser the electrons make their optical transitions cooperatively ". Gestalt psychology described Perception and Brain-events as Self-Organizing Systems spontaneously Take That on the "best" or in Simplest Arrangement Given Conditions. In 1914, Max Wertheimer introduced the principle of Prdgnanz to describe this capacity. The geometrical hallucinations Also of migraines are an example of the Complex, reiterative Patterns in time and space Which Represent the Ways such Dynamic, non-linear Systems Organize Themselves. "For a Long time Turbulence was Identified with Noise and Disorder. Today we know this is not the case. Indeed, while turbulent motion appears as irregular or chaotic on the macroscopic scale, it is, on the contrary, highly organized on the microscopic scale ... Viewed in this way, the transition from laminar flow to turbulence is a process of self-organization "(Prigogine, Order Out of Chaos, p.141) One example of a Self-Organizing River system is a system. While All water Runs downhill, both Rivers carve channels and Carry a load bed (Which silts up the channels) This results in meandering. On a geological time scale this adaptation by changing form can be described as a life-like behaviour. While the thermodynamic processes of the hydrological cycle (rainfall = evaporation plus runoff) and of the landmass are near equilibrium in local regions, they display a fairly Ergodic Character. A single River system CAN be described as Being Born, Living, and Dying. For Per Bak, Kurt Wiesenfeld, and Chao Tang, large Interactive Dynamical Systems Typically Self-Organize into a globally Correlated "Critical" State, far out of Equilibrium. See Self-org. Criticality. Molar / Molecular ---------- In mechanics, Molar Properties Those are of a mass of Matter , as Opposed to ITS parts - Atoms or Molecules. Deleuze and Guattari For, molarity is the Site of Coded wholes. It is a Productive Process: a making-the-Same. Its attractor That is State of Equilibrium STABLE. It is the mode of being, rather than becoming.The principle revolutionary objective of their writing is to break down molar aggregates in favor of molecularity, and the "microphysics of desire." They call for becoming rather than being, for becoming-other rather than being the Same. For Them, Becoming-Other is Thoroughly Political. Molar The Body is defined by ITS Self-similarity accross ITS variations, or rather, ITS invariant Self-Identity. It seeks to make an Either / or from the both / and. Oedipus , phallocentrism, and personhood are All Molar wholes. Rhizome, pack, haecceity, singularity, Event .. All are Molecular. "Becoming is an Equilibrium-Seeking system at a Crisis Point where IT Suddenly perceives a deterministic constraint, Becomes" sensitive "to it, and is catapulted into a highly unstable supermolecular state enveloping a bifurcating future. "(Massumi, p.95) "There is no Becoming Because MAN-MAN is the Molar Entity par Excellence, whereas becomings are ... Molecular Man consitutes the Majority, or rather the standard on Which the Majority is based: White, male , adult, 'rational,' etc., in short, the average European, the subject of enunciation. "(A thousand Plateaus, p 292) (See the Critique of this Concept in Elizabeth Grosz, Volatile Bodies, PP 173-183) The Bwo is a Site of Decoded Flows. It opens a zone of indeterminacy, for imagination as a new Circuit Between the actual and the virtual, in the Place of Habit. Becoming Bears on a Population, even When IT is Initiated by a single Body. http://www.christianhubert.com/hypertext/form_matter.html # 2 form / Matter - ---------- The Judeao-Christian Account of Creation is an Account of the origin of form, not of Matter. "In the Beginning .. the Earth was without form, and void." The Passage ... Deals at Length with the origin of order. For Aristotle, the Generation of each Organism was the result of a formal Cause male (Conveyed by the semen) and a Female Material Cause (the menstrual Blood.) (See epigenesis) cf Body / Soul. The "hylomorphic" model Separates a form organizes That Matter and a Matter Prepared for the form. Gilbert Simondon bases HIS Critique of the hylomorphic model on "the Existence, Between Matter and form, of a zone of Medium and Intermediary dimension. " A formalist Concept presupposes a Contrast Between Matter and form. Indeed, without this the absolutization of Distinction form Makes no sense. "When We Look upon the work of Art, the Means and the Materials are Forgotten and IT is in Satisfying Itself as form. "(Gottfried Semper) In The Life of Forms in Art, Henri Focillon argues Against the Antithesis of form and Matter. For HIM, Art is bound to Matter, and "unless and until IT Actually exists in Matter, form is little better than a vista of the mind, a mere speculation on space that has been reduced to geometrical intelligibility. "(p95) According to Focillon," matter, even in its most minute details, is always structure and activity, that is to say , form. "(p.96) and each kind of matter has a certain" formal vocation. "But the life which inhabits matter undergoes a

metamorphosis as it becomes a substance of art, and technique is a" whole poetry of action "as a Means to Achieve Metamorphoses. Konrad Fiedler described the Achievement of Classical Greek architecture as the intellectualization of Complete All the Elements of Building Material. "We CAN Speak Understanding of a Greek Building from the Great period Only When We Perceive How the Force That strives for a pure expression of form has taken command of every part of the building. "(" Observations on Architecture, in Empathy, Form, and Space, p. 134) Still, Fiedler is careful to note that "Form has no existence except in material, and the Material, to the Mind, is not the Only Means by Which form Expresses Itself but the Medium in Which form achieves Existence. " For Heinrich Wtslfflin, the force of form (Formkraft), the opposition between the tendency of matter towards formless Collapse and the Opposing Force of Will, life, or whatever, sets the Entire Organic World in Motion and the Principle is Theme of architecture. Ferdinand de Saussure For, "Language is a form and not a substance." (Cours, p.169) For Deleuze and Guattari, the Distinction Between Matter and form is Characteristic of "Royal Science", the science of a Society Divided into Governors and governed. For nomad science the relevant distinction is material-forces rather than matter-form. Materials for nomad science are not homogenous, and form is not fixed. The singularities or haecceities are already like implicit forms that are topological rather than geometrical, and that combine with the processes of deformation: for example the variable undulations and torsions of the fibers guiding the operation of splitting wood. "Energetic materiality in movement" (see Thousand Plateaus, p 408) (See transcendent / immanent) (Also See Natural form) Energy ---------- The Greek energeia Term derives etymologically from en ("within") and ergein ("to work"). THUS Energy is the actual work or Potential That is immanent within a Body. Joule's Principle of the Conservation of Energy is an example of the subsumption of Qualitative transformations into a quantifiable Entity. "Thus it is that order is maintained in the universe - nothing is deranged, nothing ever lost, but the entire machinery, complicated as it is, works smoothly and harmoniously." (Joule, quoted in Prigogine, p.108-9) In the period from 1775-1825, the Word and Concept of Energy was bruited by All About manners of literate Individuals, Often as a shorthand, or cipher for talking About organicism as a process uniting mind, nature, and art. (Stuart Peterfreund, "Organicism and the Birth of Energy" in Burwick, Approaches to Organic Form.) Freud Dynamic model of the Unconscious, Called "Hydraulic" by critics, described Energy Flows That build up and seek Release. In this model, tension is displeasure and Will Always seek an Avenue for release's Force ------- Is the Concept of Force a Technico / Scientific one or a Philosophical, even metaphysical one? By Defining Force purely as the Product of an acceleration (a purely kinematic magnitude - from Greek kineo, referring to constrained or controlled motion) and a mass (a coefficient to be determined empirically), modern science eliminated both the metaphysical terminology and psychological origins of the concept of force. (EJ Dijksterhuis) (See Machine) See Also qualititative / Quantitative. For Classical Physics, Force is really explainable Only in Terms of Bodies in Contact. "Action at a distance", which is also a feature of gravity, seems "occult." Yet from the work of Kepler and Newton, particles had a dual nature: on the one hand, a highly localized object, and on the other an influence extending through the whole of space. But is the Concept of Force to be Primarily Associated with Newtonian mechanics - as Opposed to Newton's alchemical Writings? Newton refused to define the Fundamental Nature of HIS Gravitational Force. He emphasized its heuristic importance and the possibility of mathematizing it. On the other hand, his alchemical speculations downright invited the amplification and translation of his concept of force into the realm of living things. Could magnetism and electricity, he asked, fulfill the same role for living beings that gravitation did for inanimate matter? (See mechanism / vitalism) Whether as quasi-mechanical attraction of molecules (as suggested by Buffon and Maupertius) or eventually as formative force (as suggested by the epigeneticists) the Newtonian concept of force supplied the debate about generation with a new energy in the Eighteenth century. In What is Matter?, Hermann Weyl Gives a Detailed outline of the displacement of the old "substance Theory" by the Modern "Field Theory." - a kind of tension or Stress That CAN exist in empty space in the Absence of matter. In 1907 Einstein theorized that light propagated across a gravitational field is curved. Since the trajectory of light is the shortest distance between two points and the basis for all measurement, he concluded that by slowing down the speed of light gravity warps time and space. He also hypothesized that gravity is not a force but an intrinsic curvature of a space-time continuum. It is Currently Accepted That there are four kinds of Force in the universe: Electromagnetic, gravitation, Weak, and strong. Wtslfflin For Heinrich, the principal theme of architecture is the opposition between matter and the force of form. (Formkraft) ("Prologemena to a Psychology of Architecture," p.

159-162) This dimension of architecture is experienced in sympathy with the body. Wolfflin sharply distinguishes between the lawfulness of geometric form "which does not strike us as a pleasing vital form," and the features of architectural form whose principles we understand "by virtue of our own bodily organization and expressive movements." (This is the same opposition that Wilhelm Worringer drew between abstraction and empathy.) "Matter itself, to a certain extent, longs for form .... The perfect form, for its part, presents itself as an entel echy, that is, as the actualization of the potential inherent in the matter. (cf formative force in mechanism / vitalism) For Wtslfflin, ornament is an epression of an excessive force of form. It is "the blossoming of a force that has nothing more to achieve." (p.181) See form / Matter. Classic In HIS 1917 book On Growth and Form, D'Arcy Thompson analyzed biological form in terms of physical forces such as viscosity or gravity. Biological form and engineering design were directly related as conforming to or resisting physical Force. See di Agram for Contemporary Analyses of Forces in architecture. Metaphysical The usage of the creative Force of Being found a new proponent in Henri Bergson. Bergson conceived of a life Force (ylan), radically distinct from inert matter, but passing through IT and forcing IT to Become Organized. Gilles Deleuze For the Affective Force, Level of intensity, Sire de, or conveys ideas and Affirmation Ultimately Govern Their tr UTH-value. Psychoanalytic Accounts of the in stinctual describe Drives Them in Terms of Force. See Affect Re ssentiment for Nietzsche's Account of a Force turning back on Itself. Power -------- For Foucault, every relation Between Forces is a "Power relation." Power is not Essentially repressive. It is practiced before IT is possessed, and it passes through the hands of the mastered no less than through the hands of the masters. Therefore we should ask: "How is it practiced?" It is a physics of abstract action. (Deleuze, Foucault, p. 72) "Power has its principle not so much in a person as in a certain concerted distribution of bodies, surfaces, lights, gazes; in an arrangement whose internal mechanisms produce the relations in which the individuals are caught up." (Discipline and Punish, p .202) (See Also Spectacle) "Power must be Understood ... as the multiplicity of force relations immanent in the sphere in which they operate and which constitute their own organization; as the process which, through ceaseless struggles and confrontations, transforms, strengthens, or reverses them. "(History of Sexuality, Vol 1, p . 26) According to Deleuze, is local Power Because IT is Never global, but IT is not local or localized Because IT is diffuse "(Foucault, P. 26) note relation Between Power, Knowledge, and Pleasure. For Bertrand Russell, “Scientific thought is essentially power thought - the sort of thought whose purpose, conscious or unconscious, is to give power to its possessor. " For Foucault, knowledge and truth are inseparable from power. If Linking Knowledge consists of the Visible and the articulable, Power and Knowledge presuppose and constitute each Other. http://www.christianhubert.com/hypertext/desire.html Desire -------- "Once the imagination Has Been excluded from experience as unreal, and its place has been taken by the ego cogito, the status of desire changes radically: it becomes essentially insatiable. " "The expulsion of imagination from the sphere of experience sunders what Eros united in himself: desire and need, in such a way that they cannot coincide in the same subject." (Giorgio Agamben, infancy and History, p.26) Freud's use of the Word Wunsch, Which corresponds to 'wish' does not have the connotations Same as the English Word 'Desire "or the French dysir. His clearest elucidation of the concept is in the theory of dreams. Freud does not identify need with desire. Need can be satisfied through the action which procures the adequate object. (eg. food) Wishes, on the other hand, are governed by a relationship with signs, with memory- traces of Excitation, and the Desire to Re-cathect MNEMIC Images. The Freudian conception of Desire Refers above All to Unconscious wishes, bound to Indestructible infantile signs, Organized as Phantasy. Freud's Analysis of Hysteria Showed How hysterics have Taken Their imagination for Reality, and, more fundamentally, how they have translated - according to specific laws of transposition - their desire into reality. Their own desire to seduce the father has been translated, in inverse form, into an actual scene of seduction by the father. Freud Also Compared Hysterical symptoms to monuments erected to commemorate events. THUS Anna O. 's symptoms are "MNEMIC Symbols" of the Illness and Death of her father. Freud Has Been criticized for renouncing the Seduction Theory - Which Recognized a Reality to Parental Seduction - and for Reducing the Realm of Desire to the Interior Realm of the Self and the Family. Narcissism and Desire: Narcissus expired through HIS delusory Desire "Because He Could not Lay hold of Himself, and yet Himself Perceived as Other. (Ovid) Jacques Lacan's theory of the mirror stage follows both Ovid and Freud in stressing the basically narcissistic relation of the subject to his counterpart, the specular ego. In this way Lacan also sets up the erotic attraction or aggressive tension as a relation to a counterpart ("another who is me"), who can only exist because the ego is originally another. (See "Imaginary" in Laplanche and

Pontalis, The Language of Psychoanalysis) The Collapse into the Self enacted by Narcissus is Transformed for prototypically Narcissistic Female subjects into a Fundamentally alienated Relationship to her external Image. http://www.christianhubert.com/ hypertext/desire1.jpeg Lack and Desire: In a tradition That Reaches from Plato to Lacan and Beyond, Desire Has Been Understood as Negative, Abyssal, a Lack at the Level of Ontology Itself (this was Most Clearly articulated in Hegel's Lack Understanding of the [of the object] of desire being the necessary condition for the maintenance of desire), a lack in being that strives to be filled through the (impossible) attainment of an object. Lacan calls this lack un manque-a-ktre a "Want-to-be" (or a wannabe?) For Georg Simmel, "the Possibility of Desire is the Possibility of Objects of Desire." (The Philosophy of Money, p. 66) For Simmel enjoyment is a unity similar to the state prior to the differentiation of subject and object. With the observation that "our mind has a remarkable ability to think of contents as being independent of the act of thinking, he observes that" In desiring what we do not yet own or enjoy, we place the content of our desire outselves ... . We desire objects only if they are not immediately given to us for our use and enjoyment; that is, to the extent that they resist our desire. "Value is characterized by this separation, which takes the form of a claim or demand. The purpose of establishing a distance is that it should be overcome. Withdrawal and approach are the two sides of our relationship to objects, which we call subjectively our desire and objectively their value. For Simmel, aesthetic value is the most complete projection of our feelings. "The content of the feeling is, as IT Were, absorbed by the object and confronts the subject as something autonomous Which Has Significance, Which is Inherent in the object." (p.73) Immanuel Kant Sought to separate from Desire Aesthetics (as well as concepts). For Kant, disinterest Lies at the core of Aesthetics, and while Enjoyment of the Pleasant May Satisfy specific Desires, the core of Aesthetics is a Freedom from Desire. Contemporary theorists have tried to reconceptualize Desire in positive Terms. Feminists have objected to the association of lack with the feminine, (eg. the Freudian interpretation of womens' genitals) of the phallus as the unique symbol of desire, (and power) such that desire in women can only appear as envy. Jessica Benjamin links the intersubjective realm of the holding environment and transitional experience to the experience of inner self, which she sees as enabling the experience of women's desire. ("A Desire of One's Own, in Teresa de Lauretis, ed. Feminist Studies / Critical Studies) (see Sexual pscho-space) Desire and Philosophy: The bulk of the Western tradition Has Sustained Skepticism Towards the Philosophical Possibilities of Desire. To desire the world and to know its meanings and structures have seemed conflicting enterprises. But when philosophy interrogates its own possibilities as engaged or practical knowledge, it tends to ask after the philosophical potential of desire. In Moral Actions, the relation Between subject and object is Completely Different from That of Theoretical Knowledge, for the Desire to Realize the Good Produces ITS Own Objects. Spinoza and Kant admitted Desire to Philosophy, and both Developed a Philosophy of Psychic Integration Between reason and desire. If the philosopher is not beyond desire, then (s) he is a being of rational desire who knows what he wants and wants what he knows. (Judith Butler) For Kant, the Good is the object of Will (IE a Faculty of Desire Determined by reason. (Critique of Judgement, p.43) Kant was convinced That the Human Mind Has a Natural Desire to be reminded of the Fact that it is possessed of the faculty of Reason. Although we commonly think of desire as having for its object a natural object, possession of which is desired. Kant held that there is another quite different kind of desire, viz., the desire to realize the good. (see HW Cassirer, A Commentary on Kant's Critique of Judgement, p 125) This Moral and Self-fufilling form of Desire is echoed in Contemporary Writers such as Rosi Braidotti. She calls the "desi Re for Philosophy" a Will to know that is fundamentally affective - a love of, and desire for, higher knowledge. Braidotti attributes this form of desire to both Gilles Deleuze and Luce Iragaray "... Not just libidinal desire, but rather ontological desire, the desire to be. . "(Rosi Braidotti, refs. to Lyotard, Le postmodernisme expliquy aux Enfants) For Gilles Deleuze the Affective Force, Level of intensity, Desire, or conveys ideas and Affirmation Ultimately Govern Their Truth-value. prephilosophical The moment of Desire is not Only unthought, but remains unthought at the very heart of philosophy, because it is that which sustains the very activity of philosophizing. Braidotti alludes to "the unspeakable desire for thought." "The desire to know is, like all desires, related to the problem Representing one's of Origins, of Answering the Most childish and consequently of Fundamental questions: "where did I Come from?" (Braidotti, Nomadic Subjects, p.90) Deleuze's Notes on Foucault, Published by François Rewald under the title "Desire and Pleasure , "contain an important definition. Life, Deleuze says, is" desire's variable field of immanence. "Given what we know of Deleuzian immanence, this means that the term "life" designates nothing more and nothing less than the immanence

of desire to itself. (Agamben, "Absolute Immanence", p. 235) It is clear that for Deleuze, desire implies neither alterity nor a lack. For Deleuze and Guattari, desire is a process of production. (See desiring machines) (see also BwO)

On the basis of this definition it may be said that the processes of action **are related to and influenced by the** attainment of the gratification or the avoidance of deprivations of the correlative actor, whatever they concretely be in the light of the relative personal structures that there may be. All social actions proceed from mechanism which is their ultimate source. It does not mean that these actions are solely connected with organism. They are also connected with actor's relations with other persons' social situations and culture. Philosophy and art, Badiou revisits the relationships that have been proposed between poetry and philosophy. They include the early fusion of philosophy into the sacred aura of utterance (Parmenides), the separation of poetry from philosophy in Plato, and the inclusion of poetry into the organization of knowledge in Aristotle. Badiou commends Heidegger's re-establishment of the function of poetry to produce truth, while criticizing him for restoring the sacral authority of the authentic poetic utterance. For Badiou, "poetry is the thought of the presence of the presence." (p.99) He recognizes that the "undecidable supplement" of the event **requires** poetic naming. In this sense, the poem produces truths. Yet the truth procedures of philosophy **must be subtracted** from the eventual singularity that weaves them into the real. For Badiou, philosophical truth is not about sense and the interpretation of its conditions. It must **subtract** all aura of sense, **all trembling and all pathos**, to seize truth's proving of itself as such. (p.104)

Systems of social action (Observers in natural sciences: an analogy)

Social actions are **guided by the** following three systems which may also be called as three aspects of the systems of social action Personality system: This aspect of the system of social action is responsible for the needs for fulfillment of which the man makes effort and performs certain actions. But once man makes efforts he has to meet certain conditions. These situations have definite meaning and they are distinguished by various symbols and symptoms. Various elements of the situation come to have several meanings for ego as signs or symbols which become relevant to the organization of his expectation system. For Alain Badiou, the three main strands of contemporary philosophy, the hermeneutic, the analytic, and the postmodern all claim that the ideal of truth as it was put forth by classical philosophy has come to its end and that for the idea of truth we must **substitute** the plurality of meanings. (Infinite Thought, p. 46) Instead, Badiou calls for a philosophy **founded on** singularity, a philosophy of contemporary rationality, a philosophy of the event. Badiou traces the trajectory of a truth as starting with an undecidable event, such as a creation or invention. For Badiou, the axiom of truth **opened up** by such an event is infinite. His examples include: the appearance, with Aeschylus, of theatrical tragedy; the eruption, with Galileo, **of mathematical physics**; an amorous encounter which changes a whole life, the French Revolution of 1792. The artistic truth of Greek Tragedy, opened up by Aeschylus is inexhaustible. The work **is finite**; a pure choice made in what was previously in discernable. But Tragedy itself, as an artistic truth, continues to infinity.

A possibly meaningful (but unparsimonious) answer to the Ultimate Why is that the universe exists (more precisely, is perceived to exist) roughly because it is possible. The reasoning would be as follows. Absolute impossibility -- the state of affairs in which nothing is possible -- is itself not possible, because if nothing truly were possible, then absolute impossibility would not be possible, implying that at least something must be possible. But if at least one thing is possible, then it seems the universe we perceive should be no less possible than anything else. Now, assuming that physicalism is right and that qualia and consciousness are epiphenomena, then the phenomenology of a mind and its perfect simulation are identical. So whether the universe we perceive existed or not, it as a merely possible universe would be perceived by its merely possible inhabitants no differently than our actual universe is perceived by its actual inhabitants. By analogy, the thoughts and perceptions of a particular artificial intelligence in a simulated universe would be the same across identical "runs" of the simulation, regardless of whether we bothered to initiate such a "run" once, twice -- or never. Thus, the universe might merely be the undreamed possible dream of no particular dreamer.**Brian Holtz (Human knowledge: Foundations and limits**)

Cultural system: Once the process of the social action develops the symbols and the signs acquire general meaning. They also develop as a result of systematized system and ultimately when different actors under a particular cultural system perform various social interactions, special situation develops.

Social System: A social system consists in a plurity of individual actor's interacting with each other in a situation which has at least a physical or environmental aspect actors are motivated in terms of tendency to the optimization of gratification and whose relations to the situation including each other is defined and motivated in terms of system of culturally structured and shaped symbols.

In Parson's view each of the three main type of social action systems-culture, personality and social systems has a distinctive coordinative role in the action process and therefore has some degree of causal autonomy. Thus personalities organize the total set of learned needs, demands and action choices of individual actors, no two of whom are alike.

Every social system is confronted with 4 functional problems. These problems are those of pattern maintenance, integration, goal attainment and adaptation. Pattern maintenance refers to the need to maintain and reinforce the basic values of the social system and to resolve tensions that emerge from continuous commitment to these values. Integration refers to the allocation of rights and obligations, rewards and facilities to ensure the harmony of relations between members of the social system. Goal attainment involves the necessity of mobilizing actors and resources in organized ways for the attainment of specific goals. Adaptation refers to the need for the production or acquisition of generalized facilities or resources that can be employed in the attainment of various specific goals. Social systems tend to differentiate these problems so as to increase the functional capabilities of the system. Such differentiation whether through the temporal specialization of a structurally undifferentiated unit or through the emergence of two or more structurally distinct units from one undifferentiated unit is held to constitute a major verification of the fourfold functionalist schema. It also provides the framework within which are examined the plural interchanges that occur between structurally differentiated units to provide them with the inputs they require in the performance of their functions and to enable them to dispose of the outputs they produce.

The scientist can learn about the world by observation but Kierkegaard emphatically denied that observation could reveal the inner workings of the spiritual world. In 1847 Kierkegaard described his own view of the single individual:

God is not like a human being; it is not important for God to have visible evidence so that he can see if his cause has been victorious or not; he sees in secret just as well. Moreover, it is so far from being the case that you should help God to learn anew that it is rather he who will help you to learn a new, so that you are weaned from the worldly point of view that insists on visible evidence. (...) A decision in the external sphere is what Christianity does not want; (...) rather it wants to test the individual's faith."

"The crowd is untruth. And I could weep, in every case I can learn to long for the eternal, whenever I think about our age's misery, even compared with the ancient world's greatest misery, in that the daily press and anonymity make our age even more insane with help from "the public," which is really an abstraction, which makes a claim to be the court of last resort in relation to "the truth"; for assemblies which make this claim surely do not take place. That an anonymous person, with help from the press, day in and day out can speak however he pleases (even with respect to the intellectual, the ethical, the religious), things which he perhaps did not in the least have the courage to say personally in a particular situation; every time he opens up his gullet—one cannot call it a mouth—he can all at once address himself to thousands upon thousands; he can get ten thousand times ten thousand to repeat after him—and no one has to answer for it; in ancient times the relatively unrepentant crowd was the almighty, but now there is the absolutely unrepentant thing: No One, an anonymous person: the Author, an anonymous person: the Public,

sometimes even anonymous subscribers, therefore: No One. No One! God in heaven, such states even call themselves Christian states. One cannot say that, again with the help of the press, "the truth" can overcome the lie and the error.

"O, you who say this, ask yourself: Do you dare to claim that human beings, in a crowd, **are just as quick** to reach for truth, which is not always palatable, as for untruth, which is always deliciously prepared, when in addition this **must be combined** with an admission that one has let oneself be deceived! Or do you dare to claim that "the truth" is just as quick to let itself be understood as is untruth, which requires no previous knowledge, no schooling, no discipline, no abstinence, no self-denial, no honest self-concern, and no patient labor! No, "the truth," which detests this untruth, the only goal of which is to desire its increase, is not so quick on its feet. Firstly, it cannot work through the fantastical, which is the untruth; its communicator is only a single individual. And its communication relates itself once again to the single individual; for in this view of life the single individual is precisely the truth. The truth can neither be communicated nor be received without being as it were before the eyes of God, nor without God's help, nor without God being involved as the middle term, since he is the truth. It can therefore only be communicated by and received by "the single individual," which, for that matter, every single human being who lives could be: this is the determination of the truth in contrast to the abstract, the fantastical, impersonal, "the crowd" – "the public," which excludes God as the middle term (for the personal God cannot be the middle term in an impersonal relation), and also thereby the truth, for God is the truth and its middle term." Søren Kierkegaard, Copenhagen, spring 1847

Pattern Variables

Affectivity vs affectivity neutrality: The pattern is affective when an organized action system emphasizes gratification that is when an actor tries to avoid pain and to maximize pleasure; the pattern is affectively neutral when it imposes discipline and renouncement or deferment of some gratifications in favour of other interests. Self-orientation vs collectivity orientation: This dichotomy **depends on** social norms or shared expectations which define as legitimate the pursuit of the actor's private interests or obligate him to act in the interests of the group. **Particularism vs universalism**: The former refers to standards determined by an actor's particular relations with particular relations with a particular object; the latter refers to value standards that are highly generalized. **Quality vs performance:** The choice between modalities of the social object. This is the dilemma of according primary treatment to an object **on the basis of what** it is in itself an inborn quality or what it does and quality of its performance. The former involves defining people on the basis of certain attributes such as age, sex, color, nationality etc; the **latter defines** people on the basis of their abilities. **Diffusion vs specificity**: This is the dilemma of defining the relations borne by object to actor (observer)as indefinitely wide in scope, infinitely broad in involvement morally obligating and significant in pluralistic situations or specifically limited in scope and involvement. For Nietzsche, truth is "a mobile army of metaphors, metonyms, and anthropomorphisms -- in short a sum of human relations, which have been enhanced, transposed, and embellished poetically and rhetorically and which after long use seem firm, canonical, and obligatory to a people." (In "Truth and Lie in an Extra-Moral Sense." Michel Foucault picked up Nietzsche's insight into the **"politics of truth."** Foucault's position is that intellectuals are inextricably **involved in a** struggle over "**the status of truth and the economic and political role it plays."** The option before radical intellectuals is not that of "emancipating truth from every system of power (which would be a chimera, because truth is already power) but of detaching the power of truth **from the** forms of hegemony, social, economic, and cultural, within which it operates at the present time." (Barry Smart "Politics of Truth" in Hoy, ed. Foucault Reader 165-66). Thus there is a general attribution, ascription and assignment that truth is something which need have to be different from perception and as Von Neumann hesitatingly pointed, and we point out bluntly is the consciousness.

The object of faith is the actuality of another person; its relation is an infinite interestedness. The object of faith is not a doctrine, for then the relation is intellectual, and the point is not to bungle it but to reach the maximum of the intellectual relation. The object of faith is not a teacher who has a doctrine, for when the teacher has a doctrine, then the doctrine is eo ipso more important than the teacher, and the relation is intellectual, in which the point is not to bungle it but to reach the maximum of the intellectual relation. But the object of faith is the actuality of the teacher, that the teacher actually exists. Therefore faith's answer is absolutely either yes or no. Faith's answer is not in relation to a doctrine, whether it is true or not, not in relation to a teacher, whether his doctrine is true or not, but is the answer to the question about a fact: Do you accept as fact that he actually existed? Please note that the answer is with infinite passion. In other words, in connection with a human being it is thoughtless to lay so infinitely much weight upon whether he has existed or not. Therefore, if the object of faith is a human being, the whole thing is a prank by a foolish person who has not even grasped the esthetic and the intellectual. The object of faith is therefore the god's actuality in the sense of existence. But to exist signifies first and foremost to be a particular individual, and this is why thinking must disregard existence, because the particular cannot be thought, but only the universal. The object of faith, then, is the actuality of the god in existence, that is, as a particular individual, that is that the god has existed as an individual human being. Christianity is not a doctrine about the unity of the divine and the human, about subject-object, not to mention the rest of the logical paraphrases of Christianity. In other words, if Christianity were a doctrine, then the relation to it would not be one of faith, since there is only an intellectual relation to a doctrine. Christianity, therefore, is not a doctrine but the fact that the god has existed. Faith, then, is not a lesson for slow learners in the sphere of intellectuality, an asylum for dullards. But faith is a sphere of its own, and the immediate identifying mark of every misunderstanding of Christianity is that it changes it into a doctrine and draws it into the range of intellectuality. What holds as the maximum in the sphere of intellectuality, to remain completely indifferent to the actuality of the teacher, holds in just the opposite way in the sphere of faith-its maximum is the quam maxime [in the greatest degree possible] infinite interestedness of the actuality of the teacher. The individual's own ethical actuality is the only actuality. That this seems strange to many does not surprise me. To me it seems odd that one has finished with the system and systems without asking about the ethical. Concluding Unscientific Postscript, Vol 1, p. 326-327 Hong In order to stress the element of self-determination in thinking, philosophy declares: The absolute is because I think it. But since philosophy itself perceives that free thinking is thereby designated, not the necessary thinking it usually celebrates, it substitutes another expression: namely, that my thinking of the absolute is the absolute's thinking-itself in me. This expression is by no means identical with the one preceding; it is, however, very suggestive. That is to say, my thinking is an element of the absolute, and therein lies the necessity of my thinking, therein lies the necessity with which I think it. It is otherwise with the good. The good is because I will it, and otherwise it is not at all. This is the expression of freedom, and the same is also the case with evil-it is only inasmuch as I will it. This in no way **reduces or lowers the categories** of good and evil to merely subjective categories. On the contrary, the absolute validity of these categories is declared. The good is the being-in-and-for-itself, posited by the being-in-and-for-itself, and this is freedom. It might seem dubious for me to use the expression "to choose oneself absolutely," because this might seem to imply that I chose both the good and the evil just as absolutely and that both the good and evil belonged to me just as essentially. It was to prevent this misunderstanding that I used the expression "I repent myself out of the whole existence." Repentance specifically expresses that evil essentially belongs to me and at the same time expresses that it does not essentially belong to me. If the evil in me did not essentially belong to me, I could not choose it; but if there were something in me that I could not choose absolutely, then I would not be choosing myself absolutely at all, then I myself would not be the absolute but only a product. Either/Or Part II, Hong p. 224

Continental philosophy includes the following movements: German idealism, phenomenology, existentialism (and its antecedents, such as the thought of Kierkegaard and Nietzsche), hermeneutics, structuralism, post-structuralism, French feminism, psychoanalytic theory, and the critical theory of the Frankfurt School and related branches of Western Marxism. It is difficult to identify non-trivial claims that would be common to all the preceding philosophical movements. The term "continental philosophy", like "analytic philosophy", lacks clear definition and may mark merely a family resemblance across disparate philosophical views. Simon Glendinning has suggested that

the term was originally more **pejorative than descriptive**, functioning as a label for types of western philosophy rejected or disliked by analytic philosophers Babette Babich emphasizes the political basis of the distinction, still an issue when it comes to appointments and book contracts. Nonetheless, Michael E. Rosen has ventured to identify common themes that typically characterize continental philosophy. First, continental philosophers generally reject scientism, the view that the natural sciences are the only or most accurate way of understanding phenomena. This contrasts with analytic philosophers, many of whom have considered their inquiries as continuous with, or subordinate to, those of the natural sciences. Continental philosophers often argue that science depends upon a "pre-theoretical substrate of experience" (a version of the Kantian conditions of possible experience or the phenomenological concept of the "lifeworld") and that scientific methods are inadequate to fully understand such conditions of intelligibility. Second, continental philosophy **usually considers** these conditions of possible experience as variable: determined at least partly by factors such as context, space and time, language, culture, or history. Thus continental philosophy tends toward historicism. Where analytic philosophy tends to treat philosophy in terms of discrete problems, capable of being analyzed apart from their historical origins (much as scientists consider the history of science inessential to scientific inquiry), **continental philosophy** typically suggests that "philosophical argument cannot be divorced from the textual and contextual conditions of its historical emergence". Third, continental philosophy typically holds that conscious human agency can change these conditions of possible experience: "if human experience is a contingent creation, then it can be **recreated** in other ways". Thus continental philosophers tend to take a strong interest in the **unity of theory and practice**, and tend to see their philosophical inquiries as closely related to personal, moral, or political transformation. This tendency is very clear in the Marxist tradition ("philosophers have only interpreted the world, in various ways; the point, however, is to change it"), but is also central in **existentialism and post-structuralism**. A final characteristic trait of continental philosophy is an emphasis on metaphilosophy. In the wake of the development and success of the natural sciences, continental philosophers have often sought **to redefine the** method and nature of philosophy. In some cases (such as German idealism or phenomenology), this manifests as a renovation of the traditional view that philosophy is the first, foundational, a priori science. For Kant, judgement **and** object are strictly correlative concepts, so that in the critical sense, the truth of the object is always to be grasped and substantiated only through the truth of the judgement, once the rules of judgement have been defined. In other cases (such as **hermeneutics, critical theory, or structuralism**), it is held that philosophy investigates a domain that is irreducibly cultural or practical. And some continental philosophers (such **as Kierkegaard, Nietzsche, the later Heidegger, or Derrida**) doubt whether any conception of philosophy can coherently achieve its stated goals. **(Source: Wikipedia and Stanford encyclopedia) We left the religious person in the crisis of sickness; but this sickness is not unto death. We shall now let him be strengthened by the very same conception that destroyed him, by the conception of God. First and foremost, in each generation there certainly are not many who suffer through even the beginning of the absolute religious** relationship; and next, that a beginning in the existence-medium is anything but something that is decided once and for all, because it is only on paper that one is finished with the first phase, and then has nothing more to do with it. The absolute decision in the existence-medium still is and remains only an approximation because **the eternal aims from above at the existing person, who by existing is in motion and thus at the moment the eternal touches is already a little moment away from there. The beginning of the absolute decision in the existence-medium is least of all once** and for all, something accomplished, because the existing person is not and abstract X who accomplishes something and then goes further, goes through life, if I may put it this way, undigested; but the existing person becomes **concrete in what has been experienced, and as he proceeds he has it with him and can lose it at any moment. He has it with him, not the way one has something in a pocket, but through this, this specific thing, he is what he is more specifically defined and loses his own more specific definition by losing it.** Through the decision in existence, an existing person, more specifically defined, has become what he is. If he sets it aside, it is not he who has lost something, so that he does not have himself and has lost something, but then he has lost himself and must start from the beginning. The religious person has recovered from his sickness (tomorrow there may be a relapse due to a little unjudiciousness). He perhaps fortifies himself with the upbuilding reflection that God, who created man, certainly knows best all the numerous things that to a human being **appear to be incapable of being joined together with the thought of God-all the earthly desires, all the**

confusion in which he can be trapped, and the necessity of diversion, of rest, as well as a night's sleep. It is obvious that the discussion here is not about the indulgence that is preached in the world, where one human being consoles himself through another, consoles himself reciprocally and leaves God out. Every human being is gloriously structured, but what destroys so many is this confounded talkativeness between man and man about what must be suffered but also be matured in silence, the confession before human beings instead of before God, this candid communication to this one and that one of what ought to be a secret and be before God in secret, this impatient hankering for makeshift consolation. No, in the pain of annihilation, the religious person has learned that human indulgence is of no benefit; therefore he listens to nothing from that corner, but he is before God and suffers through what it means to be a human being and then to be before God. Therefore he cannot be comforted by what the human crowd mutually knows people who have a market-town idea of what it means to be a human being, and a fluent, talkative idea at seventeenth hand of what it means to be before God. From God he must draw his comfort, lest his entire religiousness become a rumor. P. 488-490

The triad is often said to have been extended and adopted by Karl Marx and Friedrich Engels, however, Marx referred to them in The Poverty of Philosophy as speaking Greek and "Wooden trichotomies" .The classic empiricist presentation of this approach is William James's Essays in Radical Empiricism: "There is no general stuff of which appearance is made. It is made of that, or just what appears (26-27) ... what I call **'pure experience' [is] only virtually or potentially either object or subject as yet.** For the time being, it is plain, unqualified actuality, a simple that (23) ... experience as a whole is self-containing and leans on nothing (193) ... **object and subject fuse in the fact of 'presentation' or sense-perception (197) ... knower and object exist as so many ultimate thats or facts of being**" (196). As the last phrase suggests, James's "that" (or "this-that": to signal a differential, or incipient difference) is Francis Bacon's "fact," which can in turn be understood in terms of Whitehead's definition of fact as the "undifferentiated terminus of sense-awareness," the inexhaustible unthought from which thought "diversifies" and to which it "demonstratively" returns, as to its "ideal limit" (6-10, 13-15. **On the "univocity" of the coming to expression of the multiple, see Deleuze 1990, 177-180; 1994, 35-42, 303-304.** Can the nature of thinking be uncovered by thinking alone? (Cf consciousness) "My dialectical method is, in its foundations, not only different from the Hegelian, but exactly opposite to it. For Hegel, the process of thinking, which he even transforms into an independent subject, under the name of 'the idea ', is the creator of the real world, and the real world is only the external appearance of the idea. With me, the reverse is true: the ideal is nothing but the material world reflected in the mind of man, and translated into forms of thought. "Karl Marx, Capital, vol. 1, Preface to the Second Edition. (P.102) (See Ideal / Real) For Aristotle, Thinking is the one specific activity of the Human Soul Which is Capable of Independent Existence and separate from any connection to the Body. A Strenuous counterargument has been made, precisely for body thinking. Deleuze and Guattari for, "Becoming Animal" and "Becoming intense" are Modes of resistance to abstract Thinking. Alain Badiou FOLLOWS Aristotle in using thinking **to define** specifically the Human, as Opposed to Interest. For Badiou, "the capacity which is specifically human is that of thought, and thought is nothing other than that by which the path of a truth seizes and traverses the human animal." **(Infinite Thought, p. 71)** In HIS Essay "Does Consciousness Exist?" **William James** Pointed out That Consciousness is a Process, and not a substance. (Unlike Descartes, who claimed the existence of a res cogitans?) In The Principles of Psychology, **James described five properties of what he called "thought**." Every thought, he wrote, tends to be **part of personal consciousness**. We write this despite the definition that has been held for the consciousness as a storage room and mind anagrammatically acting upon it. Concomitant quantum channeling like synapses and neuron firing might take place the in the brain. And that falls under the subterranean realm and ceratoid dualism of Neurology. Thought is always changing, is sensibly continuous, and appears to deal with objects independent of it. In addition, thought focuses on some objects to the exclusion of others. In other words, it involves attention. Alfred North Whitehead claimed that, with this Inquiry, James was to the Twentieth century What Descartes was to the Seventeenth. Louis Sullivan describes "Real Thinking," Thinking in the present tense, as Organic Thought. "It is in the present, only, that you really live, therefore it is in the present, only that you can really think. And in this sense you think organically. Pseudo-thinking is inorganic. The one is living, the other dead. The present is the organic

moment, the living moment. "(Kindergarten Chats, "Thought.") **Thinking is a form of Computation? Emergence ---------- Emergence Refers to the appearance of Patterns of Organization and is one of the Key concepts of Complexity and a -life. It is sometimes referred to as a situation where the whole is greater than** the sum of the parts, because it cannot be analyzed by taking the parts apart and examining them separately. One reason for this is that in a complex phenomenon **showing** emergent properties, the parts become a determining context for each other, and these patterns of feedback **contribute to the** appearance of the emergent phenomenon. For Michael Polanyi, "**Evolution can be understood only as a feat of emergence**." For Goethe, the Whole Living was Also More than the Sum of ITS parts. "Das ist zwar Lebendige in Elemente zerlegt, aber MAN Kann ES aus Diesen nicht wieder zusammenstellen und beleben." Goethe quoted in d'Arcy Thompson, **On Growth and Form. (P.41)** But for most contemporary scientists, Goethe's meaning of emergence was a mystical one. For Francis Crick, the scientific meaning of emergent assumes that while the **whole cannot be the simple sum of the separate parts, its behavior can, at least in principle, be understood from the nature and behavior of its parts plus the knowledge of how all these parts interact. (The Astonishing Hypothesis, P.11)** (Assuming IT's Possible to have **That Laplacian Knowledge**) Stuart Kauffman Suggests That many Properties of Organisms May be Probable **emergent Collective Properties** of Their constituents. (This is precisely Kant's definition of the organic - as the result of internal interactions instead of an assembly of preexisting parts. (See Mechanism / Vitalism) "The problem of origins requires an understanding of how new levels of order emerge from complex patterns of interactions and what the properties of these emergent structures are in terms of their robustness to perturbation and their capacities for self-maintenance **"(Brian Goodwin, How the Leopard Changed its Spots, p. 181.)** Luc Steels Offers Useful definitions of emergence in Artificial Life half. He distinguishes between first order emergence, defined as a property not explicitly programmed in, and second-order emergence, defined as an emergent behavior that adds additional functionality in the system. In general a-life researchers try to create second -order emergence, then the system for use CANS ITS own emergent Properties to create an upward Spiral of Continuing evolution and emergent behaviors. For a Critique of emergence, See Peter Cariani, "Emergence and adaptivity in Organisms and Devices," World Fudtures 32 (1991):111-32. C. Lloyd Morgan's Descriptions of emergence as a Qualitative Change in direction closely resemble the definitions of bifurcations. HIS In Emergent Evolution, of 1923, he writes, "**The emergent step, although it may seem more or less saltatory, is best Regarded as a Qualitative Change of direction, or Critical turning-Point, in the Course of events**. " Daniel Stern describes the Infant's sense of Self between Birth and two months as the "emergent Self." (See subject). It is a sense of the Coming-into-Being of Organization and Remains active for the rest of life. But IT is not an overarching schema About Integrating the Self, but rather an Experience of Process. (cf Molar / Molecular) In Gestalt Perception terms, emergence refers to our ability to perceive forms that cannot be reduced to the addition of the "atomic" parts of a pattern. Ego superimposing two squares diagonally displaced **allows for** readings of an emergent square at their overlap, two emergent L-shapes in the non-Intersecting area, in addition to the two Squares. (cf. Figure / Ground) A **Fundamental property of Open Systems is that they stabilize any improbability which Serves to elicit them**. Being / Becoming --------- ---------- According to Ernst Cassirer, Plato's Philosophy recognizes two Contrary Forms of representation, one of Which is Valid for the Realm of Being and the Other for the Realm of Becoming. (Individual and Cosmos, p.125) For Cassirer, "**Form Thinking**" Belongs to Being, while "causal Thinking" Belongs to Becoming. But strict Knowledge is Only Possible of the Always-Being. Becoming that which is CAN only be described, if at All, in the language of myth. Or rather, myth is already familiar with both the question of the "what" and the question of the "whence." "It sees everything that it grasps (the world as well as the gods) under this double aspect."(The Problem of Form and the Problem of Cause, in The Logic of the Cultural Sciences, p. 87) Socrates first Discovered the Concept, or Eidos as the relation Between the Particular and the General and as a germ of a new meaning of the general question concerning being. This meaning emerged in its full purity when the Socratic eidos went on to unfold into the (transcendental) Platonic "Idea." (See transcendence / immanence) For Plato, the soul is an intermediary between being and Becoming. The Aristotelian system seemed to Promise a Different reconciliation of the Opposition between Being and Becoming. Unlike Plato and the thinkers of pure form, Aristotle wanted to restore to Becoming ITS rightful Place, Because He was convinced **That Only in this way Could Philosophy be Transformed from a Mere Theory of concepts into a Theory of Reality. Form and Matter, Being and**

Becoming, must become correlative. Peculiarly Aristotelian the Concept of the formal Cause originates from this fusion. (Cassirer) "In Nature We May Search for successfully something that endures, and in nature we may regard becoming as if any phase of it were the "reason" of a later phase and the "consequence" of an earlier one. We may even go so far as to say: Nature is the one mediate object that obeys the postulates of the rational theory of becoming. "Hans Driesch, History and Theory of Vitalism, p. 193. We are grateful to the Russian Blog that was available on Google search and Wikipedia in the consolidation and consummation of the above facts. Gilles Deluze is another person whose Pure Immanence and The Logic of sense has been helpful. Part of it expatiated and enucleated from blog Интернет версия данной статьи находится по адресу: http://www.situation.ru/app/j_art_1041.htm Copyright (c) Альманах "Восток"

One might compare Hegel's point here to that expressed by Kant in his well known claim that without concepts, those singular and immediate mental representations he calls “intuitions” are “blind.” In more recent terminology one might talk of the “concept-” or “theory-ladenness” of all experience, and the lessons of this chapter have been likened to that of Wilfred Sellars's famous criticism of the “myth of the given” (Sellars 1997). Hegel has insisted on quarantining his defensible ideas from it (e.g., Wood 1990). However, rather than be understood as a treatise in formal (or “general”) logic, it is perhaps best understood as a version of what Kant had called “transcendental logic,” and in this sense thought of as a successor to Kant's “transcendental deduction of the categories” in the Critique of Pure Reason in which Kant attempted to “deduce” a list of those non-empirical concepts, the “categories,” which he believed to be presupposed by all empirical judgments made by finite, discursive thinkers. However, whereas the latter’s is largely etymologically oriented and given to exploit-ing semantic ambiguities, traces and aporias, Deleuze’s strategy is more geared towards conceptual and functional differentiation, exploring the horizons of Ideas (in the Kantian sense) and bringingforth the machinic and operative features of the philosophies with which he engages. hile, for Lacan, Truth is this shattering experience of the Void - a sudden insight into the abyss of Being, "not a process so much as a brief traumatic encounter, or illuminating shock, in the midst of common reality" -, for Badiou, Truth is what comes afterward: the long arduous work of fidelity, of enforcing a new law onto the situation. 5 The choice is thus: "whether a vanishing apparition of the real as absent cause (for Lacan) or a forceful transformation of the real into a consistent truth. doctrine is precisely that, while never ceasing to be dialectical in pinpointing the absent cause and its divisive effects on the whole, it nevertheless remains tied to this whole itself and is thus unable to account for the latter's possible transformation. /.../ Surely anchored in the real as a lack of being, a truth procedure is that which gives being to this very lack. Pinpointing the absent cause or constitutive outside of a situation, in other words, remains a dialectical yet idealist tactic, unless this evanescent point of the real is forced, distorted, and extended, in order to give consistency to the real as a new generic truth. Bosteels describes the modality of the truth-procedure: Setting out from the void which prior to the event remains indiscernible in the language of established knowledge, a subjective intervention **names** the event which disappears no sooner than it appears; /it/ faithfully connects as many elements of the situation as possible to this name which is the only trace of the vanished event, and **subsequently forces** the extended situation from the bias of the new truth as if the latter were indeed already generally applicable. Bosteels basic reproach, according to which, psychoanalysis collapses into an instantaneous act what is in reality an ongoing and impure procedure, which from a singular event leads to a generic truth by way of a forced return upon the initial situation. Whereas for Zizek, the empty place of the real that is impossible to symbolize is somehow already the act of truth itself, for Badiou a truth comes about only by forcing the real and by displacing the empty place, so as to make the impossible possible. 'Every truth is post-evental,' Badiou writes. An example is Descartes’ celebrated phrase at the beginning of the Discourse on the Method: Good sense is the most evenly shared thing in the world. . the capacity to judge correctly and to distinguish the true from the false, which is properly what one calls common sense or reason, is naturally equal in all men . . For Deleuze, is an image of thought? Although images of thought take the common form of an ‘Everybody knows . . .’ (DR 130), they are not essentially conscious. Rather, they operate on the level of the social and the unconscious, and function, “all the more effectively in silence.” (DR 167). Relationship of philosophy to thought must have two correlative aspects, Deleuze argues: an attack on the traditional moral image of thought, but also a movement towards understanding thought as self-engendering, an act of creation, not just of what is thought,

but of thought itself, within thought (DR 147). he thought which is born in thought, the act of thinking which is neither given by innateness nor presupposed by reminiscence but engendered in its genitality, is a thought without image. But what is such a thought, and how does it operate in the world? (DR 167; cf. 132) Durkheim's concepts of the collective representation and the process of universalization, then, **correspond to** Kantian concepts, and Durkheim acknowledges the relevance of Kantian philosophy to his epistemological and moral considerations (1965: 494). It is to be noted and stated in unmistakable and unequivocal words that the singularities and the events constitute a life coexist with the accidents of life that corresponds to it, but they are neither grouped nor divided in the same way. They connect in a manner entirely different from how individuals connect by virtue of which classification takes place. A singular life might do without any individuality, without any other concomitant that individualizes it. Individual life or that of insentient life remains inseparable from empirical determinations. On the other hand, the indefinite as such is the mark of not an empirical indetermination but of a determination by immanence or a transcendental determinability. The discovery of non-Euclidean geometries radically undermined this concept of truth. (See scientific space) (**Pure Immanence Gilles Deluze: Page number 30).** Plato and Euclid developed an indissoluble partnership between geometrical and philosophical ideas of truth. The Platonic concept of the theory of ideas was possible only because Plato had continually in mind the static shapes discovered by Greek mathematics. On the other hand, Greek geometry did not achieve completion as a real system until **it adopted** Plato's manner of thinking. (**See Ernst Cassirer**, The Problem of Knowledge.) The concepts and propositions that **Euclid p**laced at the apex of his system were a prototype and pattern for what Plato called the process of synopsis in idea. What is grasped in such synopsis is not the peculiar, fortuitous, or unstable; it possesses universal necessary and eternal truth. (See transcendence / immanence) This is the space of **universal truth that** differs from the spaces of a kind of truth that functions only in the context of local pockets, a truth that is always local, **distributed haphazardly in a plurality of spaces, with regional epistemologies**. It is to be realized that a phenomenon which we think out is engaged in process of actualization following the plane that gives it its particular reality. Plane of immanence therefore is actualization, as long as the events that populate them are localised by a law or other theorem tic corollary. Another vehement and trenchant example of classification is Hegelianism and teleology thereof. Hegelianism is a collective term for schools of thought following or referring to G. W. F. Hegel's philosophy which can be summed up by the dictum that "the rational alone is real", which means that all reality is capable of being expressed in rational categories. His goal was to reduce reality to a more synthetic unity within the system of transcendental idealism. Hegel's method in philosophy consists of the triadic development (Entwicklung) in each concept and each thing. Thus, he hopes, philosophy will not contradict experience, but will give data of experience to the philosophical, which is the ultimately true explanation. If, for instance, we wish to know what liberty is, we take that concept where we first find it—the unrestrained action of the savage, who does not feel the need of repressing any thought, feeling, or tendency to act. Next, we find that the savage has given up this freedom in exchange for its opposite, the restraint, or, as he considers it, the tyranny, of civilization and law. Finally, in the citizen under the rule of law, we find the third stage of development, namely liberty in a higher and a fuller sense than how the savage possessed it—the liberty to do, say, and think many things beyond the power of the savage. In this triadic process, the second stage is the direct opposite, the annihilation, or at least the sublation, of the first. The third stage is the first returned to itself in a higher, truer, richer, and fuller form. The three stages are, therefore, styled:

In itself (An-sich)

Out of itself (Anderssein)

In and for itself (An-und-für-sich).

These three stages are found succeeding one another throughout the whole realm of thought and being, from the most abstract logical process up to the most complicated concrete activity of organized mind in the succession of states or the production of systems of philosophy.

Deleuze uses the introduction to clarify the term "repetition." Deleuze's repetition can be understood by contrasting it to generality. Both words describe events that have some underlying connections. Generality refers to events that are connected through cycles, equalities, and laws. Most phenomena that can be directly described by science are generalities. Seemingly isolated events will occur in the same way over and over again because they are governed by the same laws. Water will flow downhill and sunlight will create warmth because of principles that apply broadly. In the human realm, behavior that accords with norms and laws counts as generality for similar reasons. Science deals mostly with generalities because it seeks to predict reality using reduction and equivalence. Repetition, for Deleuze, can only describe a unique series of things or events. The Borges story in which Pierre Menard reproduces the exact text of Don Quixote is a quintessential repetition: the repetition of Cervantes in Menard takes on a magical quality by virtue of its translation into a different time and place. Art is often a source of repetition because no artistic use of an element is ever truly equivalent to other uses. (Pop Art pushes this quality to a certain limit by bringing production near the level of capitalism.) For humans, repetition is inherently Transgressive. As in Coldness and Cruelty, Deleuze identifies humor and irony as lines of escape from the generalities of society. Humor and irony are in league with repetition because they create distance from laws and norms even while re-enacting them. Deleuze describes repetition as a shared value of an otherwise rather disparate trio: Kierkegaard, Nietzsche, and Péguy. He also connects the idea to Freud's death drive. He goes on to **define** repetition as "difference without a concept". Repetition is thus reliant on difference more deeply than it is opposed. Further, profound repetition will be **characterized b**y profound difference. The Platonic idea of truth entailed a correspondence with the world of forms

Difference in Itself

Deleuze paints a picture of philosophical history in which difference has long been subordinated to four pillars of reason: identity, opposition, analogy, and resemblance. He argues that difference has been treated as a secondary characteristic which emerges when one compares pre-existing things; these things can then be said to have differences. This network of direct relations between identities roughly overlays a much more subtle and involuted network of real differences: gradients, intensities, overlaps, and so forth .The chapter contains a discussion of how various philosophers have treated the emergence of difference within Being. This section uses Duns Scotus, Spinoza, and others to make the case that "there has only ever been one ontological proposition: Being is univocal. ... A single voice raises the clamor of being". One then tries to understand the nature of differences that arise within Being. Deleuze describes how Hegel took contradiction—pure opposition—to be the principle underlying all difference and consequently to be the explanatory principle of the entire world's texture. He accuses this conception of having a theological and metaphysical slant. In an aside to his discussion of "social construction," Ian Hacking registers his own protest to the labels of "culture wars," and "science wars." For Hacking, the willingness to describe fierce disagreements in terms of the metaphors of war makes the very existence of real wars seem more natural, more inevitable, and more a part of the human condition. (While Hacking seems generally unsympathetic to the expression "social construction," he acknowledges that the term is generally invoked against apparent natural inevitability.)

Deleuze's geophilosophy is a "surface topology". (109) Hence, we can use topology to construct and deconstruct the structure of the event. If we take one singularity, one knot, one red dot in the Pascal diagram, and spread it out over a "line of ordinary points," we get the labyrinth, the pulsed string, vibrating like a wave at a frequency in tune with its number. The singularity takes this shape of extension over an "ordinary" line in the actualization of the event. (110) "A world already envelops an infinite system of singularities selected through convergence." (109) this idea of convergence is important to Deleuze. Elsewhere, he writes on hylomorphism. The preindividual transcendental field of singularities is self-organizing. This is how a shape or structure takes its shape, its form, "to be incarnated in a body; to become the state of the body." (110) Deleuze says, "An individual is always in a world as a circle of convergence." To be a circle implies that the circle is the result of an event, a convergence that results not in a knot, or some other kind of shape, but in a circle. I visualize two series coming together, two Pascal triangles, the tips of

which are interlocking in some kind of dragon-chasing-its-tail kind of shape, a yin-yang whose sum total is something like a circle, which also represents the GCD of the two series. The two Pascal triangles intersect and produce a new gyre, and because it has been lifted up into another dimension, we see the cross-section -- the circle. The circle is the stable shape of this event, and as such associated with the individual. Hegel also diagrammed the world as a series of interlocking concentric circles. Different theory, similar diagram. If you can imagine the Pascal triangle as a 3-D gyre, when the event occurs, the potential energy in the system "falls to its lowest level" -- gets pulled to the tip of the triangle (like a tornado). This idea of energy falling to its lowest level is similar to how we referenced the pachinko game earlier. Deleuze wrote extensively on Leibniz, quoting him here that "each individual monad expresses the world." Each monad would be like a singularity, and in discovering all the frequencies of that singularity (e.g. all the factors of a huge number), we gain a perspective on every other singularity. This is very similar to Borges' aleph, a point in space that reflects all the other points by which one can see everything in the universe. Page 112: Deleuze distinguished between zones of clarity and obscurity in the singularity. In other works, this implies "consistency" and "inconsistency". Here, we can think in terms of sense and nonsense. Sense is the ability for the snowflake, crystalline form to incarnate an event. The nonsense is all the other unformed points (non-red dots). Page 114: "Incompossible" worlds (e.g. a world in which Adam is a sinner and a world in which Adam is not a sinner; not a contradiction but a mutual exclusivity) imply that there must be an "ambiguous sign" or aleatory point constructing the difference between the two worlds. This aleatory point, or shifter, is the key, or machine by which the different actualizations are realized. In the case of Adam, the shifter would be something like, "to sin." Worlds are "overthrown from within by paradox", by this aleatory point or ambiguous sign, this moment of becoming and revolution. This is an exemplary case of what Deleuze deploys in the crucial pages of his Difference and Repetition: while it may seem that the two presents are successive, at a variable distance apart in the series of reals, in fact they form, rather, two real series which coexist in relation to a virtual object of another kind, one which constantly circulates and is displaced in them /.../. Repetition is constituted not from one present to another, but between the two coexistent series that these presents form in function of the virtual object (object = x). (DR-104-105) Neither the problem nor the question is a subjective determination marking a moment of insufficiency in knowledge. Problematic structure is part of objects themselves, allowing them to be grasped as signs (DR-63-4)

Hegelian counter-argument would have been: is then the "pure" virtual difference not the very name for actual self-identity? Is it not CONSTITUTIVE of actual identity? More precisely, in the terms of Deleuze's transcendental empiricism, pure difference is the virtual support/condition of actual identity: an entity is perceived as "(self-) identical" when (and only when) its virtual support is reduced to a pure difference. In Lacanese, pure difference concerns the supplement of the virtual object (Lacan's objet a); its most plastic experience is that of a sudden change in (our perception of) an object which, with regard to its positive qualities, remains the same: "although nothing changes, the thing all of a sudden seemed totally different" - as Deleuze would have put it, it is the thing's intensity which changes. Lacan himself is here not beyond reproach, since he gets sometimes seduced by the rhizomatic wealth of language beyond (or, rather, beneath) the formal structure that sustains it. It is in this sense that, in the last decade of his teaching, he deployed the notion of lalangue (sometimes simply translated as "language") which stands for language as the space of illicit pleasures that defy any normativity: the chaotic multitude of homonymies, word-plays, "irregular" metaphoric links and resonances... Productive as this notion is, one should be aware of its limitations. Many commentators have noted that Lacan's last great literary reading, that of Joyce to whom his late seminar (XXIII: Le sinthome is dedicated, is not at the level of his previous great readings (Hamlet, Antigone, and Claudel’s Coufontaine-trilogy). T (For Lacan, the theoretical problem/task is here to distinguish between the Master-Signifier and objet a which both refer to the abyssal X in the object beyond its positive properties.) As such, pure difference is closer to antagonism than to the difference between two positive social groups one of which is to be annihilated. The universalism that sustains an antagonistic struggle **is not exclusive of anyone**, which is why the highest triumph of the antagonistic struggle **is not the** destruction of the enemy, but the explosion of the "universal brotherhood" in which agents of the opposite camp change sides and join us (recall the proverbial scenes of police or military units joining the demonstrators). It is in such explosion of enthusiastic all-encompassing brotherhood from which no one is in principle excluded, that the difference between "us" and "enemy" as positive agents is reduced to

a PURE formal difference. I never pretended that one can insert reality into the past and thus work backwards in time. However, one can without any doubt inserts there the possible, or, rather, at every moment, the possible insert itself there. Insofar as inpredictable and new reality creates itself, its image reflects itself behind itself in the indefinite past: this new reality finds itself all the time having been possible; but it is only at the precise moment of its actual emergence that it begins to always have been, and this is why I say that its possibility, which does not precede its reality, will have preceded it once this reality emerges.

Subjectification (French: subjectivation) is a philosophical concept coined by Michel Foucault and elaborated by Gilles Deleuze and Félix Guattari. It refers to the construction of the individual subject. The concept has been often used in critical theory, sometimes with Louis Althusser's concept of interpellation. In Gilbert Simondon's theory of individuation, subjectification precedes the subject in the same way as the process of individuation precedes the creation of the individual. While the classical notion of a subject considers it as a term, Foucault considered the process of subjectification to have an ontological pre-eminence on the subject as a term.

For Deleuze, the presentation of absolute difference is 'an immediate and adequate expression of an absolute Being that comprises in it all beings.' To cite a phrase Deleuze uses elsewhere, it involves a 'static genesis' of the structure of the absolute. Hegel's Science of Logic, on the other hand, performs a 'dynamic genesis' of 'the logicity of being' in such a way that 'it says its own sense' (accounts for itself through the concepts it has generated) through the very movement of thought presented step-by-step in the book itself. The Logic therefore enacts the complete and immanent interpenetration of the logic of being with the logic of thought. For instance, the movement from being to nothingness and then to becoming at the start of the Logic is simultaneously a movement of thought in which the bare thought of being reveals itself to be nothing determinate. Moreover, it is also through this approach that Hegel completes his response to the Kantian critique of the ontological argument: by arguing that the notion of bare 'existence' or 'being' cannot be conceived without introducing some determinacy into it: to be is to be something. Now Hegel's articulation of the logicity of being is of course only made possible by the claim that difference must be fundamentally understood as negation. We know that Deleuze disagrees with this, but is the necessary consequence of this disagreement that he also has to give up on a determinate and genetic account of the development of thought? If so, then he will have concomitant problems defending his account of immanence against Hegel's. Hegel manages to generate a lot of determinate possibilities out of the structure of negation: it is hard to see what determinate possibilities can be strictly generated from 'difference in itself'. In the Spinozist account, there is no direct movement from the real distinction of the attributes to the position that thought and extensions are two of these attributes. The Vertigo of Philosophy: Deleuze and the Problem of Immanence Christian Kerslake

Deleuze proposes (citing Leibniz) that difference is better understood through the use of dx, the differential. A derivative, dy/dx, determines the structure of a curve while nonetheless existing just outside the curve itself; that is, by describing a virtual tangent. Deleuze argues that difference should fundamentally be the object of affirmation and not negation. As per Nietzsche, negation becomes secondary and epiphenomenal in relation to this primary force.. Repetition for Itself: The chapter describes three different levels of time within which repetition occurs. Deleuze takes as axiomatic the notion that there is no time but the present, which contains past and future. These layers describe different ways in which past and future can be inscribed in a present. As this inscription grows more complicated, the status of the present itself becomes more abstract. Basic processes of the universe have a momentum that they carry into each present moment. A 'contraction' of reality refers to the collection of a diffuse ongoing force into the present. Prior thought and behavior, all substance performs contraction. "We are made of contracted water, earth, light, and air...Every organism, in its **receptive and perceptual elements**, but also in its viscera, is a sum of contractions, of retentions and expectations. Passive synthesis is exemplified by habit. Habit incarnates the past (and gestures to the future) in the present by transforming the weight of experience into an urgency. Habit creates a multitude of "larval selves," each of which functions like a small ego with desires and satisfactions. In Freudian discourse, this is the domain of bound excitations associated with the pleasure principle. Deleuze cites Hume and Bergson as relevant to his understanding of the passive synthesis. Active synthesis: The

second level of time is organized by the active force of memory, which introduces discontinuity into the passage of time by sustaining relationships between more distant events. A discussion of destiny makes clear how memory transforms time and enacts a more profound form of repetition: Destiny never consists in step-by-step deterministic relations between presents which succeed one another according to the order of a represented time. Rather, it implies between successive presents non-localisable connections, actions at a distance, systems of replay, resonance and echoes, objective chances, signs, signals, and roles which transcend spatial locations and temporal successions. Relative to the passive synthesis of habit, memory is virtual and vertical. It deals with events in their depth and structure rather than in their contiguity in time. Where passive syntheses created a field of 'me's,' active synthesis is performed by 'I.' In **the Freudian register**, this synthesis describes the displaced energy of Eros, which becomes a searching and problematizing force rather than a simple stimulus to gratification.

Empty time

The third layer of time still exists in the present, but it does so in a way that breaks free from the simple repetition of time. This level refers to an ultimate event so powerful that it becomes omnipresent. It is a great symbolic event, like the murder to be committed by Oedipus or Hamlet. Upon rising to this level, an actor effaces herself as such and joins the abstract realm of eternal return. The me and the I give way to "the man without name, without family, without qualities, without self or I...the already-Overman whose scattered members gravitate around the sublime image" Empty time is associated with Thanatos, a desexualized energy that runs through all matter and supersedes the particularity of an individual psychic system. Deleuze is careful to point out that there is no reason for Thanatos to (eb) produce a specifically destructive impulse or 'death instinct' in the subject; he conceives of Thanatos as simply indifferent.

Lars Marcussen on Deleuze's philosophy of space and singularities

When the nomad/State opposition is applied to space, the basic principle is that nomad space is 'smooth' and heterogeneous, while State space is 'striated' and homogeneous. Deleuze illustrates these concepts with an example from technology: woven fabric is striated, that is, with the threads of warp and woof; felt is smooth, as it consists of entangled fibres; it is no accident, Deleuze comments, that the Mongolian nomads excel in using felt for their tents, clothing and even armoury. In psychogenetic terms, the difference between metrics and projection is that the straight line is mastered in two different ways: as a 'base line' that structures metric space, and as a 'line of sight' that structures projective space. In the most elementary psychological sense, the straight line of metrics is the line that connects and denotes the distance between two points. When one notes the shortest distance by means of an actual movement, one acquires the idea of the straight line as something that denotes a fixed distance. Eventually, this experience is encoded as a general mental scheme that works automatically in all situations. When we move things around in our imagination so that their relative positions are changed in a regular manner, while their metric identities remain unchanged, the idea gradually emerges of a system of co-ordinates as a regular frame of reference for a metric space that can be expanded to include bigger and bigger entities. As a matter of fact, the very spaces inhabited by nomads – steppes and deserts – are smooth, and the same is true of the ice desert inhabited by Eskimos, and of the sea roamed by seafaring peoples. In these spaces orientations, landmarks and linkages are in continuous variation, Deleuze observes, and goes on: "there is no line separating earth and sky; there is no intermediate distance, no perspective or contour; visibility is limited; and yet there is an extraordinarily fine topology that relies not on points or objects, but rather on haecceities, on sets of relations (winds, undulations of snow or sand, the song of the sand, the creaking of the ice, the tactile qualities of both)." In contrast to this fluid state, the spaces inhabited by sedentary peoples – which are State spaces – are striated with walls, enclosures and roads that exhibit constancy of orientation and metric regularity. Deleuze designates this state of affairs as 'the actual', and he wants to show that although the actual is indeed a part of reality, it is also a kind of illusion that conceals or 'covers up' another realm of reality, which he calls 'the virtual'. In order to grasp the virtual, we should think of everything as 'individuals', each of which has its own history. The word 'individual' is used here in an extended sense. In biology, for instance, species and genera as well as organisms are defined as individuals. On another spatio-temporal scale, ecological

environments in which species are embedded and evolve are individuals, and at the opposite end of the scale each molecule and atom is an individual. In every case individuals are meshed together in fuzzy aggregates without distinct borders, and are in an incessant state of 'becoming' in a sense that escapes our perception of actual processes and states of affairs(**Lars Marcussen on Deleuze's philosophy of space and singularities)**

To utilize and expand on the possibilities offered by differential calculus, Deleuze proposes an Idea in the Kantian sense insofar as it arises from and regulates its field immanently (Deleuze, Difference and Repetition, 177). "Already Leibniz had shown that calculus … expressed problems which could not hitherto be solved or, indeed, even posed **(transcendent problems)," problems such as the complete determination of a species of curve, or problems characterized by** the paradox of Achilles and the tortoise (Deleuze, Difference and Repetition, 177). As technology has come to be seen as inherently dangerous, its unbridled development is seen as potentially leading to war. For Norbert Wiener, "There is nothing more dangerous to contemplate than World War III. It is worth considering whether part of the danger may not be intrinsic in the unguarded **use of learning machines**"(p. 175). For **Wiener,** the new and real agencies of learning machines have a "**literal-mindedness**" whose dangers our prejudices may hide from us. "The mere fact that we have made a machine does not guarantee that we will have the proper information as to whether the danger point has come." Manuel De Landa's study of War in the Age of Intelligent Machines starts with the speculation that one day "robot historians" might study the technological lineages that had lead to generations of killer robots. (cf Terminator)In this point of view, humans would have served only as machines' surrogate reproductive organs until robots acquired their own self-replication abilities. (See intro pp 1-11)

Repetition and Time(Internet Encyclopedia of Philosophy)

For Deleuze, the central stake in the consideration of repetition is time. As with difference, repetition has been subjected to the law of the identical, but also to a prior model of time: to repeat a sentence means, traditionally, to say the same thing twice, at different moments. These different moments must be themselves equal and unbiased, as if time were a flat, featureless expanse. So repetition has essentially been considered as the traditional idea of difference over time understood in a common-sense way, as a succession of moments. Deleuze asks if, given a renovated understanding of difference as in-itself, we are able to reconsider repetition also. But there is also an imperative here, since, if we are to consider difference-in-itself over time, based in the traditional logic of repetition, we once again reach the point of identity. As such, Deleuze's critique of identity must revaluate the question of time.

Deleuze's argument proceeds through three models of time, and relates the concept of repetition to each of them. The first is time as a circle. Circular time is mythical and seasonal time, the repetition of the same after time has passed through its cardinal points. These points may be simple natural repetitions, like the sun rising daily, the movement of summer to spring, or the elements of tragedy, which Deleuze suggests operate cyclically. There is a sense of both destiny and theology in the concept of time as a circle, as a succession of instants which are governed by an external law.

When time is considered in this fashion, Deleuze argues (DR 70-9), repetition is solely concerned with habit. The subject experiences the passing of moments cyclically (the sun will come up every morning), and contracts habits which make sense of time as a continually living present. Habit is thus the passive synthesis of moments that creates a subject. The second model of time is linked by Deleuze to Kant (KCP vii-viii), and it constitutes one of the central ruptures that the Kantian philosophy creates in thought, for Deleuze: this is time as a straight line. In the Critique of Pure Reason, Kant liberates time from the circular model by proposing it as a form that is imposed upon sensory experience. For Deleuze, this reverses the earlier situation by placing events into time (as a line), rather than seeing the chain of events constituting time by the passing of present moments. Habit can thus no longer have any power, since in this model of time, nothing returns. In order for sense to be made of what has occurred, there must be an active process of synthesis, which makes of the past instances a meaning (DR 81). Deleuze calls this second synthesis memory. Unlike habit, memory does not relate to a present, but to a past which has never been present, since it synthesizes from passing moments a form in-itself of things which never existed before the operation. The novels of Marcel Proust are for Deleuze the most profound development of memory as the pure past, or in Proust's terminology, as time regained. (DR 122; PS passim)

In this second model of time, repetition thus has an active sense in line with the synthesis, since it repeats something, in the memory, that did not exist before – this does not save it, however, from being an operation of identity, nonetheless. These two moments, the active constitution of a pure past, and the disparate experience of a present yet to be synthesized produces a further consequence for Deleuze: as in Kant, a radical splitting of the subject into two elements, the I of memory, which is only a process of synthesis, and a self of experience, an ego which undergoes experience. (DR 85-7; KCP viii-ix)

Deleuze insists that both of these models of time press repetition into the service of the identical, and make it a secondary process with regards to time. The final model of time that Deleuze proposes attempts to make repetition itself the form of time.

In order to do this, Deleuze relates the concepts of difference and repetition to each other. If difference is the essence of that which exists, constituting beings as disparates, then neither of the first two models of time does justice to them, insisting as they do on the possibility and even necessity of synthesising differences into identities. It is only when beings are repeated as something other than their disparateness is revealed. Consequently, repetition cannot be understood as a repetition of the same, and becomes liberated from subjugation under the demands of traditional philosophy.

To give body to the conception of repetition as the pure form of time, Deleuze turns to the Nietzsche an concept of the eternal return. This difficult concept is always given a forceful and careful qualification by Deleuze whenever he writes about it (eg. DR 6;41; 242; PI 88-9; NP 94-100): that it must not be considered as the movement of a cycle, as the return of the identical. As a form of time, the eternal return is not the circle of habit, even on the cosmic level. This would only allow the return of something that already existed, of the same, and would result again in the suppression of difference through an inadequate concept of repetition.

While habit returned the same in each instance, and memory dealt with the creation of identity in order to allow experience to be remembered, the eternal return is, for Deleuze, only the repetition of that which differs-from-itself, or, in Nietzsche's terminology, only the repetition of those beings whose being is becoming: "The subject of the eternal return is not the same but the different, not the similar but the dissimilar, not the one but the many . . ." (DR 126)

As such, Deleuze tells us, repetition as the third meaning of time takes the form of the eternal return. Everything that exists as a unity will not return, only that which differs-from-itself. "Difference inhabits repetition." (DR 76). So, while habit was the time of the present, and memory the being of the past, repetition as the eternal return is the time of the future.

The superiority of this third understanding of repetition as time has two main impetuses in Deleuze's argument. The first is obviously that it keeps difference intact in its movement of differing-from-itself. The second is as significant, if for different reasons. If only what differs returns, then the eternal return operates selectively (DR 126; PI 88-9), and this selection is an affirmation of difference, rather than an activity of representation and unification based on the negative, as in Hegel.

. The Image of Thought

Chapter three of Difference and Repetition provides a novel approach to an important question in philosophy, the problem of presuppositions. Deleuze pursues this topic again later in A Thousand Plateaus (374-80), and when he writes about conceptual personae in What is Philosophy? (ch. 3); he had already written on images of thought in Nietzsche and Philosophy (103-10) and Proust and Signs (94-102).

An example is Descartes' celebrated phrase at the beginning of the Discourse on the Method:

Good sense is the most evenly shared thing in the world. . The capacity to judge correctly and to distinguish the true from the false, which is properly what one calls common sense or reason, is naturally equal in all men . .For Descartes, thought has a natural orientation towards truth, just as for Plato, the intellect is naturally drawn towards reason and recollects the true nature of that which exists. This, for Deleuze, is an image of thought.

Although images of thought take the common form of an 'Everybody knows . . .' (DR 130), they are not essentially conscious. Rather, they operate on the level of the social and the unconscious, and function, "all the more effectively in silence." (DR 167)

Deleuze undertakes a thorough analysis of the traditional philosophical image of thought, and lists eight features which, in all aspects of philosophical pursuit, imply a subordination of thought to externally imposed directives. He includes the good nature of thought, the priority of the model or recognition as the means of thought, the sovereignty of representation over supposed elements in nature and thought, and the subordination of culture to method (or learning to knowledge). These all imply an a priori nature of thought, a telos, a meaning and a logic of practice. These features,

Crush thought under an image which is that of the Same and the Similar in representation, but profoundly betrays what it means to think and alienates the two powers of difference and repetition, of philosophical commencement and recommencement. (DR167)

It is this element, in Difference and Repetition that founds Deleuze's most serious criticism of the traditional image of thought: that it fails to come to terms with the true nature of difference and repetition. As a result, it is fair to say that this moment of the book is essential for understanding the way in which Deleuze both wants to base his assessment of traditional philosophies of identity and time, and how he wishes to exceed them: his reformulation of difference and repetition is made possible by this critique (cf. N 149).

The other critical angle Deleuze supplies here is related to the first, and derives from Nietzsche's critique of Western thought:

When Nietzsche questions the most general presuppositions of philosophy, he says that these are essentially moral, since Morality alone is capable of convincing us that thought has a good nature and the thinker a good will, and that only the good can ground the supposed affinity between thought and the True. (DR132; cf. LS 3) As we saw above regarding Hegel, the real point of concern is that this image of thought is in the service of practical, political and moral forces, it is not simply a matter of philosophy, in segregation from the rest of the world.

To the question 'why do we have this image of thought?' Deleuze, along with Nietzsche, that it is a moral image, and is in the service of power, but there is also a more intrinsic problem with thinking itself, that is only fully developed in the Conclusion to What is Philosophy?, and this is that thought itself is dangerous.

In contradistinction to the natural goodness of thought in the traditional image, Deleuze argues for thought as an encounter: "Something in the world forces us to think." (DR 139) These encounters **confront us with** the impotence of thought itself (DR 147), and evoke the need of thought to create in order to cope with the violence and force of these encounters. The traditional image of thought has developed, **just as Nietzsche argues** about the development of morality in The Genealogy of Morals, as a reaction to the threat that these encounters offer. We can consider the traditional image of thought, then, precisely as a symptom of the repression of this violence. As a result, the relationship of philosophy to thought must have two correlative aspects, Deleuze argues: an attack on the traditional moral image of thought, but also a movement towards understanding thought as self-engendering, an act of creation, not just of what is thought, but of thought itself, within thought (DR 147).But what about determinations beyond a single curve? Is there a means to make "a complete determination with regard to the existence and distribution of … [regular and singular] points which depends upon a completely different instance," an instance characterized in terms of a field of vectors (Devlin 44)? The goal here is to explicitly link differential equations and vector fields. A vector field is defined, by Deleuze, as the complete determination of a problem given in terms of the existence, number and distribution of points that are its condition. This corresponds fairly well to the more or less standard mathematical definition where a vector field is defined as associating a vector to every point in the field space (Deleuze, Difference and Repetition, 177). Vector fields are used in physics to model observations, such as the movement of a fluid, which include a direction for each point of the observed space. This is particularly important for Deleuze who claims that nature is a vector field. This is what Deleuze calls a plane of immanence. Referring back to the above quote from What is Philosophy?, concepts are singular points on that vector field. They are called attractors. Deleuze and Guattari claim that they attract affects, percepts, functives, and prospects, which are their components and which constitute them, but never absolutely, since such forces only ever approach singularities infinitely without ever being identical to them. Trajectories are the movements of affects, percepts, functives, and prospects as they pass back and forth (forward and backward in time) on this plane of immanence. This is where the

speed of light enters the picture (33). How do those affects, percepts, functives, and prospects escape the concepts that attract them? Ideally, they exceed the speed of light and thereby they escape the pull of the singularity. Reflections on Time and Politics **(Nathan Widder, Reflections on Time and Politics, Pennsylvania State University Press, 2008)**

The Image of Thought

This chapter takes aim at an "image of thought" that permeates both popular and philosophical discourse. According to this image, thinking naturally gravitates towards truth. Thought is divided easily into categories of truth and error. The model for thought comes from the educational institution, in which a master sets a problem and the pupil produces a solution which is either true or false. This image of the subject supposes that there are different faculties, each of which ideally grasps the particular domain of reality to which it is most suited. In philosophy, this conception results in discourses predicated on the argument that "Everybody knows..." the truth of some basic idea. Descartes, for example, appeals to the idea that everyone can at least think and therefore exists. Deleuze points out that philosophy of this type attempts to eliminate all objective presuppositions while maintaining subjective ones. Deleuze maintains, with Artaud, that real thinking is one of the most difficult challenges there is. Thinking requires a confrontation with stupidity, the state of being formlessly human without engaging any real problems. One discovers that the real path to truth is through the production of sense: the creation of a texture for thought that relates it to its object. Sense is the membrane that relates thought to its other. Accordingly, learning is not the memorization of facts but the coordination of thought with a reality. "As a result, 'learning' always takes place in and through the unconscious, thereby establishing the bond of a profound complicity between nature and mind". Deleuze's alternate image of thought is based on difference, which creates a dynamism that traverses individual faculties and conceptions. This thought is fundamentally energetic and asignifying: if it produces propositions, these are wholly secondary to its development.

Philosophy merges with ontology; ontology merges with univocity of being; analogy has always a theological vision; not a philosophical vision; one becomes adapted to the forms of god; self and world; the univocity of being does not mean that there is one and the same being; on the contrary, beings are multiple and different they are always produced by disjunctive synthesis; and they themselves are disintegrated and disjoint and divergent; membra disjuncta.the univocity of being signifies that that being is a voice that is said and it is said in one and the same "consciousness". Everything about which consciousness is spoken about. being is the same for everything for which it is said like gravity; it occurs therefore as an unique event for everything; for everything for which it happens; even tum tan tum;it is the ultimate form for all of the forms; and all these forms are disjointed; it brings about resonance and ramification of its disjunction; the vorticity of being merges with the positive use of the disjunctive synthesis, and this is the highest affirmation of its univocity like gravity; it is the eternal resurrection or a return itself, the affirmation of all chance in a single moment, the unique cast for all throws; a simple rejoin der for Einstein's god does not play dice; one being, one consciousness, for all forms and all times. a single instance for all that exists, a single phantom for all the living; a single voice for every hum of voices or a single silence for all the silences; a single vacuum for all the vacuousness consciousness should not be said without conjuring if consciousness is one unique event in which all the events communicate with each other; univocity refers both to what occurs to what it is said. The attributable to all states of bodies and states of affairs and the expressible of every proposition. So univocity of consciousness means the identity of the noematic attribute and that which is expressed linguistically and sense-fully. Univovity.We means that it does not allow consciousness to be subsisting in a quasi state and but expresses in all pervading reality.

At the end of the chapter, Deleuze sums up the image of thought he critiques with eight attributes: the postulate of the principle, or the Cogitatio natural universalis (good will of the thinker and good nature of thought); the postulate of the ideal, or common sense (common sense as the concordia facultatum and good sense as the distribution which guarantees this concord); the postulate of the model, or of recognition (recognition inviting all the faculties to exercise themselves upon an object supposedly the same, and the consequent possibility of error in the distribution

when one faculty confuses one of its objects with a different object of another faculty); the postulate of the element or of representation (when difference is subordinated to the complimentary dimensions of the Same and the Similar, the Analogous and the Opposed; the postulate of the negative, or of error (in which error expresses everything which can go wrong in thought, but only as the product of external mechanisms); the postulate of logical function, or the proposition (designation is taken to be the locus of truth, sense being no more than the neutralized double or the infinite doubling of the proposition); the postulate of modality, or solutions (problems being materially traced from propositions or indeed, formally defined by the possibility of their being solved); the postulate of the end, or result, the postulate of knowledge (the subordination of learning to knowledge, and of culture to method.

It is to be stated that we are here reproducing the following for the bastion and stylobatishness of the necessity of classificational procedural formalities and I donot believe in whatever that has been stated. Nor is it meant as propaganda. It is purely definitional with axiomatic predications and doctrinal manifestations. In Vedas and Upanishads the classification processual formalities of Universal action is undertaken thus with a mention that the proportion of the ones mentioned would be different in different individuals and individuals themselves are investigating systems notwithstanding a wider classification which is nothing but more than a Universal Law. Manduka Upanishad divides all knowledge into two categories. The knowledge that leads to Self Realization is called Para Vidya (Great or Divine Knowledge) and everything else is called Apara Vidya or Knowledge of Material world (wordly knowledge). Shaunaka approaches sage Angiras and asks "Revered Sir, by knowing what everything will be known?" Angiras replies that two know ledges should be known, one is **Para Vidya** and other is **Apara Vidya.** Knowledge of worldly things is Apara Vidya and that by which Eternal Truth or Akshara is obtained is Para Vidya. Though Apara Vidya enables one to earn ones bread and helps one to understand each object of universe separately, it does not show the Ultimate Reality (Akshara) or Root Cause of this universe. While Para Vidya doesn't teach objects of this universe but enables one to understand underlying fabric of it. Like by knowing gold all the gold ornaments could be known, by knowing Akshara, it's another manifestation, the universe is known. This Upanishad expounds the greatness of Para. To think whether a certain thing may be eaten is a thought-form of the mind; "It is good. It is not good. It can be eaten. It cannot be eaten", discriminating notions like these constitute the discriminative intellect. Because the mind alone constitutes the root-principle manifesting as the individual, God and the world, its absorption or submergence and dissolution in the Self as pure Consciousness is the final emancipation known as Kaivalya and in the Supreme Spirit, the Brahman.(Ramana maharishi and Wikipedia). **Tamas (Darkness):** Complete delusion, ignorance, illiberality, indecision in respect of action, sleep, haughtiness, fear, cupidity, grief, censure of good acts, loss of memory, unripeness of judgment, absence of faith, violation of all rules of conduct, want of discrimination, blindness, vileness of behaviour, boastful assertions of performance when there has been no performance, presumption of knowledge in ignorance, unfriendliness (or hostility), evilness of disposition, absence of faith, stupid reasoning, crookedness, incapacity for association, sinful action, senselessness, stolidity, lassitude, absence of self-control, degradation, - all these qualities are known as belonging to Darkness **(Tamas).** Whatever other states of mind connected with delusion exist in the world, all appertain to Darkness. Frequent ill-speaking of other people, censuring the deities and the Brahmanas (priests), illiberality, vanity, delusion, wrath, unforgiveness, hostility towards all creatures, are regarded as the characteristics of Darkness. Whatever undertakings exist that are unmeritorious (in consequence of their being vain or useless), what gifts there are that are unmeritorious (in consequence of the unworthiness of the done, the unreasonableness of the time, the impropriety of the object, etc.), vain eating, - these also appertain to **Darknesss (Tamas).** Indulgence in calumny, unforgiveness, animosity, vanity, and absence of faith are also said to be characteristics of Darkness. Whatever men there are in this world who are characterised by these and other faults of a similar kind, and who break through the restraints provided by the scriptures, are all regarded as belonging to the quality of Darkness. Here Rishis and Munis (Seers and sages), and deities become deluded, desirous of pleasure. Darkness, delusion, the great delusion, the great obscurity called wrath, and death, that blinding obscurity (these are the five great afflictions). As regards wrath, that is the great obscurity (and not aversion or hatred as is sometimes included in the list). With respect then to its colour (nature), its characteristics, and its source, I have, ye learned Brahmanas, declared to you, accurately and in due order, everything about the quality of Darkness (Tamas). Who is there that truly understands it? Who is there that

truly sees it? That, indeed, is the characteristic of Darkness, viz., the beholding of reality in what is not real. The qualities of Darkness have been declared to you in various ways.

Rajas (Passion): Injuring others, beauty, toil, pleasure and pain, cold and heat, lordship (or power), war, peace, arguments, dissatisfaction, endurance, might, valour, pride, wrath, exertion, quarrel, jealousy, desire, malice, battle, the sense of meum or mineness, protection of others, slaughter, bonds, and affliction, buying and selling, lopping off, cutting, piercing and cutting off the coat of mail that another has worn, fierceness, cruelty, vilifying, pointing out the faults of others, thoughts entirely devoted to worldly affairs, anxiety, animosity, reviling of others, false speech, false or vain gifts, hesitancy or doubts, boastfulness of speech, praise and criticisms, laudation, prowess, defiance, attendance (as on the weak and the sick), obedience (to the commands of preceptors and parents), service or ministrations, harbouring of thirst or desire, cleverness or dexterity of conduct, policy heedlessness, contumely, possessions, and diverse decorations that prevail in the world among men, women, animals, inanimate things, houses, grief, incredulousness, vows and regulations, actions with expectation (of good result), diverse acts of public charity, the rites in respect of Swaha salutations, rites of Swadha and Vashat, officiating at the sacrifices of others, imparting of instruction, performance of sacrifices, study, making of gifts, acceptance of gifts, rites of expiation, auspicious acts, the wish to have this and that, affection generated by the merits of the object for which or whom it is felt, treachery, deception, disrespect and respect, theft, killing, desire of concealment, vexation, wakefulness, ostentation, haughtiness, attachment, devotion, contentment, exultation, gambling, indulgence in scandal, all relations arising out of women, attachment to dancing, instrumental music and songs – all these qualities have been said to belong to the quality of Passion (Rajas). Those men on earth who meditate on the past, present and the future, who are devoted to the aggregate of the three viz., Religion, Wealth and Pleasure, who acting from impulse of desire, exult on attaining to affluence in respect of every desire, are said to be enveloped by Passion (Rajas). These men have downward courses. Repeatedly reborn in this world, they give themselves up to pleasure. They covet what belongs to the world as also all those fruits that belong to the world hereafter. They make gifts, accept gifts, offer oblations to the Pitris, and pour libations on the sacrificial fire.

Sattwa (Goodness): Brahma (the Grandsire Prajapati) said: Sattwa is beneficial to all creatures in the world, and unblamable, and constitutes the conduct of those that are good. Joy, satisfaction, nobility, enlightenment, and happiness, absence of stinginess, absence of fear, contentment, disposition for faith, forgiveness, courage, abstention from injuring any creature, equability, truth, straightforwardness, absence of wrath, absence of malice, purity, cleverness, prowess- these appertain to the quality of Goodness (Sattwa).He who is devoted to the duty of Yoga, regarding knowledge to be vain, conduct to be vain, service to be vain, and mode of life to be vain, attains to what is highest in the world hereafter. Freedom from the idea of meum (mineness), freedom from egoism, freedom from expectations, looking upon all with an equal eye, and freedom from desire, - these constitute the eternal religion of the good. Confidence, modesty, forgiveness, renunciation, purity, absence of laziness, absence of cruelty, absence of delusion, compassion to all creatures, absence of the disposition to calumniate, exultation, satisfaction, rapture, humility, good behaviour, purity in all acts having for their object the attainment of tranquillity, righteous understanding, emancipation from attachments, indifference, Brahmacharya (celibacy), complete renunciation, freedom from the idea of meum, freedom from expectations, unbroken observance of righteousness, beliefs that gifts are vain, sacrifices are vain, study is vain, vows are vain, acceptance of gifts is vain, observance of duties is vain, and penances are vain – those Brahmanas (priests) in this world, whose conduct is marked by these virtues, who adhere to righteousness, who abide in the Vedas, are said to be wise and possessed of correctness of vision.[Note: Compare from The Mahabharata, Aswamedha Parva, and Sec.XLIV): Brahma said: The Unmanifest is the source of all the worlds as, indeed, that is the end of everything. Days end with the sun's setting and Nights with the sun's rising. The end of pleasure is always sorrow, and the end of sorrow is always pleasure. All accumulations have exhaustion for their end, and all ascent have falls for their end. All associations have dissociations for their end, and life has death for its end. All action ends in destruction and all that is born is certain to meet with death. Every mobile and immobile thing in this world is transient. Sacrifice, gift, penances, study, vows, observances, - all these have destruction for their end. Of Knowledge, there is no end. Hence, one that is possessed of a tranquil soul, that

has subjugated his senses, that is freed from the sense of meum (mineness), that is devoid of egoism, is released from all sins by pure knowledge.] Casting off all sins and freed from grief, those men possessed of wisdom attain to Heaven and create diverse bodies for themselves. Attaining the power of governing everything, self-restraint, minuteness, and these high-souled ones make operations of their own mind, like the gods themselves dwelling in Heaven. Such men are said to have their courses directed upwards. They are veritable gods capable of modifying all things. Attaining to Heaven they modify all things by their very nature. They get whatever objects they desire and enjoy them. Thus have I, ye foremost of regenerate ones, described to you what that conduct is which appertains to the quality of Goodness (Sattwa). Understanding this duly, one acquires whatever objects one desire. The qualities that appertain to Goodness have been declared particularly. The conduct which those qualities constitute has also been properly set forth. That man, who always understands these qualities, succeeds in enjoying the qualities without being attached to them. **Sattwa, Rajas & Tamas** Whatever object exists in this world, everything in it is fraught with the three qualities.

It is to be noted that the primary focus and locus of homologous repetitiveness and differentially instrumental activity we are talking about is the application of some certain Universal laws should not be mistaken for the Universalistic laws and particularistic laws themselves. All we are talking about is the application of paradigmatic relational content to specific investigating systems under consideration**.** On the other hand there certainly exist **Universalistic** Laws and Particularistic Laws. We are not in from any point of view talking about the imperative compatibilities and structural variabilities of the two different forms of laws but the application of say Einstein's Law of mass energy equivalence to some certain galaxy which is under investigation. Following essay allays and ameliorates the vestiges of doubt in the eventuality of confoundment between the two schools of thought we have essayed. It is simple application systems that are also investigatory and conforms and congruenential with the given Einsteinian Theory. **This universalism which is not one Linda M. G. Zerilli From: Diacritics Volume 28, Number 2, summer 1998 pp. 3-20 | 10.1353/dia.1998.0013 In lieu of an abstract, here is a brief excerpt of the content: Diacritics 28.2 (1998) 3-20 Review Essay Ernesto Laclau. Emancipation(s). London: Verso, 1996.** Judging from the recent spate of publications devoted to the question of the universal, it appears that, in the view of some critics, we are witnessing a reevaluation of its dismantling in twentieth-century thought. One of the many oddities about this "return of the universal" is the idea that contemporary engagements with it are more or less of a piece, and that they reflect a growing consensus that poststructuralist political theories are incapable of generating a viable alternative to the collective fragmentation that characterizes late modernity. The putative return to the universal marks, on this view, both a homecoming to Enlightenment ideals -- purified of their more poisonous elements, of course -- and a reconciliation of sorts between those who refuted these ideals and those who sought to realize them. Now that "we" all know and agree that post structuralism is critically valuable but politically bankrupt; now that we all know and agree that the "old universal" was indeed a "pseudo-universal," so the homecoming narrative goes; we can get on with the project of constructing a "new universal." This authentic universal would really be inclusive of all people, regardless of race, class, gender, sexuality, ethnicity, nationality, and whatever else attaches to the "embarrassing etcetera" that, as Judith Butler reminds us, inevitably accompanies such gestures of acknowledging human diversity. Existentialism is as much a way of life as it is a philosophy. In the existentialism proffered by atheistic philosophers it is a life-view where the individual, in a universe without God, and thus without revealed morality, must create his own system of ethics. It is a life-view where the individual is ultimately responsible for his actions. With man at the center of all things, it is up to each individual to create an essence out of the facthood of his own existence. This does not necessarily lead to a nihilistic schema void of ethics—though it certainly can—but rather an ethical system in which man builds meaning out of meaninglessness, and thus brings order out of chaos.

The theistic existentialist also emphasizes the individual and personal responsibility—but the individual before God. The existence of a moral system given to us by God in no way limits our responsibility and necessity to live our own

lives with intelligence and volition. Theistic existentialism recognizes the chaotic and ferocity in man, and presents it to God for forgiveness, healing, and strength. This is not an abdication of responsibility. In fact, God requires that we make choices, and he honors them.

Aspects of existentialism (See Kierkegaard and Wikipedia)

Peter Angeles lists several general features of existentialism.

1) EXISTENCE precedes ESSENCE. Forms do not determine existence to be what it is. Existence fortuitously becomes and is whatever it becomes and is, and that existence then makes up its "essence". 2) An individual has no essential nature, no self-identity other than that involved in the act of choosing. 3) Truth is subjectivity. 4) Abstractions can never grasp nor communicate the reality of individual existence. 5) Philosophy must concern itself with the human predicament and inner states such as alienation, anxiety, inauthenticity, dread, sense of nothingness, anticipation of death. 6) The universe has no rational direction or scheme. It is meaningless and absurd. 7) The universe does not provide moral rules. Moral principles are constructed by humans in the context of being responsible for their actions and for the actions of others. 8) Individual actions are unpredictable. 9) Individuals have complete freedom of the will. 10) Individuals cannot help but make choices. 11) An individual can become completely other than what he is (Dictionary of Philosophy, p. 88).

It should be apparent to many readers that these definitions do not all apply to Kierkegaardian existentialism, but have been included for the sake of completeness. In fact, Kierkegaard rarely used the word existential and its cognates, and not usually in the modern sense of the terms. Of the characteristics above number 1 (in part), 3, 4, 5, 9 (in part) and 10 more directly apply to Kierkegaardian dialectic.

Before signing on to this felicitous agreement about "**the necessity of universalism**," we may wish to know whether we have anything like a minimal agreement in language, that is, whether we who speak of this universal are even speaking about the same thing. Apart from the not insignificant problem of translating from a philosophical to a political idiom, the whole question of this agreement is virtually occluded by the rush to rescue politics from the virulent particularisms **that admit no** common ground or sense of collective belonging. Presented in terms of the familiar binary couple, the choice between universalism and particularism seems settled by merely pointing to global and domestic political realities. Universalism is the only alternative to social fragmentation, wild child of the collapse of communism, the rise of deadly nationalisms, and the multiculturalist romance with particularism. To invoke the name of the universal in any affirmative sense is already to sign on to the political diagnosis and its solution. One of the many virtues of Ernesto Laclau's Emancipation(s) is that it offers both an alternative to the binarisms spawned by the "return" to the universal (for example, false universalism/true universalism) and a trenchant critique of the original binary couple itself (universalism/ particularism). Demonstrating the imbrications of the universal and the particular, Laclau shows why it is a matter not of choosing one over the other but of articulating, in a scrupulously political sense, the relation between the two. He thus explicitly rejects the notion that this relation is one of mutual exclusion, and shows that the tendency to see it as just that has led to the impasse of the contemporary debate, an impasse that is glossed over in some highly visible academic cases by proclaiming the necessary return to the universal. Although the language of universalism as spoken by Laclau searches for some common ground between particularists and Universalists, it is more by way of articulating their mutual contamination, that is, how each is rendered impure by the irreducible presence of the other. Aristotle assumes that there is a plurality of individual things (substances) that are characterized by intrinsic properties (forms) each. David Lewis, an analytical philosopher, provided the thesis of Humean supervenience: at the basic level of the world, there are only local qualities in the sense of intrinsic properties instantiated by space-time points or point-sized particles or field sources at space-time points. Space-time points can qualify as individual things in the above mentioned sense. Everything there is in a world like ours supervenes on the distribution of basic intrinsic properties over all space–

time points. Whether really everything supervenes on that distribution is not relevant to the present paper. What is important here is the claim that, except for spatio-temporal relations, all the relations between the things at the basic level supervene on their intrinsic properties. We can apply the fundamental principles of mathematical proofs to locate to the True form of nature natural things: Through the systematic observation and analysis (breaking down and classification) of the natural world, in combination with rigorous logic we can make True statements about the natural world and understand: 1) The nature of essences (what something is) 2) The nature of causes (why things occur) Unlike Plato, Aristotle also believes that the other arts are very useful for helping us understand things. Relevant to our course is that he believes argument -- or dialectic -- is a key ingredient for people reaching understandings: by arguing over issues the truth and falsity of the claims becomes increasingly apparent, where Plato might have believed that honing such rhetorical arts only confuses the matter. Entities are not separately determinate individuals but rather inseparable parts of a single phenomenon. In particular, there are no preexisting-individually-determinate-entities-with-determinate-spatial-positions-communicating-instantaneously-at-some-remove-from-one-another outside of a phenomenon that determinately resolves the boundaries and properties of the entangled components in a way that gives meaning to the notion of individual. Indeed, "individual" is ontologically and semantically indeterminate in the absence of an apparatus that resolves the inherent indeterminacy in a way that makes this notion intelligible (Barad: 316)".

Deleuze's subrepresentational self-differentiation is superior to Hegel's infinite representational self-differentiation in solving the problems those self-identity causes in Kant's and Aristotle's (likewise Russell's) representational systems.

Self-identity in representational systems leads to problems in explaining the highest part, the lowest part, and the compositional principle of Kant's, Aristotle's (and Russell's) representational systems. Hegel's and Deleuze's principles of self-difference can solve them. Hegel's productive self-contradiction and sublation suffices logically but not in application to evolution. Deleuze's non-oppositional subrepresentational difference however does suffice in this regard.

A strict view of logical identity leads to problems in Kant's and Aristotle's representational systems. The unities of and between concept and intuition that enable our subject-predicate judgments of the world for Kant are based on the unity of a transcendental self. But Sartre shows this is merely a convenient assumption, because for him the unity of consciousness of the object is based on the continuous unity of the object's givenness. For Deleuze the grounds for our judgments are based on neither the unity of the subject, of the concept, nor of the object, but rather on the unity of incompossible undetermined predicates implying a subject with virtual variations. As it is made of the integration of incompossibilities, it lacks the coherence of self-unity necessary for representation. Another question regards the representational nature of the categories we use for judgment. Aristotle and Russell have hierarchies, but because they exclude self-reference and excluded middle, the very foundations (largest parts), compositions of individuals (smallest parts), and method of composition (division/class inclusion) of their representational systems are problematic and irrepresentable within the systems themselves. Hegel and Deleuze have different ways of solving this by incorporating self-difference into their systems. Hegel's productive self-opposition creates a genetic line of sublated categories. Deleuze's Bergsonian continuously integrated heterogeneous multiplicity allows a plurality of differentially related incompossible actualizations to coexist virtually. Hegel's and Deleuze's interpretations of differential calculus show us that for Hegel there are ultimately determinate parts that are not finite but make finite values when differentially related; where for Deleuze there are ultimate undetermined parts that are nonetheless determinable as extensive finite values when differentially related. For Hegel the unconditioned (dialectic) and the conditioned (categories it creates) are on the same ontological plane, but for Deleuze the unconditioned (genesis of virtual differentials) and the conditioned (actualization of extensities) are on two different planes, the virtual and the actual, so they cannot succumb to Hegel's system by being sublated. However, Hegel's dialectic can be seen within Deleuze's system as a secondary movement to the genesis of difference. Hegel's theory

cannot explain the creation of variation needed in evolutionary, but Deleuze's can. **One cannot think condition in terms of the conditioned.**

Deleuze and Hegel offer solutions to the problems of representational systems, which are philosophical systems based on self-unity, identity, and the law of excluded middle. For Kant, our judgments of things have a subject-predicate structure that is parallel to the subject-predicate structure of concepts and intuitive objects (having the subject-property structure). What unifies each them and all with each other is the a priori unity of a transcendental self. Sartre thinks the unity of objects precedes that of the self. Deleuze's transcendental empiricism posits neither the unity of the self nor of the object; things obtain something like a subject-predicate form from incompossibilities being various and indeterminate, but the thing taking these possible predicates is the subject of the judgments. A strict adherence to the principle of identity and the law of excluded middle causes problems in Aristotle's and Russell's logical systems of classification. For Aristotle, the highest genus is being or unity. But, as it has no genus above it, it cannot be defined according to the structure of this system of division (the problem of the large) and yet all beings under it are characterized by this irrepresentable classification. The species gives the essence, and under the species is the individual, which is distinguished from other individuals not by essential but by accidental traits. But in moments of change, something with one essence has contradictory accidents, and also we don't know until after the change what was essential and what was accidental. So the individual cannot be represented properly in this system (the problem of the small). Also there are cases in the natural world, ring species, which cannot be classified using Aristotle's system of division (the problem of division). Russell also has the problem of the large. He must ban self-reference from his system of set classifications so to avoid the paradox of the set of all non-self-including sets. Yet self-reference is needed to establish identity. Hegel's dialectic and Deleuze's philosophy of difference are competing solutions to these problems with representational systems. Deleuze's system is based on Bergson's continuously integrated heterogeneous multiplicity. We can understand it **using Riemann topographical space**. Deleuze's virtual/actual relation is like topographical phase space portraits, where we can see all possible actual ways a system can behave. All the incompossible actualizations are differentially related yet are continuously integrated. Hegel's internal dialectic makes use of productive contradiction: from out of a concept arises its contradiction, and from out of that opposition arises a new concept not implied in the first ones. So contraries are located within one another, and there is a genetic chain of production of the categories of understanding. Finite thought like that used in Kant's and Aristotle's systems would find such contradiction unthinkable, but Hegel's infinite thought can think these contradictions. Hegel's and Deleuze's interpretations of the differential calculus help us elaborate their positions. For Hegel, the differential is a relation between vanishing values, they are caught in the act of the sublation of the finite and infinite: the values are vanishing and hence are not finite, but their differential ratio is a finite value; and each is constitutive of the other. For Deleuze, the differential values are not determinate, yet they are determinable in differential relation to one another. For, they are subrepresentational, meaning that they are on a level where parts are not externally exclusive like in extensity, so they are not self-identifiable. The important distinction is that for Hegel the terms are represent able and determinate, but are only thinkable using infinite thought. In examining Kant's antimony of the beginning or beginningless of the world, we see that for Kant, we have this antinomy because we mistakenly think the **unconditioned is among the conditioned**, when in fact it is noumenal; for Hegel the antinomy indicates the sublation of finite (the necessity for a limit) and infinite (the necessity for all limits to be surpassed), and for him the conditioned and unconditioned are on the same level; yet for Deleuze, the conditioned (actual) is on a different level than the unconditioned (virtual) but the virtual is not outside our knowledge, rather it is only knowable outside representational thinking. Hegel could subsume Deleuze's virtual and actual by sublating them, but that would fail since they are two tendencies of the real and are not really opposites. Deleuze could subsume Hegel's dialectic by saying it is a secondary movement to genesis of pure difference that happens on the level of actuality and representation. But Hegel could say that from the perspective of logic there is no such thing as Deleuze's difference. So we test them by seeing how their theories of the composition of the organism suffice in evolutionary theory. Hegel is like Cuvier in thinking that the organism's parts are matters of how they **function in** service of the whole (teleological); organ and organism, individual and species are like **sublated** opposing terms. But this means that deformations are deteriorations in structure and **thus cause**

deficiencies in functionality. But evolutionary theory depends on a positive account of anatomical variations; natural selection needs to pick the best from a variety of mutations. Geoffrey's and Deleuze's view sees variations as different actualizations of a transcendental model, so their view is more compatible with evolutionary theory. Because Hegel's anatomy is representational, we can see one superiority in Deleuze's subrepresentational response to representational philosophies as opposed to Hegel's infinite representational approach.

Henry Somers-Hall will examine how Deleuze and Hegel respond to the shared problematic of representation in philosophy. He then pits them against one another, and applies them to the role of the structure of the organism in evolution, to evaluate them with respect to one another. Antonio Negri gives a more overtly militant dimension to truth when he claims that "The sole criterion of truth, for me, is action. Truth is itself an action.... a confrontation. When one acts, one goes beyond solitude **because (e)** to act is to search for the truth, and truth always appears in common." (Negri on Negri, p.26). This is true of quantum mechanics too. The "local" is a politically contested concept. Is it a site of resistance to globalization and the hegemonic ideological systems that go under the names of the West, McWorld, and EMPIRE? Is the local the only viable link to tradition, or is it a source of fascisms, of ethnic atavisms, of all the raging cultural fundamentalisms and defensively defined communities that Benjamin Barber calls jihad? We shall say that QM rather unmistakably **is (e) recalcitrant** towards to classical mechanics.

In the first part, we begin by seeing how the history of philosophy contains tendencies toward building theories and systems on the basis of representation. This means that they make use of principles of self-unity, identity, and the law of excluded middle. There are problems with these approaches. Deleuze and Hegel offer solutions. There are two cases under investigation. The first is the transcendental grounds of our knowledge, specifically, what principle allows us to make judgments of the world? Kant offers a representational theory. It is representational, because it is based on a self-identical a priori unified self that is represented in all inner acts, in their accompanying 'I think'. This unity unifies the empirical world into things with a subject-property structure, it unifies our concepts into subject-predicate structure, and it unifies our concepts and our intuitions into represent able judgments with the subject-predicate structure. Sartre's critical stance would say that it is really the unity of objects, and not subjects, that comes first and the unity of the self comes secondly. For Deleuze, a unity neither of consciousness, of self, nor of the objects, is what grounds our subject-predicate knowledge of things. Rather, each moment, events can go many different ways, so the same subject now has many various undetermined predicates, and they are incompossible. Because they are contradictory, the predication of a subject is not representing able, even though the subject-predicate structure is there. This is Deleuze's transcendental empiricism. Another case of representation in the history of philosophy is the use of the principle of identity and excluded middle in Aristotle's and Russell's theories of classification. For Aristotle, we define species on the basis of their differences. But the highest genus, being or unity, has nothing to differentiate from, no genus above it or species beside it, so it is indeterminate. However, it is the basic principle saying that all beings are self-unified and have identities (and thus also the system is thoroughly representational). The very representational basis of his system is not itself representational. Russell's theory of class inclusion is also representational. Things are strictly categorized and defined by their groupings. There cannot be contradiction in the system or instances when something's identity contradicts its classification. So it cannot have the paradoxical class of all non-self-inclusive classes. Such a class is meaningless; it cannot be represented in the system. And yet, such a class is based on the same notion of inclusion as all the others. Hence class inclusion, as a universal concept that forms the basis of all instances of classification in his system, is not represent able in this system. So somehow the nature of inclusion for each level is distinguishable, when in fact it is the same sort of inclusion each time. Also, identities and essences are representing able, but moments of self-contradiction do not fit into such representable systems. This means that when something is changing, we cannot represent what is happening in the phase of transition when contradictory properties are coincident (like being both wood and fire in the action of ignition). Hegel's solution is to make contradiction productive, using internal dialectic, where some concept brings about its own self-contradiction, and out of it comes a new concept not implied in the first. For Deleuze, this solution still has the problems of representation, as we will later see. Deleuze's solution is a non-oppositional concept of difference.

In the second part, we formulate Deleuze's and Hegel's alternate proposals. Deleuze's is based on Bergson's duration, which is a continuously-integrated heterogeneous multiplicity, unlike the discrete multiplicity of externally related extensive parts characterizing homogeneous space. Bergson's heterogeneous multiplicity better explains living systems.

Deleuze then uses Bergson's continuously integrated heterogeneous multiplicity to characterize the Idea, the problem, and the concept. In all cases, they are terrains of virtual differential incompossibly-actualizeable paths of developmental explication. We can understand them with the model of topographical phase space portraits. They indicate all the tendencies for a system's development using terrain features. This describes the system's behavior on a whole, but in each instantiation only one possible line of development indicated in the map is actualized, because all the lines are incompossible yet coincident in this virtual form. They explicate into extensity. And any one actualization implicates the totality of the whole 'problematic'.

Hegel's dialectical movement brings contraries within one another, and also unites them on the basis of their genetic productions of one another. This allows him to go beyond Kant's finite thought, and also to have totality to his system and an account of change, which is lacking in Aristotle's system. Kant cannot go beyond finite thought, because it cannot think contraries together, like Hegel's infinite thought can. And because Hegel's concepts are united genetically, differences are inherently linked, and thus he can have totality to a system of differences without the need of some generic category to encompass them all, which was a source of a problem for Aristotle. Rather, their genetic process of unfolding is the glue uniting the differences. Thus Hegel can also explain the process of change, as he has accounted for the process that generates and unites the diverse contradictory changes when something alters.

Now in this final part, we examine how Deleuze and Hegel propose theories that try to overcome the problems of representational theories like the ones we saw in the first part, and we pit the theories against one another and apply them to evolution so to better evaluation them. First we examined how Deleuze's and Hegel's responses to classical representationalist philosophical can be compared on the basis of their different interpretations of differential calculus and Kant's **antinomies**. We found that for Deleuze we have an unconditioned ground of sensible and intelligible things that is subrepresentational, and for Hegel it is representable, using infinite thought. The calculus differential determines the varying relation between variables that vary with respect to one another. Leibniz saw it as the relation between infinitesimal magnitudes but there are formal problems with this. Newton saw it in terms of vanishing values. Hegel regards the vanishing values as being determinate values that combine finite and infinite, and being and nothingness. This contradiction is only thinkable with infinite thought. For Deleuze the terms of the differential relation are undetermined and subrepresentational, but they are determinable in relation to one another and are the unconditioned condition of conditioned actual determinations. Kant thinks we arrive at antinomous theories regarding whether there is a temporal beginning to the world because our understanding is unable to grasp the unconditioned, the thing in itself, with its categories. Hegel thinks the antinomies go together. Together they express the genuine infinite, because they affirm both that there is a limit and also that it is surpassed. For Hegel the unconditioned, the dialectical contradiction, is representable with infinite thought. Deleuze thinks that the unconditioned is thinkable but not using representational thought but rather using the logic of incompossibility.

We then saw how Deleuze's philosophy of difference is more resilient to attack than Hegel's, when both are pitted against one another. If Hegel wanted to critique Deleuze's philosophy of difference, he would show how Deleuze's virtual and actual as contraries dialectically sublate, which collapses the basic distinction of Deleuze's ontology. However, because Deleuze's virtual and actual are two tendencies of the real and not contraries, such a Hegelian critique would not hold. From a Deleuzean perspective, Hegel's dialectic could be viewed as a false movement, with Deleuze's genesis of difference being the real movement. Yet Hegel purely from a logical standpoint might say that Deleuze's difference does not exist.

So we then applied Deleuze's and Hegel's responses to representation to evolutionary theory to see which one is more compatible and also to see if Deleuze's three criticisms of Hegel still hold: Hegel's is a false movement, Hegel's logic revolves around a single center, and Hegel's dialectic does not provide enough precision for characterizing the world. For Hegel, nature is the one totality and it externalizes into multiplicity, but these form unified systems where parts and their whole are reciprocally determining. Hegel's dialectic is not temporal, so it does not describe an evolutionary progress through time. The structure of the organism is the reciprocally determining relation between organism and organs, which are opposing dialectical pairs like the one and the many. Individuals and species bear this organ/organism relation too for Hegel. Hegel's structure of the organism is more closely tied to Cuvier's anatomy, which is **functional and teleological**, meaning that organisms' anatomical structures can be understood in terms of their functional purposes. Geoffrey's homological theory of the unity of composition does not identify anatomical parts on the basis of their functions. Rather, he looks to see if the relations between the parts are isomorphic to a transcendental model which is so abstract that it can actualize in a wide variety of forms, **such that a fin can be identified with an arm**. This is compatible with Deleuze's transcendental empiricism and theory of the virtual, which sees there being a transcendental level that is actualized in various ways. Cuvier's and Hegel's theories, as teleological, regard deformations or mutations in negative terms, as degradations of the organism's structure and thus functioning as well. But evolutionary theory needs a positive view of aberrations. Geoffrey's and Deleuze's theories see variation positively, because variations are considered novel actualizations of the virtual model. Thus Deleuze's response to the problems of representational theories is better than Hegel's at least with regard to its application in evolutionary theory. We also see that Deleuze's three criticism's hold, because Hegel's movement is a matter of (infinite) representation, but because it cannot explain novel evolutionary variations, there is no real evolutionary movement involved. Hegel's structure of the organism has a teleological unity, and so there is a 'monocentering of circles' [around the organic unity of the organism.] Hegel's account is not precise enough. Because it understands the differentiation in the natural world in terms of determinate oppositions, Hegel's dialectic too strongly divides the world rather than seeing the blur rings of boundary that allow for evolutionary variation. Somers-Hall, Henry (2012) Hegel, Deleuze, and the Critique of Representation. Dialectics of Negation and Difference. Albany: SUNY. Emphasis mine

Avoiding the Void

Poets on Poets (Romantic Circles) Saltines and Ginger Ale (Annotated minutes of Yale's Using Theory lunch series) Stanford Enclopedia of Philosophy Saturday, July 28, 2007

Dread in Kierkegaard

Dread or anxiety has a specific structure for Kierkegaard, a structure Heidegger is correct in elaborating as thrown-ness in Being and Time: in other words, Heidegger's reading of Kierkegaard (and in a footnote Heidegger indeed acknowledges that he is indebted to Kierkegaard as well as to Christian theology: "The man who has gone farthest in analyzing the phenomenon of anxiety--and again in the theological context of a 'psychological' exposition of the problem of original sin--is Soren Kierkegaard") is exact. Let's see how. The structure of dread is outlined by Kierkegaard in the following passage, taking Adam's fall into sin as its concrete instance:

...When it is related in Genesis that God said to Adam, "Only of the tree of the knowledge of good and evil thou shalt not eat," it is a matter of course that Adam did not understand this word [i.e. that he was "innocent"]. For how could he have understood the difference between good and evil, seeing that this distinction was in fact consequent upon the enjoyment of the fruit?

When one assumes that the prohibition awakens the desire, one posits knowledge instead of ignorance; for Adam would have had to have knowledge of freedom, since his desire was to ["freely"] use it [i.e. the distinction between good and evil]. The explanation therefore anticipates what was subsequent. The prohibition alarms Adam [or

induces a state of dread] because the prohibition awakens in him the possibility of freedom... [And yet this freedom] is a nothing, the alarming possibility of being able... Thus innocence is brought to its last extremity.

-The Concept of Dread, in Existentialism from Dostoevsky to Sartre, 103-104.

Kierkegaard is saying that dread is fundamentally what allows innocence--exemplified in Adam before he ate of the tree of knowledge--to be "brought to its last extremity," or, in other words, to fall into the state of sin, whose essence, Kierkegaard later says, is guilt. Let's represent this structure in this way:

Innocence → dread → guilt

In this passage **Kierkegaard p**oints out explicitly that this structure bypasses the concept of prohibition as we normally understand it completely. Thus he also implicitly indicates fundamentally that prohibition is what is re-evaluated or revalued throughout his analysis of dread's engendering guilt. Indeed, Christian theology has often focused on this particular issue (prohibition) primarily when it comes to Adam's fall in Genesis. What is prohibited for Kierkegaard, however, in contrast to this Christian tradition, is not eating of the tree, and therefore something like the desire to know. We'll specify exactly what Kierkegaard thinks is prohibited by God in this statement later, and thus what he conceives of as desire. But for now, with this brought up, the first question to put to ourselves is why he wants to bypass the issue of prohibition in this traditional way? How will looking at dread as the origin of guilt require us to revalue the concept of prohibition?

Because the key to understanding dread is to understand it as also bound up with knowledge. As Kierkegaard says, "when one assumes that the prohibition awakens the desire, one posits a knowledge instead of ignorance.", and this covers up the essential phenomenon of how guilt arises. Essentially, understanding prohibition as the thing that awakens desire covers up the **fundamental ignorant** innocence that is constitutive for the act of eating of the tree--the ignorance that is precisely what makes the story (like the story of Job) so extremely disturbing to us. Why would God punish what essentially is an act done without our ability to know that it was wrong? Positing that Adam was focused on the prohibition itself when he ate of the tree requires that he know what the tree will do when he eats of it--in short, if the prohibition awakens the desire, it is precisely because Adam would know beforehand why what is prohibited is prohibited. Obviously this is untenable: how could Adam know beforehand what the tree would give him? No, the whole instance proceeded in complete ignorance. But how, then, could Adam have become guilty of trespassing against God's word? That is where dread comes in.

Dread is what, for Kierkegaard, leads us out of innocence and ignorance and into guilt, instead of knowledge. Knowledge is not what causes the fall of Adam, but dread, and with it, freedom. Thus to the structure we outlined above, there would seem to correspond a structure that looks like this:

Ignorance → dread → knowledge

But because we do not know how dread begets knowledge, this statement as yet is empty. Furthermore, if we were to try and tie freedom into this structure by representing it as the following we would be fundamentally wrong:

Ignorance → dread → knowledge, freedom

Freedom is not a consequence of dread for Kierkegaard, but rather underlies it as what dread itself activates or actualizes in dread's becoming actual. Thus, we must ask, how does dread lead us out of innocence and ignorance by way of being connected with freedom for Kierkegaard?

It should be noted that all the foregoing has done is to outline the interconnected nature of two structures, the structure of innocence and guilt, and the structure of ignorance and knowledge, which we can represent by combining our previous representations:

Innocence → dread → guilt

Ignorance → dread → knowledge

It is also clear from the foregoing that two concepts remain to be connected to this nexus: prohibition and, most essentially, freedom. But it is not by specifying what either of these are for Kierkegaard that we get a handle on what he means by them and dread more generally. Rather, it is by proceeding with the structures that we already have outlined that we can specify their nature. In other words, though Kierkegaard has a specific nature of freedom, he does not specify what freedom is concretely and then derive his analysis of dread (not to mention his reading of Scripture) from it--this would be too abstract for him. Rather he sticks close to what he already knows and what Scripture already tells him about the transition between ignorance and innocence and guilt and knowledge. We'll do the same. Indeed, it should be obvious by bringing the two structures we have outlined that, if we posit some other term--dread--instead of the prohibition itself as the thing that leads us out of ignorance and innocence, that this third term--dread--will be responsible for relating both ignorance and innocence. In other words, from what we have already seen Kierkegaard say, it should be obvious that dread leads us out of innocence ignorantly, and that dread leads us out of ignorance innocently. It is by analyzing these relations that Kierkegaard sticks close to what he already knows and does not give in to a pressure to cover up the disturbing nature of Adam's fall--not to mention the fall of anyone innocently or ignorantly into guilt and knowledge.

Now, if one is led out of innocence, or, as we normally call it, is guilty for some reason, and yet is led out of innocence ignorantly, he still seems innocent: indeed, this is why Kierkegaard says "he who... becomes guilty is innocent" (102)--that is, if they become guilty ignorantly. Why should this be the case? Why do we normally make this provision for guilt? Or, put differently, how can one who is guilty be innocent at the same time if they are ignorant? By our normal reasoning, we say that if an innocent person is lead to guilt ignorantly, it is something other than effectuates the guilt, not the person who was lead into guilt. "They didn't really mean it," we say, "it wasn't really their fault." Guilt must be tied up with this "something other," then. Kierkegaard simply names this "something other," "dread." That is, instead of pursuing the issue of guilt by making an exception of the case in which someone is led to guilt ignorantly, as we usually do, he analyzes it as a positive phenomenon. In other words, he is not excluding this instance from the rest of our codified ways of assigning guilt, making of it a case where the "rules don't apply in the same way:" rather, he will eventually conclude that this instance **is the exemplary instance** of the way guilt arises.

Can the nature of thinking be uncovered by thinking alone? (Cf consciousness) "My dialectical method is, in its foundations, not only different from the Hegelian, but exactly opposite to it. For Hegel, the process of thinking, which he even transforms into an independent subject, under the name of 'the idea ', is the creator of the real world, and the real world is only the external appearance of the idea. With me, the reverse is true: the ideal is nothing but the material world reflected in the mind of man, and translated into forms of thought. “Karl Marx, Capital, vol. 1, Preface to the Second Edition. (P.102) (See Ideal / Real) For Aristotle, Thinking is the one specific activity of the Human Soul Which is Capable of Independent Existence and separate from any connection to the Body. A Strenuous counterargument has been made, precisely for body thinking. Deleuze and Guattari for, "Becoming Animal" and "Becoming intense" are Modes of resistance to abstract Thinking. Alain Badiou FOLLOWS Aristotle in using thinking **to define** specifically the Human, as Opposed to Interest. For Badiou, "the capacity which is specifically human is that of thought, and thought is nothing other than that by which the path of a truth seizes and traverses the human animal." **(Infinite Thought, p. 71)** In HIS Essay "Does Consciousness Exist?", William James Pointed out That Consciousness is a Process, and not a substance. (Unlike Descartes, who claimed the existence of a res cogitans?) In The Principles of Psychology, **James described five properties of what he called "thought**." Every thought, he wrote, tends to be part of personal consciousness. Thought is always changing, is sensibly continuous, and appears to deal with objects independent of it. In addition, thought focuses on some objects to the exclusion of others. In other words, it involves attention. Alfred North Whitehead claimed That, with this Inquiry, James was to the Twentieth century What Descartes was to the Seventeenth. Louis Sullivan describes "Real

Thinking," Thinking in the present tense, as Organic Thought. "It is in the present, only, that you really live, therefore it is in the present, only that you can really think. And in this sense you think organically. Pseudo-thinking is inorganic. The one is living, the other dead. The present is the organic moment, the living moment. " (Kindergarten Chats, "Thought.") Thinking is a form of Computation? Emergence ---------- Emergence Refers to the appearance of Patterns of Organization and is one of the Key concepts of Complexity and a -life. It is sometimes referred to as a situation where the whole is greater than the sum of the parts, because it cannot be analyzed by taking the parts apart and examining them separately. One reason for this is that in a complex phenomenon showing emergent properties, the parts become a determining context for each other, and these patterns of feedback contribute to the appearance of the emergent phenomenon. For Michael Polanyi, "Evolution CAN be understood Only as a feat of emergence." For Goethe, the Whole Living was Also More than the Sum of ITS parts. "Das ist zwar Lebendige in Elemente zerlegt, aber MAN Kann ES aus Diesen nicht wieder zusammenstellen und beleben." Goethe quoted in d'Arcy Thompson, **On Growth and Form. (P.41)** But for most contemporary scientists, Goethe's meaning of emergence was a mystical one. For Francis Crick, the scientific meaning of emergent assumes that while the **whole cannot be the simple sum of the separate parts, its behavior can, at least in principle, be understood from the nature and behavior of its parts plus the knowledge of how all these parts interact. (The Astonishing Hypothesis, P.11)** (Assuming IT's Possible to have **That Laplacian Knowledge**) Stuart Kauffman Suggests That many Properties of Organisms May be Probable **emergent Collective Properties** of Their constituents. (This is precisely Kant's definition of the organic - as the result of internal interactions instead of an assembly of preexisting parts. (See Mechanism / Vitalism) "The problem of origins requires an understanding of how new levels of order emerge from complex patterns of interactions and what the properties of these emergent structures are in terms of their robustness to perturbation and their capacities for self-maintenance **"(Brian Goodwin, How the Leopard Changed its Spots, p. 181.)** Luc Steels Offers Useful definitions of emergence in Artificial Life half. He distinguishes between first order emergence, defined as a property not explicitly programmed in, and second-order emergence, defined as an emergent behavior that adds additional functionality in the system. In general a-life researchers try to create second -order emergence, then the system for use CANS ITS own emergent Properties to create an upward Spiral of Continuing evolution and emergent behaviors. For a Critique of emergence, See Peter Cariani, "Emergence and adaptivity in Organisms and Devices," World Fudtures 32 (1991):111-32. C. Lloyd Morgan's Descriptions of emergence as a Qualitative Change in direction closely resemble the definitions of bifurcations. HIS In Emergent Evolution, of 1923, he writes, "**The emergent step, although it may seem more or less saltatory, is best Regarded as a Qualitative Change of direction, or Critical turning-Point, in the Course of events**. " Daniel Stern describes the Infant's sense of Self between Birth and two months as the "emergent Self." (See subject). It is a sense of the Coming-into-Being of Organization and Remains active for the rest of life. But IT is not an overarching schema About Integrating the Self, but rather an Experience of Process. (cf Molar / Molecular) In Gestalt Perception terms, emergence refers to our ability to perceive forms that cannot be reduced to the addition of the "atomic" parts of a pattern. Ego superimposing two squares diagonally displaced **allows for** readings of an emergent square at their overlap, two emergent L-shapes in the non-Intersecting area, in addition to the two Squares. (cf. Figure / Ground) A **Fundamental property of Open Systems is that they stabilize any improbability which Serves to elicit them**. Being / Becoming --------- ---------- According to Ernst Cassirer, Plato's Philosophy recognizes two Contrary Forms of representation, one of Which is Valid for the Realm of Being and the Other for the Realm of Becoming. (Individual and Cosmos, p.125) For Cassirer, "**Form Thinking**" Belongs to Being, while "causal Thinking" Belongs to Becoming. But strict Knowledge is Only Possible of the Always-Being. Becoming That Which is CAN Only be described, if at All, in the language of myth. Or rather, myth is already familiar with both the question of the "what" and the question of the "whence." "It sees everything that it grasps (the world as well as the gods) under this double aspect."(The Problem of Form and the Problem of Cause, in The Logic of the Cultural Sciences, p. 87) Socrates first Discovered the Concept, or Eidos as the relation Between the Particular and the General and as a germ of a new meaning of the general question concerning being. This meaning emerged in its full purity when the Socratic eîdos went on to unfold into the (transcendental) Platonic "Idea." (See transcendence / immanence) For Plato, the soul is an intermediary between being and Becoming. The Aristotelian system seemed to Promise a Different reconciliation of the Opposition between Being and Becoming. Unlike Plato and the thinkers of

pure form, Aristotle wanted to restore to Becoming ITS rightful Place, Because He was convinced **That Only in this way Could Philosophy be Transformed from a Mere Theory of concepts into a Theory of Reality. Form and Matter, Being and Becoming, must become correlative. Peculiarly Aristotelian the Concept of the formal Cause originates from this fusion**. (Cassirer) "In Nature We May Search for successfully something that endures, and in nature we may regard becoming as if any phase of it were the "reason" of a later phase and the "consequence" of an earlier one. We may even go so far as to say: Nature is the one mediate object that obeys the postulates of the rational theory of becoming. "Hans Driesch, History and Theory of Vitalism, p. 193. "Freedom Means That a stream, as IT Were, of undetermined Becoming of a really creative Character Runs through the Totality of individuals. "(ibid. p. 238) One of the central arguments of Philosophical Writings Influenced by Contemporary science is to favor Becoming over Being, or to call for a new Resolution of the two. Deleuze and Guattari seek a Thought-in- Becoming, Which is Entirely immanent, not to Develop a General idea (model) That Would Stand out and above (Transcend) subsumes the Bodies ITS. They Constantly Becoming Stress - Becoming Other, Becoming Animal etc. as Qualitative Change. They Reject a metaphysics of Being dependent upon representation. (See Plane of immanence) and characterize Becoming as intransitive, not having an object of Production Other than Itself. "Reality Makes and Remakes Itself, but IT is Never made something." (See Also duration) In describing nomad art, Deleuze and Guattari describe "a nomadic absolute, as a local integration moving from part to part in an infinite succession of linkages and changes in direction. It is an absolute that is one with becoming itself, with process." (Mille Plateaus, p 494) Ilya Prigogine and Isabelle Stengers Hope to Reconcile Being and Becoming in theories of irreversible Processes, State Whose system is Associated with Being, but Whose laws of Temporal Change are Associated with Becoming. (See Order Out of Chaos, P. 310) They too, rejct the Identification of Being with timelessness. (See time) Machine "Machine is Derived Via Machina and the Latin from the Greek mechane, Meaning tool or Machine - Especially an Instrument to Lift Heavy Objects, a crane, or a military engine. Perhaps the original word is mlxos, Which Means an Artificial device, Especially Used Against misfortune and Troubles. The Entire Body of Modern science Rests on Descartes' metaphor of the **World as a Machine**, Which He Introduced in Part V of the **Discourse on Method as a way of understanding organisms but then generalized as a way of thinking about the entire universe.** "I have hitherto described this earth and generally the whole visible world, as if it were merely a machine in which there was nothing at all to consider except the shapes and motions of its parts. "One essential aspect of the machine is that it consists of clearly distinguishable bits and pieces, each of which has a determined causal relationship to the movement of other bits and pieces. In science, the machine metaphor is the rationale for reductionism. The Cartesian emphasis on "clear and distinct" ideas, a conception which united rationalists and empiricists, can be seen as part of the mechanical view, because the idea of the individual part, the basic unit, is thought to be Fully Independent of any Complex Assembly. The Classic Definition by "the first Great morphologist of the Machine," Franz Reuleaux, That is a (simple) Machine is **"a combination of resistant Bodies That SO arranged by Their Means of the Mechanical Forces nature can be compelled to do work accompanied by certain determinate motions**. "(Reuleaux, the Constructor. A Handbook of Machine Design, Philadelphia, 1894) Lewis Mumford Gives a Gloss of this Definition in The Myth of the Machine, Referring to Reuleux's Definition as "a combination of resistant parts, each specialized in function, operating under human control, to utilize energy and perform work." For Mumford, "The machine was a counterfeit of nature, nature analyzed, regulated, narrowed, controlled by the mind of men."(Mumford, Technics and Civilization, P.52) Following Marx, the Mumford SAW Machine Primarily as a Social formation. HIS In Analysis of the Division of Labor in Capital, vol 1, Karl Marx developed the parallel between the different parts of a machine and the division of labor which produces those parts, as a mechanism which compels the collective worker to work with the regularity of a machine, (p.469) which divides up the individual himself in order to transform him into the "automatic motor of a detail operation. "(481) For Marx," the machine is a means for producing surplus value "(the labor which the worker gives the capitalist for nothing.) Because its raison d'etre is the accumulation of surplus labor, capitalism subordinates men to machines instead of using machines to liberate men from the burden of mechanical and repetitive work. As production becomes mechanized under capitalism, the rhythm and and work content of living labor are subordinated to the mechanical needs of machinery itself. (See clock) (See also André Leroi-Gourhan's description of the Individual's smooth Performance of "Mechanical Operational sequences" in

everyday life in Gesture and Speech, eg. p.232) "The Machine Has suffered for the Sins of capitalism; contrariwise, capitalism Often Has Taken Credit for the Virtues of the Machine. "(Mumford, P.21) Ernest Mandel Outlines Three Quantum LEAPS in the evolution of machinery under Capital: Machine Production of Steam-driven motors since 1848, machine production of electric and combustion motors since the 1890's , and machine production of electronic and nuclear-powered apparatuses since the 1940's. These correspond to the three stages of capitalism: market capitalism, monopoly or imperialist capitalism, and "postindustrial" or multinational capitalism. (Late Capitalism) (Frederic Jameson identifies postmodernism with the Last phase.) How is a Distinguished Machine from a tool? For Mumford, the essential Distinction between a Machine and a tool lies in the Degree of Independence in the Operation from the Skill and Power motive of the operator. But is the Difference Simply in the source of motive power? Although he notes that tools can be distinguished from machines on the basis of their motor forces, this is not the important element for Marx. **He describes the transformation of the instruments of labor from tools into machines as the removal of the instruments of labor from the hands of the worker. (Alienation**) The machine takes the place of the worker, not of the tool. "The machine ... is a mechanism that, after being set in motion, performs with its tools the same operations as the worker formerly did with similar tools. "(Capital, Vol 1, NY, Intl. Pub. pp 374-79, Chapt 15 pp 492 -) Marx also distinguishes the social organization of machine from that of tool-users , Primarily in ITS need for a reliable Organization of Knowledge. For Mumford, "the Machine" is shorthand for the Technological Complex, Which "Swept over Our Civilization". According to Mumford, "Men Become HAD Mechanical They perfected before Complicated Machines to Express their new bent and interest; and the will to order had appeared ... in the monastery and the army and the counting-house before it finally manifested itself in the factory. "(Mumford, Technics and Civilization, p. 3) Gilles Deleuze echoes this statement when he claims that "machines are social before they are technical." (Deleuze, Foucault, pp 39-40) According to Mumford, "If the invention of the mechanical clock heralded the new will-to-order, the use of the Cannon in the Fourteenth century enlarged the Will to Power; and the Machine as We know IT represents the Convergence and Embodiment of Systematic These two Prime Elements."(p84) For Deleuze and Guattari, the Machine is the Ontology of Production. " As for the great capitalist machines, their basic mechanisms were proliferative: first urban, then royal state machines, commercial and banking machines, navigational machines, monotheistic religious machines, deterritorialized musical and plastic machines, scientific and technical machines, and so forth. "(Fylix Guattari, "Machinic Heterogenesis") Mumford calls the invisible structure which includes all the political and economic, military, bureaucratic and royal components that make the immense work-output and grand designs of collective social organization the megamachine - its technical equipment, "megatechnics. "" The problem was to turn a random collection of human beings ... into a mechanized group that could be manipulated at command. The secret of mechanical control was to have a single mind with a well-defined aim at the head of the organization, and a method of passing messages ... (to) the smallest unit. "(The Myth of the Machine, pp 191-2) Thus all forms of social regimentation, limiting the actions and movements of human beings to their bare mechanical elements, belongs to the physiology, if not to the mechanics, of the machine age. (Technics and Civilization, p. 41) (These descriptions of disciplinary mechanisms are part of what Michel Foucault would call the "anatomo-politics of the human body ". (see biopower)" As statements are inseparable from systems, so visibilities are inseparable from machines. A machine does not have to be optical; but it is an assembly of organs and functions that makes something visible and conspicuous. "(Deleuze , Foucault, p. 58) Mumford's arguments, echoed by Deleuze and Guattari, link the Machine to universality and Homogeneity. (cf smooth / striated) All of These Descriptions Stress the Social Coordination, the Plane of Organization, the coherence of Machine Assemblies. (See Big science) EJ Dijksterhuis describes the rise of Modern science as the "Mechanization of the World Picture." (See representation) One sense in Which He interprets this idea is to Think of the physical universe as a Great Machine, Which, once it has been set into motion, by virtue of its construction performs the work for which it was called into existence, (p.495) such that God becomes a "retired engineer." (see closed / open systems) The prime artifact of the mechanical universe is the clock. Dijksterhuis distinguishes between finalist pictures of the machine and a study that only wonders by **what cause a particular part is moved, without considering the object of the motion.** He describes this analysis as the study of a mechanical system as opposed to a Machine. Dijksterhuis Gives Another sense to HIS Thesis by Describing the Tendency among physicist to Look for Hidden Mechanisms, and to Assume That These Would be Essentially the

Same kind of as the simple Instruments Used to Relieve work, That such a skilful Mechanical Engineer would be able to imitate the real course of events. For Dijksterhuis mechanization is best interpreted as "with the aid of a concept of mechanics", which would be better named (mathematical) kinetics. By defining force purely as the product of an acceleration (a purely kinematic magnitude - from Greek kineo, referring to constrained or controlled motion) and a mass (a coefficient to be determined empirically), modern science eliminated both the metaphysical terminology and psychological origins of the concept of force. "The mechanization of the world-picture during the transition from ancient to classical science meant the introduction of a description of nature with the aid of the mathematical concepts of classical mechanics; it marks the beginning of the mathematization of science, which continues at an ever-increasing pace in the twentieth century. "(V: 9) (See Also Quantitative / Qualitative) The Mechanical 'World View "Has Also Been criticized as a General Tendency to Think of events as happening to passive and Things to conceive All Causes as efficient and extrinsic. The central paradigm of this sort of action is the colliding billiard balls of mechanics, in which it is supposed that the balls have no intrinsic, active powers of motion. In this view, there is no place for a paradigm of agency, such as spontaneous human action. When the body is compared to a machine, it is in terms of energy intake and expenditure (oxidation). The comparison is also made through artfacts such as the clock - both as models and as direct influence. See, for instance " the Body is a Machine, "**in HG Wells and Julian Huxley,** The Science of Life. Another strain of Philosophical Thought is More Critical of the Mechanical World Picture. In Monadology, Leibniz refuted the Mechanical Explanation of Perception: "Supposing there were a Machine whose structure produced thought, sensation, and perception, we could conceive of it as increased in size until one was able to enter into its interior, as he would proceed into a mill. Now, on going into it he would find only pieces working upon another one, but He Would Never Find anything to explain Perception. "Ludwig Wittgenstein, in HIS Explorations of the Relations between Meaning and use, discusses the Machine as symbolizing ITS Actions. (Philosophical Investigations, Paragraphs 193-4) "We talk as if these parts could only move in this way, as if they could not do anything else. How is this - do we forget the possibility of their bending, breaking off, melting, and so on? "He concludes that" the movement of the Machine-as-Symbol is predetermined in a sense Different from That in Which the movement of any actual Given Machine is predetermined. "Wolfgang Kettler distinguished Gestalt theories from "machine theory," that is, technological conceptions of science, life, and mind that equated knowledge of nature with **its effective manipulation and control**. The gestalt theorists attempted to introduce an aesthetic dimension of inherent order, meaning, and simplicity into the evaluation of scientific theories, and into the fabric of experience and nature itself. (Mitchell G. Ash , Gestalt Psychology in German Culture, 1890-1967, P.1) Michel de Certeau describes the use of tactics, Manipulation, and Diversion creatings as a "Certain Play in the Machine" through a Stratification of Different kinds of interfering and Functioning. (Practice of Everyday Life, p.30) Another Approach to Them Machines frees from any fixed Functional objective and sees a form of Creativity in Them. For Elizabeth Grosz, Machines are "Heterogeneous, disparate, discontinuous assemblages of fragments Brought Together in conjunctions or Severed through disjunctions and breaks. "This description ressembles the forms of bricolage or tinkering described by Claude Levi-Strauss and derive from the descriptions of machines in the Anti-Oedipus. (see desiring machines) Fylix Guattari talks about the theme of the machine as an affective object, rather than in mechanistic terms. D + G distinguish the machine from the mechanism, which is rigid and unmoving. Guattari wants to tear down the "iron curtain" between being and things. Instead he sees being "differentiating itself qualitatively". . "as the very extension of the creativity of machinic vectors." and ascribes to machines a "protosubjectivity." (See "On Machines" in Andrew Benjamin, ed. JPVA no.6, issue entitled Complexity: Architecture / Art / Philosophy) "A pile of stones is not a machine, whereas a wall is already a static protomachine, manifesting virtual polarities, an inside and an outside, a high and a low, a left and a right." (Guattari, "Machinic Heterogenesis") Jacques Monod, in Chance and Necessity; contrasts machines, whose design agency is outside of them, from organisms, that are formed internally. A machine presupposes a conscious and intelligent maker who has constructed it and makes it operate to realize a particular object. (This is the "cryptoteleological" aspect of machines) See Kant's distinction in mechanism / Vitalism. See also autopoesis. The study of artifical life assumes that it is possible to organize a population of machines in such a way that their interactive dynamics is "alive. "For Richard Dawkins," Machines are the direct products of living objects, and they are diagnostic of the existence of life on a planet. "(Blind Watchmaker, p. 2)

Reversing the traditional interpretations of organism as machines, Georges Canguilhem argues that "machines can be considered as organs of the human species." (cf prosthesis) With their greater range of activity, organisms suggest a model "less bound by purposiveness and more open to potentialities." Viewed as organic products, machines can be seen as "the direct or indirect products of a technical activity that is as authentically organic as the flowering of trees," products of an ingenuity "based on instinct, and developing along Darwinian notions of variation and natural selection, as extensions of human behavior or life processes . In the structuralist framework, Machines Articulate the categories of Language and Society. THUS, if Ritual is described as a Machine for Transforming diachrony into synchrony, and Play is, conversely, a Machine for Transforming synchrony into diachrony, the two CAN be Regarded as Machine a single, a single binary system Which is articulated accross two categories Which cannot be isolated and Whose Correlation Across Difference and the very Functioning of the system is based. (Giorgio Agamben, infancy and History, p.74) For Van Doesburg, " every machine is the spiritualization of an organism. "For de Stijl, machinery hastened the spiritualization of life by separating man from nature." The machine is, par excellence, a phenomenon of spiritual discipline. "If handicraft reduced men to the level of machines, for Van Doesburg, "the proper tendency for the machine is as the unique medium of the very opposite, social liberation." (See Banham, p.151) Banham considers these statements the passage to the machine aesthetic, where, in Severini's words, "we may conclude that the effect produced on the spectator by the machine is analogous to that produced by the work of art." This is neither Rationalist materialism nor Futurist mechanolatry, but rather a bridge to a depersonalized, metropolitan, "new style. " "Essentially a Machine is the physical Embodiment of a Process Which Occurs in time." (Collins, Changing Ideals in Modern Architecture, P. 163) - Concept May this be closer to the Organic than to the Machine. nomadic / sedentary - ---------- For Deleuze and Guattari, nomads are above All CHARACTERIZED by the Fact That Their mode of Existence is antithetical to the system of the State, of cultivation, and of striation, which they describe as sedentary. Because the nomads were so decisively defeated, history has always dismissed them, and indeed "history is one with the triumph of States." (p.394) Nomadism becomes, for D + G, a revolutionary Alternative to the State, Although They are Always Careful to Distinguish Between Their "de jure" or conceptual Distinctions and All the "de facto" mixes and transitions That Actually Occur. D + G describe the nomadic occupation of smooth space as "vortical", as distributed by turbulence, "as a" distribution of heterogeneity in free space - "a" local absolute. "It was the nomadic mode of warfare that so distinguished them. Their" pack "-like movement, the formidable assemblage of man / horse / bow, their warrior way of life - animated by a fundamental indiscipline - constitute what D + G call "the war machine" - the pure form of exteriority. Deleuze and Guattari mitigate the Identification of the War Machine War with making by Claiming That the nomadic way of the War Machine is Primarily a Determination to Occupy smooth space. But When the city stands in the way of nomadic Free movement, War is the result. THUS every Operation Against the state is associated with the nomad: "insubordination, rioting, guerilla warfare, or revolution as act - it can be said that a war machine has revived, that a new nomadic potential has appeared." (p.386) "a non-subjectified Machine Assemblage with no intrinsic Properties, Only Situational ones. "(P353) For Deleuze and Guattari, the Nomad is the "Outsider." Nomadic Thought is "outside" Thought (an Expression They Borrow from Blanchot) Nomadic science is "science Minor "which is itinerant, ambulant, and follows flows in vectorial fields across which singularities are scattered like so many" accidents. "(p372) Matter, in nomad science, is never prepared and therefore homogenized matter, (see smooth / striated) but is essentially laden with singularities, which constitute a form of content. (see form / matter) In describing nomad art, Deleuze and Guattari describe "a nomadic absolute, as a local integration moving from part to part in an infinite succession of linkages and changes in direction. It is an absolute that is one with becoming itself, with process. "(Mille Plateaus, p 494) Assemblage ---------- The French Terms of both agencement and dispositive Used by Deleuze and Guattari are usually translated as assemblage. An assemblage is a "site at which a discursive formation intersects with material practices" (Jonathan Crary, Techniques of the Observor, p. 31) It is "simultaneously a machinic assemblage and an assemblage of enunciation "(D + G, Thousand Plateaus, p. 504)

But let us continue--how could one be led into guilt ignorantly, if guilt essentially lies in knowingly doing something? Where is the knowledge located in this instance? If someone is guilty innocently because of ignorance,

and thus has become guilty by "something other" than her or himself, and if this "something other" is dread, then we might say that it is precisely dread that effectuates the guilt knowingly. In other words, if the innocent person is still innocent after being made guilty through dread, it is not the person who is responsible for this guilty status: for how could he have known--being ignorant--whatever it was that rendered him guilty? Dread knows, somehow, or has some relationship to the knowledge of which the person is ignorant, and thus Kierkegaard continues the sentence just quoted ("he who through dread becomes guilty is innocent") by saying "for it was not he himself but dread, an alien power which laid hold of him" in the transition between innocence and guilt (102). Thus by specifying the relationship of dread to knowledge we get a handle on exactly the way someone innocent and ignorant can be led to guilt. If they are seized by dread, they are seized by something that has a relationship to a knowledge that they do not know. We might represent this the following way before we go on to analyze this relationship:

Innocence/ignorance → (dread ← knowledge) → guilt

Notice that this gets rid of our previous schema where knowledge was somehow a product of the dread of an innocent person along with guilt, which was represented this way:

Ignorance → dread → knowledge

Knowledge is intimately connected with dread, and not with ignorance. Thus it does not directly oppose and obviate ignorance itself; knowledge does not suddenly come along and wipe out innocence because it wipes out ignorance. Rather, knowledge has a direct relationship to dread, and only can effect ignorance through this mediation.

We can specify the relationship of knowledge to dread by asking what must necessarily be the essence of a knowledge that preserves ignorance. The answer to this is that it must be a fantasy, something that is not real in content but is real in its ability **or possibility to be real. As Kierkegaard says, at the center of dread is that which in an innocent act could have been, that** which in an act was that person's particular "I can" (104), in the sense of an ignorant "I can do this which I currently fantasize about" at the moment of one's fantasizing. This is why Kierkegaard specifies dread as a "dreaming" of the spirit: the moment at which a subject dreads is the moment in which that subject indeed "projects its own reality" even though "this reality is nothing" and "this nothing constantly sees innocence outside of it" . The reality it projects is precisely its own ability to do whatever it is that it projects.

Now it should be obvious that by "seeing innocence outside of it," Kierkegaard means that in the particular moment of dread, what is, is only that which is for someone who is ignorant, or does not know reality, and is therefore innocent in the way we determined before. Let's penetrate deeper into this phenomenon. What is real in the moment of dread is nothing, and dread itself is thus dread "about nothing" as we usually say. But this nothing is not really nothing, as Kierkegaard says. It is only nothing insofar as what the dread is about in reality is nothing. In reality, whatever is dreaded has the constitution of a dream, a wish. But the fact that the dread occurs and is about something that is not real does not make the act of dreading itself disappear. The act of dreading is precisely everything, it is so much not anything--it is and remains one of the keenest and most real psychological feelings possible. We thus might represent dread as the following, making a distinction between the act of dread and what is dreaded.

Dread → content, what is dreaded → nothing, dream

↓

Form, the act of dreading

↓

Real, reality

More appropriately, we might call the act of dread a desire, since a desire is precisely a relationship to something that is, as of yet, nothing. Indeed, if we are going to call this act a wish or a dream as well this seems legitimate. But in recharacterizing this act this way, we can see desire come into play--and it should be quite clear now that it is precisely not the desire of some particular bit of knowledge like the distinction between good and evil. What is dreaded, what is desired is something that is not, and since the act of dread or desire occurs anyway, we are left to conclude that that which it is really about is precisely unrealized possibility itself, in other words, the reality of nothing--the reality of something that, as of yet, is not real and is therefore nothing. Put differently (and perhaps more clearly), what the desire is about is (in Freudian/Lacanian terms) really the drive, the act of desiring alone. Similarly, what is dreaded is really only the possibility of the individual to realize that particular thing one dreads. Thus the above relationship can be redrawn like this:

Dread → desire, what is dreaded → nothing, dream

↓

Drive, what is really dreaded

↓

Real, reality

This continual play on the word "**nothing**" Kierkegaard uses (and which we are merely reduplicating) he uses perhaps to curb the impact of the radical thesis he is putting forth here. What he is saying is that what dread is drive itself, the desire for the realization of what is unrealized and, as the desire of someone who is ignorant, is the desire of something that is not known. Desire, as dread and therefore as drive, is desire for that of which one is ignorant, that which is nothing currently. As such, it is not fundamentally that thing that is desired, but (because one does not really know that thing, the act one could have done), it is really the possibility of the realization of that thing. This is what is meant by us saying desire is desire of "the reality of nothing:" it is fundamentally the desire for the possibility to realize what nothing, the desire of drive is now. Furthermore, as a dream or wish or desire, one possesses already in dread this possibility--it is there in one's desiring of that which does not exist. This is why it is, as Kierkegaard implies, the greatest inner possibility of a subject.

Giorgio Agamben asks how it is possible to conceive of absolute immanence in the form of desire. He points to Spinoza's theory of "striving" (conatus) - the desire to persevere in one's own being - as a possible answer. For Spinoza, all beings not only persevere in their own Being (vis inertiae) but desire to do so (vis immanentiae). The desire to persevere in one's own Being is to desire one's own desire, to constitute oneself as desiring. Desire and in conatus Being THUS coincide without residue. Subjects of Desire In, Judith Butler describes the dialectical workings of Desire in Hegel's Phenomenology of Spirit, a work she describes That as a Bildungsroman - a Narrative of Transformation and actualization, Whose "Logical motor "is desire itself. Hegel give an account of a philosophical subject that strives to know itself in the external world and in its relations to an Other. Desire, according to Hegel, is the incessant human effort to overcome external differences, a project to become a self-sufficient and self-conscious subject. Human desire articulates the subject's relationship to that which is not itself. It finds satisfaction through the transformation of difference into identity, when a relation to something external to consciousness is discovered to be constitutive of the subject itself. Thus desire is both intentional in that it is always desire of or for a given object or Other, but it is also reflexive in the sense that desire is a modality in which the subject is both discovered and enhanced. Hegel claims that "self-consciousness in general is Desire." (Phenomenology of Spirit, ¶ 167) Desire and Other In Self-Consciousness, Desire Always Reveals the Desiring Agent as intrinsically Other to Itself: Self Consciousness is an EK-static Being, outside Itself, in Search of Self-Recovery. Some years ago at a specific hour of the day, a tall, slender man could be seen walking with measured steps back and forth on the flagstones in the southern section of Overgaden over Vandet. Hardly anyone failed to

notice the peculiarity in his walks, for the distance he covered was so short that even the uninitiated were bound to become aware of him, that he did not enter shops and that he was not, like others, out for a stroll either. Anyone who observed him frequently could see in his gait an image of the force of habit. … He was, of course, well known in the whole neighborhood, but even though he was mentally disordered, he was never exposed to any insult; on the contrary, the neighbors treated him with a certain respect. Conducive to this were his wealth and also his charitableness and his attractive appearance. It is true that his countenance had the impassive expression characteristic of a certain kind of mental disorder, but his features were handsome, his figure erect and well formed, his attire very meticulous, even elegant. Moreover, his mental disorder manifested itself most clearly only in the forenoon between eleven and twelve o'clock, when he paced the flagstones between Bornehaus [Orphanage] Bridge and the south end of the street. The rest of the day he presumably spent trailing after his unhappy concern, but it did not express itself in this way. he spoke with people, went on longer strolls, involved himself in many things, but between eleven and twelve o'clock no one for all the world could stop him from walking, make him walk farther, answer any questions, or even respond to a greeting-he who otherwise was courtesy itself. … The conduct of the nearby residents toward him was almost reminiscent of the conduct of the Indians toward a mentally disordered person, whom they venerated as a wise man, in private they possibly had many conjectures as to the cause of his misfortune. It happens not infrequently that by this kind of conjecturing the so-called sagacious people betray just as much disposition to lunacy or perhaps more foolishness than anyone mentally disordered. The so-called sagacious people are often so stupid as to believe everything a lunatic says, and not infrequently stupid enough to believe that everything he says is lunacy, although many a time no one is more cunning at hiding what he wants to hide than a mentally disordered person, and although many a word from him contains a wisdom of which the wisest need not be ashamed. This no doubt explains how the same view that thinks that in the governance of existence a grain of sand or an accident determines the outcome can hold also in psychology, for it is the same view if one sees no deeper cause for insanity but regards insanity as easily explained by nothing, just as mediocre actors believe that acting the role of an intoxicated person is the easiest of tasks, which is true only if one is sure of having a mediocre audience to see the acting. Stages on Life's Way, Hong p. 278-279 (1845)The internalization of otherness is also the externalization of the subject. Difference threatens the subject with annihilation until the subject can discover that difference as an essential moment of itself. The two moments of assimilation and projection condition the irreducible ambiguity of the emerging identity of the subject. The infinite insatiability of the Endless Desire Requires Proliferation of alterity. In the section on Lordship and Bondage (Also called master and slave?) Hegel describes the Most challenging Encounter of the subject, When "Self-Consciousness is faced by Another Self-Consciousness. ' (¶ 179) The first experience of the Other's similarity is that of self-loss. It is a narcissistic project that fails through an inability to recognize the Other's freedom. Self-consciousness CAN Only supersede this otherness to Itself When IT locates Itself in reciprocal Recognition, When IT recognizes the Other as Also in the Process of retrieving Itself from estrangement in ITS Own Desire. Michael Hardt and Antonio Negri call the Desire "ontological motor" for the biopolitical Practices of Being. Ortega Y Gasset commented That Proust HAD invented "a new Distance Between Ourselves and Things" - a Distance modified by Attention and imagination, by Love and Desire. (See Also Memory) For the Surrealists, the water was of Fluidity Powered Also the Desire of Fluidity Powered Opposing the solidity of Matter. In their discussion of the French sociologist Gabriel Tardy, Deleuze and Guattari discribe a flow as both belief and desire. (Two aspects of every assemblage) "a mutant, convulsive, creative and circulatory flow." (100 Plateaux 217-8) Maurice Blanchot, Analyzing the "objective Chance" of Breton's Encounter with Nadja, Points out That "Chance is Desire." He recounts the contradiction between an amour in search of an illusory total object, incarnate in woman, (symbol of desirable reality) and a desire which is committed to whatever might respond to it - including that which maintains it as desire without satisfying it. . for the Surrealists, the city is the place of chance (and magical) encounter. "Ce qui me syduit dans une telle maniire de voir, c'est qu'a perte de vue elle est de recryatrice dysir" (André Breton, L'Amour Fou) Desire and the Body: Desire is based on a veritable cartography of the body, a rewriting of instincts and biological processes into networks of signification and meaning. (Cf Freud's theories, elaborated by Laplanche of the drives as propping on the biological instinct, translated as anaclitic for the German Anlehnung.) The erotogenic zones mark those places where vital functions (such as ingesting milk from the breast) become the source of sexual stimulation. For Freud, "There is

good reasons why a child sucking at his mother's breast has become the prototype for every relation of love. The finding of an object is in fact a re-finding of it." ("Three Essays on Sexuality) Jean Laplanche points out that a displacement has taken place nonetheless." The lost object is the object of self-preservation, and the object one seeks to refind in sexuality is an object displaced. "(Life and Death in Psychoanalysis, p. 20) Freud's use of the Word "libido," Which in Christian usage Referred to concupiscence, (Desire as cupiditas) expressed a sense of Desire, with All the Overtones of **Lust, Force, and tension That IT injected into psychical life. Freud defined libido in psychoanalysis as "in the first instance the force (thought of as quantitatively variable and measurable) of the sexual instincts directed towards an object - 'sexual' in the extended sense required by analytic theory." (A Short account of Psychoanalysis, 1924) Freud's quest** for his own desire through the prism of his dreams resulted in dreams where the voice of desire could be heard. (Didier Anzieu) Freud sought not to master his dreams (an obssessional defense) but rather to let Them Give Voice to wish. (cf imaginary / symbolic) For Deleuze and Guattari, "Freud's greatness Lies in having Determined the Essence or Nature of Desire, no longer in relation to Objects, AIMS, even sources (Territories), but as an abstract subjective essence - libido or sexuality. "But they criticize him for relating this essence to the family as" the last territoriality of private man. "(Anti-Oedipus, p. 270) For Deleuze and Guattari, Freud sets up" interiority in place of a new relationship with the outside, "while they seek to discover" beneath the familial reduction the nature of the social investments of the unconscious. "Luce Iragaray criticizes Freud for" falling back on the affirmation that the libido is necessarily male. "(This Sex Which is Not One, p. 48) and for" maintaining that there is only one libido, but that in the case of femininity it may put itself in the service of 'passive aims.' " One Might Say That Classical Greece was the Source of Iragaray's Nightmare. According to Michel Foucault, the Greek relation to the Body and ITS Pleasures was a Completely Different model from the Christian relation to Flesh and ITS Desires. The Greek View, however, was Related to a "virile Society" That was unisexual and excluded Women. (Plato's Account in the Symposium priveleges Men Seeking Their male Counterparts.) In the Anti-Oedipus, subtitled Capitalism and Schizophrenia, Volume 1, and first published in 1972, Gilles Deleuze Fylix and Guattari engaged in a radical critique of Freudianism.Like their contemporary, RD Laing, and like Wilhelm Reich before them, they linked psychic repression with social repression, and sought to recover the revolutionnary quality of desire. For Deleuze **and Guattari, desire produces reality, and they attribute to Desiring machines the productive qualities of desire. They also linked the weapon with the production of affect. (see "The War Machine" in Thousand Plateaus) and this apparent** glorification of nomad warfare may account for some of the antipathy Which They arouse in feminists. For the artist Hans Bellmer Desire is a multiplier of Objects. The Body Becomes an anagram, Pleasure Producing the reciprocal of the parts. He is, in Alain Jouffroy's the Words "tychnicien de l'impossible" who explored the What Bellmer Called the "physical Unconscious" - the Body's Underlying Awareness of Itself. Productive What is the role of Desire in Technological development? Desire and Time. Desire How does Occur in time? "One Source of the richness of Our emotional repertoire (is) rooted in the Possibility of time-indexed Desires. "(Ronald de Sousa, The rationality of Emotion, p.100)" These focus on Particular Moments or stretches of time, making Possible More Complex Forms of rationality and irrationality. " “Only microfascism provides an answer to the global question: Why does desire desire its own repression, how can it desire its own repression? "(Anti-Oedipus, p. 215) Desiring Machines ---------- ---------- In the Anti-Oedipus, Capitalism and Schizophrenia subtitled, Volume 1, and first published in 1972, Gilles Deleuze and Fylix Guattari engaged in a radical critique of Freudianism. Like their contemporary, RD Laing, and Wilhelm Reich before Like Them, They Psychic repression linked with Social repression, and Sought to Recover the revolutionary quality of Desire. Deleuze and Guattari For, Desire is not to be Identified with Lack, with the Law, or with the signifier, but rather with production, with desiring production in the social field. For them, the Oedipal myth and its institutionalization in psychoanalysis, stands directly in the way of understanding the productive unconscious. It is an obstacle to a materialist psychiatry which would introduce desire into the Mechanism and Production Introduce into Desire. Deleuze and Guattari For, Desire Produces Reality. Desiring Machines are the Production Site of That. Deleuze and Guattari For, every Machine is a **Machine connected to Another Machine. Every Machine Functions as a Break in the Flow in relation to the machine to which it is connected, but is at the same time also a flow itself, or the production of a flow. What we term libido** is the "labor" of desiring production. It is pure multiplicity, and for Deleuze and Guattari, it is anoedipal. The flow is non-personal, although investments by

desiring machines produce subjectivity alongside its components. (Guattari, "Machinic Heterogenesis") (See also bachelor machines) (For a Freudian account see play) (See also "buccal space "in psycho-Sexual space.) Rejecting the "anthropomorphic representation of Sex", Deleuze and Guattari describe the workings of the Unconscious and of the Sexual Drives as the Actions of Desiring Machines. They describe the breast as a Machine, the Mouth as a machine coupled to it, and are at pains to point out that their use of this term is not metaphorical, because for them the essence of both man and nature is production (as opposed to representation), and desiring production is the primary form Production of Human. It is the Production of Production. Desiring Production is Generation. Deleuze and Guattari Contrast "Molar" Machines - Whether Social, Technical, or Organic - with the "Molecular" microphysics of Desire. "Desiring Machines are the nonhuman sex, the molecular machinic elements, their arrangements and their syntheses, without which there would be neither a human sex specifically determined in the large aggregates, not a human sexuality capable of investing these aggregates. "(Anti-Oedipus, p. 294) The regimes of molar and molecular remain **inseparable, however. "There are no desiring-machines that exist outside the social machines that they form on a large scale; and no social machines without the desiring machines that inhabit them on a small scale." (p .340) For D + G, the Molecular Unconscious Knows nothing of castration. Its partial Objects Lack nothing and multiplicities Free form - a** microscopic transsexuality. THUS "Love making is not just Becoming as one, or even two, but as a Becoming hundred thousand "(AO, p. 296)." The schizoanalytic slogan of the desiring revolution will be first of all: to each its own sexes. "" The schizo is not a revolutionary, but the schizophrenic process ... is the potential for revolution. "(p.341) D + G propose a schizoanalysis to replace the Oedipal psychoanalysis, not so much in celebration of schizophrenia, but because the breaks and flows that they observe at the schizoid level escape some of the strictures of Oedipus and allow us to see nature as production. "Desiring production is situated at the limits of social production; the decoded flows, at the limits of the codes and the territorialities; the body without organs at the limits of the socius." (Anti-Oedipus, PP 175-6) (the Socius is the abstract Machine of Society - Massumi). Michael Hardt and Antonio Negri Deleuze and Guattari Follow in underscoring the Revolutionary Potential of Desiring Production. For Them, "the biopolitical, Seen from the Standpoint of desire, is nothing other than concrete production, human collectivity in action."(Empire, p. 397)" Empire pretends to be master of the biopolitical world because it can destroy it. What a horrible illusion! We are in Reality Masters of the World Because Our Desire and Labor REGENERATE IT Continuously. "(P.368) Some feminist Writers have approached this Claim to the non-Personal with suspicion. Iragaray Luce asks "... does not the ' desiring machine 'still partly take the place of woman or the feminine? Isn't it a sort of metaphor of her that men can use? Especially in terms of their relation to the techno-cratic? "(This Sex Which Is Not One, p.140) Alice Jardine expresses her concern that" becoming-woman, desiring machines, and other similar concepts are merely excuses for male forms of appropriation of whatever is radical and threatening about women's movements ". (How about taking it non-personally?) (Substance 44/45) quoted in Gilles Deleuze and the Theater of Philosophy. The book title is an ironic choice, since D + G refer to Oedipus as "the fantastic repression of desiring machines. It substitutes a classical theater for the unconscious as a factory. It interprets the workings of desiring machines as representations of its triangular code of mommy-daddy-me. It compromises the direct confrontation of desiring-production and social production as well as the relation of psychic repression with social repression. Production is crushed, subjected to the requirements of representation. Oedipus forces desire to take as its objects the differentiated parental persons and prohibit the ego from satisfying its desires with these persons. But the undifferentiated is created as the reverse of the differentiated. In this sense, the Oedipal is a double bind, and for Deleuze and Guattari It cannot be resolved until we do away with both the problem and the solution. "Is that not the nature of desire, that one desires the impossible?" (AO p 162.) (See Hans Bellmer and the Body) It is Worth noting That for Freud, Maternal Care is the Primary model of Seduction. Such care, in focusing on certain bodily regions, contributes to defining them as erotogenic zones, zones of exchange which demand and provoke excitation in order subsequently to reproduce it autonomously through internal stimulation. - J. Laplanche, Life and Death in Psychoanalysis, p. 44) According to Brian Massumi, the subjectivist persistent misunderstandings of the Term "Desiring Machine" as Developed in Anti-Oedipus, LED Them to substitute the Term Assemblage in A Thousand Plateaus, (for Desire VS representation, See Michel Foucault, The Order of Things, PP 208-11 and for Social Production VS representation PP 253-56) Pleasure ---------- The Basic Assumption of Freud's Pleasure Principle

That is the Most Salient and Unique Aspect of Human Experience is the Subjective experience of pleasure (tension reduction) and unpleasure (tension buildup). See Descriptions of Bateson's "Plateaus" in schismogenesis for a Culture with a seemingly Different Approach to Pleasure than one based on Release of tension as in Orgasm. For Freud the Affective and Perceptual Experiences are yoked. Without the Experience of Hedonic Tone, no Perceptions Would be registered at All. Kant describes the Pleasure We Experience When Nature Conforms to the Requirements of Our subjectivity in the Critique of Judgement. (See also unity) He describes the pleasure the transcendental philosopher takes in finding that a particular purposiveness, or conformity to law, which was not necessary, turned out nonetheless to be the case. "The sole pleasure found in disinterested judgements concerning the mere form of objects reveals that nature (feeling) and reason are capable of agreement." (Rodolphe Gashy, The Idea of Form, p.173) But Kantian Aesthetic Pleasure is not merely an Effect - an Aesthetic Manifestation of Awareness - IT is Intimately tied up with a State of Mind in Which the Powers of imagination and reason Become attuned in a judgement of taste. Taste, for Kant is "a tone of mind which is self-maintaining." (P.67) See Gashy, p. 51) For Michel Foucault, the Greek relation to the Body and ITS Pleasures was a Completely Different model from the Christian relation to Flesh Desires and ITS. If the church and the ministry stressed the principle of a morality whose precepts were compulsory and whose scope was universal, the demands of austerity in classical thought were stylizations of masculine conduct. In The Use of Pleasure, Foucault sets out to Trace the Long History of the Aesthetics of Existence and the "techniques of the Self" - the Ways in Which the Individual is summoned to Recognize Himself as an ethical subject of Sexual Conduct. Rather than strictly defining what is permitted and what is forbidden, the accent of Greek and Greco-Roman ethical thought "was placed on the relationship with the self that enabled a person to keep from being carried away by the appetites and pleasures, to maintain a mastery and superiority over them, to keep his senses in a state of tranquillity, to remain free from interior bondage to the passions, and to achieve a mode of being that could be defined by the full enjoyment of oneself, the perfect supremacy of oneself over oneself. ' (P.31) According to Foucault, foods, Wines, and Relations with Women and Boys constituted analogous ethical Material. They brought forces into play that were natural, but that always tended to be excessive. The dynamic relation of acts, pleasures, and desires was a circular one, and the ethical question of aphrodisia was not: which desires? which acts? which pleasures? but rather with what force one is transported "by the pleasures and desires." (P.43) As Aristotle expresses it, "all men enjoy in some way both savoury foods and wines and sexual intercourse, but not all men do so as they ought." (P.52) Unconscious ---------- "Is Unconscious a thing, a Place, the Unconscious, or IT is a mode, an adjective, Unconscious?" (Francoise Meltzer, in Critical Terms for Literary Study.) "The unconscious does not have a rhetoric, but is a rhetoric." (Gorgio Agamben, stanzas, p.150 n.) How CAN the unconscious be known? (In German, it is the Unbewusste, the unknown) Can it only be diagnosed, inferred through its symptoms in parapraxis? Ideas and Memories Can be Effective yet unavailable to Consciousness? Freud approached the unconscious from a number of Points of View. In his first important paper on the subject, "A Note on the Unconscious in Psycho-Analysis" (1912), Freud adopted a descriptive viewpoint and divided the psyche into conscious and unconscious. The descriptive model was the source of Freud's "topographic" or spatial geography of the mind, which divided the psychic apparatus into three systems: Ucs., Pcs., Cs. (Unconscious, preconscious, conscious) (see "The Unconscious," 1915) (See topos) Later Freud Developed a Dynamic model, Called "Hydraulic" by critics, described Which Energy Flows That build up and seek Release. In this model, tension is displeasure and will always seek an avenue for release. For Freud, these interplays of forces are of organic origin. They are represented as images or ideas with an affective charge.The dynamic model lead Freud to consider the unconscious from an economic viewpoint, that is, to attempt to quantify the relative magnitudes of the forces of excogitation. "From the economic standpoint, psycho-analysis supposses that the mental representatives of the instincts (instinctual drives) have a charge (cathexis) of definite quantities of energy. In his later publications, Freud referred to two instinctual drives: libido and agression. In the 1930's Freud Replaced the Topographical Viewpoint with the Structural one, dividing the mental apparatus into HIS Famous "tripartite" model of Agency: ID, Ego, and superego. Throughout HIS Analyses, Freud Adopted a genetic Viewpoint, Holding That any Psychological Phenomenon, in addition to its contemporary and experiential aspects, can be traced back through its ontogenesis to its psychological origin. Throughout his life, Freud remained faithful to the idea of an intrication of ontogeny and phylogeny. He was quite prepared to see the relationship between consciousness and the

unconscious in terms of an evolutive, integrative emergence of the latter from the former, and to see neurotic symptomology as a regression and freeing of early or repressed mental processes and contents. (Anzieu, Freud's Self-Analysis, p. 38) From the Beginning , the notions of the unconscious and repression (or resistance) were closely linked for Freud. The repression barrier divided consciousness from unconsciousness. The first thing that happens in repression is that idea and affect are severed. The affect can be inhibited; it can remain in consciousness but attached to another idea, or it can undergo transformation, notably into anxiety. For Freud, it was through repression that ideas, especially ones that were threatening, painful, or unpleasant, found their way into the unconscious, just as, through the Lowering of resistance, in Dreams or Therapy, They found Their way out. (What is the relation of repression to Forgetting?) Freud's Investigations into traumatic Hysteria Clearly Revealed cases where a Patient's Behaviour Could not be Explained or even Identified without Reference to operative but unconscious ideas. The fact that the same behaviour could be artificially induced by hypnosis, by the insertion of ideas into the subject's mind, strongly suggested that ideas were operative, even though the subject knew nothing of them. Freud's preferred explanation of inner conflict was in Term of the incompatibility Between Certain ideas and a mental Agency, Identified as the Ego Which exerted repression. For Freud, the residue of the "Prehistoric stage of life", from the Age of one to Year Three, is the Source of the Unconscious and contains the a etiology (the medical word for cause) of all the psychoneuroses. It is a stage normally obscured by amnesia similar to hysteria. (See screen memories for a mode of persistence to infantile memories.) Thus Freud could conclude that because small Children do not truly understand the Meaning of Death, in Our unconsious We know nothing About Death in General and Our Own Death in Particular. For Freud, the Unconscious system was CHARACTERIZED by exemption from mutual Contradiction, by Primary Process, Which is indifferent to Logical, Spatial, Temporal or order, by timelessness, and by the replacement of external by Internal Reality. Displacement and Condensation are two Forms of Primary Process That describes Freud in The Interpretation of Dreams. Gerald Edelman describes as a Primary Process That Consciousness arises from Perceptual Organization, Memory, Learning and Self / non-Self discrimination. Identity, Memory, and space compose Primary Consciousness. Primary Consciousness From this, a higher-order Consciousness evolves in MAN, with the Powers of Language, conception, and Thought. What are unconscious memories? In 1897, Freud abandoned the seduction theory, whereby hysteria (primarily in women) was a result of actual abuse. (Whatever else may be true about hysteria, it is a thoroughly gender-laden diagnosis, description, or discourse.) One reason for this shift was that for Freud there **were no criteria** in the unconscious to distinguish between an actual event and a fantasy, an issue which paved the way for the **idea of narrative truth**. Through his own self-analysis, Freud came to the conclusion that "neurosis" was caused not by an actual sexual trauma, but by incestuous fantasies. Recent critics of Freud, especially Jeffrey Masson, have claimed that Freud chose to ignore evidence of real abuse in some of his female patients, notably in the case of Emma Eckstein, Who was Also negligently Treated When Fliess left gauze up her Nose After Operating on her. For Freud "Somatic Conversion" was a link Between Repressed ideas and the Body. It was an Indication of the Body's Powers of symbolic Expression. (See Body Thinking) What is relation Between Unconscious and Automatic Behaviour? Contrary A Current Interpretation of views of the Unconscious as Enabling IT INSTEAD of custodial. Efficiency rather than repression Accounts for Unconscious mental work. Automatic mental Processes "Unconscious cerebration" do not require Comprehensive awareness, and free up consciousness to the physiological tasks for which it is best fitted. Many nineteenth-century psychologists saw the unconsious as "actively generating the processes which are integral to memory, perception and behavior. Its contents are inaccessible, not because, as in psychoanalytic theory, they are held in strenuously preventive detention, but, more interestingly, because the effective implementation of cognition and conduct does not actually require comprehensive awareness. On the contrary, if consciousness is to implement the psychological tasks for which it is best fitted, it is expedient to assign a large proportion of psychic activity to automatic control; if the situation calls for a high level management decision, the unconscious will freely deliver the necessary information to awareness. "(See Jonathan Miller," Going Unconscious "in Hidden Histories of Science, pp 1-35. This passage quoted in Crary, Suspensions of Perception, p. 40) This is Also the neurological tradition. Gestalt psychologists , Like Kurt Goldstein, rejected the Notion of the Unconscious. For Goldstein, "What Appears as the Unconscious is nothing but the Entering of a Former Reaction Pattern of the Organism present into a response, When the Situation is Suitable." (P258) The unconscious can be thought of as a step away from a unified,

centralized mind towards a decentralized one. (D + G: "molar" versus "molecular") Rather, it is everything that is left behind in a contraction of selection or sensation than Moves from one Level of Organization to Another. The Only Things are not the unconsious are present Perception and reflection. (Massumi. p.83) What is the Relationship Between the Unconscious and the "Plane of consistency of Desire" Deleuze and Guattari Which call the Body without Organs? For Deleuze "the unconscious is not a deep container of yet-unknown sources, but rather the marking of the structural non-coincidence of the subject with his / her consciousness." Rejecting the "anthropomorphic representation of sex", Deleuze and Guattari describe the workings of the Unconscious and of the Sexual Drives as the Actions of Desiring Machines. "Thinking ... is to a **very large extent Unconscious, That in the IT Expresses Desire to know, and this Desire is That Which cannot be adequately expressed in Language, Simply Because IT is That Which Language sustains."(Braidotti," Toward a New** Nomadism ") "Within the Notion of the Freudian Unconscious the Involution of the strange in the psyche Loses ITS pathological Aspect and integrates within the assumed unity of human beings an otherness that is both biological and symbolic and becomes an integral part of the same. "" Uncanny, (unheimlich) foreigness is within us: we are our own foreigners. "(Julia Kristeva, Strangers to Ourselves, p . 181) "We are in a system where there is no More Soul, no More metaphor of the Body - the Fable of the Unconscious Has lost Most of ITS resonance" (Baudrillard, The Ecstasy of Communication New York, Semiotext (e) , 1988, 50-51) to Read: Ellenberger, The Discovery of the Unconsious Thinking ---------- "Thinking ... is to a very large extent Unconscious, That in the IT Expresses Desire to know, and this Desire is That Which cannot be adequately expressed in Language, Simply Because IT is That Which sustains Language. "(Rosi Braidotti," Toward a New Nomadism ") "One thing Only do I know for Certain and That is That MAN's judgement of value follow directly his wishes for happiness - that, accordingly, they are an attempt to support his illusion with arguments. "**Sigmund Freud**, Civilization and its Discontents, penultimate and last page. For Georg Simmel," our mind has a remarkable ability to think of contents as being independent of the act of thinking, he observes that "In desiring what we do not yet own or enjoy, We Place the content of Our Desire outselves .... When the subject is constituted by Separation from the World, representation Becomes the dominant mode of Thinking and Knowing. If there Were a verb "to believe falsely," IT Would not have any significant first person, present indicative. - Ludwig Wittgenstein.

Being and Time

2.1 The Text and its Pre-History

Being and Time is a long and complex book. The reader is immediately struck by what Mulhall (2005, viii) calls the "tortured intensity of [Heidegger's] prose", although if the text is read in its original German it is possible to hear the vast number of what appear to be neologisms as attempts to reanimate the German language. According to this latter gloss, the linguistic constructions concerned—which involve hyphenations, unusual prefixes and uncommon suffixes—reveal the hidden meanings and resonances of ordinary talk. In any case, for many readers, the initially strange and difficult language of Being and Time is fully vindicated by the realization that Heidegger is struggling to say things for which our conventional terms and linguistic constructions are ultimately inadequate. Indeed, for some thinkers who have toiled in its wake, Heidegger's language becomes the language of philosophy (although for an alternative and critical view of the language of Being and Time, see Adorno 1964/2002). Viewed from the perspective of Heidegger's own intentions, the work is incomplete. It was meant to have two parts, each of which was supposed to be divided into three divisions. What we have published under the title of Being and Time are the first two divisions of (the intended) part one. The reasons for this incompleteness will be explored later in this article.

One might reasonably depict the earliest period of Heidegger's philosophical work, in Freiburg (1915–23) and Marburg (1923–6), before he commenced the writing of Being and Time itself, as the pre-history of that seminal text (although for an alternative analysis that stresses not only a back-and-forth movement in Heidegger's earliest thought between theology and philosophy, but also the continuity between that earliest thought and the later philosophy, see van Buren 1994, 2005). Viewed in relation to Being and Time, the central philosophical theme in these early years is Heidegger's complex critical relationship with Husserl's transcendental phenomenology—what Crowell (2005, p.49) calls "a dynamic of attraction and repulsion"—as driven by Heidegger's transformative reading of Aristotle. As early as a 1919 lecture course, for example, we find Heidegger arguing that Husserl's view (developed in the Logical Investigations, Husserl 1900/1973), that philosophy should renounce theory and concentrate on the things given directly in consciousness, is flawed because such givenness is itself a theoretical construct. For the young Heidegger, then, it is already the case that phenomenological analysis starts not with Husserlian intentionality (the consciousness of objects), but rather with an interpretation of the pre-theoretical conditions for there to be such intentionality. This idea will later be central to, and elaborated within, Being and Time, by which point a number of important developments (explained in more detail later in this article) will have occurred in Heidegger's thinking: the Husserlian notion of formal ontology (the study of the a priori categories that describe objects of any sort, by means of our judgments and perceptions) will have been transformed into fundamental ontology (a neo-Aristotelian search for what it is that unites and makes possible our varied and diverse senses of what it is to be); Husserl's transcendental consciousness (the irreducible thinking ego or subject that makes possible objective inquiry) will have been transfigured into Dasein (the inherently social being who already operates with a pre-theoretical grasp of the a priori structures that make possible particular modes of Being); and Husserlian intentionality (a consciousness of objects) will have been replaced by the concept of care or Being-in-the-world (a non-intentional, or perhaps pre-intentional, openness to a world).

Each of these aspects of Heidegger's framework in Being and Time emerges out of his radical rethinking of Aristotle, a **rethinking that finds its fullest and most explicit expression in a 1925–6 lecture course entitled Logik (later renamed** Logik (Aristoteles) by Heidegger's student Helene Weiß, in order to distinguish this lecture course from a later one he gave also entitled Logik; see Kisiel 1993, 559, note 23). On Heidegger's interpretation (see Sheehan 1975), Aristotle holds that since every meaningful appearance of beings involves an event in which a human being takes a being as—as, say, a ship in which one can sail or as a god that one should respect—what unites all the different modes of Being is that they realize some form of presence (present-ness) to human beings. This presence-to is expressed in the 'as' **of 'taking-as'. Thus the unity of the different modes of Being is grounded in a capacity for taking-as (making-present-to) that Aristotle argues is the essence of human existence. Heidegger's response, in effect, is to suggest that although** Aristotle is on the right track, he has misconceived the deep structure of taking-as. For Heidegger, taking-as is grounded not in multiple modes of presence, but rather in a more fundamental temporal unity (remember, it's Being and time, more on this later) that characterizes Being-in-the-world (care). This engagement with Aristotle—the Aristotle, that is, that Heidegger unearths during his early years in Freiburg and Marburg—explains why, as Sheehan (1975, 87) puts it, "Aristotle appears directly or indirectly on virtually every page" **of Being and Time. (For more on Heidegger's pre-Being-and-Time period, see e.g., Kisiel 1993, Kisiel and van Bu**ren 1994, and Heidegger's early occasional writings as reproduced in the collection Becoming Heidegger. For more on the philosophical relationship between Husserl and Heidegger, see e.g., Crowell 2001 and the review of Crowell's book by Carman 2002; Dahlstrom 1994; Dostal 1993; Overgaard 2003.)

Let's sum this up to be a little clearer. What is desired in dread is precisely that of which one is completely ignorant--to the subject it appears as nothing, making him say characteristically when asked what he is worried about, "it's nothing." Thus, what is desired is precisely not knowledge, because what is desired is precisely that of which one as of yet cannot actually know. However, desiring (and we should mention that we do not even know of this desire--desire is not like a willing)--desiring that of which one is ignorant is indeed desiring the possibility of knowing that

which is not known--this is, as it were, the side effect of desiring precisely that which one cannot know. In desiring non-knowledge, as we might put it, we **precisely desire the possibility not of knowing the non-knowledge itself, but of knowing our own possibility to know, our own possibility** to not be as ignorant as we are now. This side-effect we call drive, or the reality of the act of dread. We might represent this structure of knowledge thusly:

Dread/desire → (desire as such) that of which one is ignorant

↓ (drive)

Possibility of knowing

It should be obvious that this is just a condensed version of what we have already specified. It is this downward arrow that is active: the desire for that of which one is ignorant opens up a general, indeterminate possibility of knowing anything--i.e. not just that of which one is ignorant. Rightly, Kierkegaard does not speak of this indeterminate result as genuine knowledge yet: dread brings into view only "a nothing in lively communication with the innocence of ignorance". It is "lively communication" only that is really going on here, since what is being known is precisely only the reality of the possibility of knowing (drive). Thus, one can still speak of this knowledge of a possibility of knowing as ignorance, and we will indeed do so. It is ignorance in lively communication with only the possibility of knowing, and as such, it produces no knowledge and continues to preserve ignorance itself. We can represent this thus within the larger structure of dread:

Ignorance → {dread ← ([desire → that of which one is ignorant] → possibility of knowing (drive)/ignorance])} → guilt

This specified, we can show exactly what this type of ignorance is and does. It should be noted that grasping the role of desire has allowed us to specify the role of prohibition, but we will put this off to later when the whole phenomenon of dread comes fully into view. Now, this particular type of ignorance, this knowledge only of an open-ended possibility of knowledge this "I can," this result of a desire based in dread, Kierkegaard names "freedom." As Kierkegaard puts it, "dread is the reality of freedom as possibility anterior to possibility" . By "possibility anterior to possibility" Kierkegaard means that freedom is the reality of possibility itself as the reality of possibility in the nothing, in the unrealized, in that of which the subject is and remains ignorant. Thus we might rewrite our above structure as the following, having it retains the same meaning:

Ignorance → (dread ← freedom) → guilt

Now, it is obviously this freedom that brings one from innocence into guilt. How does this occur according to Kierkegaard--especially if it is not really knowledge still? I'll save this answer for another post.

Posted by Mike Johnduff

What is written about: Being and Time, Heidegger, Kierkegaard

3 comments:

Sean said...

Do you side with Kaufmann or Dreyfus? To what degree is Heidegger indebted to Kierkegaard? Does it really stop at thrown-ness? Or, can you trace all of Heidegger's existentialism back to 'Religiousness A' (a secularized version)? Isn't the real distinction exactly what anxiety reveals, is it the value of finite existence, or does is reveal an infinite - a sort of flipside to finite existence - as well? Also, hello I'm Sean.

February 10, 2008 at 10:48 PM

Mike said...

The relationship of Heidegger to religion is a tricky one. This precludes it being any mere "secularized version" of a text of Kierkegaard. Furthermore, this would have to deal with how, for Heidegger, Kierkegaard relates to the Aristotelian structure of being-in-the-world and the rereading of Aristotle more generally that occurs in Being and Time, which is not an easy relation to figure out. In "The Hermeneutics of Facticity," an pretty early course (1924) Heidegger is outlining much of what will become Being and Time in its basic approach to interpretation, and he there cites Kierkegaard as prominent in the sense that it inspires his hermeneutical or interpretative task or way of getting at being. I think revisiting this nexus would probably give us some directions in conceiving *exactly how* Heidegger is thinking about Kierkegaard: one could say that the interpretative method that Husserl an phenomenology gets modified into would have to be conceived as more Kierkegaard an. In other words, it is obvious Heidegger is indebted to Kierkegaard massively--the real question is how. This is all one possibility--I'm really saying that I have no clue, and thus your question opens up a great avenue to pursue. Do you have any idea? As to Kaufmann or Dreyfus, I'm not so familiar with the arguments of the former relating to Kierkegaard and their opposition to Dreyfus' readings: I think Dreyfus has a great interpretation of **Kierkegaard though**, and though I haven't listened yet to his lectures on death, guilt, and resoluteness in the Heidegger course this semester, I plan to--they should localize at least some of these problems within the Heidegger. I will say that I think Dreyfus is a little too quick perhaps to characterize Kierkegaard as merely the "existential" influence of Heidegger, as if all of Heidegger's concern with death and guilt were to stem from a concern that is able to be distinctly separated from his more Aristotelian concern in Division I. I'm not the first to criticize him in this way though: Blattner is the best at it, though perhaps going too far in the other direction and making the concerns of temporality and finitude (finitude being really what death is all about for Heidegger), the sole organizing theme of the book. Heidegger sees being as finite, and this is probably one of the central theses of the book: furthermore, he sees Aristotle saying the same thing long ago. He must elaborate how this is able to be so--and so that leads him to posit a Division I that will merge with the concern of Division II, death etc. Kierkegaard figures there also because he breaks with Hegel--that should be noted. Hegel is Heidegger's archenemy--so anyone able to say that **there is a nothingness and a freedom in nothingness, a possibility for a finite subject to relate to infinite possibility** (whether God or not), possibility en abyme (as Derrida likes to say) or, conceived probably less rigorously, as drive and repetition of desire (which Lacan says and this is how I formulate it above), would **not escape him in** his reading of Aristotle. A great helpful book on this is actually Levinas' God, Death and Time--it is really clear and outlines certain concerns about infinity and finiteness in Heidegger and elsewhere that you touched on. Whatever you think of him, too, reading the section on Kierkegaard in The Gift of Death by Derrida is helpful, because he outlines pretty much a standard reading of the suspension of the ethical in relationship to death and shows how at this point one can relate it to concerns of finiteness and infinity. But I'm talking too much--what do you think? One more thing... with regard to thrown-ness... of course it doesn't stop there, but I think that's the structure that is the best to think about with all this Kierkegaard stuff (as I say in later posts, though unsuccessfully--I stopped all this because it was getting too tough to think)... the real question is how this possibility beyond possibility that is experienced in the dread of nothingness *already exists or is factical* in Dasein. How possibility or futurity (to talk about it in temporal terms) is already within Dasein so that Dasein is thrown back upon it in its existence, so that it is thrown back upon its factical existentiality in its being-existential or projecting itself forward into nothingness, how projecting oneself forward into nothingness is throwing oneself back to the fact of **one's nothingness** (this all takes place in some remarks on nothingness that are really weird in Being and Time and have a lot to do with Hegel)--this relay, when conceived with respect to its facticity, is the real tough thing to think. **Projection is easy--thinking facticity** is the real tough stuff. And to Dreyfus' immense credit (I don't know how many people congratulate him on this--but everyone taking his course or exposed to his writings probably should, it is so vital for understanding anything in Being and Time), this is what he thinks best and why he lays so much emphasis on Division I: it's there that we get a structure that is somehow, in its being structured that way, already there (da), albeit falling and distracted, etc. I'll post this as a separate new post, so it might be easier to find--and also so that others can perhaps get in on what you asked!

February 11, 2008 at 10:14 PM

Mike said...

Also I thought of this afterwards with respect to what I was saying towards the end of my comment on the importance of facticity:

This might answer your last question--why I actually don't think that it is so much about the what anxiety reveals. It is about the nature of this revealing as such that is the key for Heidegger. Others like Levinas will change what is revealed and then go back and read it into the structure of **Dasein that Heidegger** elaborates and show how it cannot be that way (this is what he does in the book I referred you to--but of course not with the rest of his philosophy, which is much more thought out [precisely with regard to what facticity would then have to be] than this one lecture). But for Heidegger, finitude or infinitude would have to announce itself before it is revealed in the revealing itself--the real trick is not deciding then whether revealing that is done in Dasein is finite or infinite, but by trying to let Dasein itself point the way towards answering this, which only then will take us to what is revealed. This leads, however, to the hermeneutic circle Heidegger points out famously. **(From Google search, Wikipedia, Text books, and other author's pages, as cross references given in the diaspora of the text)**

Antagonism and anachronism and Quantum mechanics:

Not that always the primary focus and locus of homologous receptiveness of differentially expressive activity in the form of war would be greeted with success. Part of successful military strategy involves learning from the opponent's behaviour. For example, when Napoleon defeated the Austrians in Italy, he knew they were incapable of taking advantage of the new methods of war developed by the French Revolution. **Predictability spells defeat in war or battle**. The mongoose and the cobra, the matador and the bull, are able to time their fatal blows by learning the rhythm of their opponent's attacks. **(See discussion in Wiener, Cybernetics)** Clausewitz stressed the unpredictability of war when stated that "Everything is simple in War, but the simplest thing is difficult. These difficulties accumulate and **produce a friction** which no man can imagine exactly who has not seen War." For Clausewitz, "Friction" is "the only conception which in a general way distinguishes War from War on paper." What role does warfare play in the history of technology, especially in the West? Is it the motor force of technological development? Is it the "cutting edge" of a more general drive for domination that has motivated technological change? The drive for domination by total control lies at the heart of any military ambition. On the other hand, military strategy has increasingly had to come to grips with the **management of chaos**. Deleuze and Guattari **mitigate the** identification of the war machine with making war by claiming that the **nomadic** way of the war machine is primarily a determination to occupy **smooth space**. But when the city stands in the way of nomadic free movement, war is the result. In his study of Herodotus, **Francois Hartog** identifies the invincibility of the Scythians as the result of nomadic inderminacy. According to Herodotus, "No invader who comes against them can ever escape and none can catch them if they do not wish to be caught." Aporia has been decisively reconceived: it is no longer an absence, **a lack, a negation; it is a positive strategy--a strategy which imposes a way of life." (From Greenblatt**)

Dasein is, then, our primary 'object' of study, and our point of investigative departure is Dasein's everyday encounters with entities. But what sort of philosophical method is appropriate for the ensuing examination? Famously, Heidegger's adopted method is a species of phenomenology. In the Heideggerian framework, however, phenomenology is not to be understood (as it sometimes is) as the study of how things merely appear in experience. Rather, in a recognizably Kantian staging of the idea, Heidegger follows Husserl (1913/1983) in conceiving of phenomenology as a theoretical enterprise that takes ordinary experience as its point of departure, but which, through an attentive and sensitive examination of that experience, aims to reveal the a priori, transcendental conditions that shape and structure it. **In Heidegger's Being-centred project, these are the conditions "which, in every kind of Being that factical Dasein may possess**, persist as **determinative for the character of its Being**"

(Being and Time 5: 38). Presupposed by ordinary experience, these structures must in some sense be present with that experience, but they are not simply available to be read off from its surface, hence the need for disciplined and careful phenomenological analysis to reveal them as they are. So far so good. But, in a departure from the established Husserlian position, one that demonstrates the influence of Dilthey, Heidegger claims that phenomenology is not just transcendental, it is hermeneutic (for discussion, see e.g., Caputo 1984, Kisiel 2002 chapter 8). In other words, its goal is always to deliver an interpretation of Being, an interpretation that, on the one hand, is guided by certain historically embedded ways of thinking (ways of taking-as reflected in Dasein's preontological understanding of Being) that the philosopher as Dasein and as interpreter brings to the task, and, on the other hand, is ceaselessly open to revision, enhancement and replacement. For Heidegger, this hermeneutic structure is not a limitation on understanding, but a precondition of it, and philosophical understanding (conceived as fundamental ontology) is no exception. Thus Being and Time itself has a spiral structure in which a sequence of reinterpretations produces an ever more illuminating comprehension of Being. As Heidegger puts it later in the text:

What is decisive is not to get out of the circle but to come into it the right way… In the circle is hidden a positive possibility of the most primordial kind of knowing. To be sure, we genuinely take hold of this possibility only when, in our interpretation, we have understood that our first, last **and constant task is never to allow** our fore-having, fore-sight and fore-conception to be presented to us by fancies and popular conceptions, but rather to make the scientific theme secure by working out these fore-structures in terms of the things themselves. (Being and Time 32: 195)

On the face of it, the hermeneutic conception of phenomenology sits unhappily with a project that aims to uncover the a priori transcendental conditions that make possible particular modes of Being (which is arguably one way of glossing the project of "working out [the] **fore-structures [of understanding**] in terms of the things themselves"). And this is a tension that, it seems fair to say, is never fully resolved within the pages of Being and Time. The best we can do is note that, by the end of the text, the transcendental has itself become historically embedded. More on that below. What is also true is that there is something of a divide in certain areas of contemporary Heidegger scholarship over whether one should emphasize the transcendental dimension of Heidegger's phenomenology (e.g., Crowell 2001, Crowell and Malpas 2007) or the hermeneutic dimension (e.g., Kisiel 2002).

Martin Heidegger

(26 September 1889 – 26 May 1976)

John Phillips

National University of Singapore

Martin Heidegger's 1927 publication, Sein und Zeit (translated as Being and Time, 1962), can plausibly be considered the most influential philosophical text of the 20th century. The main focus of this work had been announced at least fifteen years earlier when Heidegger was still in his early twenties and it remained his lifelong topic until his death in 1976. He has designated this subject matter with a number of terms: life, historicity, situated being, facticity, Dasein's Sein (i.e., the being of there-being), and later in his life, das Ereignis, which is normally translated as, "the event of appropriation," and is supposed to designate the unfolding of being. The topic, in short, concerns how human beings are situated historically in a world that from the beginning has meaning for them, both making possible and setting limits to their future. Sein und Zeit is less the mighty tome or magnum opus that it appeared to be during its early publishing career, but more a seamless amalgamation of relatively independent drafts written during the years after the First World War. It thus represents the culmination of an intense period of philosophical struggle and breakthrough. It was incomplete on its publication and can be regarded as one of the stages in Heidegger's lifelong philosophical journey.

Heidegger's roots lie in deeply conservative rural Southern Germany. His lifelong suspicion of modernity, machinery, democracy, liberalism and the forms of commoditized style (like journalism and mass culture) certainly contributed to his radical critiques of the productivity and efficiency that are valorized by forms of modern life, its bland cosmopolitanism and its perpetually seductive novelty. But, alongside this, Heidegger's untiring philosophical questing was able to disclose a deep complicity between the tradition he was committed to preserve (i.e., western metaphysics) and the conditions he found around him in the modern technological world. An acknowledged and powerful alternative to predicative logic, Heidegger's thought opens with the way beings are revealed through what he called "the question of being," which he formulates as follows: why are there beings at all, and not rather nothing? The "why" in this case would not be a question about the cause of beings but something more like a celebration of the wonder that anything exists at all, thus revealing beings in hitherto unthinkable ways—for instance, opening experience up to the essential and constitutive indeterminacy of its future. Heidegger's critique of western metaphysics lies in his discovery that these constitutive aspects of experience have been systematically excluded by the metaphysical tradition, which privileges predicative logic and the correspondence theory of truth, and which reaches its ultimate form as technologizing rationality.

Heidegger was born to cellar man and church sexton Friedrich and Johanna Heidegger (née Kempf) in Messkirch, a small rural town in Baden, south-west Germany, on 26 September 1889. His sister Marie was born three years later and his brother Fritz two years after that. In 1903 Heidegger won a scholarship to the Jesuit gymnasium in Konstanz and stayed, with other scholarship children from similarly poor Catholic families, in a local Catholic boarding house. Here he prepared for a clerical career. In 1906 he transferred to Bertholds gymnasium in Freiburg, boarding at the archiepiscopal seminary of St. Georg, where the church gave him free board and lodging.

While there he received a copy of Franz Brentano's "On the Manifold Meaning of Being According to Aristotle" (1862) from Dr. Conrad Gröber, a paternal friend. Heidegger later claimed that his interest in philosophy was first aroused on studying Brentano's book, which then led him to Carl Braig's On Being: An Outline of Ontology (1896), in which he discovered excerpts from Aristotle and commentaries from Thomas Aquinas and other scholastic fathers. In 1909 he became a novice of the Society of Jesus at Tisis near Feldkirch in Austria; but on October 13 at the end of two weeks of candidature, he left, having been turned down for health reasons and possibly doubts about his spiritual vocation. In the winter semester he began studying for the priesthood at the Albert-Ludwig University in Freiburg, where he focused on philosophy and moral and natural sciences. His introduction to contemporary philosophy came when he borrowed a library copy of Edmund Husserl's Logische Untersuchungen (translated as Logical Investigations). Husserl would become a key influence in the development of Heidegger's most important publication, Sein und Zeit. The Logische Untersuchungen, which attempts a systematic enquiry into consciousness and mental processes, regarded independently of any non-mental determinations or effects, provides philosophy with a problem and a challenge. The object of thought is suspended in order to provide systematic access to the structure and origin of thought itself. Heidegger also began to read Wilhelm Dilthey, whose attempt to establish the conditions of possibility for the study of history would become a crucial reference point for Heidegger's account of historicity.

The three great early discoveries, Aristotle, Husserl and Dilthey, correspond rather neatly with the three key dimensions of Heidegger's early philosophy: ontology and the grounds of being; phenomenology and the grounds of consciousness; and historicity or the grounds of history. While it is tempting to understand these influences in terms of their content, thereby arriving at a conveniently schematic picture of Heidegger's early development, the pattern that emerges shows that his interest lies in the attempts to establish grounds rather than in any specific conclusions that might have determined Heidegger's thought. Both Husserl's and Dilthey's works are marked by some version of the Kantian procedure of critique, whereby the a priori conditions of possibility for a way of thinking or acting can be set out schematically. If Kant's Critical Philosophy addresses the questions of knowledge, action and then judgment in terms of their a priori conditions of possibility, then Husserl and Dilthey are motivated by similar aims in their attempts to establish a priori conditions for consciousness and for history respectively. For Heidegger's

developing philosophy the questions of being, consciousness and history are significant in that they each resist the tendencies of formalism, empiricism and relativism, which increasingly characterize modern science and philosophy throughout the 20th century. Already by 1913, then, Heidegger's philosophical interests reveal a highly developed pattern according to which he seizes on a basic problem of metaphysics, concerning the fundamental conditions of, for instance, knowledge, action, judgment, consciousness, language or history. The problem each time indicates a contradiction between a metaphysical question of grounds and a contemporary academic discourse. Heidegger never simply adopts the metaphysical side against the empiricist or formalist doctrines but rather he consistently establishes what the basic problems of metaphysics are, acknowledging and demonstrating the persistent and repeated failure throughout its history to achieve its aims.

So Heidegger's earliest works reveal a pattern of thought that is not strictly reducible to his early discoveries of powerful precursors. He shows a consistent interest in the covert connections between some basic problems: the increasing dominance of doctrines that favor empirical, formal or relativistic methods and conclusions; the failure of powerful metaphysical doctrines that, while allowing the criticism of empirical and formal methods, nonetheless fail to establish unquestionable grounds themselves; and already, as evident in the earliest published work, a deep interest in the implications of this connection for the social and political concerns of the modern world. The key problem concerns the relationship between the persistent failure of metaphysics and the dominance of empirical and formalist thinking in the sciences and technology, a thinking which, despite its dominance, remains vulnerable to criticism from a philosophical perspective inherited from the metaphysical tradition. It is the relationship between this key problem and his interest in contemporary and, at this time, local political and social issues that would develop into the controversial question of Heidegger's politics.

In 1910 he gave his first lecture, on Abraham a Santa Clara, in Hausen im Tal, near his home town, Messkirch. He again lectured on Abraham a Santa Clara in Kreenheinstetten, Abraham's birthplace, only ten kilometers from Messkirch. The lecture is one of the earliest works by Heidegger now published in his Collected Works (GA13). Here his philosophical interests are couched in language that ties the young Heidegger to a conservative anti-modernist tradition, which at that time favored the tough self assurance of the educated peasant classes against the fashionable bourgeoisie of modern urban life. Heidegger's topic was the court preacher Abraham a Santa Clara (1644-1709), the unveiling of whose monument in the town of his birth, Kreenheinstetten, provided the occasion for Heidegger's lecture. Abraham a Santa Clara had been adopted as a role model by those with an increasingly polemical attitude towards various liberal tendencies in the Catholic Church. At this time he began to publish articles, reviews and poems in Der Akademiker, the journal of the German Association of Catholic Graduates.

In 1911 he spent the summer semester at home because of asthma and heart problems and abandoned the theological seminary on advice of his superiors. He changed his studies to mathematics and philosophy, now concentrating on problems of logic in the history of modern metaphysics. He continued to publish reviews and poems. In 1912 he published his last articles in Der Akademiker, and wrote two articles for philosophy journals, which have since been published in revised versions in the Gesamtausgabe as part of the Frühe Schriften: "Das Realitätsproblem in der modernen Philosophie" ("The Problem of Reality in modern Philosophy") and "Neuere Forschungen über Logik" ("New Investigations of Logic").

Heidegger graduated on 26 July 1913 after his doctoral examination under Schneider with Heinrich Rickert co-examining and was awarded a Ph.D. summa cum laude on the basis of his dissertation, Die Lehre vom Urteil im Psychologismus (The Doctrine of Judgement in Psychologism), which shows the influence of Husserl in its criticism of attempts to analyze the logical notion of judgment from the perspective of human psychology. His main claim is that logic cannot be reduced to psychological processes and is marked by a developing emphasis on subjective aspects of everyday experience as well as on history and metaphysics. This is a major landmark in the important development towards the key notion of the everyday and would result two years later in what is now recognized as a major breakthrough on the way to the philosophy presented in Sein und Zeit. By 1913 his intention to "articulate the

whole region of 'being' in its various modes of reality" had been made clear and the combined resources of scholasticism, phenomenology and neo-Kantianism harnessed towards this end.

In 1915 he completed his Habilitation Dissertation, Die Kategorien und Bedeutungslehre des Duns Scotus (The Doctrine of Categories and Signification in Duns Scotus), which earned him the license to teach philosophy at the Philosophical Faculty at Freiburg. He began by assisting Father Englebert Krebs, lecturing theology students in philosophy. In his habilitation lecture, "Der Zeitbegriff in der Geschichtswissenschaft" (translated as "The Doctrine of Time in the Science of History"), Heidegger distinguishes between the notion of time as understood by modern physics and the non-uniform and qualitatively differentiated periodic time of historians, which is not reconcilable with calculable temporal duration. At this stage in his intellectual career, represented by the Habilitation Dissertation and Lecture, his focus on metaphysics is complicated by the dense fusion of scholastic philosophy, neo-Kantianism and Husserlian phenomenology. However, his central concerns yield to patient scholarship. His enquiries are already directed to what will be more fully developed in Sein und Zeit as a schematic interpretation of facticity. Facticity denotes the conditions according to which one's possibilities are determined and limited by one's circumstances, one's "being-in-the world." His interest in scholasticism focuses on categories of living language that have no correspondence with objects in the world, like marks of privation and negation and categories for non-existent entities like "Nothing." The key factor at this stage recalls the text that he claimed first aroused him to philosophy, Brentano's book on the Categories of Being in Aristotle. Categorical extremes of signification like univocity and equivocality indicate why there must be a role for a category that rescues the univocal ideal from the inevitable consequences of equivocal language use (i.e., the use of words in more than one sense—for the scholastics the paradigm was the word "God"). In the scholastic texts, following Aristotle, the analogy of being mediates between such extremes. So the empty word "being" figures at the heart of questions about the generation of sense in everyday human experience.

At this stage the problem of what he calls "the hermeneutics of facticity" lies in the self-reflexive puzzle according to which facticity must be considered, already, as a kind of hermeneutics. Everyday experience already articulates and interprets its world, so that any attempt to interpret facticity would be an attempt to interpret interpretation. Here Husserl's account of the natural or pre-theoretical attitude, an attitude that is intensified rather than exposed by theoretical procedures, indicates a way forward in the attempt to resist the philosophical tendency to objectify experience. So by 1915 Heidegger had been able to move beyond the neo-Kantian account of facticity, which posits a chaotic flux at the origin of sensation, by drawing out from the phenomenology of categorical intuition the conclusion that humans live from the first in a context, a world, which is already meaningful and which they already know interpretively.

The First World War interrupted his academic career. He published nothing between this time and 1927 but the period, especially from the end of the war in 1919, is marked by intense wide ranging study and a series of brilliant and compelling lectures. It is not possible to trace accurately his intellectual development during the four years between the Habilitation and the semester he taught immediately on his return from war duties, the so called Kriegnotsemester (war-emergency semester) of 1919. Courses on German Idealism, on Aristotle's logic, and on basic questions of logic, were announced for the 1915-16 academic year, but in 1915 he was conscripted. Because he was regarded as unsuitable for combat, he was assigned to the postal and meteorological services as military censor at the Freiburg post office. In 1915 he had met Elfride Petri, a student of political economy, and on 21 March 1917 Father Krebs presided over the first official ceremony of their lifelong marriage. Because Martin was a Catholic and Elfride a Lutheran, they were remarried a week later in a protestant ceremony with her parents. By her own account, she was never to find her faith and he would shortly lose his. In 1919 their son Jörg was born, and in 1920 a second son, Hermann, was born.

Heidegger gave a talk on Friedrich Schleiermacher, the key figure in the German Hermeneutic tradition and an important forerunner of Dilthey, on 2 August 1917. He lectured on Hegel in the summer semester, and on Plato in the winter semester. On 17 January 1918, Heidegger, then a Private in the territorial reserve, reported to barracks

for basic training, and was then sent to the front to serve at a meteorological station. He was promoted to lance corporal on 5 November, and was finally discharged eleven days later. He was never involved in any military action. In the meantime, Heidegger had written to Engelbert Krebs, breaking with "the dogmatic system of Catholicism." Heidegger continued to lecture and met with Karl Jaspers in Freiburg, where Husserl had become professor of philosophy in 1916.

On his return from the front, Heidegger became Husserl's unsalaried lecturer and assistant at Freiburg, where he taught courses on phenomenology, the foundations of logic, the history of metaphysics and the nature of the University. What marks his teaching during this war emergency semester is his forceful advocating of a kind of phenomenology that would be set apart from all hitherto existing theoretical science. The new science does not have an object as such because its subject matter is the process of life itself, within which each of us is already caught up in meaningful ways. This subject matter, before Heidegger's exceptional phenomenology, occupies the region or domain traditionally regarded—in the language of neo-Kantianism—as the transcendental, which is traditionally opposed to the empirical domain of human experience. Strictly, then, Heidegger is proposing a science of the non-objectifiable, unspeakable grounds of experience. The human being is not to be understood any longer as being derived from some extra-temporal transcendental source but is to be derived from the categories of its own facticity as a finite, situated historical being grounded in its life towards death. His fame as a teacher spread rapidly on the basis of his brilliant exposition of these original insights on the nature of the world of everyday experience and the human being. His studies of Scheler's philosophical anthropology were added to his exceptional adoption of Husserl's phenomenology, Dilthey's hermeneutics and the Christian texts of St Paul, Augustine, and Martin Luther. The role of the scholastic texts becomes clearer too, showing how they contributed to his compelling alternative to contemporary versions of ontology and epistemology still rooted in the standard readings of Greek thought. The hermeneutics of facticity—of life before theory—thus merges with the phenomenology of religious experience to provide a radical phenomenology of the everyday context or world of human experience.

In 1922 Elfride as a present had a small wood-cabin built in Todtnauberg, where Heidegger could work in solitude. It is possible, tentatively, to date the early stages of the development of the first draft of Sein und Zeit from this period. He would not publish the book until 1927, and then under considerable pressure for him to do so, but during this time he continued to write and to lecture. A letter from Husserl to Paul Natorp at the University of Marburg from 1922 describes the young Heidegger as a highly original personality but not ready to publish, because he is still "struggling, searching for himself and laboriously shaping his unique style." Heidegger himself sent Natorp an article planned as an introduction to a projected book on Aristotle, which secured him a position at Marburg in 1923. The article, Phänomenologische Interpretationem zu Aristoteles: Einfürung in die Phänomenologische Forschung ("Phenomenological Interpretations with Respect to Aristotle: Indication of the Hermeneutical Situation"), reveals something of the "laborious shaping" that Husserl describes. The vocabulary of phenomenology, the attempt to bring systematic thinking to bear on problems of philosophy, and historical considerations of philosophy, are interconnected in what remains a starkly original critical account of the human condition. At this stage the dense technical vocabulary reflected in the title encumbers the text. However, this is the first of Heidegger's extant works to introduce several of the key themes of Sein und Zeit. For the first time he makes a distinction between authentic and inauthentic existence. He also announces the project of destruction, sometimes translated as the deconstruction, of the metaphysical tradition.

In the summer of 1923 he taught a final course at Freiberg, with the title, Ontologie (Hermeneutik der Faktizität) ("Ontology: Hermeneutics of Facticity"), which is one of the core strands of Sein und Zeit. The focus is on Dasein—being there (Da, which may mean here or there)—the existing human being in its aspect as a someone who is placed somewhere and lives in a particular time. By the end of the course the notion of facticity has developed into the key notion of In-der-Welt-Sein, being in the world, Heidegger's basic subject matter, which allows him to focus on the being, the Sein of that which discloses or interprets being, that is, Dasein. As the subtitle of the course indicates, Heidegger's treatment of hermeneutics intensifies what in 1915 had already been accomplished, that is, a

radical revision of the long tradition of exegesis and commentary—hermeneutics. In Heidegger's sense, the process of disclosing one's facticity would operate only on the assumption that for Dasein this process is already an operative function of facticity—one would actually be disclosing the event of disclosure itself, drawn out from the intelligible structure of Dasein's own being.

Heidegger's time at Freiberg came to an end on July 5, when with Natorp's help he moved to the University of Marburg to take up the post of Associate Professor. Here he met the newly enrolled 18 year old Hannah Arendt. In February of 1924 they became lovers in an affair that would last until 1933. Their friendship, which she resumed after the Second World War, would last Heidegger's lifetime. In July, the lecture presented to a society of Marburg Theologians, Der Begriff der Zeit (translated as The Concept of Time), recalls the theme of his habilitation lecture, and also provides an introduction to one of the central distinctions of Sein und Zeit, between temporality in everyday human experience and time as it is conceived by the physical sciences. While his thoughts make reference to Dilthey's attempts to work out the grounds of historicity, we are also looking ahead to Sein und Zeit. Dasein is now conceived, in its most radical possibility of being, as time itself rather than a being in time. But time, contrary to how the tradition has always understood it—that is as presence—must now be understood as futural: "To have no time means to cast time into the bad present of the everyday" (14E). So futural time—the time we are yet to have—is what gives time for our experience of the present in a way that allows the past to be repeated as life. Time, therefore, will now be regarded as the possibility of repetition, which, as Heidegger points out at the conclusion of the lecture, is historicity. The possibility of repetition connects two essential concepts, in the German die Möglichkeit, meaning the possibility, and Wiederholen, meaning repetition, though an attempt to portray correctly the sense of the word in English would need to acknowledge the sense of "revision" that is also implied. The phrase "possibility of repetition" is similar in formulation to the phrase "hermeneutics of facticity." The "of" in each case is what is called a "double genitive" because it can signify in two ways. The interpretation of facticity is what happens when the attempt to interpret facticity yields to the realization that facticity is already what interprets—it is facticity's interpretation. In the same way, if Dasein is time, and if the fundamental phenomenon of time is the future, Dasein would not be separable from its future possibilities, that is, the possibilities available in the repetition of the past. The possibility of repetition, then, just is possibility.

Heidegger's father died in December 1924 at 73. His mother died less than three years later in 1927 at the age of 69, shortly after the publication of Sein und Zeit. In 1925 faculty members at Marburg proposed Heidegger as the replacement for Nicolai Hartmann in the philosophy chair, as a tenured full professor. But in January 1926 Berlin authorities rejected the proposal on the grounds of Heidegger's lack of publications. Heidegger then, in March, worked intensely to produce a substantial draft of the work he was planning at the time and completed the first and most of the second division of the first part of what was supposed to have been a work in two parts with three divisions each. Heidegger presented the manuscript to Husserl at his 67th birthday party in Todtnauberg on April 8. With the work less than half finished Heidegger found himself under increasing pressure to publish something. In June the university again proposed to appoint Heidegger to tenured professor and again in November Berlin refused on the same grounds as before. However the completed sections of Sein und Zeit went to press and were published in April 1927, first in the Yearbook for Philosophy and Phenomenological Research, edited by Husserl, and at the same time as a separate work. In the summer semester Heidegger taught the course, "Die Grundprobleme der Phänomenologie" (translated as "The Basic Problems of Phenomenology"), which is very close in subject matter to Sein und Zeit and helps to fill in some of what is missing there, although the published version did not appear until 1980. Heidegger examines the philosophical history of ontology, with an emphasis on Kant in the first half, and then examines time as temporality and its relation to being. The material covered may have been intended as part of the third division of the first part of Sein und Zeit and perhaps even looks forward to the second part. The published version does not in fact add anything new philosophically to Sein und Zeit but it does help to clarify and deepen our understanding of it. The Kantian context reveals the extent to which Heidegger's thinking had developed during the writing of Sein und Zeit, though this is not as neo-Kantianism but rather by way of a critical reading of Kant that becomes yet more explicit during the lecture courses of 1928 and 1929. In fact the notion of critical reading itself

undergoes a powerful transformation here, as it takes the form of the Destruction of metaphysics that is promised in Sein und Zeit but which in fact constitutes Sein und Zeit in its manifestation as Wiederholung—repetition/revision—of the tradition. For this reason, the texts that lead up to its publication as well as the lecture courses that immediately follow provide an indispensable complement to the study of Sein und Zeit.

Paragraph 8 of the Introduction to Sein und Zeit, which aims to address the question of the meaning of being, sets out the design of the treatise. The response to the question involves first, an interpretation of the entity for whom being is a question—Dasein; and second, because Dasein is a historical entity, an interpretation of historicity. Heidegger explains that, as the treatment of the question of being branches out into these two distinct tasks, the treatise itself must have two parts. The first part is concerned with the interpretation of Dasein in terms of temporality and "the explication of time as the transcendental horizon of the question of being." It was to have three divisions: (1) a preparatory analysis of Dasein; (2) the analysis of Dasein and temporality (Zeitlichkeit); and (3) an exposition of the relation between time and being. The third division never appeared. Part Two was supposed to have addressed the basic features of a phenomenological destruction (or deconstruction) of the history of ontology, with the problematic of temporality (Temporalität) as a guide. It was also supposed to have three divisions: (1) Kant's doctrine of schematism and time; (2) the ontological foundations of Descartes' cogito sum and medieval ontology; and (3) Aristotle's essay on time, providing a way of examining the limits of ancient ontology. None of Part Two ever appeared but he plan is clear enough in outline. The destruction would work backwards as a way of disclosing the repetitive nature of the problematic of temporality and ontology, working first from Kantian ontology, then back through the repetition of medieval ontology represented by Descartes, which itself would be revealed as a repetition of ancient ontology and the limits imposed by the phenomenal basis found in Aristotle. Indeed the groundwork of Sein und Zeit can be traced back to Heidegger's critique of Aristotle's concept of time. What Heidegger published corresponds with the design for the first two divisions. The book starts with an analysis of Dasein, as the being for whom being is an issue and a question and, therefore, as the being on whom the question returns. In order to address the meaning of being in general, the being who poses the question must first look into its own way of being. The use of the term Dasein (which is accordingly rarely translated) remains as one of Heidegger's most striking innovations. An ordinary German word, which in most philosophical contexts means something like the English existence, takes on a forceful character in Heidegger's use, in which the etymological root—Da-Sein—is emphasized. Heidegger makes a number of other innovative terminological distinctions early in Sein und Zeit, which help to distinguish Dasein and its peculiar way of being. The most important of these is the distinction between the ontological and the ontical. Paragraph 4 addresses the "ontical priority" of the question of being. The ontical denotes whatever pertains to the factual world, the world conventionally explored by the sciences. In this world a peculiar fact emerges concerning the being that Heidegger calls Dasein. It is distinguished from all other beings by the fact that, for Dasein alone, being is an issue for it. Neither rocks nor vegetables, nor even other animals, concern themselves with the question of the meaning of being. So, ontically speaking, Dasein is distinguished by being ontological, which means that it has a relation to being. However, just because Dasein is ontological that does not mean it has developed an ontology. Heidegger reserves this term for the theoretical inquiry into being. Dasein, then, is ontically distinguished from other beings, insofar as it has an understanding of being, but for the most part it remains "pre-ontological" because it has yet to work out its ontology. Heidegger adds to this distinction between the ontological and the ontical a further distinction, which is, in the context of the years leading up to the publication of Sein und Zeit, quite surprising. Existence is the term that Heidegger uses to denote the kind of being towards which Dasein comports itself, reaching both for future possibilities and back into the past. Heidegger points out that because Dasein has its being "to be," (i.e., to come) its way of being cannot be designated by a subject matter—a what. Rather, Dasein understands itself in terms of its possibilities—it can either be itself or not itself, he writes, thus turning from the question of what to the question of who. It is the ability to make decisions about the future that characterizes Dasein's existence. Accordingly, Heidegger uses the terms existentiell (existenziell) and existential (existenzial) to distinguish between the everyday decisions made by the particular Dasein and the a priori structures and modes of existence itself (the terms are surprising because they seem to recall the contemporary and fashionable versions of existentialism that Heidegger despised). Ontology must begin with

the pre-ontological understanding of being because this is the definitive characteristic of Dasein. We operate in the everyday world and must deal with other entities in practical ways. To that extent we are existentiell, which implies practical competence and an implicit understanding of our existence. On the basis of this implicit understanding it should be possible to develop an explicit theory of being, an ontology. In this way, Heidegger claims, the question of being aims at the a priori conditions of possibility not only of the objective sciences but also of the philosophical ontologies that lie at the basis of all sciences.

Having defined Dasein in terms of existence Heidegger develops the implications, which effectively overturn a number of metaphysical doctrines. Dasein cannot be considered as a particular (belonging to the genus Man, for instance). Rather, as an entity considered as its own possibility, Dasein is in each case its own being. Its possibility or existence thus always overflows or outstrips its actuality. This possibility—or "mineness"—implies two alternative conditions: Eigentlichkeit or Uneigentlichkeit. The latter condition (which is usually translated as inauthenticity, though not without controversy) should be regarded as the normal condition for Dasein most of the time. It implies that in this modality Dasein is not properly its own being. To be authentic is to be one's own self rather than to belong to some other—a social group, family, friend, institution or whatever. Heidegger's term for these others, das Man, can be translated as the "one" or the "they," and it implies a condition according to which Dasein exists without making its own decisions, conforming instead to habits, customs and practices that determine relative cultural conditions. The authentic condition, on the contrary, implies resoluteness in assuming the ability to make one's own decisions.

The authentic and inauthentic ways of being that are grounded in Dasein's existence correspond to the existentialia that characterize Dasein's everydayness. By existentialia Heidegger means the a priori conditions of possibility for being in the world. Because the world, in this sense, is a world only for Dasein (there is no world independent of Dasein, although there are certainly beings that are) then the other kinds of entity that populate it must be considered in terms of how Dasein engages with them. Dasein's world is populated not only by others but also by tools and equipment, which are not simply vorhanden (present-at-hand), inert and without practical significance, but they are zuhanden (ready-to-hand), available for work or leisure, and without our needing to deliberate unduly over them.

Dasein can also be characterized in terms of the moods it finds itself in—its throwness. Befindlichkeit (the state I find myself in) can be grasped as a kind of attunement. Heidegger's examples, Angst and boredom, illustrate what is meant by moods: they are conditions of being that, if one is responsive to them, disclose the world in its ontological dimension. Neither Angst (anxiety over nothing in particular) nor boredom can be said to have a specific object or external cause. Rather they are moods capable of disclosing the world in its everyday ordinariness and can lead to an understanding of it.

Dasein is constantly dealing with entities whose meanings are implicitly understood in the way Dasein makes use of them and in the way they are related to each other. This everyday concern, which Heidegger characterizes in terms of the way the craftsman operates in his workspace, can also be made into the topic of philosophical reflection, so that the mode of being of entities in the world is shown to be special to the mode of being of Dasein, revealing the a priori nature of Dasein's world as a "for the sake of which," grounded in Dasein's own existentiality—possibility itself. Similarly, Dasein's understanding of the world is disclosed as an active interpretation, projecting an a priori condition, Dasein's own possibility, onto the entities that populate its world. The significance of the tools or equipment that Dasein habitually uses turns out to be a projection of Dasein's mode of existence. Dasein understands its environment in terms of possibilities offered and is therefore always active, always meddling in some way. This understanding is grounded in kinds of interpretation, involving a priori a fore-structure: a Vorhabe, a Vorsicht and a Vorgriff (a fore-having, a fore-sight and a fore-conception). Heidegger points out that we never encounter equipment that is ready-to-hand in a way that involves having to work out its significance before using it. We do not first encounter a door, for instance, as an inert piece of wood and then derive its use from contemplation. We encounter it first of all as already a door. The scientific attitude (typified by Descartes and his world of res extensa) must have been derived from and abstracted from the more primordial one, in which the "as" that

constitutes the mere understanding of a table as a table (or a door as a door) is given in advance. So whenever something is interpreted as something (which is the normal state) the interpretation must be taken as having been given in advance.

These considerations of mood, understanding and interpretation emphasize the active role of being-in, the verbal aspect of being-in-the-world. They also indicate why Heidegger privileges temporality. Dasein's thrownness, revealed through moods like Angst and boredom, show that Dasein's existence involves both having been and being to come. Dasein can either exist in an inauthentic way, governed by das Man and the fashionable currency of the everyday, or it can exist in an authentic way, projecting its own possibilities into a future that it can have a say in determining for itself. Through the experience of anxiety Dasein learns that it is thrown in a world that is already marked by significance; this is its facticity. It also learns that the character of this thrownness is possibility and, therefore, that its existence is its own. Finally, it learns that it is fallen among other beings whose being is not its own. The fundamental unifying structure that supports these related dimensions is Sorge (usually translated as care). Care must use time and must reckon with time. The "time" that that Dasein experiences, however, is the phenomenal aspect of temporality—the proximal experience of the present. And this, Heidegger argues, is not only what arises as the everyday understanding of time, but also what has evolved into the traditional ontological understanding of time, according to which being is determined as presence. Having derived the ontological meaning of care, according to which temporality involves both running ahead and reaching back, Heidegger moves on to Division Two.

Division Two begins by acknowledging that the "Preparatory Fundamental Analysis" of Division One has only dealt with being in the world in a partial way. The structure of care has revealed a depth and complexity that demands a different approach. If being in the world involves reaching back into the past and running ahead into the future then Division One has only managed to bring into view the everyday happenings between birth and death. The structural roles of the end and the beginning remain obscure. Furthermore, the character of "mine-ness," which forms the ground of both authentic and inauthentic existence, has been brought into view only in terms of the inauthentic, average way of being. It will thus be the task of Division Two to reveal in the structure of care the deepest ground of the whole of Dasein's being. Division Two begins, accordingly, with the questions of how Dasein can be a whole and how Dasein can wholly be.

For this reason, Heidegger argues, the analysis must engage with the most extreme possibilities of Dasein's existence, the possibilities of birth and death (the ends of being) and the possibility of authentic existence, in which Dasein can be wholly its own. The first problem emerges with the terms of the question itself: in what sense can a being, who is to be understood in terms of its finitude and freedom, also be considered as a whole? The essence of Dasein's care is its active interpreting, which follows from an essential constitutional structure of Dasein, the temporal phenomenon of the future, the ahead-of-itself, according to which there is "eine ständige Unabgeschlossenheit," constantly something to be settled. So Dasein cannot be considered a finished whole in the conventional sense, a sum of attributes or a living thing. If Dasein is to be considered as a whole then this will be through an interpretation by Dasein that can disclose the whole of its own being—its own disclosing character. Dasein's being as a whole can only be disclosed by Dasein. The first serious problem thus emerges. The two extremes of life—birth and death—remain outside the temporal structure of Dasein's thrownnness. Dasein can never reach back behind its thrown facticity to the moment of birth nor run ahead so far as to experience its death before ceasing to be. This problem rules out the method operational in Division One, where the ontical and existentiell aspects of existence served as an essential starting point for approaching the ontological and existential problem. Here, Heidegger points out, the problem of not being able to eradicate the "ahead-of-itself" as an essential structural aspect of care need only be regarded as insurmountable from a perspective that posits Dasein as a present to hand just waiting for a not yet present to hand to be added onto it. Rather, if it is going to be possible to make Dasein accessible in its wholeness, it must be disclosed (as it must disclose itself) in an existential and ontological way.

Heidegger accomplishes this through an extended and detailed series of analyses, which link the problems of death, guilt and history to the meaning of care as time; and he concludes by arguing that Dasein's being is always and from the beginning being towards an end. The fore-structure, or a priori given-ness, of the end of its existence forces Dasein to relate to this end in definite, authentic or inauthentic, ways.

Heidegger uses the Greek term ekstasis (which has etymological connections with the Greek ek-sistence) to denote what he calls "the ecstatic unity of temporality," according to which it is possible for us to experience other beings. The unity describes the way that time is constantly "outside-itself," in so far as the present must be regarded as derived from a combination of having-been and the future. Ekstasis designates how a unity of future, having-been and present allows the human being to be "cleared." It provides a spatial and temporal present that is only authentically arrived at through this state of Ekstasis. Modern life increases the demand on existential functions like Ekstasis, provoking an escalation of the repetitiveness and intensity of the acts or performances of presencing (which allows beings to present themselves in particular times and spaces). Heidegger's radio broadcast from 1934 defending his decision not to come back to Berlin to take up the prestigious post offered to him gives us a sense of how he felt about this: "The world of the city," he said, "runs the risk of falling into destructive error," and he identifies "a very loud and very active and very fashionable obtrusiveness." This echoes his remarks from 1924 in the "Begriff der Zeit" lecture, according to which we can have time only by maintaining ourselves along side our past and running ahead (i.e., historicity). Alternatively: "All idle talk, that in which idle talk maintains itself, all restlessness, all busyness, all noise and all racing around breaks down" (14 E). It would seem, then, that from Heidegger's point of view, repeating and revising the past while running ahead into our future is the only way for us to have time, which for Heidegger, is to have ourselves.

Shortly after the publication of Sein und Zeit Heidegger's fame as a philosopher quickly spread internationally, and consequently he was promoted to full professor at Marburg on October 19. In 1928 he was offered the chair of philosophy at Freiberg, which had been vacated by its founder, the retired Husserl. The new post represented the highest honor possible in the field of phenomenological philosophy and Heidegger naturally accepted the invitation and took up the position in time for the winter semester.

In 1929 he published Kant und das Problem der Metaphysik (translated as Kant and the Problem of Metaphysics), the so called Kantbuch, which explicitly sets out the conditions according to which the futural dimension of time makes the kind of interpretation possible that would move things forward. In Sein und Zeit Heidegger had argued that the historicity of human existence, or Dasein, implies a legacy that we inherit. Historicity is not the same as history. Historicity is the condition of being historical. And as Sein und Zeit teaches, the aim of locating and understanding conditions cannot necessarily be achieved by simply studying the phenomena (i.e., historical events) that they make possible. In other words Heidegger is interested in the connection that Aristotle had suggested between what is necessary and what is possible (but not always necessary). In a strange reversal the necessary turns out to be possibility, i.e., the possibility of repetition, which overflows all the ways in which possibilities can be actualized. The consequence for Heidegger is that the best way forward, in any of the dimensions that humans operate in, implies types of response to the traditions, legacies and histories that make us what we are. That response (our responsibility—the ability to respond with a modicum of freedom to a tradition) always implies some minimal unthought-—the undetermined futural aspect of time.

The notion of repetition, Wiederholung, which was introduced in the lecture Der Begriff der Zeit, and provisionally analyzed in Sein und Zeit, is yet more fully worked out in the Kantbuch. This is his definition: "By the Wiederholung of a basic problem, we understand the opening-up of its original, long concealed possibilities, through the working-out of which it is transformed. In this way it first comes to be preserved in its capacity as a problem. To preserve a problem, however, means to free and keep watch over those inner forces which make it possible, on the basis of its essence, as a problem." The revision operates, then, on the possibilities of repetition. What are preserved are the "inner forces" that make a problem possible. What is affirmed is the problematic. The strategy thus involves the location of a persistent problem (for instance, the problem of time or of being), the working out of

its conditions and the repetition of those conditions in a response that perpetuates the problematic (i.e., the future). A basic (grounding or founding) problem of metaphysics can always be interpreted as a possibility of its future. In this way the legacy or tradition can be read in terms of its Geschick—not its destiny but its open possibilities of destination; not as fate in any of the traditional senses but as sending forth in a sense closer to that of telecommunication.

Heidegger's inaugural lecture at Freiburg, Was ist Metaphysik (translated as "What is Metaphysics?"), on its publication in the same year, attracted considerable attention in the Anglo-American analytic tradition, for whom it confirmed the suspicion that Heidegger, and by extension other fashionable continental trends in philosophy, were hopelessly unclear and muddled. This suspicion hardened into prejudice in the years that followed and led to almost wholesale scorn towards Heidegger's thought among the English speaking philosophical community. At the same time the lecture attracted hyperbolic praise among existentialist philosophers and led to a long term association of Heidegger's thought with French existentialism. The lecture can be read as a challenge to philosophize. It takes up issues familiar from Sein und Zeit, particularly the analysis of anxiety and nothingness, but presents them in a new and provocative way. Heidegger's language seemed to some to be contorted and unwieldy. The notorious phrase from the lecture, "the nothing noths," draws attention to a characteristic that had long been a part of Heidegger's style—the active verbalizing of noun phrases. Just as being is (re)verbalized in its context in Sein und Zeit as Da-Sein, the active be-ing of being-in-the-world, so too the nothing is put to work as an activity constitutive also of Dasein's world. The nothing helps to disclose being in its previously concealed strangeness, showing how the transcendence of Dasein is held out over the nothing from the strictly finite and limited vantage of being, which can now be revealed in contrast as temporarily meaningful, but bounded by meaninglessness.

Heidegger's rise to fame between the end the of the first war through the years immediately after the publication of Sein und Zeit coincided with the period of the Weimar Republic, a period of turbulent political activity, economic depression and extraordinary cultural vibrancy. By September 1930, Adolf Hitler's National Socialist German Workers Party (NSDAP or the Nazi Party) had become one of the most powerful and popular political organizations in Germany. Heidegger's political leanings had been unfocused and largely localized in his youth and had not been evident at all during the intensely productive years leading up to Sein und Zeit. Nevertheless, during the early 1930s, he became sympathetic to Nazism. He had read Hitler's Mein Kampf and, along with many of his generation, had sought in the political world an answer to the global problems caused by what he had started to describe as the fulfillment of metaphysics in technology. On 30 January 1933 Hitler was appointed Chancellor of the winning coalition of right wing parties and, after 27 February, absolute power was conferred on the NSDAP. Heidegger's enthusiasm for the new regime led, in a series of swiftly changing circumstances, to his appointment as rector of the University of Freiburg on 22 April, and on 1 May he became an official member of NSDAP.

On 27 May Heidegger gave his rectoral address, Die Selbstbehauptung der deutschen Universität (translated as The Self-Assertion of the German University), which provoked ironic responses from some commentators of the time, who suggested substituting Selbstenthauptung (Self-Beheading) for Selbstbehauptung, having observed that self-assertion, in Heidegger's view, meant adherence to National Socialism. In a move that is actually entirely in line with his long term philosophical preoccupations, he condemns contemporary versions of academic freedom as arbitrary and lacking constraint. He sees the occasion as an opportunity to establish a firm ground at last for knowledge, which would spell the end for the looseness of the various ungrounded empirical sciences and the domination of relativity amongst the disciplines. Despite his clear alignment with the party, however, his sense of what this would mean has less to do with concrete Nazi politics—there is no discussion of racial superiority or of global domination—but more with his own philosophy, such that the confrontation with being would act as a ground and unifying force for the German University in its role in the destiny of the German Volk and in what he calls its spiritual mission. Students will henceforth fulfill their mission in three ways: in labor, in the military and in knowledge. Furthermore, Heidegger sees this mission as one that will restore to the west its spiritual strength before what he calls the "moribund pseudo civilization" collapses into itself. On 11 November he gave a radio broadcast

supporting Hitler's withdrawal of Germany from the League of Nations, revealing both an idealism regarding international relations and apparent ignorance about the nature of National Socialism itself.

During 1933 some of the Jewish people most closely associated with Heidegger were forced to leave the country. His ex lover Hannah Arendt and his lifelong correspondent Elisabeth Blochmann would both renew relations with Heidegger after the war but for the moment their contacts were broken off. Heidegger's contacts with Husserl also ceased at this time. Relations between the two had anyway become strained when Husserl had digested the extent to which Sien und Zeit departed from his own project of phenomenological science. But their break during this time contributed to the unfounded rumor that Heidegger had banned his friend and ex-teacher from the university. While Heidegger did little or nothing to discourage the official dissociation of Jewish colleagues and students from the university during his time as rector, there remains scant evidence of any overt anti-Semitism. Heidegger's relations with the party were neither comfortable nor mutually supportive, despite his consistently expressed enthusiasm for the opportunities the regime seemed to him to represent for his ideals. Conflicts with party officials as well as with students and faculty led him to resign as rector on 21 April 1934, after which he played no further part in political affairs, though his thought would continue to engage on an increasingly more developed political level. In June Hitler had a number of his rivals murdered, including Ernst Röhm's Storm Troopers and a number of party members, like Gregor Strasser, whose politics were not sharply focused on opposition to Jews and to communism. The so called Röhm putsch turned out to be a thinly veiled pretext for clearing the party of potential rivals and on 1 August Hitler became "Fürher of the German Reich." Heidegger later claimed that he became disillusioned with National Socialism after the Röhm putsch but there exist both witnesses and documents **suggesting that** Heidegger's support of Nazism continued at least until 1936.

Heidegger did not publish very much during the remainder of the war, though he continued to lecture. In 1934 he began a series of lectures on Friedrich Hölderlin, which would occupy him until the early 1940s. Hölderlin, the enigmatic German poet and contemporary of Hegel, emerges in Heidegger's discussions as a resource for his radically alternative exposition of the place of human being in the world. In 1935 he lectured on "Der Ursprung des Kunstwerkes" (translated as "The Origin of the Work of Art), three successively different versions of which emerged as articles. While the earliest version is characterized by the same preoccupations as his lecture courses of that year, by the time the final version **was published these had been refined** and the final version reveals how Heidegger's thought had been developing. The key distinction in the article is between world and earth and the article works through three rather different kinds of artworks. His reading of Van Gogh's Shoes shows that art can be regarded neither as merely an aesthetic object designed to give pleasure or to portray beauty, nor as a kind of thing with the addition of aesthetic beauty. Rather, art discloses the nature of things. Equipment, unlike art, disguises its status as a thing. Its material nature is absorbed in its function. The artwork, however, draws attention to the materials from which it was formed. The artwork thus draws attention to the struggle between form and matter. Van Gogh's painting of shoes shows what would not normally be evident. The involvement of the shoes in the world of the peasant disappears from view while they are being worn; but put into view, as in Van Gogh's painting, the shoes reveal that world itself and the relation of that world to the earth. But what is the earth? Heidegger shifts his attention to a Greek Temple, partly to underline the fact that his reading of artworks is not based on a model of representation. The shoes in the Van Gogh example do the work of explication not because of the painting's naturalistic or evocative qualities. Rather the role of the shoes as equipment is revealed in an artwork that takes as its topic the medium between the world of the peasant and the earth on which the peasant treads. A famous reading of Heidegger's artwork essay takes him to task rather pedantically by pointing out that the shoes in Van Gogh's painting were actually his own. The temple shows how a work of art not only opens up a world but also unifies and structures the world of a historical people. In doing so it contextualizes the earth upon which it stands, instituting a particular interpretation of the relationship between the cultural contrivances of Dasein and the natural world with which those contrivances are engaged. Finally, Heidegger turns to what he calls Dichtung, which in its fully etymological sense means invention. The ordinary and the everyday is made strange in art, revealing the struggle between the newness of art and the state of things out of which it had to have emerged. The meaning of

Dichtung in the normal sense is, of course, poetry. Because the matter or earth of poetry is language and because language is what gives Dasein names for beings, then poetry has the power of addressing the possibility of human communications and relations. The relationship between world and earth, when it takes the form of linguistic innovation, reveals the torturous ways in which the relations between concepts and words are formed and form each other. Poetry can thus be grasped as the most essential kind of artwork because it performs an absolutely singular intervention that is also a form of disclosing. Poetry reveals the conditions on which not only artworks but all other kinds of communicating and all other kinds of thing are possible at all. For this reason Heidegger increasingly privileges Dichtung in his works of this period and later. The disclosing of being—if it is to be achieved in any way that eludes the classifying, calculating procedures of modernity—must be an evidently singular event each time.

The changes in focus that Heidegger's philosophy passes through during this time have been described as a Kehre (a "turn"), though the nature and implications of this Kehre have since been a topic of dispute and debate and remain undecided. When Heidegger uses the term himself he tends to refer to a stage or procedure that allows the passage from Division Two to Division Three (which did not appear) of Sein und Zeit. In 1935 Heidegger's thought is still dominated by the distinction between everydayness and what had become "creative self-assertion," a version of the distinction in Sein und Zeit between inauthentic and authentic existence. If Die Selbstbehauptung der deutschen Universität had transferred the topic of fundamental ontology from individual Dasein to a people or a state, then the resoluteness of one's ownmost being becomes the creative self-assertion of a people's spirit. What this implies is that inauthentic everyday distraction by the present should be overcome in the authentic disclosure of being as historical destiny. The lecture course on Hölderlin makes this explicit. The task is to learn how to listen to the poet. The approach is similar to that which begins Sein und Zeit, insofar as it begins with the need to clear away the obstacles to asking the question, "Who is Dasein?" But in the Hölderlin course, the everyday world and its distracting forms must be discarded before Dasein can be disclosed as the authentic gathering of individuals in a community. The argument suggests that Heidegger was locked in a struggle both with himself and with the politics of the German nation. The critical, arguably moral, attitude towards the everyday—which is not evident in Sein und Zeit—relates explicitly to the everyday life of the Nazi regime, which Heidegger found himself in the midst of: the political subordination of thought, art, history and writing to the urgency of everyday political interests; the busy cultural activities; the explicit biologism of racist dogma; and the ubiquity of bureaucratic procedures and hierarchies. Heidegger thus sought to transcend this everyday Nazism in an authentic version that would disclose the innermost historicity of a people. An authentic historicity would be the domain of only a few creative individuals, specifically, poets, thinkers and founders of states (thus evoking the triumvirate: Hölderlin, Heidegger and Hitler). The poet institutes the truth of a people's Dasein. The thinker explicates the poet's disclosure of being. The creator of states can manage the state in such a way as to adjust it to the essence revealed and explicated by the poet and thinker respectively.

Einführung in die Metaphysik (translated as Introduction to Metaphysics, 1959), from the lecture course given in the summer of 1935, reveals beneath Heidegger's compromising adherence to a spurious kind of geopolitical propriety, a degree of insight that ultimately renders his political ideals untenable. He gives an account of Europe, "in its unholy blindness," as always on the point of cutting its own throat. In the pincers between Russia on the one side and America on the other, Europe struggles against the mundane technologization it has itself spawned as the outcome of its metaphysics. Russia and America, seen metaphysically, Heidegger explains, are both the same: "the same hopeless frenzy of unchained technology and of the rootless organization of the average man." Heidegger's characterization of Europe—with Germany at its core—is quite complex. Europe would always have been the heir of the historicity of metaphysics, which as Heidegger teaches, completes itself in technology. Europe's "unholy blindness" would therefore be the inability of the modern subject to see its own essence. The "darkening of the world" thus described by Einführung in die Metaphysik implies a fulfillment of Europe's metaphysics in Stalin's Russia and in capitalist America, now returning as the ends of the pincers and surrounding their own source. Thus Europe would be at once the hinge and the ends of the pincers. What Heidegger is most concerned about here is the instrumentalization of spirit. The tripartite geographical distinction represents three types of reduction of spirit to

the mere instrument of intelligence: the arranging of material relations (in Stalinism); the ordering and explanation of the present to hand (of Positivism); and management of vital resources and race (everyday National Socialism). So spirit relates to the dimension in which the gods might again become thinkable: that is, to the holy. The realm of the holy [das Heiligen] and its cognate Heilen or "healing" (resonating at once with notions of salvation, well-being and wholeness) signal both Heidegger's theological heritage and his absorption of ethics into his fundamental ontology. The questioning embodies a mode of thinking through which, as Heidegger explains elsewhere, the truth of Being becomes a distress. The questions, "what for?—where to?—and what then?" represent the echo of Heidegger's question of being, provoked by circumstances which by the mid 1930s had become commonplace as the systematic global urbanism of modernity spearheaded by mass communications and the rapid expansion of military powers.

In 1936 Heidegger went to Rome to lecture on Hölderlin. He met his former pupil, Karl Lowith there, who has since insisted that Heidegger's adherence to Nazism was still evident at that time. In that year Heidegger also began his series of lectures on Nietzsche, which ran until 1942 and which were eventually published in 1961 in 2 volumes. The first volume, which covers the years 1936-39, interprets Nietzsche approvingly as foreshadowing Heidegger's analytic of Dasein and is consistent with Heidegger's other writings of the mid 1930s. But the second volume departs both in style and content from the first, in that it is marked by a polemical tone directed against Nietzsche and characterizes Nietzsche as the fulfillment of the tradition in what turns out to be an account of western metaphysics that finally seems to be doing the work of destruction advertised in Sein und Zeit.

The argument is foreshadowed by the book-length work that Heidegger wrote between 1936-38, probably for his private use, and which he did not publish in his lifetime, Beitrage zur Philosophie (Vom Eriegnis) (translated as Contributions to Philosophy (From Enowning), 1999). It contains a horrified account of the abandonment of being in modernity and suggests the extent to which Heidegger's thought was undergoing some alterations in emphasis. Being is not anything that is. So being withdraws in granting to beings their presence. Everything that exists does so because we as Dasein stand in a relation to being. But being will be discovered in its most essential form, time and time again—and in this repetition—as nothing. "There is nothing to being." The statement is enacted in the enunciation of increasingly empty phrases and achieved through this recitative mode, which springs from the various "moods"—deep awe, wonder, boredom, etc—that disclose beings in relation to their being. Heidegger's lifelong philosophical project manifests here as the desire to empty all ontic statements of their content in order to open up the relationship between the beings that appear and their secret and thus un-thematizable emergence. This emergence is explicated by the phrase, "Being essentially unfolds as appropriation [Ereignis]." The subtitle, (Vom Ereignis), offers a deliberate and, Heidegger says, proper alternative to the conventional academic banality of the main title. Ereignis is difficult to translate into English. The translators of the Beiträge coin a new word, enowning, in an attempt to indicate how it works in Heidegger's German, stressing both the active, verbal aspect as well as the sense of appropriating that it suggests etymologically. Elsewhere the phrase "event of appropriation" and the catachresis "event of propering" (this with some irony) have both been used. The word, in its conventional sense, might have been faithfully transposed as "event." Heidegger claims that his use of it cannot be translated at all; it is for him a key term, like the Greek logoV or the Chinese Tao. It does not any more mean a happening or occurrence in the ordinary sense, but it does retain the sense that what happens does so uniquely and in the singular. This resistance to translation thus remains faithful to the sense of the term itself. It supposedly resists much more than simple translation. The singular and unique unfolding of being as Ereignis could hardly fall under a conception. Furthermore, it could hardly come into experience at all, if all the forms of representation, predication, classification, calculation and conceptualization, which characterize modernity from Plato to Nietzsche in an intensification of the obliviousness to the unfolding of being, now dominate, as Heidegger claims they do. The difficulty is not simply a matter of forgetfulness. It is a structural condition, according to which the singular event always and immediately drops out of sight, withholds itself in what it grants.

The fact that this withholding can only be experienced as a lack is what leads Heidegger to characterize technology as the perpetual organization of this lack—where technology disguises the emptiness of being beneath an insistent functionality. In his Überwidung der Metaphysik (published in 1954 but based on notes made during the 1936-40 Nietzsche lectures and translated as "Overcoming Metaphysics," 1959) he suggests that the animal (or sub-human) instinct has become indistinguishable from calculating rationality. Heidegger's reflections here turn to the phenomenon, which for him fulfils technology, of armament, as the inevitable consequence of "the ordered use of beings which become the opportunity and the material for feats and their escalation." What remains concealed by the escalating consumption of beings, is that the aim of technology is purely aimless. To put it succinctly, the end of armament is the end of man. In this way the groundlessness of an ethical ground—its unconditionality—shows that self-guaranteeing armament seems designed to remove the power of decision and responsibility entirely from the human sphere. Das Man has been entirely captured by the machination of a technics that he fails to grasp in its historicity, and so is in this way enchanted by his disenchantment. The emergence of this distress would nevertheless still be found as the historicity—or emergence—of technology itself. A profound and disturbing truth beyond truth emerges, which inevitably imposes a powerful deconstruction on Heidegger's adherence to the spirit of the German Volk still inherent in his arguments of the mid to late thirties. Heidegger's philosophy begins increasingly to emphasize the enigmatic withdrawal of that which grants presence. Rather than maintain the strict division between the authentic and the inauthentic in terms of the voluntarism that marks the Hölderlin lectures and Einführung in die Metaphysik, according to which Dasein discloses its own truth, Heidegger increasingly characterizes truth as a mode of un-concealment to which humans are exposed, and to which they belong. Whether this is what should be understood by Kehre remains debatable. There is no clear distinction between the two kinds of emphasis; rather the difference takes the character of a refinement. Heidegger's philosophy develops through what is most characteristic in his style as well as in his subject matter—the repetition/revision that also characterizes the work of deconstruction. The entity which in Sein und Zeit anticipates the end of its existence and is thus capable of freely taking over its own being, now belongs together with being on the basis of an unfolding that withholds itself. In each case the relationship between Dasein and being eludes the conceptualizations that dominate metaphysics.

The Nietzsche lectures have been read as a covert criticism of the Nazi regime's appropriation of Nietzsche in support of its racist practices, especially the second volume, which includes the lectures given during the early years of the Second World War. Heidegger was under Gestapo surveillance at this time, according to the confession of an official infiltrator who claimed that he rejected Nazism on the strength of Heidegger's way of thinking. The arguments of the period are characterized by regular references to the contemporary political situation—the events leading up to and after the breaking out of war—and each time related to the metaphysical foundations of technology and instrumental reason and to the forgetting of being. Not the self assertion of individuals, dictators or authoritarian states (a clear if still covert reference to the Nazi regime), but rather "the metaphysical essence of modernity," or, echoing Nietzsche, the will to power over nature, leads to the emergence of powerful imperialist movements. Nietzsche's doctrines of the will-to-power and the eternal return of the same are now read as the culmination (and not the overcoming) of western metaphysics in nihilism. Nihilism would thus be the culmination of an increasing forgetfulness of being. Heidegger returns to the earliest documented sources of Ancient Greek philosophy, the pre-Socratics, whose term al-ētheia is normally translated as truth. In Heidegger's reading it designates a process in which beings appear yet what grants them presence reserves itself, withholds itself in what it gives. Heidegger argues that this indicates an experience of being in which withdrawal or withholding is experienced simultaneously with what appears. For the earliest philosophers, then, thinking would be a matter of remaining receptive to the withholding of being. In Plato's philosophy, however, the mere effects of the process—the appearing of things, which are regarded as the manifestation of ideas, then get abstracted from the process of unconcealment and set apart as clarity and light. Thus, for Heidegger, the essence of truth as idea in Plato signals the beginning of metaphysics and onto-theology, and can be grasped as the first stage in the forgetting of being and the obliteration of alētheia. Truth now means conformity of the mind to essence. A further step in the forgetting of being occurs with the scholastic philosophy of the Middle Ages where truth as adequatio (adequacy of the mind to

the thing) is grounded in the concept of divine (i.e., the mind of God). At the beginning of the so called modern age a further step in the oblivion of being occurs with the establishment of the human subject, and with the rationality characteristic of Descartes and Galileo. In Descartes the cogito sets itself up as the basis on which objects reveal themselves. Heidegger traces the term hupokeimenon (the Greek for basis) through its translation into the Latin subiectum to the institution of the human cogito as subject of philosophy and the one true basis upon which things can appear.

So, according to Heidegger's reading of the tradition, at the beginning of the modern era a technological interpretation of the world was instituted in which the whole of nature was regarded mechanically. Heidegger traces Nietzsche's notion of will to power back to this beginning, insisting that Descartes and those who followed had already anticipated it. Nietzsche's philosophy, which claims to break from past philosophy, actually brings it to its climax as the ultimate accomplishment of onto-theology in nihilism. The relation between subject and object discovered in Descartes has been pushed to its extremes in the notion of will to power. The will regards everything as an object to be stored and manipulated and, in an extreme form of subjectivity, reduces everything to the values it projects upon them in its drive for power over them. The meaning of being is now as far from the Ancients' conception is it could be. Being has become as nothing.

By the 1940s Heidegger had began to express his belief that the route taken by Sein und Zeit had been in danger of unintentionally repeating the intensification of the subjective turn of modern philosophy, thus reinforcing the era of calculability and technological domination. Since the Beiträge he had been using the term Ereignis in opposition to the term Gestell—the en-framing that subordinates beings to the interminable conceptualizing, evaluating, storing, manipulating and calculating of the technological era. In the Nietzsche lectures the discussion of Ereignis recapitulates the work of the Beiträge. Ereignis designates that which thinking can meditate upon in resistance to nihilism, in which thinking has been replaced by calculation and instrumentality. The notion of the frame plays a crucial role in Heidegger's discussions because it allows him to show how the technological era has long been anticipated by metaphysics and that to move forward it will be necessary to meditate on the emergence of the frame itself. Framing has taken specific forms, as the Nietzsche lectures teach, but each time the determination of being as presence divides the transcendental from the empirical, giving priority to the modes of framing by which philosophy designates its subject matter. In Ereignis the possibility arises that it might overcome the merely dominant character of the frame and transform the frame itself into the Ereignis, the singular and unique unfolding which gives rise to the belonging together of mind and being. In other words the possibility arises of meddling with the transcendental itself. If the Ereignis just is that possibility then it will be a matter of repeating the Ereignis in a way that is each time unique. The special character of Wiederholung is once more activated, as a repetition that is at the same time something new.

Towards the end of the war, late in 1944, Heidegger was drafted into the Volkssturm, the people's army, as one of the "least indispensable" members of the University, to join a labor force whose work was to dig trenches by the Rhine. The following year, as Germany approached collapse, he managed to return to Messkirch to put his manuscripts in order and to arrange their security. A manuscript dated May 1945, but rediscovered long after Heidegger had died, provides a clear sense of how Heidegger felt about Germany and its collapse by the end of the war. "Abendgespräch in einem Kriegsgefangenenlager in Rusland zwischen einem Jüngeran und einem Älteren" (Evening Dialogue in a Prisoner of War Camp in Russia between a Younger and an Older Man") uses the fictional premise of the dialogue (though both his sons were at that time in camps, as the title describes) to express the conclusion that the Nazi regime had been a catastrophe for Germany. It goes on to develop a thesis already typical in Heidegger's writings, such that Germany in defeat is not so much responsible for the evils of the war but more a victim of global devastation consequent on the metaphysical oblivion of being. In June, after Germany's capitulation, he was forced to attend a hearing on his political activities by a University denazification committee in French occupied Freiburg, to his documented indignation. The hearing considered testimony from a number of important figures, some of whom implicated him while others defended his actions. One of the most decisive

testimonies came from his former friend Karl Jaspers, who volunteered instances of Heidegger's active support of National Socialism during the years 1933-34. In 1946 the hearing concluded that Heidegger should be banned from teaching. Jaspers had suggested that the German youth after years growing up in an authoritarian state were not ready for critical responses towards a teaching that had once been associated with the Nazis. Heidegger was, however, allowed to keep his library and was granted the title of emeritus professor.

Shortly after this, experiencing emotional crisis, Heidegger sought psychiatric help from Medard Boss who not only helped him regain his emotional stability but who came to admire and to be influenced by his patient's way of thinking. Heidegger held regular seminars for Boss's circle, made up primarily of professionals trained in the thinking of the sciences and social sciences. Boss would, with Ludwig Binswanger, go on to develop what became known as existential psychiatry.

During his time in French occupied Freiburg, Heidegger had been introduced to the French intellectual, Jean Beaufret and they became friends. Beaufret helped to raise awareness of and interest in Heidegger's thought in France, where a new generation of academics like others around the globe turned to Heidegger despite official disapproval. The global successes of the 1920s were thus repeated in a second wave, which coincided with the revival of Heidegger's publishing and lecturing career. In 1946 he spoke at an occasion commemorating the 20th anniversary of Rilke's death. And in 1947 he published what turned out to be one of his most influential articles, "Brief den Humanismus" (translated as "Letter on Humanism," 1962), an "open letter," which responds to questions that Beaufret had put to Heidegger regarding Jean Paul Sartre's brief but renowned statement of existentialism, L'existentialism est un humanisme (translated as Existentialism and Humanism). True to the genre, Heidegger's anecdotal and meandering meditation nonetheless refuses to accept an alignment of his own thought with either existentialism or humanism, and asserts that no ethics would need to be added on to a philosophy that begins with the question concerning the truth of being.

In December 1949 Heidegger lectured to the Bremen Club. The ban on his teaching was lifted and he resumed correspondence with both Hannah Arendt and Karl Jaspers. In 1950 the first of a series of major publications appeared, Holzwege (wrong Paths), which collected revised versions of some of his most important lectures and seminars on art, technology and language. Throughout the 1950s he continued to publish what were for the most part revised versions of older works, largely related to his teaching. In 1951 he taught a lecture series, Was Heisst Denken? (translated as What is Called thinking, 1968). This was his last course before retiring from the University and the first he had been allowed to teach since 1944, when he was drafted into the Volkssturm. He revisits previous engagements with Parmenides, Plato and Kant as well as Hölderlin and Nietzsche in his pursuit of that which gives thought its topic by withdrawing, a withdrawal that by absenting itself is both more present than everything present and yet infinitely exceeds the actuality of everything actual. He observes that the possibility of thinking is given by what is not yet thought and that what is most thought provoking is the fact that we are not yet thinking.

In 1955-6 he returned as emeritus professor for a series of lectures, which were published in 1957 as Der Satz vom Grund (translated as The Principle of Reason, 1992). These were followed by a series on Hegel's Wissenschaft der Logik ("Science of Logic"). These lecture series are closely linked to a short text, "Der Satz der Identität" ("The Principle of Identity"), which he presented on Faculty Day, 27 June 1957 on the occasion of the 500th anniversary of the University of Freiburg. A significant factor of these texts is that the current milieu is now characterized as "the atomic age" and described in Heidegger's usual grim humor as the "mushrooming" of instrumental rationality. The lecture course of 1955-56 culminates in an address, with the same title, which condenses the careful readings of the thirteen lectures into a sustained thesis outlining the conditions and the destiny of modernity in terms of its paradoxical grounds, indicated by the phrase nihil est sine ratione: "nothing is without reason." Here, as with many of the works of this period, the motif being, always supposedly the central motif, the single thought, is glimpsed not as the fundamental ontological category, but rather as just one term or motif in a kind of loosely organized dissemination of such terms in the history of thinking. Strictly even the word term will no longer adequately capture

the kind of verbal mark or unit of signification that being must be approached as in these texts. Emphasis moves away from the category of being to what he calls its Geschick, its destiny/destination or its sending-forth. What saves this Geschick (a destiny that all philosophical determinations of being share) from the fragmentation and chaotic dissemination it appears to have in its history is what is not determined in it each time—this alone captures its sameness and simplicity. The critique of the metaphysics of presence that always animates Heidegger's works leads him to evoke the constancy of being's Geschick in terms of presencing, lingering and light. Being, of course, means different things for different philosophers in different epochs, to the extent that one epoch can be distinguished from another, in which case these heterogeneous determinations of being might actually be a helpful historiographical guide. What is significant turns out to be the sameness that remains, lies hidden, beneath these determinations: "But it does not run between the epochs like a band linking them; rather the legacy always comes from what is concealed in the Geschick, just as if from one source various streamlets arise that feed a stream that is everywhere and nowhere." One might as well talk of absencing as presencing. The difference, repeating the ontical-ontological difference of Sein und Zeit, gives us a vision of the ontological as that which is nowhere. The being-nowhere animates beings wherever they are (everywhere). The two figures, the sprouts and streamlets, indicate the full Geschick in terms of its absence, its being nowhere. The condensed version of this point in the address is yet more powerful. Taking the phrase that Leibniz coins for the principle of reason, nihil est sine ratione, Heidegger proceeds (as he has done throughout the lecture course) to uncover the hidden grounds of the principle itself—the principle of the principle. The two negatives (nihil, "nothing" and sine, "without") cancel each other out to give a positive, the belonging together of being (est) and reason (ratione).

The principle of this belonging together is identified in terms of a kind of sounding, a Klang, which draws our attention to the musical belonging together of the two terms, being and reason (est and ratione): "Being and Reason now ring in unison. In this Klang being and reason ring out as belonging together in one. The principle of reason now has a different ring and says: ground/reason belongs to being." In the unfolding of this logic of the ring, the repetition of reason in ground (in Der Satz vom Grund) makes ground and reason equivalent but, in order to save their sense in this equivocation, a further ground (and reason) is posited beneath them—Heidegger's being, to which they belong (der Satz/Grund vom Sein). But this belonging has a further ground (the belonging that being is) in the Klang, the ringing that one must hear while seeing. In this sense being as ground/reason (being/ground/reason) has no ground, reason or even being, because it is what grounds, gives reasons to and allows beings to be beings. Being is thus not a ground beneath the various forms of determination that metaphysics variously privileges, like ground and reason. Rather, by occupying a place alongside these other terms, being reveals itself and all such terms to be grounded only in the indeterminate klang of their belonging together as Geschick. Being thus cannot be represented as a being. This means first that being cannot be represented at all. There is nothing here that departs from the pattern that all metaphysical concepts fall into. Their various means of representation inevitably fall short of representation strictly speaking, for their otherworldly, supersensible status must remain infinitely outside the spatial and temporal conditions of representation, outside experience. But it means, second, that the reason why being cannot be represented at all has nothing to do with any atemporal status that might be supposed of it as a metaphysical concept; rather, being cannot be represented because the meaning of "being" is the power that makes representation possible before and beyond any actual representations. It cannot be determined because it is what gives itself to determination. It has no reason itself because it gives reasons. And so the ground of reason is not a ground but a kind of grounding potential, something in the determination that is not given but nonetheless gives itself to possible determinations as Geschick—the sending forth that makes it possible to respond to each epoch as legacy, tradition, inheritance.

In this respect Heidegger's text gains an extra dimension, for the argument takes the exact form of the response, activating the Wiederholung (repetition/retrieval) that was already thematized as early as 1915 before emerging in a more developed form in Sein und Zeit and in the Kantbuch. The legacy or tradition can be read in terms of its Geschick—it's destination (destiny) as sending forth. The concept of the same in Heidegger's text turns out to be a quite precise Wiederholung of a problem that persists in Hegel's Greater Logic. Distinguishing the same from the

empty repetition of identities, Heidegger, like Hegel, is led to a notion of identity that maintains the contradiction of difference within it. What is (est) the same—not itself a being—holds what belong together in a kind of radical difference. What is the same is their radical difference. Hegel's Logic indeed returns rather forcefully in these pages. Heidegger even affirms that text as the place where "relation" [verhältnis] most purely comes to be thought as "what is to be thought." This relation, or holding together, of radical differences is yet to be thought—it is a not yet. This, then, is what Geschick sends forth in the legacy of metaphysical thinking, the un-thought—the thought of the un-, or not- that gives itself to be thought, handing down what is not determined for an event of appropriation, that is, a singular determination.

Heidegger's late thought can be focused on a subtle and elusive distinction. In a lecture from 1962, whose title "Zeit und Sein" recapitulates the missing third division of Sein and Zeit, Heidegger gives his audience "a little hint" on how to listen. "The point is not to listen to a series of propositions," he explains, "but rather to follow the movement of showing." This distinction typifies Heidegger's philosophy. It implies a fundamental difference between propositions and the conditions enacted when a person makes them, and can be put in the following way. A proposition would be a statement that asserts something about some object. For instance, a predicate is denied or affirmed of something or someone ("this table is white"). The proposition is thus the main building block of the logic of assertion. It supports the notion of truth as correspondence, according to which assertions can be tested against established facts, and it forms the ground of predicative logic, which, Heidegger teaches, dominates the theoretical attitude. In order to follow "the movement of showing" one would focus not on the validity of assertions but on their intelligibility, on the performative or enunciative modalities of a discourse and thus on the way a modality of discourse allows certain kinds of being to come into view in certain ways. In other words, one would focus on the constitution of the world's intelligibility. An alternative notion of truth thus emerges, alētheia, which would not be reducible to questions of validity but would rather concern the ways in which given modes of discourse disclose beings. Predicative logic discloses beings in such a way as to distinguish the objects of knowledge from the subject who knows and who can thus make systematic pronouncements on things that are. An analysis of the non-objectifiable grounds of predicative discourse would thus bring to light the ways in which the subject of philosophy and science comes to being and comes to knowledge. The effect is to get beneath the subject of knowledge to the grounds of experience and knowledge itself and thus, for us situated historical beings, to an engagement with the historicity of the ways of being in the modern world.

In the 1960s and 70s Heidegger's work activities became less intense. Most of his time was spent at his home in Freiburg or at the solitary mountain cottage in Todtnauberg. In 1966 he gave an interview with Der Spiegel, which was recorded, on the condition that it would be published only after he had died. Heidegger died peacefully in 1976. His last word was "Thanks." He had made plans for the publication of all his writings in 1972, a huge undertaking which began in 1974 with around 100 volumes projected. The project is expected to go on until the first decades of the 21st century. The Gesamtausgabe is a complete edition "in the last hand," so includes only Heidegger's final revisions, which he prefaced with the motto, "ways not works." Heidegger's publishers have taken his request literally, to publish the works with no scholarly apparatuses whatsoever. The motto has with unintentional irony been reversed. It is more difficult than ever to assess his works as "ways," because readers only have access to the final versions, many of which include revisions made to the earlier manuscripts that are not reflected in the published texts. For this reason the Gesamtausgabe cannot be regarded as the definitive edition for Heidegger scholarship.

The Spiegel interview was published in May 1976 with the title "Nur noch ein Gott ka nuns retten" (translated as "Only a God Can Save Us"), which echoes a line from Hölderlin. In it he responds to questions about his involvement with the Nazis in 1933-4. Those who were looking for stronger expressions of regret than Heidegger had hitherto offered were disappointed. Their disappointment contributed to misleading debates about Heidegger's silence on the matter. The most disturbing aspect of Heidegger's involvement has been less a matter of what he did not say and more a matter of the little he did say. Nazism was, for Heidegger, one more creation of modern

metaphysics, no worse than the hydrogen bomb, mechanized agriculture or automated kitchens. He admitted during the postwar years that he had been wrong in his support of the Nazis but he never acceded to the demand to regard the Holocaust as a more horrific event than others. He once defended his attitude by drawing attention to the victimization of East Germans, who were, he said, victims no less than Jews. Furthermore he tended to minimize the extent of his involvement in Nazi politics, emphasizing instead the covert resistance of his lectures. The key concerns of the debates about Heidegger's politics, which began immediately after the war and which continued throughout the remainder of the 20th century, have been about the connections between Heidegger's philosophy and his involvement with Nazism. Works by Victor Farías and Hugo Ott during the 1980s revealed the actual extent of Heidegger's involvement and ambitions, leaving no doubt that Heidegger's self representations misrepresented his involvement in crucial respects. The question of the connection between his political involvement and his philosophy has been considerably clarified during the final decade of the 20th century, largely owing to painstaking research and patient scholarship.

Heidegger stands out as one of the greatest philosophers of the century. The impact of his writing not only in Europe and America but also in Asia is hardly calculable. He has been described as the hidden master of modern thought and as a man tormented both by philosophy and by his ambiguous relationship to his faith. His impact has been felt not only in philosophy but also in literary theory, psychology, psychiatry, psychoanalysis, theology, and ecology. A growing interest in Continental Philosophy in America has led to a wider understanding and appreciation of his work, highlighting his role in the development of hermeneutics and deconstruction and his influence on theorists like Hans-Georg Gadamer and Jürgen Habermas as well on French thinkers like Jacques Derrida, Michel Foucault, Gilles Deleuze and Pierre Bourdieu.

Heidegger's influence on modern critical and cultural thought can be assessed in two ways. If his philosophy has led to significant developments, especially in the analysis of modernity, historicity and technology, it is also the case that he develops his mode of thinking independently of a number of coterminous thinkers who by radically different routes achieve positions with comparable implications for cultural and critical thought. For instance, Michel Foucault's notion of the enunciative modality of the statement owes as much to Ferdinand de Saussure's distinction between la langue and parole (the language system and the speech event) as it does to Heidegger's notion of alētheia, though his mode of radical revision is reminiscent of Heidegger's Destruction. Here it is no longer possible to distinguish between direct influence and independent thought. The notion of destruction, or deconstruction, implies at once the repetition and the revision of the thought of a prior thinker. The hinge between repetition and revision operates at the modality of enunciation and indicates a potential for singular decisions. If Heidegger never expressed this thought as such, his philosophy nevertheless leads in this direction. The historicity of modern thought implies that beyond its systematic calculations there remains the undecidable. It is only on the basis of the undecidable that decisions can be made. Otherwise we would be following the preprogrammed plan of an auto-telic universe. The startling fact about the 1924 lecture "The Concept of Time" is not just its foreshadowing of difficult and hard to understand concepts that would emerge in the 1927 masterpiece Being and Time: namely Dasein, Care, Being-in-the-World, Historicality, Temporality, Resoluteness, etc. That great work of 1927, which changed continental European philosophy forever, practically had to invent a whole new set of terms within an already flexible German language and its philosophical resources. Yet in the 1924 lecture we get a straight forward presentation by arguably the most significant figure in twentieth century philosophy, which certainly puts him in the canon of the history of Western philosophy with other greats who pondered the mystery of time, namely Heraclitus, Parmenides, Plato, Aristotle, Descartes, Leibniz, Kant, Hegel and Nietzsche. Heidegger for his part says he is not revolutionizing philosophy by providing a revolutionary answer to the age-old question- "What is time?" Rather, his undeniable and breathtaking originality comes from the simple fact that he has utterly changed the very structure of the question by restructuring our relation to it so that it is no longer like any other question- says a question in science - what is an atom? - or a question in the social sciences- 'how do social movements arise and occur?' Rather, we humans become the question in so far as the question of time inhabits our very essence in as much as we inhabit the question. And so pursuing the **question of time separates philosophy from any other branch of human**

inquiry, say the natural and social sciences. By asking about time, we are asking about ourselves, but without either metaphysics (a concept or notion like Spirit or the Absolute) or theology (the revelation of God) as a readily available answer to the question. Hence revolutionary interrogations begin to flow such as 'Are we ourselves time? and "Am I time?' in the 1924 lecture. The great shift that Heidegger's thought catalyzes (at least in the history of Western thought) is the creation of this distinction. The history of metaphysics (Plato to Hegel) separated the human subject from the question and thought of time as an object to be probed, and therefore 'what' becomes the proper subject of the question's object, namely time; whereas in Heidegger, 'who' is substituted for 'what' and so by asking about the very 'who' of time it forces a reorientation in the very way philosophy must proceed with its greatest question (what will become the question of the meaning of being in Being and Time) and a way for this project to ground itself with its own rationale for turning away from the history of metaphysics. This is precisely the future project of 'Dasein's analytic,' or that entity that has to pose the question of Being anew (arising from the destruction of the history of metaphysics and how the latter tried to answer what time is through various great thinkers' efforts). Furthermore, Dasein does so with a 'radical individuation' in which the 'who' of Dasein itself is at root the mystery of time; but the root is one of radical transcendence, i.e. beyond anything ever conceived or intuited. Time therefore has to be the horizon for any understanding of Being, and Dasein is not like any other subject (man, God, angels or animals) asking or revealing what time 'is.' The very Being of Time is what Heidegger's thought captivates precisely by freeing it from any previous relation between beings and time, how beings relate to time and how time has been conceived by beings-- be it gods or men like the tragedies of ancient Greece. The great thing about "The Concept of Time" is that it is arguably the clearest and most concise indicator of where all the early works of Heidegger was tending prior to his massive breakthrough articulation of a problem that had haunted him along: to understand being one must reckon the mystery of time... Heidegger's fundamental analysis of Dasein from Being and Time points to temporality as the primordial meaning of Dasein's being. Dasein is essentially temporal. Its temporal character is derived from the tripartite ontological structure: existence, thrownness, and fallenness by which Dasein's being is described. Existence means that Dasein is potentiality-for-being (Seinkönnen); it projects its being upon various possibilities. Existence represents thus the phenomenon of the future. Then, as thrownness, Dasein always finds itself already in a certain spiritual and material, historically conditioned environment; in short, in the world, in which the space of possibilities is always somehow limited. This represents the phenomenon of the past as having-been. Finally, as fallenness, Dasein exists in the midst of beings which are both Dasein and not Dasein. The encounter with those beings, "being-alongside" or "being-with" them, is made possible for Dasein by the presence of those beings within-the-world. This represents the primordial phenomenon of the present. Accordingly, Dasein is not temporal for the mere reason that it exists "in time," but because its very being is rooted in temporality: the original unity of the future, the past and the present. Temporality cannot be identified with ordinary clock time – with simply being at one point in time, at one "Now" after another—which for Heidegger is a derivative phenomenon. Neither does Dasein's temporality have the merely quantitative, homogeneous character of the concept of time found in natural science. It is the phenomenon of original time, of the time which "temporalizes" itself in the course of Dasein's existence. It is a movement through a world as a space of possibilities. The "going back" to the possibilities that have been (the past) in the moment of thrownness, and their projection in the resolute movement "coming towards" (the future) in the moment of existence, which both take place in "being with" others (the present) in the moment of fallenness, provide for the original unity of the future, the past, and the present which constitutes authentic temporality.

As authentically temporal, Dasein as potentiality-for-being comes towards itself in its possibilities of being by going back to what has been; it always comes towards itself from out of a possibility of itself. Hence, it comports itself towards the future by always coming back to its past; the past which is not merely past but still around as having-been. But in this "going back" to what it has been which is constitutive together with "coming towards" and "being with" for the unity of Dasein's temporality, Dasein hands down to itself its own historical "heritage," namely, the possibilities of being that have come down to it. As authentically temporal, Dasein is thus authentically historical. The repetition of the possibilities of existence, of that which has been, is for Heidegger constitutive for the phenomenon of original history which is rooted in temporality.

It should be no surprise that Carl von Clausewitz first heard the term of "Center of Gravity" at a physics conference in 19th Century Germany (pg 110) .The physics at the time, Newtonian Physics, did well to explain most observable phenomena then in existence. Newton's take on gravity was tied to a mass's body: that physical point within an object at which the gravitational attraction to other bodies is exactly equal in every direction. **Clausewitz,** it is postulated, heard the concept and immediately imagined a wrestler losing his balance due to his opponent finding a center of gravity and exerting a force against it. He further imagined this concept being used in the military context: as a general would find that point at which to concentrate effects so that the opposing army would likewise lose balance and "fall". Of course, Newtonian physics suffered from a few issues, the main ones being that it is only accurate as long as one is dealing with sufficiently large objects and relatively low speeds. As one approaches the speed of light, for instance, Newtonian, or "classical" physics does not explain observable phenomena very well at all. The military, of course, taking its cues from Clausewitz's metaphor, have a very distinct take on the military "Center of Gravity**". (Replacing the Center of Gravity Concept with a Quantum Mechanics Military Theory and Its Implications by Grant Martin)**

Scheme of division of the paper is given in the following:

SECTION ONE

Worm Holes And Time Travel: Saltus In Demonstrando-Salva Veritate Model

SECTION TWO

Negatively Curved Space-Time And Flat Space-Time: Non In Legendo Sed In Intelligendo Legis Consistent Model

SECTION THREE

"Roman Ring": Nemo Mortalium Omnibus Horis Sapit-Nemo Saltat Sobrius Model

SECTION FOUR

String Landscape: Naturam Expellas Furca, Tamen Usque Recurret Model

SECTION FIVE

Philosophy As The Invention Of Concepts (Deluze And Guattari): Saltus In Demonstrando-Salva Veritate Model

SECTION SIX

Creation Of Concepts: Acta Deos Nunquam Mortalia Fallunt Model

SECTION SEVEN

Mathematicised Ontology (Badiou And Deleuze): Ab Aeterno Model

SECTION EIGHT

Subject And Event (Badiou And Deleuze): Sed Ipse Spiritus Postulat Pro Nobis, Gemitibus Inenarrabilibus Model

SECTION NINE

Being And Event: Nosce Te Ipsum - Omnia Mutantur Nos Et Mutamur In Illis Model

SECTION TEN

A Thousand Plateaus: Omnia Causa Fiunt - Perfer Et Obdura-Dolor Hic Tibi Proderit Olim Model

SECTION ELEVEN

Truth Of An Event: Acta Deos Nunquam Mortalia Fallunt Model

SECTION TWELVE

Identity And Contradiction: Actus Non Facit Reum Nisi Mens Sit Rea Model

SECTION THIRTEEN

Pure Difference: Dictum Sapienti Sat Est Model

SECTION FOURTEEN

"Minimal Difference" - The Purely Parallax Gap: Quod Me Nutrit Me Destruit. Qui Tacet Consentire Model

SECTION FIFTEEN

Dark Matter: Nemo Dat Quod Non Habet - Non Est Ad Astra Mollis E Terris Via

SECTION SIXTEEN

Anapole Dark Matter: Nosce Te Ipsum - Omnia Mutantur Nos Et Mutamur In Illis Model

SECTION SEVENTEEN

Majorana Fermions And Anapole: Quod Me Nutrit Me Destruit. Qui Tacet Consentire Model

SECTION EIGHTEEN

Zeno’s Paradox And High-Fidelity Quantum Computation: Non Causa Pro Causa-Non Bis In Idem Model

SECTION TWENTY

Numerous Landscapes Of Eternal Inflation And Lambda: Sed Ipse Spiritus Postulat Pro Nobis, Gemitibus Inenarrabilibus Model

SECTION TWENTY ONE

Regime Of Eternal Inflation: Non Obstante Veredicto Model

SECTION TWENTY TWO

Quantum Dynamics And Casimir Energy: Eheu Fugaces Labuntur Anni

SECTION TWENTY THREE

Stylobate Problem In Many-Worlds Theories: Saltus In Demonstrando-Salva Veritate Model

SECTION TWENTY FOUR

Coherent Quantum Control: Non Est Ad Astra Mollis E Terris Via Model

SECTION TWENTY FIVE

Quantum Formalism And Holographic Renormalization Group: Non Causa Pro Causa-Non Bis In Idem Model

SECTION TWENTY SIX

Holographic Quantum Fields: Naturam Expellas Furca, Tamen Usque Recurret

SECTION TWENTY SEVEN

Artificial Intelligence And Uncertainty Modeling: Non In Legendo Sed In Intelligendo Legis Consistent

SECTION TWENTY SEVEN

Quantum Weirdness: Mutantur Omnia Nos Et Mutamur In Illis

SECTION TWENTY NINE

God Does Not Play Dice: Quod Me Nutrit Me Destruit. Qui Tacet Consentire

SECTION THIRTY

Nagarjuna's Tetra Lemma: Legum Servi Sumus Ut Liberi Esse Possimus - Mutantur Omnia Nos Et Mutamur In Illis

SECTION THIRTY ONE

Cyclotomic Quinary: Fiat Lux Model

SECTION THIRTY TWO

Pan-Hylism And Pan-Psychism: A Uncomfortable Menage A Trios

SECTION THIRTY THREE

Consciousness And Quantum Computing: Veritas Vos Liberabit Model

SECTION THIRTY FOUR

Pastry Chef And Contemporary Computer Science: Sacrosuchus Imperator-Bonvivavt Model

SECTION THIRTY FIVE

Organizational Invariance And Conscious Experience Aeì Koloiòs Parà Koloiôi Hizánei Model

SECTION THIRTY SIX

Holographic And Anthropic Multiverse: Aei Ho Theos Geōmetreî

SECTION THIRTY SEVEN

Hamiltonian Engineering: Lucus A Non Lucendo-Lux Et Veritas--Lux Hominum Vita

SECTION THIRTY EIGHT

Quantum Control Systems: Labor Omnia Vincit-Leges Humanae Nascuntur, Vivunt, Et Moriuntur

SECTION THIRTY NINE

Hamiltonian Quantum Systems: Nanos Gigantum Humeris Insidentes

SECTION FORTY

Atomic Engineering: Cato Esse Quam Videri Bonus Male Bat

SECTION FORTY ONE

Nature Nanotechnology: Perfer Et Obdura; Dolor Hic Tibi Proderit Olim

SECTION FORTY TWO

A Single-Atom Transistor: Exitus Acta Probat Errare Humanum Est

SECTION FORTY TWO A

Kane Computer: Una Hirundo Non Facit Ver- Usus Est Magister Optimus

SECTION FORTY THREE

Quantum Control: Saltus In Demonstrando Model

SECTION FORTY FOUR

Quantum Transport: A Gesamtbedeutung Model

SECTION FORTY FIVE

Electric Dipole Control: Tanquam Ex Ungue Leonem-Technica Impendi Nationi

SECTION FORTY SIX

Quantum Dots: A Poindexter Poinsettia

SECTION FORTY SEVEN

Spintronics: Gotterdammerung Model

SECTION FORTY EIGHT

Griffith Program: Sapientia Et Doctrina Model

SECTION FORTY NINE

Spintronics And Quantum Computation: Natura Non Contristatur-Natura Non Facit Saltum Ita Nec Lex

SECTION FIFTY

Holographic Quantum Computation: Naturam Expellas Furca, Tamen Usque Recurret

SECTION FIFTY ONE

Quantum Information Theory In The Field Of Quantum Gravity: A Hauptvermutung

SECTION FIFTY TWO

Interference Fringes Of Electrons In Backscattering: A Posse Ad Esse

SECTION FIFTY THREE

Non-Abelian Anyons: Ad Augusta Per Angusta

SECTION FIFTY FOUR

Monolayer Doping (MLD): Victoria Concordia Crescit

SECTION FIFTY FIVE

Quantum Geometry And Black Hole Entropy: Aeì Ho Theòs Ho Mégas Geōmetreî Tò Sýmpan

SECTION FIFTY SIX

Laws Of Black Hole Mechanics: Deus Ex Machina

SECTION FIFTY SEVEN

Correlation Space And Extremal Entanglement: Labor Omnia Vincit-Leges Humanae Nascuntur, Vivunt, Et Moriuntur

SECTION FIFTY EIGHT

Extreme Spin Squeezing: Crede Quod Habes, Et Habes

SECTION FIFTY NINE

Quantum Feedback Control: Multum In Parvo

SECTION SIXTY

Measurement And Control For Quantum Computing: Affirmanti Non Neganti Incumbit Probatio

SECTION SIXTY ONE

Quantum Control Engineering: Omnia Causa Fiunt

SECTION SIXTY TWO

General Relativity And Quantum Cosmology Computational Complexity: Generalia Specialibus Non Derogant

SECTION SIXTY THREE

Quantum Transport: Hannibal is at the Gate

SECTION SIXTY FOUR

Quantum Topological Computation: Multum In Parvo: Much In Little

SECTION SIXTY FIVE

LQG And Quantum Geometry: Dona Clandestina Sunt Semper Suspiciosa Clandestine Gifts And Sybaritic, Luxuriant Trade Offs

SECTION SIXTY SIX

Light Harvesting And Quantum Computation: Noscitur A Sociis: Meaning Of A Word Is From The Context

SECTION SIXTY SEVEN

Quantum Computation In Neurobiological Framework: Nosce Te Ipsum: Know Thyself

SECTION SIXTY EIGHT

Cognitive Informatics: Mutantur Omnia Nos Et Mutamur In Illis :All Things Change, And We Change With Them

SECTION SIXTY NINE

Mahavairocana (Dainichi-Nyorai 大日如来) Mandala: Constituents of Dependent Arising

SECTION SEVENTY

Vajrasattva (Kongō-Satta 金剛薩埵) Mandala: Adamantine Being

SECTION SEVENTY ONE

Nagarjuna (Ryūju-Bosatsu 龍樹菩薩) Mandala: Proof of Convention

SECTION SEVENTY TWO

Nagabodhi (Ryūchi-Bosatsu 龍智菩薩) Mandala: Hymn to the Absolute Reality

SECTION SEVENTY THREE

Exotic matter: Rocky Razzmatazz

SECTION SEVENTY FOUR

Correlation Phenomenon: Omnia Causa Fiunt: Everything Happens For A Reason

SECTION SEVENTY FIVE

Quantum Phase transition: minima maxima sunt: The smallest things are most important

SECTION SEVENTY SIX

Quantum Floating Point: Autumn Sonata

SECTION SEVENTY SEVEN

Decoherence Engineering: Deux Ex machina

SECTION SEVENTY EIGHT

Perception: Nosce Te Ipsum: Know Thyself

SECTION SEVENTY NINE

Motion Inducers: Isotropy-Antistrophe

SECTION EIGHTY

Change Of Mind: Rocky Razzmatazz And Autumn Sonata

SECTION EIGHTY ONE

Change Of Mind: Rocky Razzmatazz And Autumn Sonata

SECTION EIGHTY TWO

Living On The Edge: Abyssus Abyssum Invocat: Deep Calls Unto Deep

SECTION EIGHTY THREE

Dasein: Cygnus Inter Anates: Swan Among Ducks

SECTION EIGHTY FOUR

Pure Phenomenology: Adversus Solem Ne Loquitor: Donot Speak Against The Sun

SECTION EIGHTY FIVE

Being And Time: Agere Sequitur Esse: Action Follows Being

SECTION EIGHTY SIX

"Ontologico-Existential Concept Of Worldhood": Welträtsel: Nature Of The Universe And Meaning Of Life

SECTION EIGHTY SEVEN

Cartesianism: Adequatio Intellectus Nostri Cum Re: Conformity Of Minds To The Facts

SECTION EIGHTY EIGHT

Spatio Temporal Actualization: Wertfreiheit: Value Neutrality

SECTION EIGHTY NINE

Objective, Intersubjective And Subjective Dimensions: Altagsgeschichte: Micro History

SECTION NINETY

Living Lives At The Limits Of Itself…: Fingerspitzengefuhl: Intuitive

SECTION NINETY ONE

Worm Holes And Time Travel: Freiwirtschaft: Natural Order

SECTION ONE

Worm Holes And Time Travel: Saltus In Demonstrando-Salva Veritate Model

INTRODUCTION—VARIABLES USED

Wormholes and Time Travel Shortcut through Spacetime - If Traversable Wormholes Exist, They Could Allow Time Travel

(1) A short-cut through space time that exists because of the general relativity. Traveling through a wormhole is faster than traveling at the speed of light, **which (eb) allows** the possibility of time travel.
(2) In physics and fiction, a wormhole is a hypothetical topological feature of spacetime that would be, fundamentally, a "shortcut"(e: distance reducing aisle, artery, avenue, beat, beaten path, boulevard, byway, crosscut, direction, drag,footpath, groove , highway, lane, line, pass, passage, pathway, procedure, rail, road,roadway, route, rut, shortcut , str eet, stroll, terrace, thoroughfare, track, trail, walk,walkway) through spacetime.
(3) A proposed time-travel machine **using a** traversable wormhole would (hypothetically) work in the following way: One end of the wormhole **is (eb)accelerated** to some significant fraction of the speed of light, perhaps with some advanced propulsion system, and then brought back to the point of origin.
(4) Alternatively, another way is to take one entrance of the wormhole and **move it to (eb) within** the

gravitational field of an object that has higher gravity than the other entrance, and then return it to a position near the other entrance.

(5) For both of these methods, time dilation **causes (eb)** the end of the wormhole that has been moved to have aged less than the stationary end, as seen by an external observer;

(6) However, time **connects** differently through the wormhole than outside it, so that synchronized clocks at either end of the wormhole will always remain synchronized as seen by an observer passing through the wormhole, no matter how the two ends move around.

(7) This means that an observer **entering (eb)** the accelerated end would exit the stationary end when the stationary end was the same age that the accelerated end had been at the moment before entry

(8) for example, if prior to entering the wormhole the observer noted that a clock at the accelerated end read a date of 2007 while a clock at the stationary end read 2012, then the observer **would (eb)exit the** stationary end when its clock also read 2007, a trip backwards in time as seen by other observers outside.

NOTATION

Module One

A short-cut through space time that exists because of the general relativity. Traveling through a wormhole is faster than traveling at the speed of light, **which (eb) allows** the possibility of time travel.

G_{13} : Category one of **short**-cut through space time that exists because of the general relativity. Traveling through a wormhole is faster than traveling at the speed of light

G_{14} : Category two of short-cut through space time that exists because of the general relativity. Traveling through a wormhole is faster than traveling at the speed of light

G_{15} : Category three of short-cut through space time that exists because of the general relativity. Traveling through a wormhole is faster than traveling at the speed of light

T_{13} : Category one of possibility of time travel

T_{14} : Category two of possibility of time travel

T_{15} : Category three of possibility of time travel

Module Two

In physics a wormhole is a hypothetical topological feature of spacetime that would be, fundamentally, a "shortcut"(e: distance reducing aisle, artery, avenue, beat, beaten path, boulevard, byway, crosscut, direction, drag,footpath, groove, highway, lane, line, pass, passage, pathway, procedure, rail, road,roadway, route, rut, shortcut , street, stroll, terrace, thoroughfare, track, trail, walk,walkway) through spacetime

G_{16} : Category one of through spacetime

G_{17} : Category two of through spacetime

G_{18} : Category three of through spacetime

T_{16} : Category one of wormhole is a hypothetical topological feature of spacetime

T_{17} : Category two of wormhole is a hypothetical topological feature of spacetime

T_{18} : Category three of wormhole is a hypothetical topological feature of spacetime

Module three

A proposed time-travel machine **using a** traversable wormhole would (hypothetically) work with one end of the wormhole **is (eb)accelerated** to some significant fraction of the speed of light, perhaps with some advanced propulsion system, and then brought back to the point of origin

G_{20} : Category one of proposed time-travel machine **using a** traversable wormhole would (hypothetically) work with one end of the wormhole

G_{21} : Category two of proposed time-travel machine **using a** traversable wormhole would (hypothetically) work with one end of the wormhole

G_{22} : Category three of proposed time-travel machine **using a** traversable wormhole would (hypothetically) work with one end of the wormhole

T_{20} : Category one of significant fraction of the speed of light, perhaps with some advanced propulsion system, and then brought back to the point of origin

T_{21} : Category two of significant fraction of the speed of light, perhaps with some advanced propulsion system, and then brought back to the point of origin

T_{22} : Category three of significant fraction of the speed of light, perhaps with some advanced propulsion system, and then brought back to the point of origin

Module four

Alternatively, another way is to take one entrance of the wormhole and **move it to (eb) within** the gravitational field of an object that has higher gravity than the other entrance, and then return it to a position near the other entrance.

G_{24} : Category one of one entrance of the wormhole

G_{25} : Category two of one entrance of the wormhole

G_{26} : Category three of one entrance of the wormhole

T_{24} : Category one of **move it to (eb) within** the gravitational field of an object that has higher gravity than the other entrance, and then return it to a position near the other entrance.

T_{25} : Category two of **move it to (eb) within** the gravitational field of an object that has higher gravity than the other entrance, and then return it to a position near the other entrance.

T_{26} : Category three of **move it to (eb) within** the gravitational field of an object that has higher gravity than the other entrance, and then return it to a position near the other entrance.

Module five

For both of these methods, time dilation **causes (eb)** the end of the wormhole that has been moved to have aged less than the stationary end, as seen by an external observer

G_{28} : Category one of time dilation

G_{29} : Category two of time dilation

G_{30} : Category three of time dilation

T_{28} : Category one of end of the wormhole that has been moved to have aged less than the stationary end, as seen by an external observer

T_{29} : Category two of end of the wormhole that has been moved to have aged less than the stationary end, as seen by an external observer

T_{30} : Category three of end of the wormhole that has been moved to have aged less than the stationary end, as seen by an external observer

Module six

However, time **connects** differently through the wormhole than outside it, so that synchronized clocks at either end of the wormhole will always remain synchronized as seen by an observer passing through the wormhole, no matter how the two ends move around.

G_{32} : Category one of time

G_{33} : Category two of time

G_{34} : Category three of time

T_{32} : Category one of **connects** differently through the wormhole than outside it, so that synchronized clocks at either end of the wormhole will always remain synchronized as seen by an observer passing through the wormhole, no matter how the two ends move around

T_{33} : Category two of **connects** differently through the wormhole than outside it, so that synchronized clocks at either end of the wormhole will always remain synchronized as seen by an observer passing through the wormhole, no matter how the two ends move around

T_{34} : Category three of **connects** differently through the wormhole than outside it, so that synchronized clocks at either end of the wormhole will always remain synchronized as seen by an observer passing through the wormhole, no matter how the two ends move around

Module seven

G_{36} : Category one of **connection** through the wormhole than outside it, so that synchronized clocks at either end of the wormhole will always remain synchronized as seen by an observer passing through the wormhole, no matter how the two ends move around

G_{37} : Category two of **connection** through the wormhole than outside it, so that synchronized clocks at either end of the wormhole will always remain synchronized as seen by an observer passing through the wormhole, no matter how the two ends move around

G_{38} : Category three of **connection** through the wormhole than outside it, so that synchronized clocks at either end of the wormhole will always remain synchronized as seen by an observer passing through the wormhole, no matter how the two ends move around

T_{36} : Category one of time

T_{37} : Category two of time

T_{38} : Category three of time

Module eight

This means that an observer **entering (eb)** the accelerated end would exit the stationary end when the stationary end was the same age that the accelerated end had been at the moment before entry

G_{40} : Category one of time **connects** differently through the wormhole than outside it, so that synchronized clocks at either end of the wormhole will always remain synchronized as seen by an observer passing through the wormhole, no matter how the two ends move around

G_{41} : Category two of time **connects** differently through the wormhole than outside it, so that synchronized clocks at either end of the wormhole will always remain synchronized as seen by an observer passing through the wormhole, no matter how the two ends move around

G_{42} : Category three of time **connects** differently through the wormhole than outside it, so that synchronized clocks at either end of the wormhole will always remain synchronized as seen by an observer passing through the wormhole, no matter how the two ends move around

T_{40} : Category one of observers **entering (eb)** the accelerated end would exit the stationary end when the stationary end was the same age that the accelerated end had been at the moment before entry

T_{41} : Category two of observers **entering (eb)** the accelerated end would exit the stationary end when the stationary end was the same age that the accelerated end had been at the moment before entry

T_{42} : Category three of observers **entering (eb)** the accelerated end would exit the stationary end when the stationary end was the same age that the accelerated end had been at the moment before entry

Module Nine

prior to entering the wormhole the observer noted that a clock at the accelerated end read a date of 2007 while a clock at the stationary end read 2012, then the observer **would (eb)exit the** stationary end when its clock also read 2007, a trip backwards in time as seen by other observers outside. There are many such systems

G_{44} : Category one of prior to entering the wormhole the observer noted that a clock at the accelerated end read a date of 2007 while a clock at the stationary end read 2012, then the observer

G_{45} : Category two of prior to entering the wormhole the observer noted that a clock at the accelerated end read a date of 2007 while a clock at the stationary end read 2012, then the observer

G_{46} : Category three of prior to entering the wormhole the observer noted that a clock at the accelerated end read a date of 2007 while a clock at the stationary end read 2012, then the observer

T_{44} : Category one of stationary end when its clock also read 2007, a trip backwards in time as seen by other observers outside. There are many such systems. See classification remarks in the introduction.

T_{45} : Category two of stationary end when its clock also read 2007, a trip backwards in time as seen by other

observers outside. There are many such systems

T_{46} : Category three of stationary end when its clock also read 2007, a trip backwards in time as seen by other observers outside. There are many such systems

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}),t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$
$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$
$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$
$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$
$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$
$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**
$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}), t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}), t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition

coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)}\ \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\ \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\ \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for

category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad 78$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

are fourth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{16}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{17}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{18}'')^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{20}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{21}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{22}'')^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{46}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{45}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{44}'')^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{28}')^{(5)}\boxed{+(a_{28}'')^{(5)}(T_{29},t)} & \boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)} & \boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a_{29}')^{(5)}\boxed{+(a_{29}'')^{(5)}(T_{29},t)} & \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)} & \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{30}')^{(5)}\boxed{+(a_{30}'')^{(5)}(T_{29},t)} & \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)} & \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad 81$$

Where $\boxed{+(a_{28}'')^{(5)}(T_{29},t)}, \boxed{+(a_{29}'')^{(5)}(T_{29},t)}, \boxed{+(a_{30}'')^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3
And $\boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3
$\boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3
$\boxed{+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3
$\boxed{+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3
$\boxed{+(a_{40}'')^{(8,8\,,8,8,8,8)}(T_{41},t)}, \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3
$\boxed{+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)} & +(a_{36}'')^{(7)}(T_{37},t) & +(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)} & +(a_{37}'')^{(7)}(T_{37},t) & +(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)} & +(a_{38}'')^{(7)}(T_{37},t) & +(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a_{36}'')^{(7)}(T_{37},t)$, $(a_{37}'')^{(7)}(T_{37},t)$, $(a_{38}'')^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)} \boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} \boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)} G_{44}$$
$$- \left[\begin{array}{ccc} (a'_{45})^{(9)} \boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)} G_{45}$$
$$- \left[\begin{array}{ccc} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)} T_{45} -$$
$$\left[\begin{array}{ccc} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{13}$$
$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ \ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(A) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(B) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(C) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(D)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(E) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(F) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{16})^{(2)}$, $T_i^0 \leq (\hat{Q}_{16})^{(2)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^{0} + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^{0} + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^{0} + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{36})^{(7)}$, $T_i^0 \leq (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(a) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$sup_i\{max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}ds_{(16)} +$

$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)};\ (G_{23})^{(2)}, (T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \qquad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \qquad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \qquad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \qquad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq\frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \qquad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \qquad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq-(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq-(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq-(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq-(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(a) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)} G_{14}^{*}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)} G_{14}^{*}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)} T_{14}^{*}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)} T_{14}^{*}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)} G_{17}^{*}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)} G_{17}^{*}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)} T_{17}^{*}}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)} T_{17}^{*}}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)} G_{21}^{*}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)} G_{21}^{*}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)} T_{21}^{*}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)} T_{21}^{*}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

(586 A)

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWO

Negatively Curved Space-Time And Flat Space-Time : Non In Legendo Sed In

Intelligendo Legis Consistent Model

INTRODUCTION—VARIABLES USED

Science News:

(1) A new paper released by Professor Skenderis and Dr Marco Caldarelli from the University of Southampton, Dr Joan Camps from the University of Cambridge and Dr Blaise Goutéraux from the Nordic Institute for Theoretical Physics, Sweden published in the Rapid Communication section of Physical Review D, makes connections between negatively curved space-time **and (e&eb)** flat space-time.
(2) Space-time is usually understood **to describe** space existing in three dimensions, with time playing the role of a fourth dimension and all four coming together to form a continuum, or a state in which the four elements can't be distinguished from each other.
(3) Flat space-time and negative space-time **describe an** environment in which the Universe is non-compact, with space extending infinitely, forever in time, in any direction.
(4) The gravitational forces, such as the ones produced by a star, **are best described** by flat-space time.
(5) Negatively curved space-time **describes a** Universe filled with negative vacuum energy. The mathematics of holography is best understood for negatively curved space-times.
(6) Professor Skenderis has developed a mathematic model which finds striking similarities between flat space-time **and** negatively curved space-time, with the latter however formulated in a negative number of dimensions, beyond our realm of physical perception.
(7) He comments: "According to holography, at a fundamental level the universe has one less dimension than we perceive in everyday life **and is (e&eb) governed** by laws similar to electromagnetism.
(8) The idea is similar to that of ordinary holograms where a three-dimensional image **is (e) encoded in** a two-dimensional surface, such as in the hologram on a credit card, but now it is the entire Universe that is encoded in such a fashion.

"Our research is ongoing, and we hope to find more connections between flat space-time, negatively curved space-time and holography. Traditional theories about how the Universe operates go some way individually to describing its very nature, but each fall short in different areas. It is our ultimate goal to find a new combined understanding of the Universe, which works across the board."

(9) AdS/Ricci-flat correspondence **and t**he Gregory-Laflamme instability specifically explains what is known as the Gregory Laflamme instability
(10) AdS/Ricci-flat correspondence **and t**he Gregory-Laflamme instability specifically explains what is known as the Gregory Laflamme instability and according to which certain types of black hole **break(e&eb) up into** smaller black holes when disturbed – rather like a thin stream of water breaking into little droplets when you touch it with your finger.
(11) This black hole phenomenon has previously been **shown (eb) to exist** through computer simulations and this work provides a deeper theoretical explanation.

NOTATION

Module One

A new paper released by Professor Skenderis and Dr Marco Caldarelli from the University of Southampton, Dr Joan Camps from the University of Cambridge and Dr Blaise Goutéraux from the Nordic Institute for Theoretical Physics, Sweden published in the Rapid Communication section of Physical Review D, makes

connections between negatively curved space-time **and (e&eb)** flat space-time

G_{13} : Category one of negatively curved space-time

G_{14} : Category two of negatively curved space-time

G_{15} : Category three of negatively curved space-time

T_{13} : Category one of flat space-time

T_{14} : Category two of flat space-time

T_{15} : Category three of flat space-time

Module Two(Case two)

G_{16} : Category one of flat space-time

G_{17} : Category two of flat space-time

G_{18} : Category three of flat space-time

T_{16} : Category one of negatively curved space-time

T_{17} : Category two of negatively curved space-time

T_{18} : Category three of negatively curved space-time

Module three

Space-time is usually understood **to describe** space existing in three dimensions, with time playing the role of a fourth dimension and all four coming together to form a continuum, or a state in which the four elements can't be distinguished from each other.

G_{20} : Category one of Space-time

G_{21} : Category two of Space-time

G_{22} : Category three of Space-time

T_{20} : Category one of space existing in three dimensions, with time playing the role of a fourth dimension and all four coming together to form a continuum, or a state in which the four elements can't be distinguished from each other.

T_{21} : Category two of space existing in three dimensions, with time playing the role of a fourth dimension and all four coming together to form a continuum, or a state in which the four elements can't be distinguished from each other.

T_{22} : Category three of space existing in three dimensions, with time playing the role of a fourth dimension and all four coming together to form a continuum, or a state in which the four elements can't be distinguished from each other.

Module four

Flat space-time and negative space-time **describe an** environment in which the Universe is non-compact, with space extending infinitely, forever in time, in any direction

G_{24} : Category one of flat space-time and negative space-time

G_{25} : Category two of flat space-time and negative space-time

G_{26} : Category three of flat space-time and negative space-time

T_{24} : Category one of environment in which the Universe is non-compact, with space extending infinitely, forever in time, in any direction

T_{25} : Category two of environment in which the Universe is non-compact, with space extending infinitely, forever in time, in any direction

T_{26} : Category three of environment in which the Universe is non-compact, with space extending infinitely, forever in time, in any direction

Module five

The gravitational forces, such as the ones produced by a star, **are best described** by flat-space time.

G_{28} : Category one of flat-space time

G_{29} : Category two of flat-space time

G_{30} : Category three of flat-space time

T_{28} : Category one of gravitational forces, such as the ones produced by a star

T_{29} : Category two of gravitational forces, such as the ones produced by a star

T_{30} : Category three of gravitational forces, such as the ones produced by a star

Module six

Professor Skenderis has developed a mathematic model which finds striking similarities between flat space-time **and** negatively curved space-time, with the latter however formulated in a negative number of dimensions, beyond our realm of physical perception

The Geometry of the Universe (Blog on Wikipedia)

The most profound insight of General Relativity was the conclusion that the effect of gravitation could be reduced to a statement about the geometry of spacetime. In particular, Einstein showed that in General Relativity mass caused space to curve, and objects travelling in that curved space have their paths deflected, exactly as if a force had acted on them.

Curvature of Space in Two Dimensions

The idea of a curved surface is not an unfamiliar one since we live on the surface of a sphere. More generally, mathematicians distinguish 3 qualitatively different classes of curvature, as illustrated in the following image (Source):

These are examples of surfaces that have two dimensions. For example, the left surface can be described by a coordinate system having two variables (x and y, say); likewise, the other two surfaces are each described

by two independent coordinates. The flat surface at the left is said to have zero curvature, the spherical surface is said to have positive curvature, and the saddle-shaped surface is said to have negative curvature.

Curvature of 4-Dimensional Spacetime

The preceding is not too difficult to visualize, but General Relativity asserts that space itself (not just an object in space) can be curved, and furthermore, the space of General Relativity has 3 space-like dimensions and one time dimension, not just two as in our example above. This IS difficult to visualize! Nevertheless, it can be described mathematically by the same methods that mathematicians use to describe the 2-dimensional surfaces that we can visualize easily.

The Large-Scale Geometry of the Universe

Since space itself is curved, there are three general possibilities for the geometry of the Universe. Each of these possibilities is tied intimately to the amount of mass (and thus to the total strength of gravitation) in the Universe, and each implies a different past and future for the Universe:

If space has negative curvature, there is insufficient mass to cause the expansion of the Universe to stop. The Universe in that case has no bounds, and will expand forever. This is termed an open universe.

If space has no curvature (it is flat), there is exactly enough mass to cause the expansion to stop, but only after an infinite amount of time. Thus, the Universe has no bounds in that case and will also expand forever, but with the rate of expansion gradually approaching zero after an infinite amount of time. This is termed a flat universe or a Euclidian universe (because the usual geometry of non-curved surfaces that we learn in high school is called Euclidian geometry).

If space has positive curvature, there is more than enough mass to stop the present expansion of the Universe. The Universe in this case is not infinite, but it has no end (just as the area on the surface of a sphere is not infinite but there is no point on the sphere that could be called the "end"). The expansion will eventually stop and turn into a contraction. Thus, at some point in the future the galaxies will stop receding from each other and begin approaching each other as the Universe collapse on itself. This is called a closed universe.

Which of these scenarios is correct is still unknown because we have been unable to determine exactly how much mass is in the Universe.

Is the Universe Open, Flat, or Closed?

The Density Parameter of the Universe

Source Value

Baryons (BB nucleosynthesis) (0.013 +/- 0.005) h-2

Stars in Galaxies 0.004

Intergalactic Stars <0.04

Rich Clusters 0.01

Dynamics (r < 10 h-1 Mpc) ~0.05 - 0.2

Dynamics (r > 30 h-1 Mpc) ~0.05 - 1

Source: P. J. E. Peebles, Principles of Physical Cosmology

The geometry of the Universe is often expressed in terms of the density parameter, which is defined to the the ratio of the actual density of the Universe to the critical density that would just be required to cause the expansion to stop. Thus, if the Universe is flat (contains just the amount of mass to close it) the density parameter is exactly 1, if the Universe is open with negative curvature the density parameter lies between 0 and 1, and if the Universe is closed with positive curvature the density parameter is greater than 1.

The density parameter determined from various methods is summarized in the adjacent table. In this table, BB nucleosynthesis refers to constraints coming from the synthesis of the light elements in the big bang, +/- denotes an experimental uncertainty in a quantity, and the parameter h lies in the range 0.5 to 0.85 and measures the uncertainty in the value of the Hubble parameter.

Although most of these methods (which we will not discuss in detail) yield values of the density parameter far below the critical value of 1, we must remember that they have likely not detected all matter in the Universe yet. The current theoretical prejudice (because it is predicted by the theory of cosmic inflation) is that the Universe is flat, with exactly the amount of mass required to stop the expansion (the corresponding average critical density that would just stop this is called the closure density), but this is not yet confirmed. Therefore, the value of the density parameter and thus the ultimate fate of the Universe remains one of the major unsolved problems in modern cosmology.

G_{32} : Category one of flat space-time

G_{33} : Category two of flat space-time

G_{34} : Category three of flat space-time

T_{32} : Category one of negatively curved space-time, with the latter however formulated in a negative number of dimensions, beyond our realm of physical perception

T_{33} : Category two of negatively curved space-time, with the latter however formulated in a negative number of dimensions, beyond our realm of physical perception

T_{34} : Category three of negatively curved space-time, with the latter however formulated in a negative number of dimensions, beyond our realm of physical perception

Module seven

He comments: "According to holography, at a fundamental level the universe has one less dimension than we perceive in everyday life **and is (e&eb) governed** by laws similar to electromagnetism.

G_{36} : Category one of universe has one less dimension than we perceive in everyday life according to holography. Characterstics of the systems that are under consideration are taken in to account for classification. It is to be understood that there are perceived perceivable and unobserved and expanding universes and multiverses too. The prediction model gives the expression for all these counts.

G_{37} : Category two of universe has one less dimension than we perceive in everyday life according to holography. Characterstics of the systems that are under consideration are taken in to account for classification. It is to be understood that there are perceived perceivable and unobserved and expanding

universes and multiverses too. The prediction model gives the expression for all these counts

G_{38} : Category three of universe has one less dimension than we perceive in everyday life according to holography. Characterstics of the systems that are under consideration are taken in to account for classification. It is to be understood that there are perceived perceivable and unobserved and expanding universes and multiverses too. The prediction model gives the expression for all these counts

T_{36} : Category one of **governed** by laws similar to electromagnetism

T_{37} : Category two of **governed** by laws similar to electromagnetism

T_{38} : Category three of **governed** by laws similar to electromagnetism

Module eight

The idea is similar to that of ordinary holograms where a three-dimensional image **is (e) encoded in** a two-dimensional surface, such as in the hologram on a credit card, but now it is the entire Universe that is encoded in such a fashion.

G_{40} : Category one of **encoded in** a two-dimensional surface, such as in the hologram on a credit card, but now it is the entire Universe that is encoded in such a fashion. Systems analogy is given here. There are various systems with two dimensional enclosures.

G_{41} : Category two of **encoded in** a two-dimensional surface, such as in the hologram on a credit card, but now it is the entire Universe that is encoded in such a fashion.

G_{42} : Category three of **encoded in** a two-dimensional surface, such as in the hologram on a credit card, but now it is the entire Universe that is encoded in such a fashion.

T_{40} : Category one of ordinary holograms where a three-dimensional image

T_{41} : Category two of ordinary holograms where a three-dimensional image

T_{42} : Category three of ordinary holograms where a three-dimensional image

Module Nine

AdS/Ricci-flat correspondence **and t**he Gregory-Laflamme instability specifically explains what is known as the Gregory Laflamme instability

G_{44} : Category one of AdS/Ricci-flat correspondence **and t**he Gregory-Laflamme instability

G_{45} : Category two of AdS/Ricci-flat correspondence **and t**he Gregory-Laflamme instability

G_{46} : Category three of AdS/Ricci-flat correspondence **and t**he Gregory-Laflamme instability

T_{44} : Category one of Gregory Laflamme instability

T_{45} : Category two of Gregory Laflamme instability

T_{46} : Category three of Gregory Laflamme instability

SECTION THREE

"Roman Ring": Nemo Mortalium Omnibus Horis Sapit-Nemo Saltat Sobrius Model

INTRODUCTION—VARIABLES USED

Wormholes and Time Travel Shortcut through Spacetime - If Traversable Wormholes Exist, They Could Allow Time Travel

(1) One significant limitation of such a time machine is that it is only possible to go as far back in time as the initial creation of the machine; in essence, it is more of a path **through (e&eb**) time than it is a device that itself moves through time, and it would not allow the technology itself to be moved backwards in time. This could provide an alternative explanation for Hawking's observation: a time machine will be built someday, but has not yet been built, so the tourists from the future cannot reach this far back in time.

Albert Einstein: Worm Hole Is time travel possible?

(2) According to current theories on the nature of wormholes, construction of a traversable wormhole **would (e) require the** existence of a substance with negative energy (often referred to as "exotic matter").

(3) More technically, the wormhole spacetime **requires a** distribution of energy that violates various energy conditions, such as the null energy condition along with the weak, strong, and dominant energy conditions.

(4) However, it is known that quantum effects **can lead to** small measurable violations of the null energy condition, and many physicists believe that the required negative energy may actually be possible due to the Casimir effect in quantum physics.

(5) physical forces **arising (e) from** a quantized field

(6) Although early calculations suggested a very large amount of negative energy **would be required**, later calculations showed that the amount of negative energy can be made arbitrarily small. In 1993, Matt Visser argued that the two mouths of a wormhole with such an induced clock difference could not be brought together **without** inducing quantum field and gravitational effects that would either make the wormhole collapse or the two mouths repel each other.

(7) Because of this, the two mouths could not be brought close enough **for (e)** causality violation to take place.

(8) However, in a 1997 paper, Visser hypothesized that a complex "Roman ring" (named after Tom Roman) configuration of an N number of wormholes arranged in a symmetric polygon could still

act as a time machine**, although he concludes** that this is more likely a flaw in classical quantum gravity theory rather than proof that causality violation is possible.

NOTATION

Module One

AdS/Ricci-flat correspondence **and t**he Gregory-Laflamme instability specifically explains what is known as the Gregory Laflamme instability and according to which certain types of black hole **break(e&eb) up into** smaller black holes when disturbed – rather like a thin stream of water breaking into little droplets when you touch it with your finger. This black hole phenomenon has previously been **shown (eb) to exist** through computer simulations and this work provides a deeper theoretical explanation

G_{13} : Category one of AdS/Ricci-flat correspondence **and t**he Gregory-Laflamme instability specifically explains what is known as the Gregory Laflamme instability and according to which certain types of black hole

G_{14} : Category two of AdS/Ricci-flat correspondence **and t**he Gregory-Laflamme instability specifically explains what is known as the Gregory Laflamme instability and according to which certain types of black hole

G_{15} : Category three of AdS/Ricci-flat correspondence **and t**he Gregory-Laflamme instability specifically explains what is known as the Gregory Laflamme instability and according to which certain types of black hole

T_{13} : Category one of smaller black holes when disturbed – rather like a thin stream of water breaking into little droplets when you touch it with your finger.

T_{14} : Category two of smaller black holes when disturbed – rather like a thin stream of water breaking into little droplets when you touch it with your finger.

T_{15} : Category three of smaller black holes when disturbed – rather like a thin stream of water breaking into little droplets when you touch it with your finger.

Module Two

One significant limitation of such a time machine is that it is only possible to go as far back in time as the initial creation of the machine; in essence, it is more of a path **through (e&eb**) time than it is a device that itself moves through time, and it would not allow the technology itself to be moved backwards in time. This could provide an alternative explanation for Hawking's observation: a time machine will be built someday, but has not yet been built, so the tourists from the future cannot reach this far back in time.

G_{16} : Category one of limitation of such a time machine is that it is only possible to go as far back in time as the initial creation of the machine; in essence, it is more of a path

G_{17} : Category two of limitation of such a time machine is that it is only possible to go as far back in time as the initial creation of the machine; in essence, it is more of a path

G_{18} : Category three of limitation of such a time machine is that it is only possible to go as far back in time as the initial creation of the machine; in essence, it is more of a path

T_{16} : Category one of device that itself moves through time, and it would not allow the technology itself to be moved backwards in time. This could provide an alternative explanation for Hawking's observation: a time machine will be built someday, but has not yet been built, so the tourists from the future cannot reach this far back in time.

T_{17} : Category two of device that itself moves through time, and it would not allow the technology itself to be moved backwards in time. This could provide an alternative explanation for Hawking's observation: a time machine will be built someday, but has not yet been built, so the tourists from the future cannot reach this far back in time.

T_{18} : Category three of device that itself moves through time, and it would not allow the technology itself to be moved backwards in time. This could provide an alternative explanation for Hawking's observation: a time machine will be built someday, but has not yet been built, so the tourists from the future cannot reach this far back in time.

Module three

More technically, the wormhole spacetime **requires a** distribution of energy that violates various energy conditions, such as the null energy condition along with the weak, strong, and dominant energy conditions

G_{20} : Category one of distribution of energy that violates various energy conditions, such as the null energy condition along with the weak, strong, and dominant energy conditions

G_{21} : Category two of distribution of energy that violates various energy conditions, such as the null energy condition along with the weak, strong, and dominant energy conditions

G_{22} : Category three of distribution of energy that violates various energy conditions, such as the null energy condition along with the weak, strong, and dominant energy conditions

T_{20} : Category one of wormhole spacetime

T_{21} : Category two of wormhole spacetime

T_{22} : Category three of wormhole spacetime

Module four

However, it is known that quantum effects **can lead to** small measurable violations of the null energy condition, and many physicists believe that the required negative energy may actually be possible due to the Casimir effect in quantum physics.

G_{24} : Category one of quantum effects

G_{25} : Category two of quantum effects

G_{26} : Category three of quantum effects

T_{24} : Category one of small measurable violations of the null energy condition, and many physicists believe that the required negative energy may actually be possible due to the Casimir effect in quantum physics.

T_{25} : Category two of small measurable violations of the null energy condition, and many physicists believe that the required negative energy may actually be possible due to the Casimir effect in quantum physics.

T_{26} : Category three of small measurable violations of the null energy condition, and many physicists believe that the required negative energy may actually be possible due to the Casimir effect in quantum physics.

Module five

physical forces **arise (e) from** a quantized field

G_{28} : Category one of quantized field

G_{29} : Category two of quantized field

G_{30} : Category three of quantized field

T_{28} : Category one of physical forces

T_{29} : Category two of physical forces

T_{30} : Category three of physical forces

Module six

Although early calculations suggested a very large amount of negative energy **would be required**, later calculations showed that the amount of negative energy can be made arbitrarily small. In 1993, Matt Visser argued that the two mouths of a wormhole with such an induced clock difference could not be brought together **without** inducing quantum field and gravitational effects that would either make the wormhole collapse or the two mouths repel each other.

G_{32} : Category one of **without** inducing quantum field and gravitational effects that would either make the wormhole collapse or the two mouths repel each other.

G_{33} : Category two of **without** inducing quantum field and gravitational effects that would either make the wormhole collapse or the two mouths repel each other.

G_{34} : Category three of **without** inducing quantum field and gravitational effects that would either make the wormhole collapse or the two mouths repel each other.

T_{32} : Category one of Visser argued that the two mouths of a wormhole with such an induced clock difference could not be brought together

T_{33} : Category two of Visser argued that the two mouths of a wormhole with such an induced clock difference could not be brought together

T_{34} : Category three of Visser argued that the two mouths of a wormhole with such an induced clock difference could not be brought together

Module seven

Because of this, the two mouths could not be brought close enough **for (e)** causality violation to take place.

G_{36} : Category one of two mouths could not be brought close enough **for (e)** causality violation to take place.

G_{37} : Category two of two mouths could not be brought close enough **for (e)** causality violation to take place.

G_{38} : Category three of two mouths could not be brought close enough **for (e)** causality violation to take place.

T_{36} : Category one of Visser argued that the two mouths of a wormhole with such an induced clock difference could not be brought together **without** inducing quantum field and gravitational effects that would either make the wormhole collapse or the two mouths repel each other.

T_{37} : Category two of Visser argued that the two mouths of a wormhole with such an induced clock difference could not be brought together **without** inducing quantum field and gravitational effects that would either make the wormhole collapse or the two mouths repel each other.

T_{38} : Category three of Visser argued that the two mouths of a wormhole with such an induced clock difference could not be brought together **without** inducing quantum field and gravitational effects that would either make the wormhole collapse or the two mouths repel each other.

Module eight

However, in a 1997 paper, Visser hypothesized that a complex "Roman ring" (named after Tom Roman) configuration of an N number of wormholes arranged in a symmetric polygon could still act as a time machine**, although he concludes** that this is more likely a flaw in classical quantum gravity theory rather than proof that causality violation is possible.

G_{40} : Category one of Visser hypothesized that a complex "Roman ring" (named after Tom Roman) configuration of an N number of wormholes arranged in a symmetric polygon could still act as a time machine**,**

G_{41} : Category two of Visser hypothesized that a complex "Roman ring" (named after Tom Roman) configuration of an N number of wormholes arranged in a symmetric polygon could still act as a time machine**,**

G_{42} : Category three of Visser hypothesized that a complex "Roman ring" (named after Tom Roman) configuration of an N number of wormholes arranged in a symmetric polygon could still act as a time machine**,**

T_{40} : Category one of **although he concludes** that this is more likely a flaw in classical quantum gravity theory rather than proof that causality violation is possible.

T_{41} : Category two of **although he concludes** that this is more likely a flaw in classical quantum gravity theory rather than proof that causality violation is possible.

T_{42} : Category three of **although he concludes** that this is more likely a flaw in classical quantum gravity theory rather than proof that causality violation is possible.

Module Nine

Traversable Lorentzian Wormholes: An Overview Adam Getchell

An overview of the history and relevance of wormholes is given. Morris and Thorne's description of a traversable wormhole is revisited, and the requirement for exotic matter is derived. The null, weak, strong, dominant, averaged null, averaged weak, and averaged strong energy conditions and The ways that traversable wormholes violate them is discussed. The connection between topological censorship and the averaged null energy condition is shown, with the result that wormholes are not topologically censored. The existence of a method for extracting a sufficient amount of exotic matter from the vacuum stress energy to build a wormhole with an arbitrarily large throat is proven. The connection between wormholes and time machines is explored, and how theorems related to closed timelike curves (chronology protection) or global hyperbolicity (cosmic censorship) might prevent their creation. Finally, it is stated how physical theories

might be reconciled to the notion of traversable Lorentzian wormholes**.**

G_{44} : Category one of theorems related to closed timelike curves (chronology protection) or global hyperbolicity (cosmic censorship)

G_{45} : Category two of theorems related to closed timelike curves (chronology protection) or global hyperbolicity (cosmic censorship)

G_{46} : Category three of theorems related to closed timelike curves (chronology protection) or global hyperbolicity (cosmic censorship)

T_{44} : Category one of connection between wormholes and time machines

T_{45} : Category two of connection between wormholes and time machines

T_{46} : Category three of connection between wormholes and time machines

SECTION FOUR

String Landscape: Naturam Expellas Furca, Tamen Usque Recurret Model

INTRODUCTION—VARIABLES USED

String landscape distributive in the universe? And might it manifest a Peano cosmic consciousness? Richard D. Ruquist, PhD

(1) A consequence of 10d supersymmetric string theory **is(eb) the** existence of a universal subspace called the Calabi-Yau Compact Manifold (CYCM) which is composed of a 3D array of discrete units of 6d compactified space.
(2) The recent astronomical evidence that Sommerfeld's fine-structure constant varies slightly across the visible universe (from north to south in an Earth perspective) **suggests that** the flux windings in the discrete 10^-30 cm diameter CYCM units may vary similarly across the universe.
(3) String theorists estimate that the flux has 10 quantum states while winding through the 500 or so holes in each CYCM unit, **so that there(eb) are about** 10^500 possible windings, the so-called string landscape. Such a large number is more than enough possible windings to fill many good size universes with distinct CYCM units.
(4) If the discrete and distinct CYCM units are also numerable, they may have the properties of Peano arithmetic and possibly **manifest (eb) an** invisible emergent collective cosmic consciousness that permeates the entire universe, separate from human physical consciousness.
(5) string theory and string landscape,
(6) fine-structure-constant and Calabi-Yau compact manifold
(7) Peano arithmetic and cosmic consciousness
(8) mind/brain duality and Calabi-Yau compact manifold
(9) cosmic consciousness and string theory

NOTATION

Module One

A consequence of 10d supersymmetric string theory **is(eb) the** existence of a universal subspace called the Calabi-Yau Compact Manifold (CYCM) which is composed of a 3D array of discrete units of 6d compactified space.

G_{13} : Category one of consequence of 10d supersymmetric string theory

G_{14} : Category two of consequence of 10d supersymmetric string theory

G_{15} : Category three of consequence of 10d supersymmetric string theor

T_{13} : Category one of existence of a universal subspace called the Calabi-Yau Compact Manifold (CYCM) which is composed of a 3D array of discrete units of 6d compactified space.

T_{14} : Category two of existence of a universal subspace called the Calabi-Yau Compact Manifold (CYCM) which is composed of a 3D array of discrete units of 6d compactified space.

T_{15} : Category three of existence of a universal subspace called the Calabi-Yau Compact Manifold (CYCM) which is composed of a 3D array of discrete units of 6d compactified space.

Module Two

The recent astronomical evidence that Sommerfeld's fine-structure constant varies slightly across the visible universe (from north to south in an Earth perspective) **suggests that** the flux windings in the discrete 10^-30 cm diameter CYCM units may vary similarly across the universe.

G_{16} : Category one of astronomical evidence that Sommerfeld's fine-structure constant varies slightly across the visible universe (from north to south in an Earth perspective)

G_{17} : Category two of astronomical evidence that Sommerfeld's fine-structure constant varies slightly across the visible universe (from north to south in an Earth perspective)

G_{18} : Category three of astronomical evidence that Sommerfeld's fine-structure constant varies slightly across the visible universe (from north to south in an Earth perspective)

T_{16} : Category one of flux windings in the discrete 10^-30 cm diameter CYCM units may vary similarly across the universe.

T_{17} : Category two of flux windings in the discrete 10^-30 cm diameter CYCM units may vary similarly across the universe.

T_{18} : Category three of flux windings in the discrete 10^-30 cm diameter CYCM units may vary similarly across the universe.

Module three

String theorists estimate that the flux has 10 quantum states while winding through the 500 or so holes in each CYCM unit, **so that there(eb) are about** 10^500 possible windings, the so-called string landscape. Such a large number is more than enough possible windings to fill many good size universes with distinct CYCM units

G_{20} : Category one of String theorists estimate that the flux has 10 quantum states while winding through the 500 or so holes in each CYCM unit

G_{21} : Category two of String theorists estimate that the flux has 10 quantum states while winding through the 500 or so holes in each CYCM unit

G_{22} : Category three of String theorists estimate that the flux has 10 quantum states while winding through

the 500 or so holes in each CYCM unit

T_{20} : Category one of 10^500 possible windings, the so-called string landscape. Such a large number is more than enough possible windings to fill many good size universes with distinct CYCM units

T_{21} : Category two of 10^500 possible windings, the so-called string landscape. Such a large number is more than enough possible windings to fill many good size universes with distinct CYCM units

T_{22} : Category three of 10^500 possible windings, the so-called string landscape. Such a large number is more than enough possible windings to fill many good size universes with distinct CYCM units

Module four

If the discrete and distinct CYCM units are also numerable, they may have the properties of Peano arithmetic and possibly **manifest (eb) an** invisible emergent collective cosmic consciousness that permeates the entire universe, separate from human physical consciousness

G_{24} : Category one of discrete and distinct CYCM units are also numerable, they may have the properties of Peano arithmetic and possibly

G_{25} : Category two of discrete and distinct CYCM units are also numerable, they may have the properties of Peano arithmetic and possibly

G_{26} : Category three of discrete and distinct CYCM units are also numerable, they may have the properties of Peano arithmetic and possibly

T_{24} : Category one of **manifest (eb) an** invisible emergent collective cosmic consciousness that permeates the entire universe, separate from human physical consciousness

T_{25} : Category two of **manifest (eb) an** invisible emergent collective cosmic consciousness that permeates the entire universe, separate from human physical consciousness

T_{26} : Category three of **manifest (eb) an** invisible emergent collective cosmic consciousness that permeates the entire universe, separate from human physical consciousness

Module five

string theory and string landscape

G_{28} : Category one of string theory

G_{29} : Category two of string theory

G_{30} : Category three of string theory

T_{28} : Category one of string landscape

T_{29} : Category two of string landscape

T_{30} : Category three of string landscape

Module six

fine-structure-constant and Calabi-Yau compact manifold

G_{32} : Category one of fine-structure-constant

G_{33} : Category two of fine-structure-constant

G_{34} : Category three of fine-structure-constant

T_{32} : Category one of Calabi-Yau compact manifold

T_{33} : Category two of Calabi-Yau compact manifold

T_{34} : Category three of Calabi-Yau compact manifold

Module seven

Peano arithmetic and cosmic consciousness(nature's general ledger)

G_{36} : Category one of Peano arithmetic

G_{37} : Category two of Peano arithmetic

G_{38} : Category three of Peano arithmetic

T_{36} : Category one of cosmic consciousness(nature's general ledger)

T_{37} : Category two of cosmic consciousness(nature's general ledger)

T_{38} : Category three of cosmic consciousness(nature's general ledger)

Module eight

mind/brain duality and Calabi-Yau compact manifold

G_{40} : Category one of mind/brain duality (mind is essentially an anagrammatic field as assumed by us acting on consciousness the storage of information in all three states producing and fabricating results which are dreams, pathological states of mind due to gratification and deprivation states)

G_{41} : Category two of mind/brain duality (mind is essentially an anagrammatic field as assumed by us acting on consciousness the storage of information in all three states producing and fabricating results which are dreams, pathological states of mind due to gratification and deprivation states)

G_{42} : Category three of mind/brain duality (mind is essentially an anagrammatic field as assumed by us acting on consciousness the storage of information in all three states producing and fabricating results which are dreams, pathological states of mind due to gratification and deprivation states)

T_{40} : Category one of Calabi-Yau compact manifold

T_{41} : Category two of Calabi-Yau compact manifold

T_{42} : Category three of Calabi-Yau compact manifold

Module Nine

cosmic consciousness(nature's general ledger) and string theory

G_{44} : Category one of cosmic consciousness(nature's general ledger)

G_{45} : Category two of cosmic consciousness(nature's general ledger)

G_{46} : Category three of cosmic consciousness(nature's general ledger)

T_{44} : Category one of string theory

T_{45} : Category two of string theory

T_{46} : Category three of string theory

SECTION FIVE

Philosophy As The Invention Of Concepts(Deluze And Guattari): Saltus In Demonstrando-Salva Veritate Model

INTRODUCTION—VARIABLES USED

The World Seen From Within: Deleuze and the Philosophy of Events Paul Patton from: Theory & Event Volume 1, Issue 1, 1997 | 10.1353/tae.1991.0006 .In lieu of an abstract, here is a brief excerpt of the content:

(1) It is easy to decry the abstraction and sophistication of political theory **and to (eb) advocate** a return to a more immediate engagement with real events.
(2) It is more difficult to specify the nature of this theoretical engagement or **to (eb) define** the 'real events' with which theory is summoned to engage.
(3) One response modeled upon the sciences would suggest that events **serve (e) to** confirm or refute particular theories.
(4) However, like experimental results, events require (e) interpretation and it is impossible to disentangle their meaning if not their very nature as events of a certain kind from the theories which inform their description.
(5) These points to a different response, according to which theory **engages (e&eb)** with events at the level of their identification and specification: what kinds of events are there? What kind of event is this? Foucault provides many examples of this form of engagement with events.
(6) For example, by showing that the prison and its associated discourse of reform have always co-existed, Discipline and Punish **transforms (e&eb)** a typical political response to prison protests into a non-event: the intention to make prisons more effective mechanisms of rehabilitation turns out to be a continuation of the modern apparatus of criminal punishment rather than a break with its past.
(7) Deleuze and Guattari's conception of philosophy as the invention of concepts which **'express'(eb)** events might also be understood in terms of the identification and specification of events.
(8) On their view, the task of philosophy when it creates concepts is to extract an event from things, **to**

(e) give them a new event.

(9) The aim is to engage with everyday social and historical reality in a manner which would **challenge (e&eb)** received ideas about the nature of events.

NOTATION

Module One

It is easy to decry the abstraction and sophistication of political theory **and to (eb) advocate** a return to a more immediate engagement with real events.

G_{13} : Category one of abstraction and sophistication of political theory

G_{14} : Category two of abstraction and sophistication of political theory

G_{15} : Category three of abstraction and sophistication of political theory

T_{13} : Category one of immediate engagement with real events

T_{14} : Category two of immediate engagement with real events

T_{15} : Category three of immediate engagement with real events

Module Two

It is more difficult to specify the nature of this theoretical engagement or **to (eb) define** the 'real events' with which theory is summoned to engage. We assume the relationship between the nature of theoretical engagement and the real event and give a model

G_{16} : Category one of nature of this theoretical engagement

G_{17} : Category two of nature of this theoretical engagement

G_{18} : Category three of nature of this theoretical engagement

T_{16} : Category one of 'real events'

T_{17} : Category two of 'real events'

T_{18} : Category three of 'real events'

Module three

One response modeled upon the sciences would suggest that events **serve (e) to** confirm or refute particular theories

G_{20} : Category one of confirmation or refutation of particular theories

G_{21} : Category two of confirmation or refutation of particular theories

G_{22} : Category three of confirmation or refutation of particular theories

T_{20} : Category one of One response modeled upon the sciences would suggest that events

T_{21} : Category two of One response modeled upon the sciences would suggest that events

T_{22} : Category three of One response modeled upon the sciences would suggest that events

Module four

However, like experimental results, events require (e) interpretation and it is impossible to disentangle their meaning if not their very nature as events of a certain kind from the theories which inform their description.

G_{24} : Category one of interpretation and it is impossible to disentangle their meaning if not their very nature as events of a certain kind from the theories which inform their description.

G_{25} : Category two of interpretation and it is impossible to disentangle their meaning if not their very nature as events of a certain kind from the theories which inform their description.

G_{26} : Category three of interpretation and it is impossible to disentangle their meaning if not their very nature as events of a certain kind from the theories which inform their description.

T_{24} : Category one of like experimental results, events require

T_{25} : Category two of like experimental results, events require

T_{26} : Category three of like experimental results, events require

Module five

These points to a different response, according to which theory **engages (e&eb)** with events at the level of their identification and specification: what kinds of events are there? What kind of event is this? Foucault provides many examples of this form of engagement with events

G_{28} : Category one of theories

G_{29} : Category two of theories

G_{30} : Category three of theories

T_{28} : Category one of events at the level of their identification and specification: what kinds of events are there? What kind of event is this? Foucault provides many examples of this form of engagement with events

T_{29} : Category two of events at the level of their identification and specification: what kinds of events are there? What kind of event is this? Foucault provides many examples of this form of engagement with events

T_{30} : Category three of events at the level of their identification and specification: what kinds of events are there? What kind of event is this? Foucault provides many examples of this form of engagement with events

Module six (case two)

G_{32} : Category one of events at the level of their identification and specification: what kinds of events are there? What kind of event is this? Foucault provides many examples of this form of engagement with events

G_{33} : Category two of events at the level of their identification and specification: what kinds of events are there? What kind of event is this? Foucault provides many examples of this form of engagement with events

G_{34} : Category three of events at the level of their identification and specification: what kinds of events are there? What kind of event is this? Foucault provides many examples of this form of engagement with events

T_{32} : Category one of theories

T_{33} : Category two of theories

T_{34} : Category three of theories

Module seven

Deleuze and Guattari's conception of philosophy as the invention of concepts which **'express'(eb)** events might also be understood in terms of the identification and specification of events

G_{36} : Category one of conception of philosophy as the invention of concepts

G_{37} : Category two of conception of philosophy as the invention of concepts

G_{38} : Category three of conception of philosophy as the invention of concepts

T_{36} : Category one of identification and specification of events

T_{37} : Category two of identification and specification of events

T_{38} : Category three of identification and specification of events

Module eight

For example, by showing that the prison and its associated discourse of reform have always co-existed, Discipline and Punish **transforms (e&eb)** a typical political response to prison protests into a non-event: the intention to make prisons more effective mechanisms of rehabilitation turns out to be a continuation of the modern apparatus of criminal punishment rather than a break with its past.

G_{40} : Category one of showing that the prison and its associated discourse of reform have always co-existed, Discipline and Punish **transforms**

G_{41} : Category two of showing that the prison and its associated discourse of reform have always co-existed, Discipline and Punish **transforms**

G_{42} : Category three of showing that the prison and its associated discourse of reform have always co-existed, Discipline and Punish **transforms**

T_{40} : Category one of typical political response to prison protests into a non-event: the intention to make prisons more effective mechanisms of rehabilitation turns out to be a continuation of the modern apparatus of criminal punishment rather than a break with its past.

T_{41} : Category two of typical political response to prison protests into a non-event: the intention to make prisons more effective mechanisms of rehabilitation turns out to be a continuation of the modern apparatus of criminal punishment rather than a break with its past.

T_{42} : Category three of typical political response to prison protests into a non-event: the intention to make prisons more effective mechanisms of rehabilitation turns out to be a continuation of the modern apparatus of criminal punishment rather than a break with its past.

Module Nine

On their view, the task of philosophy when it creates concepts is to extract an event from things, **to (e) give** them a new event.

G_{44} : Category one of extraction of an event from things, **to (e) gives** them a new event.

G_{45} : Category two of extraction of an event from things, **to (e) give** them a new event

G_{46} : Category three of extraction of an event from things, **to (e) give** them a new event

T_{44} : Category one of creation of concepts

T_{45} : Category two of creation of concepts

T_{46} : Category three of creation of concepts

SECTION SIX

Creation Of Concepts: Acta Deos Nunquam Mortalia Fallunt Model

INTRODUCTION—VARIABLES USED

The World Seen From Within: Deleuze and the Philosophy of Events Paul Patton from: Theory & Event Volume 1, Issue 1, 1997 | 10.1353/tae.1991.0006 .In lieu of an abstract, here is a brief excerpt of the content:

(1) The conceptualization of new events **enables (eb) us** to become conscious of processes and forces at work in the present, those which we might seek to advance as well as those we might oppose.
(2) Philosophy understood in this manner is intended **to (e&eb) orient** thought and action.
(3) Deleuze and Guattari insist upon the utopian vocation of philosophy, arguing that the creation of new concepts **may (eb) contribute** to the emergence of 'a new earth and people that do not yet exist'.
(4) In order to clarify their stated aim of a philosophy which would be 'worthy' of the event, it may be helpful to examine Deleuze's concept of the event and to ask how this might apply to present social and political events. Throughout his work, Deleuze has always drawn the key elements of his concept of the event **from (e) the** Stoics.
(5) In The Logic of Sense, he argues that they were the first **to create** a philosophical concept of the event, describing this as '... an incorporeal, complex and irreducible entity, at the surface of things, a pure event which inheres or subsist in the proposition.'
(6) The Stoics drew a fundamental distinction between two realms of being, a material realm of bodies **and** states of affairs and an incorporeal realm of events.
(7) Events are **expressed (e) by** means of language, in statements, but they are attributes of bodies and physical states of affairs.
(8) Thus, the knife opening up a wound in flesh **is (=)** an attribute of the interpenetration of bodies, but

the event of 'being cut' is what is expressed by the statement 'He was cut with the knife'.

(9) The fact of being cut **is (=)** a property of neither the flesh nor the knife; it is an incorporeal attribute of the flesh.

(10) It is an event which may be expressed in a variety of ways, for example in the statement that he has a wound. On this account, events are the epiphenomena **of (e)** corporeal causal interactions: they do not affect bodies and states of affairs but they do affect other events, such as the responses and actions of agents. Pure events are both the expressed of statements and the 'sense' of what happens.

NOTATION

Module One

The conceptualization of new events **enables (eb) us** to become we conscious of processes and forces at work in the present, those which we might seek to advance as well as those might oppose.

G_{13} : Category one of conceptualization of new events

G_{14} : Category two of conceptualization of new events

G_{15} : Category three of conceptualization of new events

T_{13} : Category one of conscious of processes and forces at work in the present, those which we might seek to advance as well as those might oppose.

T_{14} : Category two of conscious of processes and forces at work in the present, those which we might seek to advance as well as those might oppose.

T_{15} : Category three of conscious of processes and forces at work in the present, those which we might seek to advance as well as those might oppose.

Module Two

Philosophy understood in this manner is intended **to (e&eb) orient** thought and action

G_{16} : Category one of such an understanding of the conceptualization

G_{17} : Category two of such an understanding of the conceptualization

G_{18} : Category three of such an understanding of the conceptualization

T_{16} : Category one of **orientation of** thought and action

T_{17} : Category two of **orientation of** thought and action

T_{18} : Category three of **orientation of** thought and action

Module three

Deleuze and Guattari insist upon the utopian vocation of philosophy, arguing that the creation of new concepts **may (eb) contribute** to the emergence of 'a new earth and people that do not yet exist'.

G_{20} : Category one of creation of new concepts

G_{21} : Category two of creation of new concepts

G_{22} : Category three of creation of new concepts

T_{20} : Category one of emergence of 'a new earth and people that do not yet exist'.(We interpret this as newer

thoughts and actions)

T_{21} : Category two of emergence of 'a new earth and people that do not yet exist'.(We interpret this as newer thoughts and actions)

T_{22} : Category three of emergence of 'a new earth and people that do not yet exist'.(We interpret this as newer thoughts and actions)

Module four

In The Logic of Sense, he argues that they were the first **to create** a philosophical concept of the event, describing this as '... an incorporeal, complex and irreducible entity, at the surface of things, a pure event which inheres or subsist in the proposition.'

G_{24} : Category one of conceptualization of event, thought, and phenomenological correlates

G_{25} : Category two of conceptualization of event, thought, and phenomenological correlates

G_{26} : Category three of conceptualization of event, thought, and phenomenological correlates

T_{24} : Category one of creation of concept of the event, describing this as '... an incorporeal, complex and irreducible entity, at the surface of things, a pure event which inheres or subsist in the proposition.'

T_{25} : Category two of creation of concept of the event, describing this as '... an incorporeal, complex and irreducible entity, at the surface of things, a pure event which inheres or subsist in the proposition.'

T_{26} : Category three of creation of concept of the event, describing this as '... an incorporeal, complex and irreducible entity, at the surface of things, a pure event which inheres or subsist in the proposition.'

Module five

In order to clarify their stated aim of a philosophy which would be 'worthy' of the event, it may be helpful to examine Deleuze's concept of the event and to ask how this might apply to present social and political events. Throughout his work, Deleuze has always drawn the key elements of his concept of the event **from (e) the** Stoics.

G_{28} : Category one of stoic philosophy

G_{29} : Category two of stoic philosophy

G_{30} : Category three of stoic philosophy

T_{28} : Category one of Deleuze's concept of the event

T_{29} : Category two of Deleuze's concept of the event

T_{30} : Category three of Deleuze's concept of the event

Module six

The Stoics drew a fundamental distinction between two realms of being, a material realm of bodies **and** states of affairs and an incorporeal realm of events.

REALER THAN REAL The Simulacrum According to Deleuze and Guattari Brian Massumi.

Originally published in Copyright no.1, 1987.Links to other works by Brian Massumi on the Web

There is a seductive image of contemporary culture circulating today. Our world, Jean Baudrillard tells us, has been launched into hyperspace in a kind of postmodern apocalypse. The airless atmosphere has asphyxiated the referent, leaving us satellites in aimless orbit around an empty center. We breathe an ether of floating images that no longer bear a relation to any reality whatsoever.1 That, according to Baudrillard, is simulation: the substitution of signs of the real for the real.2 In hyperreality, signs no longer represent or refer to an external model. They stand for nothing but themselves, and refer only to other signs. They are to some extent distinguishable, in the way the phonemes of language are, by a combinatory of minute binary distinctions.3 But postmodernism stutters. In the absence of any gravitational pull to ground them, images accelerate and tend to run together. They become interchangeable. Any term can be substituted for any other: utter indetermination.4 Faced with this homogeneous surface of syntagmatic slippage, we are left speechless. We can only gape in fascination.5 For the secret of the process is beyond our grasp. Meaning has imploded. There is no longer any external model, but there is an immanent one. To the syntagmatic surface of slippage there corresponds an invisible paradigmatic dimension that creates those minimally differentiated signs only in order for them to blur together in a pleasureless orgy of exchange and circulation. Hidden in the images is a kind of genetic code responsible for their generation.6 Meaning is out of reach and out of sight, but not be cause it has receded into the distance. It is because the code has been miniaturized. Objects are images, images are signs, signs are information, and information fits on a chip. Everything reduces to a molecular binarism. The generalized digitality of the computerized society.7

And so we gape. We cannot be said to be passive exactly, because all polarity, including the active/passive dichotomy, has disappeared. We have no earth to center us, but we ourselves function as a ground--in the electrical sense.8 We do not act, but neither do we merely receive. We absorb through our open eyes and mouths. We neutralize the play of energized images in the mass entropy of the silent majority.

It makes for a fun read. But do we really have no other choice than being a naive realist or being a sponge?

Deleuze and Guattari open a third way. Although it is never developed at length in any one place, a theory of simulation can be extracted from their work that can give us a start in analyzing our cultural condition under late capitalism without landing us back with the dinosaurs or launching us into hypercynicism.

A common definition of the simulacrum is a copy of a copy whose relation to the model has become so attenuated that it can no longer properly be said to be a copy. It stands on its own as a copy without a model. Fredric Jameson cites the example of photorealism. The painting is a copy not of reality, but of a photograph, which is already a copy of the original.9 Deleuze, in his article "Plato and the Simulacrum," takes a similar definition as his starting point, but emphasizes its inadequacy. For beyond a certain point, the distinction is no longer one of degree. The simulacrum is less a copy twice removed than a phenomenon of a different nature altogether: it undermines the very distinction between copy and model.10 The terms copy and model bind us to the world of representation and objective (re)production. A copy, no matter how many times removed, authentic or fake, is defined by the presence or absence of internal, essential relations of resemblance to a model. The simulacrum, on the other hand, bears only an external and deceptive resemblance to a putative model. The process of its production, its inner dynamism, is entirely different from that of its supposed model; its resemblance to it is merely a surface effect, an illusion.11 The production and function of a photograph has no relation to that of the object photographed; and the photorealist painting in turn envelops an essential difference. It is that masked difference, not the manifest resemblance, that produces the effect of uncanniness so often associated with the simulacrum. A copy is made in order to stand in for its model. A simulacrum has a different agenda, it enters different circuits. Pop Art is the example Deleuze uses for simulacra that have successfully broken out of the copy mold:12 the multiplied, stylized images take on a life of their own. The thrust of the process is not to become an equivalent of the "model" but to turn against it and its world in order to open a new space for the simulacrum's own mad proliferation. The simulacrum affirms its own difference. It is not an implosion, but a differentiation; it is an index not of absolute proximity, but of galactic distances.

The resemblance of the simulacrum is a means, not an end. A thing, write Deleuze and Guattari, "in order to become apparent, is forced to simulate structural states and to slip into states of forces that serve it as masks. . . . underneath the mask and by means of it, it already invests the terminal forms and the specific higher

states whose integrity it will subsequently establish."13 Resemblance is a beginning masking the advent of whole new vital dimension. This even applies to mimickry in nature. An insect that mimics a leaf does so not to meld with the vegetable state of its surrounding milieu, but to reenter the higher realm of predatory animal warfare on a new footing. Mimickry, according to Lacan, is camouflage.14 It constitutes a war zone. There is a power inherent in the false: the positive power of ruse, the power to gain a strategic advantage by masking one's life force.

Ridley Scott's film Blade Runner shows that the ultimate enemy in this war of ruse is the so-called "model" itself. The off-world replicants return to earth not to blend in with the indigenous population, but to find the secret of their built-in obsolescence so they can escape their bondage and live full lives, and on their own terms. Imitation is an indication of a life force propelling the falsifier toward the unbridled expression of its uniqueness. The dominant replicant makes a state ment to the man who made his eyes that can be taken as a general formula for simulation: if only you could see what I have seen with your eyes. If they find out how to undo their pre-programed deaths, the replicants will not remain on earth as imitation humans. They will either take over or flee back to their own vital dimension of interplanetary space to see things no human being ever has or will. Their imitation is only a way-station en route to an unmasking and the assumption of difference. As Eric Alliez and Michel Feher observe, the best weapon against the simulacrum is not to unmask it as a false copy, but to force it to be a true copy, thereby resubmitting it to representation and the mastery of the model: the corporation that built the rebellious replicants introduces a new version complete with second-hand human memories.15

I said earlier that the simulacrum cannot adequately be discussed in terms of copy and model, and now I find myself not only talking about a model again, but claiming that it is in a life and death struggle with the simulacrum. The reality of the model is a question that needs to be dealt with. Baudrillard sidesteps the question of whether simulation replaces a real that did indeed exist, or if simulation is all there has ever been.16 Deleuze and Guattari say yes to both. The alternative is a false one because simulation is a process that produces the real, or, more precisely, more real (a more-than-real) on the basis of the real. "It carries the real beyond its principle to the point where it is effectively produced."17 Every simulation takes as its point of departure a regularized world comprising apparently stable identities or territories. But these "real" entities are in fact undercover simulacra that have consented to feign being copies. A silent film by Louis Feuillade illustrates the process.

Vendémiaire takes place in the final days of World War I. The plot is simple: members of a well-to-family from the north of France who cannot fight in the war flee to unoccupied territory in the south to contribute their efforts to the wine harvest. There they meet one of the daughters' husband-to-be and a sinister pair of German prisoners of war who have obtained identity papers by killing two Belgians and try to pass themselves off as Allies until they can get enough money to flee to Spain. The Germans' plan is to steal from the vineyard owners and pin the theft on a gypsie woman who is also working on the harvest. The plan fails when one of the Germans, about to be found out, jumps into an empty grape storage tank. He is killed by poisonous gases produced by grapes fermenting in the next tank. His corpse is found still clutching the loot, and the gypsie woman is saved. His lonely comrade later betrays himself by getting drunk and speaking in German.

The film is bracketted by grapes. The grape harvest supplies the initial motivation that sets up the situation of the plot, and the grapes themselves rather than any human hero resolve the dilemma. The film is not only bracketted by grapes; it swims in wine as its very element. Every crucial moment is expressed in terms of wine: love is expressed by the scintillating image of the faraway wife dancing in the husband's wine cup; the German menace in its highest expression is one of the escapees stomping on the grape vine; heroism is exemplified by an altruistic trooper who braves death to bring wine back to the trenches to give his comrades a taste of the homeland that will revive their will to victory; when victory does come, it is toasted to with wine, and the movie ends with a sentimental tableau of the vines and a final intertitle saying that from these vineyards a new nation will be reborn. "Simulation," Deleuze and Guattari write, "does not replace reality . . . but rather it appropriates reality in the operation of despotic overcoding, it produces reality on the new full body that replaces the earth. It expresses the appropriation and production of the real by a quasi-cause."18 The undivided, abstract flow of wine is the glorified body of the nation. It arrogates to itself the power of love, victory and rebirth. It presents itself as first and final cause. But the war was obviously not won with

wine. Its causality is an illusion. But it is an effective illusion because it is reinjected into reality and sets to work: it expresses love, and thereby motivates the man to be a good husband and give sons to the nation rising; it expresses patriotism, and thereby spurs the soldiers to victory. That is why it is called a quasi-cause. It abstracts from bodies and things a transcendental plane of ideal identities: a glorious wife, a glorious family, a glorious nation. ("It carries the real beyond its principle...") Then it folds that ideal dimension back down onto bodies and things in order to force them to conform to the distribution of identities it lays out for them. ("...to the point where it is effectively produced.") It creates the entire network of resemblance and representation. Both copy and model are the products of the same fabulatory process, the final goal of which is the recreation of the earth, the creation of a new territory.

The power of the quasi-cause is essentially distributive. It separates the good bodies from the bad, in other words the bodies that agree to resemble the glorious illusion it presents them as a model from those that do not; and it polices for renegade copies operating with a different agenda. The quasi-cause enables the French patriots to unmask the conniving Germans,19 and it shows up the gypsie for the true, hard-working Frenchwoman that she is despite her apparent otherness.

This account overcomes the polarity between the model and the copy by treating them both as second-order productions, as working parts in the same machine; but it seems to leave intact the dichotomy between the real and the imaginary--until it is realized that the bodies and things that are taken up by this fabulatory process are themselves the result of prior simulation-based distributions operating on other levels with different quasi-causes. Simulation upon simulation. Reality is nothing but a well-tempered harmony of simulation. The world is a complex circuit of interconnected simulations, in which Feuillade's own film takes its place. It was made in 1919, just after the war. Every war, especially one of those dimensions, has a powerful deterritorializing effect: the mobilization of troops and supplies, refugees from other countries, refugees to other countries, families broken, entire regions levelled... The film itself is a simulation meant to insert itself into that disjointed situation to help induce a unifying reterritorialization, to contribute to the rebirth of the nation. Vendémiare is the first month of the Republican calendar.

So what we are left with is a distinction not primarily between the model and the copy, or the real and the imaginary, but between two modes of simulation. One, exemplified in Feuillade's film, is normative, regularizing, and reproductive. It selects only certain properties of the entities it takes up: hard work, loyalty, good parenting, etc. It creates a network of surface resemblances. They are surface resemblances because at bottom they not resemblances at all but standardized actions: what those entities do when called upon (the gypsie in this respect is as French as the French). What bodies do depends on where they land in a abstract grid of miraculated identities that are in practice only a bundle of normalized and basically reproductive functions. It is not a question of Platonic copies, but of human replicants. Every society creates a quasi-causal system of this kind. In capitalist society the ultimate quasi-cause is capital itself,20 which is described by Marx as a miraculating substance that arrogates all things to itself and presents itself as first and final cause. This mode of simulation goes by the name of "reality."

The other mode of simulation is the one that turns against the entire system of resemblance and replication. It is also distributive, but the distribution it effects is not limitative. Rather than selecting only certain properties, it selects them all, it multiplies potentials: not to be human, but to be human plus. This kind of simulation is called "art." Art also recreates a territory, but a territory that is not really territorial. It is less like the earth with its gravitational grid than an interplanetary space, a deterritorialized territory providing a possibility of movement in all directions. Artists are replicants who have found the secret of their obsolescence.

In A Thousand Plateaus, Deleuze and Guattari invent a vocabulary enabling them to discuss both modes of simulation without lapsing into the terminology of representation. The key concept is double becoming. There are always at least two terms swept up in a fabulous process that transforms them both.21 David Cronenberg's film, The Fly, presents an instance of this, although a failed one. A scientist named Brundle accidentally splices himself with a fly as he is experimenting with a machine that can dematerialize objects and transport them instantly to any chosen location, in defiance of gravity and Newtonian physics generally. When the accident occurs, Brundle does not so much become fly, nor the fly human. Rather, certain properties or potentials of both combine in a new and monstrous amalgation: a Brundle-Fly that can walk on walls and think and speak well enough to describe itself as the world's first "insect politician." It tries to

purify itself of the fly in it by repeating the process backwards, but only succeeds in combining with the machinery itself. In limitative or negative becoming as portrayed in Vendémiare, one of the terms is an abstract identity and the body in question must curtail its potentials in order to fit into the grid, or at least appear to. In nonlimitative or positive becoming, as in The Fly, both terms are on the same level: rather than looking perpendicularly up or down, one moves sideways toward a another position on the grid for which one was not destined, toward an animal, a machine, a person of a different sex or age or race, an insect, a plant. The fabulatory process, though as abstract as subatomic physics, is immanent to the world of the things it affects, and is as real as a quark.22 .The transporting machine is on the same plane as the terms it combines. Its operating principle dips into that world's quantum level, into its pool of virtuality, to create an as yet unseen amalgamation of potentials. It produces a new body or territory from which there is no turning back. The only choice is to keep on becoming in an endless relay from one term to the next until the process either makes a breakthrough or exhausts its potential, spends its fuel, and the fabulous animal dies. Likening this to interplanatary space can be misleading: there is nothing farther from free-floating weightlessness than this. There is no such thing as total indetermination. Every body has its own propulsion, its own life force, its own set of potentials defining how far it can go. And it moves in a world filled with the obstacles thrown down by sedimentations of preexisting simulations of the "real" persuasion. There is no generalized indetermination, but there are localized points of undecidability where man meets fly. The goal is to reach into one's world's quantum level at such a point and, through the strategic mimickry of double becoming, combine as much potential as possible. Deleuze and Guattari, of course, are not suggesting that people can or should "objectively" become insects. It is a question of extracting and combining potentials, which they define as abstract relations of movement and rest, abilities to affect and be affected: abstract yet real. The idea is to build our own transporting machine and use it to get a relay going and to keep it going, creating ever greater and more powerful amalgamations and spreading them like a contagion until they infect every identity across the land and the point is reached where a now all-invasive positive simulation can turn back against the grid of resemblance and replication and overturn it for a new earth. Deleuze and Guattari insist on the collective nature of this process of becoming, even when it is seemingly embodied in a solitary artist. Revolutionary or "minor"23 artists marshal all of the powers of the false their community has to offer. They create a working simulation that may then reinject itself into society like Feuillade's wine assemblage, but to very different, though perhaps equally intoxicating, effect.

Returning to The Fly, the former scientist's only hope for a breakthrough is to convince his former girlfriend to have a child by him and the fly. His hope, and her fear, is that he will infect the human race with Brundle-Flies, and a new race with superhuman strength will rise up to replace the old. The overman as superfly.24 Reproduction, and the forging of a new ethnic identity, is aspects of this process of simulation, but they are not the goal. The goal is life, a world in which the New Brundle can live without hiding and repressing his powers. That possibility is successfully squelched by the powers that be. Brundle-Fly is deprived of an escape route. The original formula, as inscribed in the bodies of Brundle and the fly, was apparently flawed. They did the best they could do, but only reached obsolescence.

How does all of this apply to our present cultural condition? According to Deleuze, the point at which simulacrum began to unmask itself was reached in painting with the advent of Pop Art. In film, it was Italian neo-Realism and the French New Wave.25 Perhaps we are now reaching that point in popular culture as a whole. Advanced capitalism, Deleuze and Guattari argue, is reaching a new transnational level that necessitates a dissolution of old identities and territorialities and the unleashing of objects, images and information having far more mobility and combinatory potential than ever before.26 As always, this deterritorialization is effected only in order to make possible a reterritorialization on an even grander and more glorious land of worldwide capital reborn. But in the meantime, a breach has opened. The challenge is to assume this new world of simulation and take it one step farther, to the point of no return, to raise it to a positive simulation of the highest degree by marshaling all our powers of the false toward shattering the grid of representation once and for all.

This cannot be done by whining. The work of Baudrillard is one long lament. Both linear and dialectical causality no longer function, therefore everything is indetermination. The center of meaning is empty; therefore we are satellites in lost orbit. We can no longer act like legislator-subjects or be passive like slaves, therefore we are sponges. Images are no longer anchored by representation; therefore they float weightless in hyperspace. Words are no longer univocal; therefore signifiers slip chaotically over each other. A circuit has

been created between the real and the imaginary, therefore reality has imploded into the undecidable proximity of hyperreality. All of these statements make sense only if it is assumed that the only conceivable alternative to representative order is absolute indetermination, whereas indetermination as he speaks of it is in fact only the flipside of order, as necessary to it as the fake copy is to the model, and every bit as much a part of its system. Baudrillard's framework can only be the result of a nostalgia for the old reality so intense that it has difformed his vision of everything outside of it. He cannot clearly see that all the things he says have crumbled were simulacra all along: simulacra produced by analyzable procedures of simulation that were as real as real, or actually realer than real, because they carried the real back to its principle of production and in so doing prepared their own rebirth in a new regime of simulation. He cannot see becoming, of either variety. He cannot see that the simulacrum envelops a proliferating play of differences and galactic distances. What Deleuze and Guattari offer, particularly in A Thousand Plateaus, is a logic capable of grasping Baudrillard's failing world of representation as an effective illusion the demise of which opens a glimmer of possibility. Against cynicism, a thin but fabulous hope--of ourselves becoming realer than real in a monstrous contagion of our own making.

G_{32} : Category one of material realm of bodies

G_{33} : Category two of material realm of bodies

G_{34} : Category three of material realm of bodies

T_{32} : Category one of states of affairs and an incorporeal realm of events(Deluze represents this by transcendental surface)

T_{33} : Category two of states of affairs and an incorporeal realm of events(Deluze represents this by transcendental surface)

T_{34} : Category three of states of affairs and an incorporeal realm of events(Deluze represents this by transcendental surface)

Module seven

Events are **expressed (e) by** means of language, in statements, but they are attributes of bodies and physical states of affairs.

G_{36} : Category one of language, in statements, but they are attributes of bodies and physical states of affairs.

G_{37} : Category two of language, in statements, but they are attributes of bodies and physical states of affairs.

G_{38} : Category three of language, in statements, but they are attributes of bodies and physical states of affairs.

T_{36} : Category one of Events

T_{37} : Category two of Events

T_{38} : Category three of Events

Module eight

Thus, the knife opening up a wound in flesh **is** an attribute of the interpenetration of bodies, but the event of 'being cut' is what is expressed by the statement 'He was cut with the knife'.

G_{40} : Category one of attribute of the interpenetration of bodies, but the event of 'being cut' is what is expressed by the statement 'He was cut with the knife'.

G_{41} : Category two of attribute of the interpenetration of bodies, but the event of 'being cut' is what is expressed by the statement 'He was cut with the knife'.

G_{42} : Category three of attribute of the interpenetration of bodies, but the event of 'being cut' is what is expressed by the statement 'He was cut with the knife'.

T_{40} : Category one of knife opening up a wound in flesh

T_{41} : Category two of knife opening up a wound in flesh

T_{42} : Category three of knife opening up a wound in flesh

Module Nine

The fact of being cut **is (=)** a property of neither the flesh nor the knife; it is an incorporeal attribute of the flesh.

G_{44} : Category one of property of neither the flesh nor the knife; it is an incorporeal attribute of the flesh.

G_{45} : Category two of property of neither the flesh nor the knife; it is an incorporeal attribute of the flesh.

G_{46} : Category three of property of neither the flesh nor the knife; it is an incorporeal attribute of the flesh.

T_{44} : Category one of fact of being cut

T_{45} : Category two of fact of being cut

T_{46} : Category three of fact of being cut

SECTION SEVEN

Mathematicised Ontology(Badiou And Deleuze): Ab Aeterno Model

INTRODUCTION—VARIABLES USED

(1) That which is more abstract **is (=) that** which incorporates a larger, that is, more multiple, sample, considering not only human but also animal, inorganic and other inhuman processes.
(2) According to Badiou, of course, this is not true abstraction. What Deleuze discounts as the mere formality of pure quantity **is (=) actually** wherein its power rests for Badiou, and for two reasons: multiplicity must be pure quantity (no quality) if it is to generate the paradoxes that condition the possibility of an event; and it must be homogeneous and empty if it is to be the basis for a truly universalist politics.
(3) Conversely, according to Badiou, Deleuze's antipathy towards a wholly mathematicised ontology (Deleuze does, admittedly, embrace the differential calculus in his work, but more on account of its kinship with vital processes rather than its inherent formalism),**is due (e) in** part to a poorly thought-out polemic against sets that erroneously reduces them to number (whereas number actually reduces to sets).
(4) Before turning next to possibly the most important idea Badiou shares with Deleuze (the idea of the event), let me say a few short words about two topics on which they appear to diverge most, the Subject and Truth. According to Badiou, truth and subjectivity are **co-engendered (e)** through the 'event', which must be understood conceptually, that is, through reason.

Being and Event

(5) Being and event is that 'a subject is **nothing (=) other than** an active fidelity to the event of truth'.
(6) Where Deleuze gives the sensation(of) matter the transcendental function of constituting the subject, for Badiou, the event of truth and the subject come together **so (eb) that** neither supervenes over the other, as objectivism or subjectivism would respectively entail.
(7) There is an identity of being and thinking, but only **via (e&eb)** events and the decisions immanent to them – they are co-engendered through one process.
(8) Qua a non-processual state, the subject is indeed **transcended (eb)by** truth –'every truth is transcendent to the subject'
(9) Qua process, it is through subjectivization that **manifestation of** truth is possible.

NOTATION

Module One

That which is more abstract **is (=) that** which incorporates a larger, that is, more multiple, sample, considering not only human but also animal, inorganic and other inhuman processes

G_{13} : Category one of incorporates a larger, that is, more multiple, sample, considering not only human but also animal, inorganic and other inhuman processes

G_{14} : Category two of incorporates a larger, that is, more multiple, sample, considering not only human but also animal, inorganic and other inhuman processes

G_{15} : Category three of incorporates a larger, that is, more multiple, sample, considering not only human but also animal, inorganic and other inhuman processes

T_{13} : Category one of abstract

T_{14} : Category two of abstract

T_{15} : Category three of abstract

Module Two

According to Badiou, of course, this is not true abstraction. What Deleuze discounts as the mere formality of pure quantity **is actually** wherein its power rests for Badiou, and for two reasons: multiplicity must be pure quantity (no quality) if it is to generate the paradoxes that condition the possibility of an event; and it must be homogeneous and empty if it is to be the basis for a truly universalist politics

G_{16} : Category one of power rests for Badiou, and for two reasons: multiplicity must be pure quantity (no quality) if it is to generate the paradoxes that condition the possibility of an event; and it must be homogeneous and empty if it is to be the basis for a truly universalist conceptualization

G_{17} : Category two of power rests for Badiou, and for two reasons: multiplicity must be pure quantity (no quality) if it is to generate the paradoxes that condition the possibility of an event; and it must be homogeneous and empty if it is to be the basis for a truly universalist conceptualization

G_{18} : Category three of power rests for Badiou, and for two reasons: multiplicity must be pure quantity (no quality) if it is to generate the paradoxes that condition the possibility of an event; and it must be homogeneous and empty if it is to be the basis for a truly universalist conceptualization

T_{16} : Category one of According to Badiou, of course, this is not true abstraction. What Deleuze discounts as the mere formality of pure quantity. Characterstics of the systems studied are taken in to consideration in the classification schema and doxa. For extremal details kindly refer to Introduction. There are many systems that satisfy the criterion presupposed by both the bulwarks.

T_{17} : Category two of According to Badiou, of course, this is not true abstraction. What Deleuze discounts as the mere formality of pure quantity

T_{18} : Category three of According to Badiou, of course, this is not true abstraction. What Deleuze discounts as the mere formality of pure quantity

Module three

Conversely, according to Badiou, Deleuze's antipathy towards a wholly mathematicised ontology (Deleuze does, admittedly, embrace the differential calculus in his work, but more on account of its kinship with vital processes rather than its inherent formalism),**is due (e) in** part to a poorly thought-out polemic against sets that erroneously reduces them to number (whereas number actually reduces to sets).

G_{20} : Category one of poorly thought-out polemic against sets that erroneously reduces them to number (whereas number actually reduces to sets).

G_{21} : Category two of poorly thought-out polemic against sets that erroneously reduces them to number (whereas number actually reduces to sets).

G_{22} : Category three of poorly thought-out polemic against sets that erroneously reduces them to number (whereas number actually reduces to sets).

T_{20} : Category one of Deleuze's antipathy towards a wholly mathematicised ontology (Deleuze does, admittedly, embrace the differential calculus in his work, but more on account of its kinship with vital processes rather than its inherent formalism),

T_{21} : Category two of Deleuze's antipathy towards a wholly mathematicised ontology (Deleuze does, admittedly, embrace the differential calculus in his work, but more on account of its kinship with vital processes rather than its inherent formalism),

T_{22} : Category three of Deleuze's antipathy towards a wholly mathematicised ontology (Deleuze does, admittedly, embrace the differential calculus in his work, but more on account of its kinship with vital processes rather than its inherent formalism),

Module four

Before turning next to possibly the most important idea Badiou shares with Deleuze (the idea of the event), let me say a few short words about two topics on which they appear to diverge most, the Subject and Truth. According to Badiou, truth and subjectivity are **co-engendered (e)** through the 'event', which must be understood conceptually, that is, through reason.

G_{24} : Category one of 'event', which must be understood conceptually, that is, through reason.

G_{25} : Category two of 'event', which must be understood conceptually, that is, through reason.

G_{26} : Category three of 'event', which must be understood conceptually, that is, through reason.

T_{24} : Category one of truth and subjectivity. Truth could be relativistic, contradictory and there are much subjectivity.

T_{25} : Category two of truth and subjectivity

T_{26} : Category three of truth and subjectivity

Module five

Being and event is that 'a subject is **nothing other than** an active fidelity to the event of truth'.

G_{28} : Category one of active fidelity to the event of truth'

G_{29} : Category two of active fidelity to the event of truth'

G_{30} : Category three of active fidelity to the event of truth'

T_{28} : Category one of Being and event is that 'a subject is

T_{29} : Category two of Being and event is that 'a subject is

T_{30} : Category three of Being and event is that 'a subject is

Module six

Where Deleuze gives the sensation(of) matter the transcendental function of constituting the subject, for Badiou, the event of truth and the subject come together **so (eb) that** neither supervenes over the other, as objectivism or subjectivism would respectively entail

G_{32} : Category one of sensation(of) matter the transcendental function of constituting the subject, for Badiou, the event of truth and the subject come together

G_{33} : Category two of sensation(of) matter the transcendental function of constituting the subject, for Badiou, the event of truth and the subject come together

G_{34} : Category three of sensation(of) matter the transcendental function of constituting the subject, for Badiou, the event of truth and the subject come together

T_{32} : Category one of neither supervenes over the other, as objectivism or subjectivism would respectively

entail

T_{33} : Category two of neither supervenes over the other, as objectivism or subjectivism would respectively entail

T_{34} : Category three of neither supervenes over the other, as objectivism or subjectivism would respectively entail

Module seven

There is an identity of being and thinking, but only **via (e&eb)** events and the decisions immanent to them – they are co-engendered through one process

G_{36} : Category one of identity of being and thinking

G_{37} : Category two of identity of being and thinking

G_{38} : Category three of identity of being and thinking

T_{36} : Category one of events and the decisions immanent to them – they are co-engendered through one process

T_{37} : Category two of events and the decisions immanent to them – they are co-engendered through one process

T_{38} : Category three of events and the decisions immanent to them – they are co-engendered through one process

Module eight (case two)

G_{40} : Category one of events and the decisions immanent to them – they are co-engendered through one process

G_{41} : Category two of events and the decisions immanent to them – they are co-engendered through one process

G_{42} : Category three of events and the decisions immanent to them – they are co-engendered through one process

T_{40} : Category one of identity of being and thinking

T_{41} : Category two of identity of being and thinking

T_{42} : Category three of identity of being and thinking

Module Nine

Qua a non-processual state, the subject is indeed **transcended (eb)by** truth –'every truth is transcendent to the subject'

G_{44} : Category one of subject is indeed **transcended (eb)by** truth –'every truth is transcendent to the subject'

G_{45} : Category two of subject is indeed **transcended (eb)by** truth –'every truth is transcendent to the subject'

G_{46} : Category three of subject is indeed **transcended (eb)by** truth –'every truth is transcendent to the subject'

T_{44} : Category one of Qua a non-processual state

T_{45} : Category two of Qua a non-processual state

T_{46} : Category three of Qua a non-processual state

SECTION EIGHT

Subject And Event(Badiou And Deleuze): Sed Ipse Spiritus Postulat Pro Nobis, Gemitibus Inenarrabilibus Model

INTRODUCTION—VARIABLES USED

Badiou and Deleuze John Mullarkey

(1) In Badiou's eyes, this(Qua a non-processual state, the subject is indeed **transcended (eb)by** truth – 'every truth is transcendent to the subject') reverses the problematic of subjectivity: it is no longer the question of how an ineluctable subject is **made (=) to tally** with Being(how does the Cartesian subject represent the world?), nor how Being generates subjectivity or some kind of naturalized self is produced by matter,('the brain becomes subject', 'souls are everywhere in matter', says Deleuze)
(2) How subject and event (or truth), come into being together as **(=) precisely** what ruptures being through a process of naming.
(3) This return to the subject, though clearly not reinstalling it as a 'self-identical substance', nonetheless **refuses(e)to** diffuse the subject (as Deleuze does) by immersion ina differential or libidinal ontology.
(4) The subject is immanently co-authored **with (e&eb) t**he truth that it names and remains faithful to.
(5) The subject is now a 'fragment of the process of truth'; a post-Cartesian and post-Lacanian thesis that **distances (eb)** the subject from any substance, but without either reducing or eliminating it.
(6) Deleuze, by contrast, must be regarded as part of the post-structuralist paradigm that did its utmost **to(e) the centre** the self, though in his case by looking to the differential forces of matter rather than language as most others did (Barthes, Derrida et al)
(7) 'One does not ask how the subject gains its experience', Deleuze wrote, 'but how experience **gives (eb) us a** subject'.

Logics of Worlds

(8) To speak more directly about truth – and against the litany of attacks on truth that constitutes most of Twentieth-Century philosophy according to Badiou – the central idea for him is, again, that truths **and** subjects come **together so that** the truth of a multiple is produced immanently to this multiple, with a subject's intervention.
(9) That is why there is no need for a dogmatism of static correspondence (between a representing subject

and 'the' world) to re-assert the claims of truth. For Badiou, then, truth is its own standard (it 'is the proof of itself'), for as new, the true simply **cannot (e)** correspond to anything (to any previously existing standard), but 'depends on its own production'.

NOTATION

Module One

In Badiou's eyes, this(Qua a non-processual state, the subject is indeed **transcended (eb)by** truth –'every truth is transcendent to the subject') reverses the problematic of subjectivity: it is no longer the question of how an ineluctable subject is **made to tally** with Being(how does the Cartesian subject represent the world?), nor how Being generates subjectivity or some kind of naturalized self is produced by matter,('the brain becomes subject', 'souls are everywhere in matter', says Deleuze)

G_{13} : Category one of **tally** with Being(how does the Cartesian subject represent the world?), nor how Being generates subjectivity or some kind of naturalized self is produced by matter,('the brain becomes subject', 'souls are everywhere in matter', says Deleuze)

G_{14} : Category two of **tally** with Being(how does the Cartesian subject represent the world?), nor how Being generates subjectivity or some kind of naturalized self is produced by matter,('the brain becomes subject', 'souls are everywhere in matter', says Deleuze)

G_{15} : Category three of **tally** with Being(how does the Cartesian subject represent the world?), nor how Being generates subjectivity or some kind of naturalized self is produced by matter,('the brain becomes subject', 'souls are everywhere in matter', says Deleuze)

T_{13} : Category one of Qua a non-processual state, the subject is indeed **transcended (eb)by** truth –'every truth is transcendent to the subject') reverses the problematic of subjectivity: it is no longer the question of how an ineluctable subject

T_{14} : Category two of Qua a non-processual state, the subject is indeed **transcended (eb)by** truth –'every truth is transcendent to the subject') reverses the problematic of subjectivity: it is no longer the question of how an ineluctable subject

T_{15} : Category three of Qua a non-processual state, the subject is indeed **transcended (eb)by** truth –'every truth is transcendent to the subject') reverses the problematic of subjectivity: it is no longer the question of how an ineluctable subject

Module Two

How subject and event (or truth), come into being together as **precisely** what ruptures being through a process of naming.

G_{16} : Category one of **precisely** what ruptures being through a process of naming.

G_{17} : Category two of **precisely** what ruptures being through a process of naming.

G_{18} : Category three of **precisely** what ruptures being through a process of naming.

T_{16} : Category one of subject and event (or truth), come into being together

T_{17} : Category two of subject and event (or truth), come into being together

T_{18} : Category three of subject and event (or truth), come into being together

Module three

This return to the subject, though clearly not reinstalling it as a 'self-identical substance', nonetheless **refuses(e)to** diffuse the subject (as Deleuze does) by immersion ina differential or libidinal ontology

An ethics of the pre-individual O'Donnell, Aislinn (2001) An ethics of the pre-individual. PhD thesis, University of Warwick. We quote the abstract of the thesis which clearly expresses the necessity of development of an alternative ontology of becoming corresponding to a philosophy of difference.

Deleuze opposes ethics to Morality. He claims that an ethics develops immanent criteria to evaluate modes of existence, while a Morality imposes transcendent principles. This thesis explores the question of ethics, and I investigate the possibilities of an ethics of the pre-individual. Consequently this enterprise involves the development of an alternative ontology of becoming corresponding to a philosophy of difference. By taking this trajectory, I seek to show that an anti-human humanism is possible and demonstrate how this might work.

Deleuze and Guattari always emphasise the practical and concrete nature of philosophy; therefore, in order to situate their concepts I begin the thesis with an examination of different theoretical approaches to the question of difference. However, I suggest that difference and heterogeneity cannot be simply affirmed in and of themselves **since new forms of domination** also affirm difference. My next chapter follows up on this idea by interrogating the allegation that philosophies of difference have made a political covenant with global capitalism. I draw on a distinction between power (Potentia) and Power (Potestas) in order to explain how different modes of **social organisation and domination** can minimise the creative and transformational capacities of humans. By analysing a number of theoretical accounts of capitalism I demonstrate how and why it differs from other social formations. Nonetheless, I conclude that philosophy can indeed be distinguished from capitalism. Philosophy, as the art of inventing concepts, develops the conditions for real experimentation and new ways of thinking, being and existing.

By turning to Spinoza's Ethics I propose that by thinking about the human differentially, as a part of nature, we can develop an imminent ethics. I explain how Spinoza's ontology operates especially in terms of its renovated conception of the human. In generating an ontology that is not centred on the individuated individual but grasps instead the individual as both relational and a degree of power, the pre-individual and transindividual dimensions of the human are emphasised and she is opened up to her non-human becomings.
Simondon's account of metastable being explores this in greater detail. He argues that we have tended to extrapolate from the individuated individual in order to try to understand its conditions of existence. Alternatively, we have relied upon a principle of individuation that pre-exists the process of individuation. By intertwining his focus on the process of individuation with his idea that being is more than unity, more than identity and fundamentally incompatible with itself, I present Simondon's account of ontology of becoming and his correlative conception of a pre-individual field. Residing at the core of his endeavour is a theory of difference and disparateness that understands identity to be emergent, partial, relative and derivative.
Simondon's emphasis on disparateness recurs in Deleuze's work 'Difference and Repetition' mobilise this idea in order to distinguish between a created possible and a realisable possible, and to elucidate the ethico-political implications of this distinction. The concept of the 'image of thought' that rests on a series of non-philosophical pre-suppositions helps us to critique dominant modes of thinking and acting. In addition to critique, I seek to construct other ways of thinking and existing. Once again I focus upon the preindividual and transindividual dimensions of the human when in my concluding chapter I map the different conceptions of ethics and subjectivity that emerge once we transform our understanding of ontology. An ethics of the pre-individual relies on immanent criteria for evaluating modes of existence, does not fetishise the human, and ultimately constructs the possibility, of things being otherwise.

Library of Congress Subject Headings (LCSH): Deleuze, Gilles, 1925-1995, Guattari, Félix, 1930-1992, Spinoza, Benedictus de, 1632-1677, Ethics, Comparative, Philosophy, Comparative, Immanence (Philosophy)Supervisor(s)/Advisor: Ansell-Pearson, Keith, 1960-

G_{20} : Category one of diffuse the subject (as Deleuze does) by immersion ina differential or libidinal ontology

G_{21} : Category two of diffuse the subject (as Deleuze does) by immersion ina differential or libidinal ontology

G_{22} : Category three of diffuse the subject (as Deleuze does) by immersion ina differential or libidinal ontology

T_{20} : Category one of return to the subject, though clearly not reinstalling it as a 'self-identical substance'

T_{21} : Category two of return to the subject, though clearly not reinstalling it as a 'self-identical substance'

T_{22} : Category three of return to the subject, though clearly not reinstalling it as a 'self-identical substance'

Module four

The subject is immanently co-authored **with (e&eb) t**he truth that it names and remains faithful to

G_{24} : Category one of subject is immanently co-authored

G_{25} : Category two of subject is immanently co-authored

G_{26} : Category three of subject is immanently co-authored

T_{24} : Category one of truth that it names and remains faithful to (say like Higgs Boson being searched by subjects at CERN)

T_{25} : Category two of truth that it names and remains faithful to

T_{26} : Category three of truth that it names and remains faithful to

Module five

The subject is now a 'fragment of the process of truth'; a post-Cartesian and post-Lacanian thesis that **distances (eb)** the subject from any substance, but without either reducing or eliminating it.

Note: reader's attention is drawn to "Futures Research Methodology – V 2.0" report. Sense making is presented as a possible framework for achieving this objective, and supplements the efforts already invested in this direction by the "Futures Research Methodology – V 2.0". The rationale for this supplement will also be presented. When applied the "Futures Research Methodology – V 2.0" shows, in general, some interesting features characteristic of the implementation of knowledge systems. In particular, the emergence of effects that is beyond the initial – but not the final – scope of the system. Sense making addresses the challenge of understanding and retrieving such dissipation from the knowledge system, and aims to bring it within a manageable range. In this way the article particularly addresses the shift that results from understanding and managing complexity. It puts forth a demonstration of these effects on futures research methods, and shows how foresight and hindsight may be used to gain insight into the present How Do We Make Sense of the Future? **An Analysis of Futures Research Methodology – V2.0 Mika Aaltonen**

G_{28} : Category one of subject is now a 'fragment of the process of truth'; a post-Cartesian and post-Lacanian thesis

G_{29} : Category two of subject is now a 'fragment of the process of truth'; a post-Cartesian and post-Lacanian thesis

G_{30} : Category three of subject is now a 'fragment of the process of truth'; a post-Cartesian and post-

Lacanian thesis

T_{28} : Category one of **distances (eb)** the subject from any substance, but without either reducing or eliminating it.

T_{29} : Category two of **distances (eb)** the subject from any substance, but without either reducing or eliminating it.

T_{30} : Category three of distances **(eb)** the subject from any substance, but without either reducing or eliminating it.

Module six

Deleuze, by contrast, must be regarded as part of the post-structuralist paradigm that did its utmost **to(e) the centre** the self, though in his case by looking to the differential forces of matter rather than language as most others did (Barthes, Derrida et al)

Note: **Back to Enquiry Learning Back to Home Addressing Methodologies. Tracing the Self In Significant Slips: Shadow Dancing. Volume 2 Addendum Jill Schostak 2005. This beautiful work throws light on some of the problems that are being encountered by quantum physics. It is reproduced in extenso. Emphasis mine. Some paragraphs and references are deleted due to spatial constraints.**

Addendum reviews the literature on "intertextuality" and "meaning-slips" with, beside and against Jacques Derrida's notion of "écriture" **to open up a space to** address the doing, the theorizing and the effects of the performativity of 'intertextuality' and "meaning-slips". It thus elaborates one of a number of exploratory resources for the reading and re-writing of Covers – in the de-signer space and In-Between Slips that constitute the experiment in writing. However, this elaboration is not at all to suggest a) a preface/pre-figuration function of Addendum, nor that b) a review of intertextuality and meaning-slips is the only approach – others will be alluded to as appropriate and picked up again in the concluding discussion of Addendum and finally nor is it c) to suggest it is the Last word on the experiment in writing. Like Derrida 'I absolutely refuse a discourse that would assign me a single code, a single language game, a single context, a single situation; and I claim this right not simply out of caprice or because it is to my taste, but for ethical and political reasons' (Derrida. 1996: 81).

Firstly then, an initial brief look at écriture is helpful before turning to review and discuss intertextuality and meaning-slips. Écriture is writing in a broader sense than the script produced on paper by whatever means, hand or other. It is a metaphor, a figure for 'an entire structure of investigation, not merely "writing in the narrow sense," graphic notation on a tangible material' (Spivak. 1976: ix-lxxxix). Rather it is 'the constitution of a thick space where the play of **hiding/revealing** may take place' (Lyotard. 1971: 75) , the space of the secret and the imperceptible. Neither irreducible to a series of rules on the investigations of graphic systems nor a simple opposition to speech in order to invert a binary opposition, it announces a rhetoric of identity situated in some physical context (Wolfreys, 1998). Not only does the notion of writing refer to speech and thought as forms of writing, but it is also expanded along further horizons to include the writing, the written-ness, of the subject's identity (Wolfreys. 1998)

Applying this academic/writer has to distance her/himself and others not just from a language, a particular discourse, but from the language itself, **as a** pre-existing classification of the world. Thus to intervene in the world through writing is not to just add to the existing mass of linguistic production. Rather a working-over of language will detach it from prevailing codes **and stop it from reproducing** a pre-given structuring of the world, and thereby set going certain in directedness, since the world is intervened in only through an

intervention in language itself. Thus, Barthes views the intertext as

What comes to me, not what I summon up; not an "authority," simply a circular memory. Which is what the inter-text is: the impossibility of living outside the infinite text?

Barthes. 1976: 36

The issue then is not that of "authority", but rather that of 'a circular memory' and of 'the impossibility of living outside' of context. Instantly within this 'infinite text', echoes between Barthes's comments in the citation above and Derrida's 'il n'y a pas de hors de texte' [see Addendum: 67] are set circulating. What-is-more, the English Oxford Reference Dictionary interestingly ascribes to the word 'cite' meanings of 'frequent' and 'set moving'. Barthes himself employs a phrase incorporating a notion of set moving [see Endnote 15] and I deploy this as a Kristevan-like minimal textual unit [see Addendum: 13] to mark the iterability of Barthes's circular memory in Derridean fashion: as inter-text, no less and the impossibility of 'living outside' of context. Thus Barthes's statement provides a kind of 'preface', a parergon as Derrida might call it, a Lyotard-like thick space for the work of Addendum which focuses on four key "primary" authors, namely, Roland Barthes, Julia Kristeva, Jacques Derrida and Michael Riffaterre. Some additional reference is made to the work of Jean-François Lyotard as a resource for discussing issues of meaning-slippage and to further explore what is at stake in the debates within the review. Each author is the occasion of a circular reading, as Barthes's 'circular memory' comes into force illustrating the circularity, in the sense of circulation/set going rather than circling, involved in reading and hence in writing.

Moreover, it is also informed by the works of Roth (1988), (Buchanon, (2000), Rabaté, (2002), Baugh, (2003) and Davis, (2004), who point out that, for the most part, the work of French post-structuralists can be viewed as a response or reaction to Hegel, and also to structuralism which eschewed Hegelianism in its search for a scientific model. In referring to Hegel within the context of Addendum, I speak of a "French Hegel" who emerged from the interpretations given to Hegel's Phenomenology of Spirit by Kojève in his famous lectures and also by Hyppolite, both of whom, incidentally, might be described as primary "authors" for the career trajectories of many of the French post-structuralists (for Foucault, Lacan, and Derrida, to name but three). These **post-structuralists respond and react by reading with, beside and against a) the works of Hegel, and also b) that of structuralism and scientism both of which are totalizing systems.**

Despite the diversity of the projects of the various post-structuralists, commonly held purposes and strategies can be identified. Their collective avant-garde intention **was to undermine and disturb** Authority and to challenge any form of totalizing system that might **give rise to** Absolute Knowledge and Absolute Reason or Science. They set out to develop revolutionary ways of thinking that would keep systems open-ended, heterogeneous and dynamic, **thus clearing the way** to alternative playful interpretations, alternative ways of seeing, being and working. **This led to** notions of reading/writing undergoing significant changes. Traditional notions of the Author, of the reader, and of the Text, and thus by implication of truth and body, were called into question. For example, Kristeva comments in Oliver's The Portable Kristeva, For us, structuralism … was already accepted knowledge. […] From the outset, however, our task was to take this acquired knowledge and immediately do something else.

Oliver. 1997: 9

Setting out to 'do something else' led to strategies involving

Desire/ intent to disturb, disrupt, destabilize, unsettle, to shock

Desire/ intent to create and maintain the fluently elusive and/or the fluidly ephemeral

Desire/ intent for open and dynamic systems

The danger Kristeva, Barthes, Derrida and others sought to avoid was **that of "interiorizing" conflict** in order to reduce it and stabilize it, which in turn, thus effectively opposed and resisted change. In other words, all three set out to do something other than sign up to following the model of a Hegelian strategy which led to Absolute Reason. It is thus within this set of concerns that the key terms of this Addendum take their significance. Intertextuality is one way of thinking how to undermine the totalitarian grip of Reason.

In Barthes's words, every text holds the intertextual, itself being 'the text-between' of another text, quite different to its "sources", but nevertheless, **marking the "influences**", falling in with 'the myth of filiation', even whilst of and in citations that are 'anonymous, untraceable and yet already read: they are quotations without inverted commas' (Barthes. 1977: 160). Earlier echoes between Barthes and Derrida were made [Addendum: 3]. Similarly with Riffaterre who posits a hidden inter-text which gives meaning to the (poetic) text currently being read, as will be discussed in detail later [Addendum: 73-81]. The 'circular memory' is already at play (Barthes. 1976: 36). But first, a distinction is required in order to describe further what is at stake in intertextuality.

Is intertextuality distinct from source studies and from influence studies? Traditionally, in literary studies, the scholar had continued to search after the 'reliable text' despite knowing that there is no fixed meaning since the historical appropriation of meaning cannot be halted. Indeed, the drive for 'actual' meaning remains over-ridingly powerful and insistent (Mai, 1991). Kristeva, however, envisioned a '**new kind of hermeneutics'** (Mai. 1991: 43), one which **propogated numerous conceptual frameworks and models (Ruprecht, 1991), through the juxta-position of its intertexts, which create forces of tension and conflict between the "source" and the new (Muller, 1991).** Kristeva is adamant that intertextuality is distinct and does not elide/compete with influence or sources, not by imitation nor by quotation. In fact, so keen is she to avoid the reduction of intertextuality to the traditional notions of influence, source-study and simple 'context' that, for these very reasons, she introduced the term 'transposition' (Kristeva. 1984).

The term inter-textuality denotes the transposition of one (or several) sign system[s] into another; but since this term has often been understood in the banal sense of "study of sources", we prefer the term transposition because it specifies that the passage from one signifying system to another demands a new articulation of the thetic – **of enunciative and denotative positionality**.

Bakhtin's emphasis on the historical and the social aspects of the utterance opens up a space for the reader/writer and stands in stark contrast to the sign system of **Saussure's structural linguistics**. Saussurean structuralism effaces the writer/reader through a system of arbitrary signs that are also differential 'without positive terms' (Saussure, 1974: 120 cited in Allen, 2000). This linguistic system pre-exists any speaker signaling that the social focus of language **simply does not exist for** Saussure, for whom words are only relational with respect to their position within abstract systems.

For Kristeva, a speaking being is crucial for 'understanding oral and written literature, politics, national identity, sexuality, culture and nature' (McAfee. 2004: 1). Thus it is crucial for a Kristevan understanding of intertextuality. Far from being separate domains, 'the speaking being is a "strange fold" between them all', where inner drives interject with language, sexuality **plays with** thought, where body and culture meet, (McAfee. 2004: 1). Kristeva calls this speaking being the **"speaking subject**". It is split between the conscious and the unconscious, reason and desire, the communicable and the incommunicable. Such a positioning begins to open up that magical milieu, invoking that mystical space wherein 'a fact is narrated no longer with a view to acting directly on reality but intransitively' (Barthes. 1977: 142). This milieu, this sudden transformative space, wherein writing does not act directly on reality, since it is 'the very practice of the symbol itself', sets going a disconnection, voice loses its origin, and 'intertextuality' later emerges

(Barthes. 1977: 142). Saussurean language after Kristeva is 'as a system' that is 'articulated through the signifier **which exceeds the** consciousness (and therefore systematization) of the speaking subject' (Kristeva. 1985: 210).

Wherever 'the subject is implicated, whether body or history, or symbolic order, reason, intelligibility', desire arises (Kristeva. 1980: 116). Kristeva borrows the symbolic order from Lacan, but then turns aside from the Lacanian Imaginary, looking instead to Freud's work on 'primary process' and the pre-symbolic stage of the child. A notion 'of the semiotic emerges, characterized by pre-symbolic drives, impulses, bodily 'pulsions' (rhythms and movements). Kristeva's semiotic is '**an articulation of unconscious processes which fracture the common idealization of those images and signs which secure the status quo, and guarantee the establishment' (Smith. 1998: 16).** It constantly and subversively threatens 'the symbolic order of things which Kristeva herself stresses', in case we should have assumed otherwise, '**is no monolithic structure, but an illusion of stability'** (Smith. 1998: 16). Although Kristeva's semiotic is inextricably linked to the body it is at the point of the thetic phase that the human subject enters the social world/the real/the symbolic/father/ phallus.

Poetic analysis.

And it is here at this very intersection that I anacoluthonally locate the parerga-figural [see Endnotes 1, 5, 6] of a) the 'Fig' with, beside and against the other 'Figs' of In-Between Slips, b) Covers – in the de-signer space with, beside and against In-Between Slips and c) In-Between Slips with, beside and against Covers – in the de-signer space and therein/of set going an "**oblique translation**", a SlanTr a poetic analysis – an enmeshing, the experiment in writing, no less. Hence poetic analysis and its relation to intertextuality and meaning-slips is central to Addendum in its function as one of a number of exploratory resources reading with, beside and against Covers – in the de-signer space and In-Between Slips.

With Kristevan semiotics and her subject split between two signifying fields, (the semiotic and the symbolic [to be elaborated below]) the mediation at work here through the dynamic and through heterogeneity transforms and transposes, opening up rather than forcing closure. The semiotic is likened to pre-speech infant babble that continues with us even though on another level we [apparently] lose [sight of] it when we enter the symbolic signifying field. **Central focuses of the semiotic is the chora, and through it, our 'infant' pre-speech fluidity of self and poetic language bubble up to ceaselessly disturb the monologic order of the symbolic. Poetic language whilst sited in the symbolic thus remains forever shot through with traces of the semiotic. Kristeva argues for more than grammar, syntax, or vocabulary, since 'sensation will leave its indelible stamp and that this imprint** of the body in language is readable' (Smith. 1998: 14). What might such an event look like? A brief recourse to Lyotard's notion of a reading event is helpful here.

The semiotic, Kristeva insists, **is not an extension of the language system but transversal to and coextensive with it. It is through the semiotic that we can connect language as a formal system to something outside this, in the realm of the psycho-somatic, to a body** and a bodily subject structuring and de-structuring identity' (Smith. 1998: 18).

In other words, notions of inside/outside are unsettled as slippage and meaning-slips may pass, or traverse or transform. But are there times when they may crash? Are meaning-slips now trapped in performing closure or are they keeping the system open? What about the imprint of language in body? Aware that her theory of semiotics is a paradox, **an aporia, as is her notion of the speaking subject**, I turn now to Kristeva's notion of the subject on trial/in process - the sujet-en-procès - in order to explore further the possible implications for the effects of such slippages.

Since discourse, for Kristeva, is a practice between speaking subjects and a system (language), 'the field of sense and signification beyond linguistics' is opened up towards the sociological and the psychological such that systems of meaning emerge based on the socio-historical and the intersubjective (Kristeva. 1985: 211). For the most part, we are locked into a semiotics of "ideologies" – as myths, rituals, and moral codes become sign systems comprising the law governing any social practice. Every social practice is therefore embedded in an order of language and thus has a double articulation (signifier/signified), a duality which stands in an arbitrary relation to the referent (R) such that 'all social functioning is marked by the split between the referent and the symbolic and by the shift from signified to signified coextensive with that' (Kristeva. 1985: **212). In other words, the speaking subject is now a process, 'simultaneously a unity and the transition to zero of this unity' (Kristeva. 1985: 213). Moreover, for Kristeva, the speaking subject is always already in a state of crisis as this 'is a permanent state of functioning' (Kristeva. 1989: 37). Not as anarchist in a negative or pejorative sense as it seems at first glance, Kristeva explains that, for her, the word 'anarchy' can have** creative possibilities and that is exactly what is in play here.

This state of crisis **between unity and zero of that unity then, is inherent in the speaking subject and necessary to postulate that the position of conscious mastery of meaning by the subject constitutes and guarantees the subject's unity, not so much as a fixed totalizing totality but rather an** 'if only' for the moment. The notion of mastery pre-supposes relativity between various types of signifying experiences and the limit of this is the zero state. However, the relationship between the referent (R) as object/situation and the signifiable 'is never a relation of identity, but rather one of displacement', of 'contradiction' (Kristeva. 1985: 214). Indeed, this is a 'relationship of emptiness, of the unnameable, of non-sense and of opaque experience which we would designate by the word increasingly haunting semiological theories – the body' (Kristeva. 1985: 214).

Thus the Kristevan speaking subject is a traversal beyond both the bounds of the Saussurean abstract system, and those aspects **of Bakhtin/Volosinov, which privilege the social specificity of language and thus view it 'always in a 'ceaseless flow of becoming', (Allen. 2000: 18). Invested with poetic language, Kristevan intertextuality embodies otherness** and thus situates itself against monologism, rejoicing in the poetic.

When she discussed her reading with Roland Barthes, her supervisor, he was 'quite fascinated' and invited her to do a seminar on Bakhtin in 1966, (Waller. 1985: web-site: 19 Dec 2003). Thus Bakhtin is eliding gradually but ineluctably into a Kristevan Bakhtin. Which is not the same as saying the meaning slips have reached monumental proportions, whatever that might mean, rather it is to say that Bakhtin-as-reading/writing undergoes a transposition? the meaning-slips inherent in the transposition, what **is the nature of this being 'faithful'? It can not be a faithful copy, but rather of the ilk of one of Dunkelsbuhler's SlanTrs and/or 'wandering translation[s] [traduction errante]' (Dunkelsbuhler. 2002: 59), constituting a parergon as creating a framing - boundaries of honouring and paying homage to a master, whose work is worthy** of being elevated to a position deserving of due care and attention. Yet also within this ilk is Derrida's 'la tournette d'une derive interne' – the letter which does not always arrive at its destination and the fact that it holds to this possibility as part of its structure afflicts it with an internal misdirection (Derrida 1987) or destinerrance (see In-Between Slips: Fig 2: 286). And yet, as seen already above, intertextuality problematizes the notion of Author and of origin.

Kristeva warms to her theme. Although Bakhtin's work exhibits two axes, 'dialogue and ambivalence', these are not 'clearly distinguished' (Kristeva. 1980: 66). The horizontal axis, dialogue, refers to the subject-addressee; the vertical axis, ambivalence, to the text-context, and it is here that the intersection for transposition opens up to a Kristevan poetic analysis. **The tensions of author-reader/reader-writer remain clearly evident.**

However, Kristeva would maintain that what she brings to the reading of Bakhtin is psychoanalysis. Indeed, she confirms that psychoanalysis – 'the only continent that we had never left' (Oliver. 1997: 19) - remains at the heart of her project. Thus, in the Waller interview, Kristeva comments that 'the notion', her notion that is, 'that the participation of different **texts at different levels reveals a particular mental activity' was 'unique' (Waller. 1985: web-site: 19 Dec. 2003).**

Picking up the loose thread of the carnival à la poetic word, the 'carnival participant is both actor and spectator' passing '**through a zero point of carnivalesque activity' and** splitting into 'a subject of the spectacle and an object of the game', and 'sees itself created as self and other, as man and mask' (Kristeva. 1980: 78). Signifiers refer to signifiers in the Saussurean system of language, such that 'this empty and yet significant place called the zero degree of the sign' (Kristeva. 2000: 214) where writing has become 'absence' is called **'the zero degree of writing'** (Barthes. 1967: 11). However, in the process of signification, as if in a game of chess, signifier checks signified to zero point, and the body falls out as a remainder – the letter kills. Via flows and transformative forces the game/play unfolds where-in/of designs form the basis of evoked self/other relationships. Or to take a Barthesian slant 'the pleasure of writing' makes its debut mediated by the body that desires thereby opening up 'the infinite pleasure of the text' as discussed later in Addendum (Kristeva. 2000: 215). The transition **to zero and thus 'loss of identity, afflux of drive and a return of symbolic capacities' marks the collapse of the unified "Transcendental ego" (Kristeva. 1985: 217). This shift allows the speaking subject the capacity to renew** the order in which s/he is trapped and for the subject this constitutes the capacity of enjoyment. Rather, intertextuality announces 'an intrapsychic or psychoanalytic finding', 'concerning the status of the "creator," the one who produces a text by placing himself or herself at the intersection of this plurality of texts on their very different levels' (Waller. 1985: web-site: 19 Dec. 2003). Kristeva has moved from recognising Bakhtin as author[ity] to Bakhtin-reader to Bakhtin-rewriter. But is she reader-rewriter-cum-author[ity] of her own conceptualization?

Continuing her story, she explains, 'I myself speak of a "subject-in-process" which makes possible my attempt to articulate' 'a logic' 'between identity and unity' mediated through 'its reduction to zero', that 'moment of crisis, of emptiness' that is the pre-lude to 'the reconstitution of a new plural identity' – this dynamic holds for both writer and reader (Waller. 1985: web-site: 19 Dec. 2003) [my italics].

Reference to Freud's speechlessness constitutes the very heart of Covers – in the de-signer space and In-Between Slips, with the emphasis on reading, on writing, on the written mark rather than the phonetic mark, tracing a sensate shadow dancing in its undecidability both-or-and as-open[ing]. 'Open', its link to the Barthesian body and/or desire is palpable. Kristeva writes of how 'Barthes threw himself into interpretative writing', 'with a sense of going against the natural' thereby revealing 'the body beneath meaning' which, for him, '**always shimmered on the horizon' as glistening 'mirage' (Kristeva. 2000: 190). In a society driven by capitalist notions, production and consumption are separated, hence author and reader became divided. But Barthes adamantly proclaims bodies desire to write, and in particular they desire to write the scriptible, not** least because the ensuing nausea from the reading of lisible scripts shows one how things could be otherwise. Closed, 'oppressed in the order of sign itself', vomit spews forth (Barthes. 1975: 139). Nevertheless, the body is not unproblematic, since as will be discussed later a notion of warfare in language is a central motif for Barthes. The relation of the body to culture is thus highly complex. To desire or to be desired is always to be trapped in a process of repetition and quotation – '[w]ithout Book, without the Code … no desire' (Barthes. 1975: 73). Desire is always already compromised. Desire where dictated by culture as doxa results in nothing but a strangulating process of repetition, suffocation by the stereotype. **The body collapses the anti-thesis of the symbolic, divorces desire from culture, but in so doing pays the price of death of the subject – ghost-like, it falls out as remainder as in Kristeva's**

game-play.

The move' (Spivak. 1976: lxxi). Forever in a vocabulary on the move, and weaving in the movement of supplementarity of Derrida's identity with the **Saussure of differences without positive terms, for Derrida '[s]pacing designates nothing,' but yet 'of a movement**, a displacement that indicates an irreducible alterity' (Derrida. 1981: 81).

'[F]eet on a tight-rope' (Cixous. 1998: 57), well aware of running the risk of "mastering" 'deconstruction' and/or différance and/or spacing and others in the nonsynonomous chain of substitution Derrida suggests the 'concept of spacing by itself thus cannot account for anything, any more than any other concept' (Derrida. 1981: 81). Thus through the play of writing and of différance, Derrida's dividing line splits and splits again; strands spin apart and then together, and weave in other threads; threads of other binaries such as presence/absence, inside/outside, same/other, us/them are called into question. 'Forever unable to saturate a context, what reading will ever master the "on" of living on?' (Derrida. 1979: 76-7). '**This is not word play, not on your life'** (Derrida. 1979: 77). A choreography of presencing/absencing, of spell-binding spaces and lace-ings shadow dancing - splitting, spinning apart, spinning together, self-ing - enmeshes the telling spaces of Covers – in the de-signer space and In-Between Slips – **'"I" hold life in my hands', I write in Slipping into Beginnings: 38**.

Although we are still "same" and "self" as previously, suddenly, apparently same and self are uncertain, foreign and other, inaccessible to ourselves. How are we now positioned in relating to the other when, in fact, we discover we are foreign to ourselves? Derrida suggests lending an ear and listening to the other such that one is taught/attempts '"to hear with [one's] eyes" too' (Derrida. 1982: xiii). Physiologically speaking, the tympanic membrane, that 'figure of the oblique', positioned between the outer and middle ear functions as both limit and passage for transmission of vibrations (Derrida. 1982: xiv). The fact **that it is oblique significantly** increases the surface area for drumming thereby facilitating the act of listening. The effect is to render one available-yet-vulnerable since the other's drumming will resonate within us or fall on deaf ears.

The former **unified system, now equivocal, suggestive of hierarchical configurations, undergoing incessant struggle, gives rise to a double reading and a double writing, thus a-voiding any fencing-in dialectically constructed synthesis as "One". Pirouetting, it 'turns around the points of a ballerina, analyzes "the syntax of point [none**] and step [not] [la syntaxe du point et du pas]' to tell 'how "each pair, in this circuit will always have referred to some other, signifying too the operation of signifying"' (Derrida. 1987: 264).

'Determination' of différance.

I turn now to set différance working to illustrate its 'determination'. Différance, neither word nor concept 'is not, does not exist' having 'neither existence nor essence' (Derrida. 1982: 6), being 'neither simply active nor simply passive' (Derrida. 1982: 12), '**is no more an effect than it has a cause'** (Derrida. 1982: 12). Différance thus announces itself as 'unthinkable' (Derrida. 1982: 19), and 'unnameable' (Derrida. 1982: 26) since any words which could conceivably describe it already involve concepts such as being and presence, and these are the very concepts it seeks to undermine. '[A]rchi-themes' (Culler. 1983: 212), that are not so much a structure of a theme, but rather pertain to a textual structure/process/ a structurality emerge parergonally. **Thus a theme of supplementarity cannot itself be substituted because 'it happens that this theme describes the chain itself, the being-chain of a textual chain, the structure of substitution' (Derrida. 1976: 163; underlined in the original**). Elevation to totalization is thus impossible – différance

lacks a 'master-word' as previously discussed (Derrida. 1981: 40) being middle but no halfway house (Llewelyn, 1986, Gasché, 1986). **This is a beautiful stability analysis of the problem Deleuzean analysis poses us.**

Gasché (1986) refers to the chain of **substitutions** as an '**infrastructure'** as well as to différance, supplement, arche-writing and so on. However, the term 'infrastructure' has connotations of 'hierarchy [which] detract[s] from its pertinence' (Hobson. 1998: 239n: 31). Gasché, himself, (1986, 1995) notes the difficulties that follow from his choice of term. Yet other words such as 'quasi-transcendentals (Derrida, 1988, 1991, 1996; Gasché, 1986) and 'lexemes' (Hobson, 1998) also seem inappropriate. In Addendum, however, I am continuing to use the word 'infrastructure' keeping to its sense of 'structurality of structure' rather than its sense of 'basic' structure. Infrastructures are 'instances of an intermediary discourse concerned with a middle in which the differends are suspended and preserved' (Gasché. 1986: 151). This signifying **structure, structure signifiante (Derrida. 1976: 158), knots together the discrepancies and differences in question toward their synthesizing infrastructure that is intrinsic to the text, whatever that might be. Thus there is 'an interlacing, a weaving, or a web, which would allow the different threads and different lines of sense or force to separate again, as being ready to bind others together' (Derrida. 1973: 132).**

Infrastructurally, Derrida's dividing lines then, divide, deferring**, differentiating spatially and temporally. 'To each his Other, which is his Same. Or the I is two: by definition'** (Derrida. 1982: 288). Deferring, differentiating, rather like an ekphrastic impulse, a rhythm drives Derrida's thoughts, textual through and through, 'the figure is never one. Not only is it the Other, but there is no unity or stability of the figural', the 'subject desists' and 'its only chance of 'grasping itself' lies in introducing itself and oscillating between figure and figure' (Lacoue-Labarthe. 1998: 175), appearing/disappearing, hiding/revealing.

A textual system then is always already contaminated by the traces of other discourses and languages. Already the discourse resonates with previous discussion in this review of intertextuality and this resonance continues to build in strength. **The system is one of difference rather than one of presence. Infrastructurally, différance makes possible not only binary opposition and the double but also determines oneness since 'the subject is constituted only in being divided from itself'** (Derrida. 1981:29). For instance, let us take as our subject the present moment, which is neither present to itself, but related to a past or a future. For the present to be present to itself, an interval must separate present from itself dividing it in and of itself (Derrida, 1982). This interval of différance is the singular moment of presence; thus the supposedly singular is always already divided**. The movement which produces seeming unity is the same movement as those effecting binary oppositions. Différance works repeatedly to count as the same. It also receives the double mark becoming différance of différance**, the 'trace of the trace' (Derrida. 1982: 24).

A systematic questioning of the fixed, the separated, the totalitarian, and the programmatic discourses and figures takes place. Thus **algorithmic discourses** are put under the spotlight, as are discourses of mass consumerism that render one the 'plaything' of the desire of the other. For example, consumer discourses around the figure of the Barbie doll and 'her' iconicity are also discussed both in relation to her 'birth'/debut and her de-signer function as displayed in 'Dolly Mixture'. Whilst all the above criteria, in a sense, can be summarized and rendered explicit as being anti-totalitarian criteria, the algorithmic cannot. However, that is not to say **that algorithmic systems simply stand in opposition to non-algorithmic systems in Covers – in the de-signer space and In-Between Slips such that** dialectical logic brings the conflicts under Reason. Rather, it is to say, that, in a 'turning operation' (Bloom. 1979: 20), a reading from the bias and over-the-shoulder is set going, through différance, to 'turn[s] the system-maintaining and **system-generating** dialectic against the system itself' in a Derrida-like deconstructive manner (Zima. 2002: 7). Hence, reading from the

margins to unsettle and disturb [see all above criteria], maintaining an alert attention to totalitarian closure, by engaging with, beside and against such **struturalististic-like algorithms in a matrix-like/ parergonal/figural fashion, constitutes a criterion that provides a way of ensuring this list of criteria remains open. There is throughout the thesis a questioning of totalitarian discourses and** figures that can be seen in the example of Barbie as 'eidos' as 'ideal' as perfect repeatability of the 'same'. I show how marketing ploys by Mattel cash in on '[t]he ideality and invisibility of the eidos [that] are it's power-to-be-repeated. Now, law is always a law of repetition, and repetition is always submission to a law' (Derrida. 1981: 123); the law of 'must have', the law of 'must be like'. The issue then is to subvert the closure wrought by the law of the Master signifier. However there remains the **danger of 'falling back' into binaries whether willingly, cynically, reluctantly or kicking and screaming. This criterion speaks to the dynamics of an open system and the ever-present risks of an illusory yet powerful totalization**. The last 'criterion' leads to the problematic that there is no language, no text, and no discourse that is fully 'external' to Western Metaphysics that can 'successfully' avoid being drawn back into its totalitarian binaries.

This circle is unique. It describes **the form of the relation between the history of metaphysics and the destruction of the history of metaphysics. There is no sense in doing without the concepts of metaphysics in order to shake metaphysics. We have no language** – no syntax and no lexicon – which is foreign to this history …Derrida. 1978: 280

Conclusion.

Intertextuality, as we have seen, **unsettles by de-totalizing and de-unifying**. It is violence. It decentres and problematizes notions of identity **by moving boundaries**. Whilst it still "frames", it does so in a most perturbing manner. It still creates margins/verges/edges/frayings but in unpredictable ways that constantly surprise. Has it, in fact, rendered traditional criticism a thing of the past? Barthes thinks so. Certainly notions of Authorship, of origin, of unity and of identity, and of traditional critique are all challenged and unsettled. Hence the examples of 'criteria' for writing/ reading that emerged from Addendum as outlined in the preceding section. These explored the basis for 'assessing'the'successes of the experiment in writing.

Such criteria for 'judgement' then open up a space for a reading/writing to the effect that there exists a horizon of potential readings **that transcends** mimesis/literality but only if the text itself opens up in-between spaces that thus provide a set of indefinite positions in a matrix where readers can jostle for making a reading and an 'assessment' theirs as against the readings and 'assessments' of others. The telling space of the experiment in writing is the place where these potential meanings emerge as speaking subjects or as face to face positions. It is the space where what counts as a successful 'reading/writing' can be contested.

Glimmer or no, the reader/writer is that impossible but necessary **combination of interiority and exteriority. There is something narcissistic about the act of reading, something be-witching. Rather like watching a film where the eye perceives a continuum but is in point of fact actually 'seeing' a set of still frames/point capiton whilst the brain over-rides the gaps between the frames suturing it in hymen-like fashion to give a continuity without visible or noticeable scar tissue.**

G_{32} : Category one of **centre** the self, though in his case by looking to the differential forces of matter rather than language as most others did (Barthes, Derrida et al)

G_{33} : Category two of **centre** the self, though in his case by looking to the differential forces of matter rather than language as most others did (Barthes, Derrida et al)

G_{34} : Category three of **centre** the self, though in his case by looking to the differential forces of matter

rather than language as most others did (Barthes, Derrida et al)

T_{32} : Category one of post-structuralist paradigm

T_{33} : Category two of post-structuralist paradigm

T_{34} : Category three of post-structuralist paradigm

Module seven

'One does not ask how the subject gains its experience', Deleuze wrote, 'but how experience gives (eb) us a subject'.

G_{36} : Category one of 'One **does** not ask how the subject gains its experience', Deleuze wrote, 'but how experience

G_{37} : Category two of 'One does not ask how the subject gains its experience', Deleuze wrote, 'but how experience

G_{38} : Category three of 'One does not ask how the subject gains its experience', Deleuze wrote, 'but how experience

T_{36} : Category one of subjects

T_{37} : Category two of subjects

T_{38} : Category three of subjects

Module eight

Logics of Worlds

To speak more directly about truth – and against the litany of attacks on truth that constitutes most of Twentieth-Century philosophy according to Badiou – the central idea for him is, again, that truths **and** subjects come **together so that** the truth of a multiple is produced immanently to this multiple, with a subject's intervention.

G_{40} : Category one of truths and subjects come together

G_{41} : Category two of truths and subjects come together

G_{42} : Category three of truths and subjects come together

T_{40} : Category one of truth of a multiple is produced immanently to this multiple, with a subject's intervention.

T_{41} : Category two of truth of a multiple is produced immanently to this multiple, with a subject's intervention.

T_{42} : Category three of truth of a multiple is produced immanently to this multiple, with a subject's intervention.

Module Nine

That is why there is no need for a dogmatism of static correspondence (between a representing subject and 'the' world) to re-assert the claims of truth. For Badiou, then, truth is its own standard (it 'is the proof of itself'), for as new, the true simply **cannot (e)** correspond to anything (to any previously existing standard), but 'depends on its own production'.

G_{44} : Category one of correspondence to anything (to any previously existing standard), but 'depends on its own production'.

G_{45} : Category two of correspondence to anything (to any previously existing standard), but 'depends on its own production'.

G_{46} : Category three of correspondence to anything (to any previously existing standard), but 'depends on its own production'.

T_{44} : Category one of truth is its own standard (it 'is the proof of itself'), for as new, the true

T_{45} : Category two of truth is its own standard (it 'is the proof of itself'), for as new, the true

T_{46} : Category three of truth is its own standard (it 'is the proof of itself'), for as new, the true

SECTION NINE

Being And Event : Nosce Te Ipsum - Omnia Mutantur Nos Et Mutamur In Illis Model

INTRODUCTION—VARIABLES USED

Badiou and Deleuze John Mullarkey

(1) As such, truth is irrepresentable, for it is what happens just once and so is not repeatable. Strangely enough, when it is put this way, Badiou's notion of truth is not entirely alien to Deleuze. Indeed, in Cinema , Deleuze **links(e&eb) the** problematic of truth to that of the event explicitly

(2) Time has always put the notion of truth into crisis….it is the form or rather the pure force of time that **puts (eb)** truth into crisis.

(3) Consequently, Deleuze **prioritizes (e&eb)** the 'powers of the false'; but we should not let his terminology confuse us: it is the correspondence (or representational) theory of truth that Deleuze has it in his mind **to attack here** such that this power of the false has much in common with Badiou's evental truth (as Badiou must partly concede when he allows that 'times truth itself' for Deleuze, only what time essentially is being different for the two).

(4) For, like Badiou, Deleuze does not see thought **as an access** to being – as representational – either**, but (=) as a** quasi-being or event in itself. Truth itself is not entirely absent from Deleuze, then, but becomes its own, immanent standard (again, following Spinoza).

(5) Where Deleuze and Badiou continue to differ on this immanent truth is in the fact that, for Badiou, such immanence must be entirely actual if it is to be its own standard. Hence, Badiou thinks of Deleuze's dualism of virtual and actual as representational: actualities **are(=)** ontical beings, or 'simulacra of Being', that is, representations, and so in a relationship of inequality with Being (equivocal),rather than being both productive and genuinely univocal as Deleuze claims.

(6) Whether this is a fair assessment of the virtual-actual dyad is contentious, however, **because(e)** Badiou ignores what is univocal in the Virtual, namely difference, the formula

A Thousand Plateaus

(7) 'Monism = pluralism', conveying the fact that the only thing that is always the same, is difference.

(8) Besides the concept of the multiple, if there is one other key concept shared between Deleuze and Badiou – that must thereby be contested – it is that of the event

Being and Event

(9) Deleuze's are common and natural (they **occur (eb) in** both the human and inhuman realms). Badiou's events **have(e) to be** thought,

(10) Deleuze's **have (e) to be** felt. Going against the idea of a 'pure empiricism of what happens' (and Badiou most likely has Deleuze in mind here)

NOTATION

Module One

As such, truth is irrepresentable, for it is what happens just once and so is not repeatable. Strangely enough, when it is put this way, Badiou's notion of truth is not entirely alien to Deleuze. Indeed, in Cinema , Deleuze **links(e&eb) the** problematic of truth to that of the event explicitly

G_{13} : Category one of **problematic** of truth (Truth is problematized like events are. Say there is crime syndicate that is following you. They are problematizing the events and your reactionary potential. High if it is, more flagitious, flagrant consequences and repercussions would it have.

G_{14} : Category two of problematic of truth

G_{15} : Category three of problematic of truth

T_{13} : Category one of Problematization of the event. One may be dashed against, rammed with a car, or even killed. These are the various ways of Problematization. One event which law oversees is the repetitive exposure to particular scenario, scimitar, schottische and that disturbs the mind of the subject which essentially is an analogue computer.

T_{14} : Category two of Problematization of the event

T_{15} : Category three of Problematization of the event

Module Two

Time has always put the notion of truth into crisis….it is the form or rather the pure force of time that **puts (eb)** truth into crisis. Note quantum mechanics speaks of infinite possibilities and the measurement problem persists. This is of paramount importance in the **background of EPR of Einstein**.

G_{16} : Category one of Time has always put the notion of truth into crisis….it is the form or rather the pure force of time

G_{17} : Category two of Time has always put the notion of truth into crisis….it is the form or rather the pure force of time

G_{18} : Category three of Time has always put the notion of truth into crisis….it is the form or rather the pure force of time

T_{16} : Category one of truth into crisis

T_{17} : Category two of truth into crisis

T_{18} : Category three of truth into crisis

Module three

Consequently, Deleuze **prioritizes (e&eb)** the 'powers of the false'; but we should not let his terminology confuse us: it is the correspondence (or representational) theory of truth that Deleuze has it in his mind **to attack here** such that this power of the false has much in common with Badiou's evental truth (as Badiou must partly concede when he allows that 'times truth itself' for Deleuze, only what time essentially is being different for the two). In essence, the **more you follow the substance the more you chase the shadow**

G_{20} : Category one of 'powers of the false'

G_{21} : Category two of 'powers of the false'

G_{22} : Category three of 'powers of the false'

T_{20} : Category one of correspondence (or representational) theory of truth

T_{21} : Category two of correspondence (or representational) theory of truth

T_{22} : Category three of correspondence (or representational) theory of truth

Module four

For, like Badiou, Deleuze does not see thought **as an access** to being – as representational – either**, but (=) as a** quasi-being or event in itself. Truth itself is not entirely absent from Deleuze, then, but becomes its own, immanent standard (again, following Spinoza). In this connection it is to be stated that modern technologies try to read thought which is of utmost importance. Unless the mind is under the feeling of being under surveillance like in a office, it is bound to go hunky dory. Creativity certainly is not about any ill feeling about world or human beings. Even despondency and desolation is poetic as is seen in the romanticist poems.

G_{24} : Category one of quasi-being or event in itself (see the theory that thought is also an action)

G_{25} : Category two of quasi-being or event in itself

G_{26} : Category three of quasi-being or event in itself

T_{24} : Category one of thought is not **an access** to being nor as representational (We have already said that sensory organs are responsible for thought and it is just a line like in a computer. GD axiomatic predication and postulation alcovishness notwithstanding.)

T_{25} : Category two of thought is not **an access** to being nor as representational

T_{26} : Category three of thought is not **an access** to being nor as representational

Module five

Where Deleuze and Badiou continue to differ on this immanent truth is in the fact that, for Badiou, such immanence must be entirely actual if it is to be its own standard. Hence, Badiou thinks of Deleuze's dualism of virtual and actual as representational: actualities **are(=)** ontical beings, or 'simulacra of Being', that is, representations, and so in a relationship of inequality with Being (equivocal),rather than being both

productive and genuinely univocal as Deleuze claims.

Meaning of ontical from Wikipedia - in philosophy, ontic (from the Greek grc, ὄν, genitive grc, ὄντος: 'of that 'i differentiate the 'ontic' ('ontical' etc.)

- In being and time , martin Heidegger made the distinction between ontic al and properly speaking ontical, and ontology ought to designate - 'world' is used as an ontical concept, and signifies the totality of things which can be present-at-hand within the world. - Of Dasein that 'is ontically distinctive in that it 'is' ontological references: 'the ontical priority of the question of being.' - About the ontic state of the system at that time must equal the amount of knowledge one lacks this is called the 'knowledge balance principle.' - In the work being and time, the word existential is used to describe an ontic understanding of beings in the world. Precondition for - In certain situations where there is a balance of ontic good s and ontic evil s (ontic evils are those that are not immoral but merely - Critical ontology, ontic relationalism, and being-principle: critical ontology, ontic relationalism, and being-principle are the three major - characterizations/ definitions: ontic aspect: the ontic constituency of biological mechanisms includes entities and activities. - Feminine present participle of grc , εἶναι (to be); it is analogous to the English participle being , and the modern philosophy adjectival ontic

G_{28} : Category one of ontical beings, or 'simulacra of Being', that is, representations, and so in a relationship of inequality with Being (equivocal),rather than being both productive and genuinely univocal as Deleuze claims.

G_{29} : Category two of ontical beings, or 'simulacra of Being', that is, representations, and so in a relationship of inequality with Being (equivocal),rather than being both productive and genuinely univocal as Deleuze claims.

G_{30} : Category three of ontical beings, or 'simulacra of Being', that is, representations, and so in a relationship of inequality with Being (equivocal),rather than being both productive and genuinely univocal as Deleuze claims.

T_{28} : Category one of dualism of virtual and actual is representational: actualities

T_{29} : Category two of dualism of virtual and actual is representational: actualities

T_{30} : Category three of dualism of virtual and actual is representational: actualities

Module six

Whether this is a fair assessment of the virtual-actual dyad is contentious, however, **because(e)** Badiou ignores what is univocal in the Virtual, namely difference,

G_{32} : Category one of **because(e)** Badiou ignores what is univocal in the Virtual, namely difference

G_{33} : Category two of **because(e)** Badiou ignores what is univocal in the Virtual, namely difference

G_{34} : Category three of **because(e)** Badiou ignores what is univocal in the Virtual, namely difference

T_{32} : Category one of virtual and actual is representationalitiesof : actualities ontical beings, or 'simulacra of Being', that is, representations, and so in a relationship of inequality with Being (equivocal),rather than being both productive and genuinely univocal as Deleuze claims is contentious

T_{33} : Category two of virtual and actual is representationalitiesof : actualities ontical beings, or 'simulacra of Being', that is, representations, and so in a relationship of inequality with Being (equivocal),rather than being both productive and genuinely univocal as Deleuze claims is contentious

T_{34} : Category three ofvirtual and actual is representationalitiesof : actualities ontical beings, or 'simulacra of Being', that is, representations, and so in a relationship of inequality with Being (equivocal),rather than being both productive and genuinely univocal as Deleuze claims is contentious

Module seven

'Monism = pluralism', conveying the fact that the only thing that is always the same, is difference

G_{36} : Category one of monism

G_{37} : Category two of monism

G_{38} : Category three of monism

T_{36} : Category one of pluralism

T_{37} : Category two of pluralism

T_{38} : Category three of pluralism

Module eight

Deleuze's are common and natural (they **occur (eb) in** both the human and inhuman realms). Badiou's events **have(e) to be** thought

G_{40} : Category one of thought as an event

G_{41} : Category two of thought as an event

G_{42} : Category three of thought as an event

T_{40} : Category one of events(Deleuzean) common and natural

T_{41} : Category two of events(Deleuzean) common and natural

T_{42} : Category three of events(Deleuzean) common and natural

Module Nine

Deleuze's **have (e) to be** felt. Going against the idea of a 'pure empiricism of what happens' (and Badiou most likely has Deleuze in mind here)

G_{44} : Category one of feelings of gratification and deprivation(AP-GP dogma holds here too)

G_{45} : Category two of feelings of gratification and deprivation

G_{46} : Category three of feelings of gratification and deprivation

T_{44} : Category one of Thought(Deleuzean)

T_{45} : Category two of Thought(Deleuzean)

T_{46} : Category three of Thought(Deleuzean)

SECTION TEN

A Thousand Plateaus: Omnia Causa Fiunt - Perfer Et Obdura-Dolor Hic Tibi Proderit Olim Model

INTRODUCTION—VARIABLES USED

Badiou and Deleuze John Mullarkey

Being and Event

(1) Also, for Badiou the event is co-engendered with the subject – subject and event are **tied (e&eb) together** in the decision of the former to name the latter.
(2) The subject is not connected to the event as the one thatpre-exists it, but as the one **that comes into existence (eb)** with that nomination.
(3) For Deleuze, alternatively, the event **is (=) a** subjective.
(4) The event comes before and **generates the** human subject out of 'larval' selves. His events are ubiquitous in nature. Cartier-Bresson's adage that there is nothing in this world that **does (e) not have** a decisive moment is worth recapitulation in this context.

The Logic of Sense

(5) We get an image of the event as an incorporeal transformation that **gathers (e) up** everyday objects and individuals for its own purpose. A genuine event doesn't happen to someone, but in and through someone (that makes him or her part of the event).
(6) Again, as it so often does logic of world **ramifies (e) the** account in Being and Event
(7) Here rendering change in four types (modifications, facts, weak singularities, and strong singularities); yet the most radical form of change **remains (eb)** the event (or strong singularity), one that still lies ill with Deleuze's conception of the event. At one level at least, though, Badiou is close to Deleuze: for both, the event is about change. But it is an odd temporality for each of them.
(8) The only temporality, therefore, **is (=)** future anterior – purely intellectual and non-durational.
(9) An event is 'decoupled' **from (e) t**emporality.

NOTATION

Module One

Also, for Badiou the event is co-engendered with the subject – subject and event are **tied (e&eb) together** in the decision of the former to name the latter.

G_{13} : Category one of event is co-engendered with the subject – subject and event are **tied (e&eb) together**

G_{14} : Category two of event is co-engendered with the subject – subject and event are **tied (e&eb) together**

G_{15} : Category three of event is co-engendered with the subject – subject and event are **tied (e&eb) together**

T_{13} : Category one of decision of the former to name the latter

T_{14} : Category two of decision of the former to name the latter

T_{15} : Category three of decision of the former to name the latter

Module Two

The subject is not connected to the event as the one thatpre-exists it, but as the one subject **that comes into existence (eb)** with that nomination.

G_{16} : Category one of subject(individual general ledger in all three states with thoughts also as actions and concomitant GD with ASCII characters as in a computer) is not connected to the event as the one thatpre-exists it

G_{17} : Category two of subject(individual general ledger in all three states with thoughts also as actions and concomitant GD with ASCII characters as in a computer) is not connected to the event as the one thatpre-exists it

G_{18} : Category three of subject(individual general ledger in all three states with thoughts also as actions and concomitant GD with ASCII characters as in a computer) is not connected to the event as the one thatpre-exists it

T_{16} : Category one of subject that **comes into existence (eb)** with that nomination.

T_{17} : Category two of subject that **comes into existence (eb)** with that nomination.

T_{18} : Category three of subject that **comes into existence (eb)** with that nomination.

Module three

Event **is (=) a** subjective (we interpret this statement as quantum mechanical. It can be treated on the same lines as we treat the gratification,deprivations,thoughts actions in all three states)

G_{20} : Category one of subjectivity

G_{21} : Category two of subjectivity

G_{22} : Category three of subjectivity

T_{20} : Category one of event

T_{21} : Category two of event

T_{22} : Category three of event

Module four

The event comes before and **generates the** human subject out of 'larval' selves. His events are ubiquitous in nature. Cartier-Bresson's adage that there is nothing in this world that **does (e) not have** a decisive moment is worth recapitulation in this context. **This reminds one of Heisenberg's principles of uncertainty, collapse of Schrodinger's wave function, manifestation of one certainty out of infinite possibilities and also the existence of all the possibilities in multiverses**.

G_{24} : Category one of decisive moment

G_{25} : Category two of decisive moment

G_{26} : Category three of decisive moment

T_{24} : Category one of everything in this world. Note observable, non observable and even extra sensory

T_{25} : Category two of everything in this world

T_{26} : Category three of everything in this world

Module five

We get an image of the event as an incorporeal transformation that **gathers (e) up** everyday objects and individuals for its own purpose. A genuine event doesn't happen to someone, but in and through someone (that makes him or her part of the event). **There exists shades of observer-potentialities interaction**

G_{28} : Category one of everyday objects and individuals for its own purpose. A genuine event doesn't happen to someone, but in and through someone (that makes him or her part of the event).

G_{29} : Category two of everyday objects and individuals for its own purpose. A genuine event doesn't happen to someone, but in and through someone (that makes him or her part of the event).

G_{30} : Category three of everyday objects and individuals for its own purpose. A genuine event doesn't happen to someone, but in and through someone (that makes him or her part of the event).

T_{28} : Category one of images of the event as an incorporeal transformation

T_{29} : Category two of images of the event as an incorporeal transformation

T_{30} : Category three of images of the event as an incorporeal transformation

Module six

Again, as it so often does logic of world **ramifies (e) the** account in Being and Event

G_{32} : Category one of account in Being and Event

G_{33} : Category two of account in Being and Event

G_{34} : Category three of account in Being and Event

T_{32} : Category one of logic of world

T_{33} : Category two of logic of world

T_{34} : Category three of logic of world

Module seven

Here rendering change in four types (modifications, facts, weak singularities, and strong singularities); yet the most radical forms of change **remains** the event (or strong singularity), one that still lies ill with

Deleuze's conception of the event. At one level at least, though, Badiou is close to Deleuze: for both, the event is about change. But it is an odd temporality for each of them

G_{36} : Category one of event (or strong singularity)

G_{37} : Category two of event (or strong singularity)

G_{38} : Category three of event (or strong singularity

T_{36} : Category one of radical forms of change

T_{37} : Category two of radical forms of change

T_{38} : Category three of radical forms of change

Module eight

The only temporality, therefore, **is (=)** future anterior – purely intellectual and non-durational

G_{40} : Category one of future anterior – purely intellectual and non-durational

G_{41} : Category two of future anterior – purely intellectual and non-durational

G_{42} : Category three of future anterior – purely intellectual and non-durational

T_{40} : Category one of temporality

T_{41} : Category two of temporality

T_{42} : Category three of temporality

Module Nine

An event is 'decoupled' **from (e) t**emporality.

G_{44} : Category one of **t**emporality.

G_{45} : Category two of **t**emporality.

G_{46} : Category three of **t**emporality.

T_{44} : Category one of event is 'decoupled'

T_{45} : Category two of event is 'decoupled'

T_{46} : Category three of event is 'decoupled'

SECTION ELEVEN

Truth Of An Event: Acta Deos Nunquam Mortalia Fallunt Model

INTRODUCTION—VARIABLES USED

Badiou and Deleuze John Mullarkey

(1) The truth (of and in an event) for Badiou **is (=)** neither temporal nor atemporal but, he claims, the 'interruption' of 'a time …of the situation'.
(2) Likewise, for Deleuze, the event **is (=)** understood in terms of multiplicity rather than process: be it the 'original time' in his Proust and Signs or the ' eventum tantum' in The Logic of Sense – in each case there is ade-temporalisation of time.
(3) Why? Because, as with Badiou, time must be **contained (eb) in** eternity. But Deleuze's eternity is less futural and intellectual and more Proustian and affective, the event being is what **makes(eb)** the present past because it is always what has been, a 'past in general', the Virtual or Aion
(4) Unlike Deleuze, Badiou says that he cannot 'think that the new is a fold of the past …. This is why I conceptualize absolute beginnings (which require a theory of the void)….'
(5) Deleuze's Platonism: Ideas **as(=) Objective reality**

Deleuze and the Lacanian Real

(6) Recall the Old Catholic strategy to guard men against the temptation of the flesh: when you see in front of you a voluptuous feminine body, imagine how it will look in a couple of decades - the dried skin, sagging breasts... (Or, even better, imagine what lurks now already beneath the skin: raw flesh and bones, inner fluids, half-digested food and excrements...) Far from enacting a return to the Real destined to break the imaginary spell of the body, such a procedure **equals (=) the** escape from the Real, the Real announcing itself in the seductive appearance of the naked body.
(7) That is to say, in the opposition between the spectral appearance of the sexualized body and the repulsive body in decay, it is the spectral appearance with is the Real, and the decaying body which is reality - we take recourse to the decaying body in order **to (e) avoid** the deadly fascination of the Real which threatens to draw us into its vortex of jouissance.
(8) A "raw" Platonism would have claimed here that only the beautiful body fully **materializes(eb)** the Idea

(9) A body in its material decay simply **falls (e) off from** its Idea, is no longer its faithful copy.

NOTATION

Module One

The truth (of and in an event) for Badiou **is (=)** neither temporal nor atemporal but, he claims, the 'interruption' of 'a time …of the situation'

G_{13} : Category one of neither temporal nor atemporal but, he claims, the 'interruption' of 'a time …of the situation'
G_{14} : Category two of neither temporal nor atemporal but, he claims, the 'interruption' of 'a time …of the

situation'

G_{15} : Category three of neither temporal nor atemporal but, he claims, the 'interruption' of 'a time …of the situation'

T_{13} : Category one of truth (of and in an event)

T_{14} : Category two of truth (of and in an event)

T_{15} : Category three of truth (of and in an event)

Module Two

Likewise, for Deleuze, the event **is (=)** understood in terms of multiplicity rather than process: be it the 'original time' in his Proust and Signs or the ' eventum tantum' in The Logic of Sense – in each case there is ade-temporalisation of time.

G_{16} : Category one of terms of multiplicity rather than process: be it the 'original time' in his Proust and Signs or the 'eventum tantum' in The Logic of Sense – in each case there is ade-temporalisation of time.

G_{17} : Category two of terms of multiplicity rather than process: be it the 'original time' in his Proust and Signs or the ' eventum tantum' in The Logic of Sense – in each case there is ade-temporalisation of time.

G_{18} : Category three of terms of multiplicity rather than process: be it the 'original time' in his Proust and Signs or the ' eventum tantum' in The Logic of Sense – in each case there is ade-temporalisation of time.

T_{16} : Category one of event

T_{17} : Category two of event

T_{18} : Category three of event

Module three

Why? Because, as with Badiou, time must be **contained (eb) in** eternity. But Deleuze's eternity is less futural and intellectual and more Proustian and affective, the event being is what **makes(eb)** the present past because it is always what has been, a 'past in general', the Virtual or Aion

G_{20} : Category one of time

G_{21} : Category two of time

G_{22} : Category three of time

T_{20} : Category one of eternity

T_{21} : Category two of eternity

T_{22} : Category three of eternity

Module four

Why? Because, as with Badiou, time must be **contained (eb) in** eternity. But Deleuze's eternity is less futural and intellectual and more Proustian and affective, the event being is what **makes(eb)** the present past because it is always what has been, a 'past in general', the Virtual or Aion

G_{24} : Category one of eternity is less futural and intellectual and more Proustian and affective, the event

G_{25} : Category two of eternity is less futural and intellectual and more Proustian and affective, the event

G_{26} : Category three of eternity is less futural and intellectual and more Proustian and affective, the event

T_{24} : Category one of present past because it is always what has been, a 'past in general', the Virtual or Aion

T_{25} : Category two of present past because it is always what has been, a 'past in general', the Virtual or Aion

T_{26} : Category three of present past because it is always what has been, a 'past in general', the Virtual or Aion

Module five

Deleuze's Platonism: Ideas **as** Objective reality. Note that knowledge which is anagrammatic as the one we have kills the Truth

G_{28} : Category one of Objective reality (Truth)

G_{29} : Category two of Objective reality (Truth)

G_{30} : Category three of Objective reality (Truth)

T_{28} : Category one of ideas (knowledge)

T_{29} : Category two of ideas (knowledge)

T_{30} : Category three of ideas (knowledge)

Module six

Recall the Old Catholic strategy to guard men against the temptation of the flesh: when you see in front of you a voluptuous feminine body, imagine how it will look in a couple of decades - the dried skin, sagging breasts... (Or, even better, imagine what lurks now already beneath the skin: raw flesh and bones, inner fluids, half-digested food and excrements...) Far from enacting a return to the Real destined to break the imaginary spell of the body, such a procedure **equals (=) the** escape from the Real, the Real announcing itself in the seductive appearance of the naked body.

G_{32} : Category one of escape from the Real, the Real announcing itself in the seductive appearance of the naked body.

G_{33} : Category two of escape from the Real, the Real announcing itself in the seductive appearance of the naked body.

G_{34} : Category three of escape from the Real, the Real announcing itself in the seductive appearance of the naked body.

T_{32} : Category one of repelation of return to the Real destined to break the imaginary spell of the body

T_{33} : Category two of repelation of return to the Real destined to break the imaginary spell of the body

T_{34} : Category three of repelation of return to the Real destined to break the imaginary spell of the body

Module seven

That is to say, in the opposition between the spectral appearance of the sexualized body and the repulsive body in decay, it is the spectral appearance with is the Real, and the decaying body which is reality - we take recourse to the decaying body in order **to (e) avoid** the deadly fascination of the Real which threatens to draw us into its vortex of jouissance.

G_{36} : Category one of the deadly fascination of the Real which threatens to draw us into its vortex of jouissance.

G_{37} : Category two of the deadly fascination of the Real which threatens to draw us into its vortex of jouissance.

G_{38} : Category three of the deadly fascination of the Real which threatens to draw us into its vortex of jouissance.

T_{36} : Category one of recourse to the decaying body. There are many who take recourse. Characterstics of the individual systems that are investigated are taken in to consideration in the schema and doxa of stratification.

T_{37} : Category two of recourse to the decaying body

T_{38} : Category three of recourse to the decaying body

Module eight

A "raw" Platonism would have claimed here that only the beautiful body fully **materializes(eb)** the Idea

G_{40} : Category one of beautiful bodies

G_{41} : Category two of beautiful bodies

G_{42} : Category three of beautiful bodies

T_{40} : Category one of materialization of ideas

T_{41} : Category two of materialization of ideas

T_{42} : Category three of materialization of ideas

Module Nine

A body in its material decay simply **falls (e) off from** its Idea, is no longer its faithful copy.

G_{44} : Category one of stickler punctiliousness to the idea and is no longer the faithful copy

G_{45} : Category two of stickler punctiliousness to the idea and is no longer the faithful copy

G_{46} : Category three of stickler punctiliousness to the idea and is no longer the faithful copy

T_{44} : Category one of body in its material decay simply

T_{45} : Category two of body in its material decay simply

T_{46} : Category three of body in its material decay simply

SECTION TWELVE

Identity And Contradiction: Actus Non Facit Reum Nisi Mens Sit Rea Model

INTRODUCTION—VARIABLES USED

Badiou and Deleuze John Mullarkey

(1) From a Deleuzian (and, here, Lacanian) view, on the contrary, the specter that attracts us **is (=) the** Idea of the body as Real. This body is not the body in reality, but the virtual body in Deleuze's sense of the term: the incorporeal/immaterial body of pure intensities.
(2) Deleuze's most radical anti-Hegelian argument concerns pure difference: Hegel is unable to think pure difference which is outside the horizon of identity/contradiction; Hegel **conceives (eb) a** radicalized difference as contradiction which, then, through its dialectical resolution, is again subsumed under identity. (Here, Deleuze is also opposed to Derrida who, from his perspective, remains caught within the vicious cycle of contradiction/identity, merely postponing resolution indefinitely.)
(3) And insofar as Hegel is the philosopher of actuality/actualization, insofar as, for him, the "truth" of a potentiality is revealed in its actualization, Hegel's inability to think pure difference **equals his** inability to think the virtual in its proper dimension, as a possibility which already qua possibility possesses its own reality
(4) pure difference is not actual, it **does(e) not** concern different actual properties of a thing or among things, its status is purely virtual,
(5) It is a difference which takes place at its purest precisely when nothing changes in actuality, when, in actuality, the SAME thing **repeats (e&eb)** itself.
(6) Effectively, it may appear that it is only Deleuze who formulates the truly post-Hegelian program of thinking difference: the Derridean "opening" which **emphasizes (eb)** the endless difference, the dissemination that cannot ever be sublated/reappropriated, etc., remains within the Hegelian framework, merely "opening" it up...
(7) But, here, the Hegelian counter-argument would have been: is then the "pure" virtual difference **not the** very name for actual self-identity?
(8) "Pure" virtual difference CONSTITUTIVE (e) of actual identity?
(9) More precisely, in the terms of Deleuze's transcendental empiricism, pure difference **is the** virtual support/condition of actual identity: an entity is perceived as "(self-) identical" when (and only when) its virtual support is reduced to a pure difference.

NOTATION

Module One

From a Deleuzian (and, here, Lacanian) view, on the contrary, the specter that attracts us **is the** Idea of the body as Real. This body is not the body in reality, but the virtual body in Deleuze's sense of the term: the incorporeal/immaterial body of pure intensities.

G_{13} : Category one of Idea of the body as Real. This body is not the body in reality, but the virtual body in Deleuze's sense of the term: the incorporeal/immaterial body of pure intensities.

G_{14} : Category two of Idea of the body as Real. This body is not the body in reality, but the virtual body in

Deleuze's sense of the term: the incorporeal/immaterial body of pure intensities.

G_{15} : Category three of Idea of the body as Real. This body is not the body in reality, but the virtual body in Deleuze's sense of the term: the incorporeal/immaterial body of pure intensities.

T_{13} : Category one of specter that attracts us

T_{14} : Category two of specter that attracts us

T_{15} : Category three of specter that attracts us

Module Two

Deleuze's most radical anti-Hegelian argument concerns pure difference: Hegel is unable to think pure difference which is outside the horizon of identity/contradiction; Hegel **conceives (eb) a** radicalized difference as contradiction which, then, through its dialectical resolution, **is again(eb) subsumed** under identity. (Here, Deleuze is also opposed to Derrida who, from his perspective, remains caught within the vicious cycle of contradiction/identity, merely postponing resolution indefinitely.)

G_{16} : Category one of radicalized difference as contradiction which, then, through its dialectical resolution

G_{17} : Category two of radicalized difference as contradiction which, then, through its dialectical resolution

G_{18} : Category three of radicalized difference as contradiction which, then, through its dialectical resolution

T_{16} : Category one of identity. (Here, Deleuze is also opposed to Derrida who, from his perspective, remains caught within the vicious cycle of contradiction/identity, merely postponing resolution indefinitely.)

T_{17} : Category two of identity. (Here, Deleuze is also opposed to Derrida who, from his perspective, remains caught within the vicious cycle of contradiction/identity, merely postponing resolution indefinitely.)

T_{18} : Category three of identity. (Here, Deleuze is also opposed to Derrida who, from his perspective, remains caught within the vicious cycle of contradiction/identity, merely postponing resolution indefinitely.)

Module three

And insofar as Hegel is the philosopher of actuality/actualization, insofar as, for him, the "truth" of a potentiality is revealed in its actualization, Hegel's inability to think pure difference **equals his** inability to think the virtual in its proper dimension, as a possibility which already qua possibility possesses its own reality

G_{20} : Category one of inability to think the virtual in its proper dimension, as a possibility which already qua possibility possesses its own reality

G_{21} : Category two of inability to think the virtual in its proper dimension, as a possibility which already qua possibility possesses its own reality

G_{22} : Category three of inability to think the virtual in its proper dimension, as a possibility which already qua possibility possesses its own reality

T_{20} : Category one of in capacitance of thought of pure difference

T_{21} : Category two of in capacitance of thought of pure difference

T_{22} : Category three of in capacitance of thought of pure difference

Module four

pure difference is not actual, it **does(e) not** concern different actual properties of a thing or among things, its status is purely virtual

G_{24} : Category one of concernment of different actual properties of a thing or among things, its status is purely virtual

G_{25} : Category two of concernment of different actual properties of a thing or among things, its status is purely virtual

G_{26} : Category three of concernment of different actual properties of a thing or among things, its status is purely virtual

T_{24} : Category one of pure difference is not actual

T_{25} : Category two of pure difference is not actual

T_{26} : Category three of pure difference is not actual

Module five

Pure difference is a difference which takes place at its purest precisely when nothing changes in actuality, when, in actuality, the SAME thing **repeats (e&eb)** itself.

G_{28} : Category one of Pure difference is a difference which takes place at its purest precisely when nothing changes in actuality, when, in actuality, the SAME thing

G_{29} : Category two of Pure difference is a difference which takes place at its purest precisely when nothing changes in actuality, when, in actuality, the SAME thing

G_{30} : Category three of Pure difference is a difference which takes place at its purest precisely when nothing changes in actuality, when, in actuality, the SAME thing

T_{28} : Category one of thing that **repeats (e&eb)** itself. There are many things which could be repeated. We reiterate that the axiomatic predications of the classificational scheme are the characterstics of the investigating systems. It may be a shot beamed on the screen. It may be a colour like red shown continuously. Note that this is what makes pathological and that is how mind works associating "red' with danger and further concocting and fabricating association of tomato juice for blood for murder of say a close relative. This is one of the methods people use to instigate and incite anger in people. Repetitive sentences usually called "nagging" repetitive sign language" is what fosters indignation and high reactionary potential in the subject. This is also misused by vested interests. Subject starts off fusillades and filibuster at the very sight of such a sign or colour.

T_{29} : Category two of thing that **repeats (e&eb)** itself.

T_{30} : Category three of thing that repeats **(e&eb)** itself.

Module six(case two)

G_{32} : Category one of thing that repeats **(e&eb)** itself.

G_{33} : Category two of thing that repeats **(e&eb)** itself.

G_{34} : Category three of thing that repeats **(e&eb)** itself.

T_{32} : Category one of Pure difference is a difference which takes place at its purest precisely when nothing changes in actuality, when, in actuality, the SAME thing

T_{33} : Category two of Pure difference is a difference which takes place at its purest precisely when nothing changes in actuality, when, in actuality, the SAME thing

T_{34} : Category three of Pure difference is a difference which takes place at its purest precisely when nothing changes in actuality, when, in actuality, the SAME thing

Module seven

But, here, the Hegelian counter-argument would have been: is then the "pure" virtual difference **not (=) the** very name for actual self-identity?
Note: Identity formation, also called **individuation**, is the development of the **distinct personality** of an individual regarded as a persisting entity (known as personal continuity) in a particular stage of life in which individual characteristics are possessed and by which a person is recognised or known (such as the establishment of a reputation). This process **defines** individuals to others and themselves. Pieces of the person's actual identity include a sense of continuity, a sense of uniqueness from others, and a sense of affiliation. Identity formation **leads to** a number of issues of personal identity and an identity where the individual has some sort of comprehension of him or herself as a **discrete and separate entity**. This may be through individuation whereby the undifferentiated individual tends to become unique, or undergoes stages through which differentiated facets of a person's life tend toward **becoming a** more indivisible whole.

Identity is often described as finite and consisting of separate and distinct parts (family, cultural, personal, professional, etc.), yet according to Parker J. Palmer, it is an ever evolving core within where our genetics (biology), culture, loved ones, those we cared for, people who have harmed us and people we have harmed, the deeds done (good and ill) to self and others, experiences lived, and choices made come together to form **who we are at** this moment (Wikipedia)

G_{36} : Category one of actual self-identity

G_{37} : Category two of actual self-identity

G_{38} : Category three of actual self-identity

T_{36} : Category one of "pure" virtual difference

T_{37} : Category two of "pure" virtual difference

T_{38} : Category three of "pure" virtual difference

Module eight

"Pure" virtual difference CONSTITUTIVE (e) of actual identity

Note: A virtual function makes its class a polymorphic base class. Derived classes can override virtual functions. Virtual functions called through base class pointers/references will be resolved at run-time. That is, the dynamic type of the object is used instead of its static type:


```
Derived d;

Base& rb = d;

// if Base::f() is virtual and Derived overrides it, Derived::f() will be called

rb.f();
```

A pure virtual function is a virtual function whose declaration ends in =0:

```
Class Base {

 // ...

 Virtual void f() = 0;

 // ...
```

A pure virtual function makes the class it is defined for abstract. Abstract classes **cannot be** instantiated. Derived classes **need to** override/implement all inherited pure virtual functions. If they do not, they too will become abstract.
In C++, a class can define a pure virtual function that has an implementation. (What that's good for is debatable.)

Virtual functions must be implemented in the base class, but they need not be implemented in their derived classes. Derived classes will automatically inherit the base class implementations of all virtual functions, but can override those functions by providing their own implementations. The base class methods can still be called explicitly to provide common functionality. A function that is declared virtual is expected to be implemented by derived classes, but it is not a requirement.

Pure-virtual functions need not be implemented in the base class, but they must be implemented in the derived classes. Any class that declares a pure-virtual function is automatically rendered an ADT (abstract data type) and cannot be instantiated other than through derivation. Derived classes do not inherit any of the abstract base class implementations, but they can call them explicitly. Derived classes that do not implement all the pure-virtual functions inherited from their abstract base classes are automatically rendered abstract themselves; however any implementations they do provide are subsequently treated as virtual methods (and are therefore inherited by derivatives). A concrete class is a class that inherits or implements the sum total of all pure-virtual methods inherited from all abstract base classes. Only concrete classes can be instantiated without being derived from.

G_{40} : Category one of actual identity

G_{41} : Category two of actual identity

G_{42} : Category three of actual identity

T_{40} : Category one of Pure" virtual difference

T_{41} : Category two of Pure" virtual difference

T_{42} : Category three of Pure" virtual difference

Module Nine

More precisely, in the terms of Deleuze's transcendental empiricism, pure difference **is the** virtual support/condition of actual identity: an entity is perceived as "(self-) identical" when (and only when) its virtual support is reduced to a pure difference

G_{44} : Category one of virtual support/condition of actual identity: an entity is perceived as "(self-) identical" when (and only when) its virtual support is reduced to a pure difference

G_{45} : Category two of virtual support/condition of actual identity: an entity is perceived as "(self-) identical" when (and only when) its virtual support is reduced to a pure difference

G_{46} : Category three of virtual support/condition of actual identity: an entity is perceived as "(self-) identical" when (and only when) its virtual support is reduced to a pure difference

T_{44} : Category one of pure difference

T_{45} : Category two of pure difference

T_{46} : Category three of pure difference

SECTION THIRTEEN

Pure Difference: Dictum Sapienti Sat Est Model

INTRODUCTION—VARIABLES USED

Deleuze's Platonism: Ideas as Real: Slavoj Zizek Deleuze and the Lacanian Real

Deleuze's most radical anti-Hegelian argument concerns pure difference: Hegel is unable to think pure difference which is outside the horizon of identity/contradiction; Hegel conceives a radicalized difference as contradiction which, then, through its dialectical resolution, is again subsumed under identity. (Here, Deleuze is also opposed to Derrida who, from his perspective, remains caught within the vicious cycle of contradiction/identity, merely postponing resolution indefinitely.) And insofar as Hegel is the philosopher of actuality/actualization, insofar as, for him, the "truth" of a potentiality is revealed in its actualization, Hegel's inability to think pure difference equals his inability to think the virtual in its proper dimension, as a possibility which already qua possibility possesses its own reality: pure difference is not actual, it does not concern different actual properties of a thing or among things, its status is purely virtual, it is a difference which takes place at its purest precisely when nothing changes in actuality, when, in actuality, the SAME thing repeats itself. - Effectively, it may appear that it is only Deleuze who formulates the truly post-Hegelian program of thinking difference: the Derridean "opening" which emphasizes the endless difference, the dissemination that cannot ever be sublated/reappropriated, etc., remains within the Hegelian framework, merely "opening" it up...

PLASTIC DIFFERENCE IN QUANTUM COMPUTATION:

One major difference between silicon and silicone is their commercial use. Pure silicon is an ideal semi-conductor, like the more expensive element germanium. It is possible to heat silicon into a molten state and

then form it into wafers. These semi-conductive wafers can then be used as a base for the intricate and microscopic electronic circuitry required for computer processing. This relationship between silicon and the technology industry inspired the nickname for the California high-tech region known as Silicon Valley. Silicone, on the other hand, is used for a number of other industrial applications. Liquid silicone is often used as a lubricant for machine parts, for example. A thicker form of rubberized silicone can also be used as a waterproof sealant around windows and other gaps. Cured silicone is flexible, moisture-resistant and transparent. Lattice spacing increases with tensile stress till it reaches its elastic limit.

The LED consists of a chip of semiconducting material doped with impurities to create a p-n junction. As in other diodes, current flows easily from the p-side, or anode, to the n-side, or cathode, but not in the reverse direction. Charge-carriers—electrons and holes—flow into the junction from electrodes with different voltages. When an electron meets a hole, it falls into a lower energy level, and releases energy in the form of a photon. The wavelength of the light emitted, and thus its color depends on the band gap energy of the materials forming the p-n junction. In silicon or germanium diodes, the electrons and holes recombine by a non-radiative transition, which produces no optical emission, because these are indirect band gap materials. The materials used for the LED have a direct band gap with energies corresponding to near-infrared, visible, or near-ultraviolet light.

LED development began with infrared and red devices made with gallium arsenide. Advances in materials science have enabled making devices with ever-shorter wavelengths, emitting light in a variety of colors. LEDs are usually built on an n-type substrate, with an electrode attached to the p-type layer deposited on its surface. P-type substrates, while less common, occur as well. Many commercial LEDs, especially GaN/InGaN, also use sapphire substrate. Most materials used for LED production have very high refractive indices. This means that much light will be reflected back into the material at the material/air surface interface. Thus, light extraction in LEDs is an important aspect of LED production, subject to much research and development.

Badiou and Deleuze John Mullarkey

(1) In Lacanese, pure difference **concerns(e&eb)** the supplement of the virtual object (Lacan's object)
(2) Its most plastic experience is that of a sudden change in (our perception of) an object which, with regard to its positive qualities, remains the same: "although nothing changes, the thing all of a sudden **seemed (e) totally** different" - as Deleuze would have put it, it is the thing's intensity which changes. (For Lacan, the theoretical problem/task is here to distinguish between the Master-Signifier and object which both refer to the abyssal X in the object beyond its positive properties.)
(3) As such, pure difference is closer to antagonism **than to (eb) the closeness to the** difference between two positive social groups one of which is to be annihilated.
(4) The universalism that sustains an antagonistic struggle is not exclusive of anyone, which is why the highest triumph of the antagonistic struggle **is not** the destruction of the enemy, **but the (eb) explosion** of the "universal brotherhood" in which agents of the opposite camp change sides and join us (recall the proverbial scenes of police or military units joining the demonstrators).
(5) It is in such explosion of enthusiastic all-encompassing brotherhood from which no one is in principle excluded, that the difference between "us" and "enemy" as positive agents **is (eb) reduced to** a PURE formal difference.
(6) This brings us to the topic of difference, repetition, and change (in the sense of the rise of something really new). Deleuze's thesis according to which New and repetition are not opposed, i.e., according to which New **arises (e) only** from repetition, is to be read against the background of the difference between the Virtual and the Actual.
(7) To put it directly: changes which concerns only the actual aspect of things **are(=) only** changes

within the existing frame, not the emergences of something really New –

(8) New only emerges when the virtual support of the Actual changes, and this change **occurs (eb)** precisely in the guise of a repetition in which a thing remains the same in its actuality.

(9) In other words, things really change not when A transforms itself into B, but when, while A remains exactly the same with regard to its actual properties, **it e(e) imperceptibly** "totally changes"...

NOTATION

Module One

In Lacanese, pure difference **concerns(e&eb)** the supplement of the virtual object (Lacan's object)

G_{13} : Category one of pure difference

G_{14} : Category two of pure difference

G_{15} : Category three of pure difference

T_{13} : Category one of supplement of the virtual object (Lacan's object)
T_{14} : Category two of supplement of the virtual object (Lacan's object)
T_{15} : Category three of supplement of the virtual object (Lacan's object)

Module Two

Its most plastic experience is that of a sudden change in (our perception of) an object which, with regard to its positive qualities, remains the same: "although nothing changes, the thing all of a sudden **seemed (e) totally** different" - as Deleuze would have put it, it is the thing's intensity which changes. (For Lacan, the theoretical problem/task is here to distinguish between the Master-Signifier and object which both refer to the abyssal X in the object beyond its positive properties.)

G_{16} : Category one of **totally** different" - as Deleuze would have put it, it is the thing's intensity which changes. (For Lacan, the theoretical problem/task is here to distinguish between the Master-Signifier and object which both refer to the abyssal X in the object beyond its positive properties.)

G_{17} : Category two of **totally** different" - as Deleuze would have put it, it is the thing's intensity which changes. (For Lacan, the theoretical problem/task is here to distinguish between the Master-Signifier and object which both refer to the abyssal X in the object beyond its positive properties.)

G_{18} : Category three of **totally** different" - as Deleuze would have put it, it is the thing's intensity which changes. (For Lacan, the theoretical problem/task is here to distinguish between the Master-Signifier and object which both refer to the abyssal X in the object beyond its positive properties.)

T_{16} : Category one of plastic experience is that of a sudden change in (our perception of) an object which, with regard to its positive qualities, remains the same: "although nothing changes, the thing all of a sudden

T_{17} : Category two of plastic experience is that of a sudden change in (our perception of) an object which, with regard to its positive qualities, remains the same: "although nothing changes, the thing all of a sudden

T_{18} : Category three of plastic experience is that of a sudden change in (our perception of) an object which, with regard to its positive qualities, remains the same: "although nothing changes, the thing all of a sudden

Module three

As such, pure difference is closer to antagonism **than to (e) the closeness to the** difference between two positive social groups one of which is to be annihilated.

G_{20} : Category one of **closeness to the** difference between two positive social groups one of which is to be annihilated.

G_{21} : Category two of **closeness to the** difference between two positive social groups one of which is to be

annihilated.

G_{22} : Category three of **closeness to the** difference between two positive social groups one of which is to be annihilated.

T_{20} : Category one of pure difference is closer to antagonism

T_{21} : Category two of pure difference is closer to antagonism

T_{22} : Category three of pure difference is closer to antagonism

Module four

The universalism that sustains an antagonistic struggle is not exclusive of anyone, which is why the highest triumph of the antagonistic struggle **is not** the destruction of the enemy, **but the (eb) explosion** of the "universal brotherhood" in which agents of the opposite camp change sides and join us (recall the proverbial scenes of police or military units joining the demonstrators).

CLOSENESS: A MATHEMATICAL PERSPECTIVE

In topology and related areas in mathematics closeness is one of the basic concepts in a topological space. Intuitively we say two sets are close if they are arbitrarily near to each other. The concept can be defined naturally in a metric space where a notion of distance between elements of the space is defined, but it can be generalized to topological spaces where we have no concrete way to measure distances

Definition

Given a metric space (X, d) a point p is called close or near to a set A if

$$d(p, A) = 0,$$

Where the distance between a point and a set is defined as

$$d(p, A) := \inf_{a \in A} d(p, a).$$

G_{24} : Category one of universalism that sustains an antagonistic struggle is not exclusive of anyone, which is why the highest triumph of the antagonistic struggle **is not** the destruction of the enemy

G_{25} : Category two of universalism that sustains an antagonistic struggle is not exclusive of anyone, which is why the highest triumph of the antagonistic struggle **is not** the destruction of the enemy

G_{26} : Category three of universalism that sustains an antagonistic struggle is not exclusive of anyone, which is why the highest triumph of the antagonistic struggle **is not** the destruction of the enemy

T_{24} : Category one of **explosion** of the "universal brotherhood" in which agents of the opposite camp changes sides and join us (recall the proverbial scenes of police or military units joining the demonstrators).

T_{25} : Category two of **explosion** of the "universal brotherhood" in which agents of the opposite camp change sides and join us (recall the proverbial scenes of police or military units joining the demonstrators).

T_{26} : Category three of **explosion** of the "universal brotherhood" in which agents of the opposite camp

change sides and join us (recall the proverbial scenes of police or military units joining the demonstrators).

Module five

It is in such explosion of enthusiastic all-encompassing brotherhood from which no one is in principle excluded, that the difference between "us" and "enemy" as positive agents **is (eb) reduced to** a PURE formal difference

G_{28} : Category one of enthusiastic all-encompassing brotherhood from which no one is in principle excluded, that the difference between "us" and "enemy" as positive agents

G_{29} : Category two of enthusiastic all-encompassing brotherhood from which no one is in principle excluded, that the difference between "us" and "enemy" as positive agents

G_{30} : Category three of enthusiastic all-encompassing brotherhood from which no one is in principle excluded, that the difference between "us" and "enemy" as positive agents

T_{28} : Category one of PURE formal difference Such "formal differences" can be found in linguistic models of Quantum mechanical computation. Described here is a quantum computational **linguistic model**. It is based on the fact that formal grammars have representations in the operators on a Hilbert space. These operators form the set of unitary transformations on an n level quantum system. A simplified version of English is cast as a group and basic properties of that language are **probed algebraically**. It is shown that unitary operators **will not provide** a sufficient space for a representation of a realistic, natural language. The space of superoperators is suggested as the next step. There are no calculations of the matrices which act as representations of any algebraic structure which may model natural language. Instead the paper simply gathers together some disparate areas of mathematics and linguistics so that, maybe, a few practitioners of those areas might then actually **provide** the final representations.(**A QUANTUM COMPUTATIONAL LINGUISTIC MODEL BEN SPROTT)**

T_{29} : Category two of PURE formal difference The primary vehicle of calculus and other higher mathematics is the function. Its "input value" is its argument, usually a point ("P") expressible on a graph. The difference between two points, themselves, is known as their Delta (ΔP), as is the difference in their function result, the particular notation being determined by the direction of formation:

Forward difference: $\Delta F(P) = F(P + \Delta P) - F(P)$;

Central difference: $\delta F(P) = F(P + \frac{1}{2}\Delta P) - F(P - \frac{1}{2}\Delta P)$;

Backward difference: $\nabla F(P) = F(P) - F(P - \Delta P)$.

The general preference is the forward orientation, as F (P) is the base, to which differences (i.e., "ΔP"s) are added to it. Furthermore,

If |ΔP| is finite (meaning measurable), then ΔF (P) is known as a finite difference, with specific denotations of DP and DF(P);

If |ΔP| is infinitesimal (an infinitely small amount—ℓ—usually expressed in standard analysis as a limit: $\lim_{\Delta P \to 0}$), then ΔF(P) is known as an infinitesimal difference, with specific denotations of dP and dF(P) (in calculus graphing, the point is almost exclusively identified as "x" and F(x) as "y").

The function difference divided by the point difference is known as the difference quotient (attributed

to Isaac Newton, it is also known as Newton's quotient):

$$\frac{\Delta F(P)}{\Delta P} = \frac{F(P+\Delta P) - F(P)}{\Delta P} = \frac{\nabla F(P+\Delta P)}{\Delta P}.$$

T_{30} : Category three of PURE formal difference

Module six

This brings us to the topic of difference, repetition, and change (in the sense of the rise of something really new). Deleuze's thesis according to which New and repetition are not opposed, i.e., according to which New **arises (e) only** from repetition, is to be read against the background of the difference between the Virtual and the Actual.

G_{32} : Category one of repetition, is to be read against the background of the difference between the Virtual and the Actual.

G_{33} : Category two of repetition is to be read against the background of the difference between the Virtual and the Actual.

G_{34} : Category three of repetition is to be read against the background of the difference between the Virtual and the Actual.

T_{32} : Category one of New and repetition are not opposed, i.e., according to which New **arises**

T_{33} : Category two of New and repetition are not opposed, i.e., according to which New **arises**

T_{34} : Category three of New and repetition are not opposed, i.e., according to which New **arises**

Module seven

To put it directly: changes which concern only the actual aspect of things **are only** changes within the existing frame, not the emergences of something really New.

G_{36} : Category one of changes within the existing frame, not the emergences of something really New.

G_{37} : Category two of changes within the existing frame, not the emergences of something really New.

G_{38} : Category three of changes within the existing frame, not the emergences of something really New.

T_{36} : Category one of changes which concern only the actual aspect of things

T_{37} : Category two of changes which concern only the actual aspect of things

T_{38} : Category three of changes which concern only the actual aspect of things

Module eight

New only emerges when the virtual support of the Actual changes, and this change **occurs (eb)** precisely in the guise of a repetition in which a thing remains the same in its actuality.

G_{40} : Category one of New only emerges when the virtual support of the Actual changes

G_{41} : Category two of New only emerges when the virtual support of the Actual changes

G_{42} : Category three of New only emerges when the virtual support of the Actual changes

T_{40} : Category one of. changes in the Actual **occurs (eb)** precisely in the guise of a repetition in which a thing remains the same in its actuality

T_{41} : Category two of changes in the Actual **occurs (eb)** precisely in the guise of a repetition in which a thing remains the same in its actuality

T_{42} : Category three of changes in the Actual **occurs (eb)** precisely in the guise of a repetition in which a thing remains the same in its actuality

Module Nine

In other words, things really change not when A transforms itself into B, but when, while A remains exactly the same with regard to its actual properties, **it e(e) imperceptibly** "totally changes"...

G_{44} : Category one of things really change not when A transforms itself into B, but when, while A remains exactly the same with regard to its actual properties

G_{45} : Category two of things really change not when A transforms itself into B, but when, while A remains exactly the same with regard to its actual properties

G_{46} : Category three of things really change not when A transforms itself into B, but when, while A remains exactly the same with regard to its actual properties

T_{44} : Category one of **imperceptible** "totally changes"...

T_{45} : Category two of **imperceptible** "totally changes"...

T_{46} : Category three of **imperceptible** "totally changes"...

SECTION FOURTEEN

"Minimal Difference" - The Purely Parallax Gap: Quod Me Nutrit Me Destruit. Qui

Advances in Physics Theories and Applications
ISSN 2224-719X (Paper) ISSN 2225-0638 (Online)
Vol 14, 2013

Tacet Consentire Model

INTRODUCTION—VARIABLES USED

Badiou and Deleuze John Mullarkey

(1) Lacanian Real, in its opposition to the Symbolic, has ultimately nothing whatsoever to do with the standard empiricist (or phenomenological, or historicist, or Lebensphilosophie, for that reason) topic of the wealth of reality that resists formal structures that cannot be reduced to its conceptual determinations - language is grey, reality is green... The Lacanian Real is, on the contrary, even more "**reductionist**"(eb) that any symbolic structure

(2) We touch it when we subtract from a symbolic field all the wealth of its differences, **reducing (eb) it to a** minimum of antagonism.

(3) Lacan himself is here not beyond reproach, since he gets sometimes **seduced (eb) by** the rhizomatic wealth of language beyond (or, rather, beneath) the formal structure that sustains it.

(4) It is in this sense that, in the last decade of his teaching, he deployed the notion of lalangue (sometimes simply translated as "language") which stands for language **as(=) the** space of illicit pleasures that defy any normativity the chaotic multitude of homonymies, word-plays, "irregular" metaphoric links and resonances..

(5) Productive as this notion is, one should be aware of its limitations. Many commentators have noted that Lacan's last great literary reading, that of Joyce to whom his late seminar (XXIII: Le sinthome) is dedicated, is not at the level of his previous great readings (Hamlet, Antigone, Claudel's Coufontaine-trilogy). There is effectively something fake in Lacan's fascination with late Joyce, with Finnegan's Wake as the latest version of the literary Gesamtkunstwerk with its endless wealth of lalangue in which not only the gap between singular languages, but the very gap between linguistic meaning and jouissance seems **overcome(e)** and the rhizome-like jouis-sense (enjoyment-in-meaning: enjoy-meant) proliferates in all directions.

(6) The true counterpart to Joyce is, of course, Samuel Becket: after his early period in which he more or less wrote some variations on Joyce, the "true" Becket constituted himself through a true ethical act, a CUT, a **rejection(e)** of the Joycean wealth of enjoy-meant, and the ascetic turn towards a "minimal difference," towards a minimalization, "subtraction," of the narrative content and of language itself (this line is most clearly discernible in his masterpiece, the trilogy Molloy - Malone Dies - L'innomable).

(7) So what is the "minimal difference" - the purely parallax gap - that sustains Becket's mature production? One is tempted to propose the thesis that it is the very difference between French and English: as is known, Becket wrote most of his mature works in French (not his mother tongue), and the, desperate at the low quality of translations, translated them himself into English, and these translations are not mere close translations, but effectively a different text. It is because of this "minimalist" - purely formal and insubstantial - status of the Real that, for Lacan, repetition **precedes (e)** repression - or, as Deleuze put it succinctly: "We do not repeat because we repress, we repress **because** (e) we repeat." (DR-105)

(8) It is not that, first, we repress some traumatic content, and then, since we are unable to remember it and thus to clarify **our relationship (e&eb)** to it, this content continues to haunt us, repeating itself in disguised forms.

(9) If the Real **is(=) a** minimal difference, then repetition (that establishes this difference) **is(=)** primordial

NOTATION

Module One

Lacanian Real, in its opposition to the Symbolic, has ultimately nothing whatsoever to do with the standard

empiricist (or phenomenological, or historicist, or Lebensphilosophie, for that reason) topics of the wealth of reality that resists formal structures that cannot be reduced to its conceptual determinations - language is grey, reality is green... The Lacanian Real is, on the contrary, even more "**reductionist**"(eb) that any symbolic structure

G_{13} : Category one of topics of the wealth of reality that resists formal structures that cannot be reduced to its conceptual determinations - language is grey, reality is green... The Lacanian Real is, on the contrary, even more

G_{14} : Category two of topics of the wealth of reality that resists formal structures that cannot be reduced to its conceptual determinations - language is grey, reality is green... The Lacanian Real is, on the contrary, even more

G_{15} : Category three of topics of the wealth of reality that resists formal structures that cannot be reduced to its conceptual determinations - language is grey, reality is green... The Lacanian Real is, on the contrary, even more

T_{13} : Category one of symbolic structure

T_{14} : Category two of symbolic structure

T_{15} : Category three of symbolic structure

Module Two

We touch it when we subtract from a symbolic field all the wealth of its differences, **reducing (eb) it to a** minimum of antagonism

FIELDS:

In abstract algebra, a field **is a** commutative ring which **contains a** multiplicative inverse for every nonzero element, equivalently a ring whose nonzero elements **form an** abelian group under multiplication. As such it is an algebraic structure with notions of addition, subtraction, multiplication, and division satisfying the appropriate Abelian group equations and distributive law. The most commonly used fields are the field of real numbers, the field of complex numbers, and the field of rational numbers, but there are also finite fields, fields of functions, various algebraic number fields, p-adic fields, and so forth. To avoid confusion with other uses of the word "field", the term "corpus" may also be used.

Any field may **be used** as the scalars for a vector space, which is the standard general context for linear algebra. The theory of field extensions (including Galois Theory) involves the roots of polynomials with coefficients in a field; among other results, this theory leads to impossibility proofs for the classical problems of angle trisection and squaring the circle with a compass and straightedge, as well as a proof of the Abel–Ruffini theorem on the algebraic insolubility of quintic equations. In modern mathematics, the theory of fields (or field theory) plays an essential role in number theory and algebraic geometry.

As an algebraic structure, every field is a ring, but not every ring is a field. The most important difference is that fields allow for division (though not division by zero), while a ring need not possess multiplicative inverses; for example the integers form a ring, but 2x = 1 has no solution in integers. Also, the multiplication operation in a field is required to be commutative. A ring in which division is possible but commutativity is not assumed (such as the quaternions) is called a division ring or skew field. (Historically, division rings were sometimes referred to as fields, while fields were called commutative fields.)

As a ring, a field may be classified as a specific type of integral domain, and can be characterized by the following (not exhaustive) chain of class inclusions:

Commutative rings ⊃ integral domains ⊃ integrally closed domains ⊃ unique factorization domains ⊃ principal ideal domains ⊃ Euclidean domains ⊃ fields' ⊃ finite fields (Note the model is applicable here also. Wikipedia).

Beckett with Lacan part 1 Slavoj Zizek(emphasis mine)

One can understand James Joyce, with all the obscenities that permeate his writings, as the ultimate Catholic author, "the greatest visionary of the dark underground of Catholicism, an underground embodying a pure transgression, but one which is nevertheless a profoundly Catholic transgression." Catholicism is legalistic, and, as Paul knew it so well, the Law **generates its** own transgression; consequently, the staging of the obscene underground of the Law, the travesty of the Black Mass (or, in Joyce's case, the elevation of Here Comes Everybody into Christ who has to die in order to be reborn as the eternal Life-Goddess, from Molly Bloom to Anna Livia Plurabelle), is the supreme Catholic act.

This achievement of Joyce simultaneously signals his limit, the limit which pushed Beckett to break with him. If there ever was a kenotic writer, the writer of the utter self-emptying of subjectivity, **of its (eb) reduction to** a minimal difference, it is Beckett.

We touch the Lacanian Real when **we (e) subtract** from a symbolic field all the wealth of its differences, reducing it to a minimum of antagonism. Lacan gets sometimes seduced by the rhizomatic wealth of language beyond (or, rather, beneath) the formal structure that sustains it. It is in this sense that, in the last decade of his teaching, he deployed the notion of lalangue (sometimes simply translated as "llanguage") which stands for language as the space of illicit pleasures **that (e) defy** any normativity: the chaotic multitude of homonymies, word-plays, "irregular" metaphoric links and resonances… Productive as this notion is, one should be aware of its limitations. Many commentators have noted that Lacan's last great literary reading, that of Joyce to whom his late seminar (XXIII: Le sinthome) is dedicated, is not at the level of his previous great readings (Hamlet, Antigone, Claudel's Coufontaine-trilogy). There is effectively something fake in Lacan's fascination with late Joyce, with Finnegan's Wake as the latest version of the literary Gesamtkunstwerk with its endless wealth of lalangue in which not only the gap between singular languages, but the very gap between linguistic meaning and jouissance **seems(e) overcome** and the rhizome-like jouis-sense (enjoyment-in-meaning: enjoy-meant) **proliferates in** all directions. The true counterpart to Joyce is, of course, Samuel Becket: after his early period in which he more or less wrote some variations on Joyce, the "true" Becket constituted himself through a true ethical act, a CUT, a rejection of the Joycean wealth of enjoy-meant, and the ascetic turn towards a "minimal difference," towards a minimalization, "subtraction," of the narrative content and of language itself (this line is most clearly discernible in his masterpiece, the trilogy Molloy – Malone Dies – L'innomable). Beckett is effectively the literary counterpart of Anton Webern: both are authors of extreme modernist minimalism, of subtracting a minimal difference from the wealth of material.

Beckett's Texts for Nothing (first published in French in 1955 as Nouvelles et texts pour rien) is the fourth term which supplements the trilogy Molloy – Malone Dies – The Unnamable – Beckett himself referred to Texts as "the grisly afterbirth of L'innomable," the "attempt to get out of the attitude of disintegration /of the trilogy/ but it failed." [3] The obvious link is that the first line of the first text ("Suddenly, no, at last, long last, I couldn't any more") echoes the famous last line of The Unnamable ("you must go on, I can't go on, I'll go on"), a true Kantian imperative, a paraphrase of Kant's Du kannst, denn du sollst ("You can, because you must."). The voice of conscience tells me "you must go on," I reply, referring to my weakness, "I can't go on," but as a Kantian, I know this excuse doesn't count, so I nonetheless decide that "I'll go on," doing the impossible.

Since, for Beckett, what "must go on" is ultimately writing itself, the Lacanian version of the last line of The

Unnamable is something that ne cesse pas a s'ecrire, that doesn't cease writing itself – a necessity, the first term in the logical square which also comprises impossibility (that which ne cesse pas a ne pas s'ecrire, doesn't cease not writing itself), possibility (that which cesse a s'ecrire, ceases to write itself), and contingency (that which cesse a ne pas s'ecrire, ceases not writing itself). It is crucial to note here the clear distinction between possibility and contingency: while possibility **is the** opposite of necessity, contingency **is the** opposite of impossibility. In Badiou's terms of the attitudes towards a Truth-Event, necessity stands for the fidelity to Truth, impossibility for a situation with no truth, possibility for the possibility of a truth-procedure to exhaust its potentials and to stop, and contingency for the beginning of a new truth-procedure.

So what do Texts for Nothing register, a possibility or a contingency? A possibility, definitely – a possibility to "cease writing," to betray fidelity, to cease going on. The failure of Texts is thus good news: Texts are failed betrayals, failed attempts to get rid of the ethical injunction. They are a comical supplement to the great triad – an opportunist's attempt to squeeze out of the call of duty, somehow like Kierkegaard's "sickness unto death," where a mortal human being attempts to escape immortality, its unbearable ethical burden/injunction. In this sense, Texts are an optimistic work – their message is that one cannot but "go on" as an immortal bodiless drive, as a subject without subjectivity: "No, no souls, or bodies, or birth, or life, or death, you've got to go on without any of that junk"…

Jonathan Boulter thus got it right – on condition that we strictly distinguish between subject and subjectivity. The whole of the trilogy can be read as a gradual getting rid of subjectivity, a gradual reduction of subjectivity **to the** minimum of a subject without subjectivity – a subject which is no longer a person, whose objective correlative is no longer a body (organism), but only a partial object (organ), a subject of DRIVE which is Freud's name for immortal persistence, "going on." Such a subject is a living dead – still alive, going on, persisting, but dead (deprived of body) – undead. Texts are a comical attempt to resubjectivize this subject – among other things, to provide him with a body, to travel back the road from Cheshire cat's smile to its full body. Boulter is right to correct Alvarez who claimed that Texts are written in the same "breathless, bodiless style" as The Unnamable:

"One of the things the reader notices about Texts from its outset is that the body (of the narrator/narrated) has made an uncanny return from its near obliteration in The Unnamable: the narrator of The Unnamable is disembodied (it may be that "he" is merely a brain in an urn). At the very least, the issue of subjectivity is a complex one in the trilogy because the relation between voice (of narrator) **and** body (of narrator) is continually called into question. We may in fact argue that the trilogy in toto is about the dismantling of the physical body: in Molloy, the body is ambulatory but weakening; in Malone Dies the body is on its last legs, immobile and dying; in The Unnamable the physical body may in fact have ceased to be an issue as the narrator floats between personalities and subject positions. All of which is to indicate that in Texts, the body has made /…/ an unexpected comeback."(333-334)

The subject without subjectivity, this "living dead," **is also** timeless – when we reach this point, "time has turned **into** space and there will be no more time till I get out of here" (note how Beckett repeats here Wagner's precise formula of the sacred space of the Grail's castle from Parsifal "time become here space," which Claude Lévi-Strauss quotes as the most succinct definition of myth). The subject we thus reach, a subject without subjectivity, is a subject which "cannot maintain with any certainty that the experiences he describes are in fact his own; we have a narrating subject who cannot discern if his voice is his own; we have a subject who cannot tell if he has a body; and most crucially, we have a subject **who has no** sense of personal history, no memory. We have, in short, a subject whose ontology denies the viability of mourning and trauma, yet who seems to display the viability of mourning and trauma."(337)

Is this subject deprived of all substantial content not the subject as such, at its most radical, the Cartesian cogito? Boulter's idea is that, for Freud, trauma presupposes a subject to whom it happens and who then tries

to narrativize it, to come to terms with it, in the process of mourning. In the case of the Beckettian narrator, on the contrary, "there is no hope of establishing a link between his own present condition and the trauma that is its precondition. Instead of having a story seemingly given to him unawares – as in the case of the victim of trauma who cannot recognize his past as his own – the Beckettian narrator can only hope (without hope /…/) for a story that will reconnect his present atemporal /…/ condition to his past."(341)

This is the division of the subject at its most radical: the subject is reduced to $ (the barred subject), even its innermost self-experience is taken from it. This is how one should understand Lacan's claim that the subject is always "decentered" - his point is not that my subjective experience is regulated by objective unconscious mechanisms that are decentered with regard to my self-experience and, as such, beyond my control (a point asserted by every materialist), but, rather, something much more unsettling: I am deprived of even my most intimate subjective experience, the way things "really seem to me," that of the fundamental fantasy that constitutes and guarantees the core of my being, since I can never consciously experience it and assume it. One should counter Boulter's question "To what extent do trauma and mourning require a subject?"(337) with a more radical one: to what extent does (the very emergence of) a subject require trauma and mourning? The primordial trauma, the trauma constitutive of the subject, **is the** very gap that bars the subject from ITS OWN "inner life."

Scenes from a Happy Life. This inner and constitutive link between trauma **and** subject is the topic of what is undoubtedly Beckett's late masterpiece: Not I, a twenty-minute dramatic monologue written in 1972, an exercise in theatric minimalism: there are no "persons" here, intersubjectivity is reduced to its most elementary skeleton, that of the speaker (who is not a person, but a partial object, a faceless MOUTH speaking) and AUDITOR, a witness of the monologue who says nothing throughout the play (all the Auditor does is that, in "a gesture of helpless compassion"(Beckett), he four times repeats the gesture of simple sideways raising of arms from sides and their falling back. (When asked if the Auditor is Death or a guardian angel, Beckett shrugged his shoulders, lifted his arms and let them fall to his sides, leaving the ambiguity intact – repeating the very gesture of the Auditor.) Beckett himself pointed to the similarities between Not I and The Unnamable with its clamoring voice longing for silence, circular narrative and concern about avoiding the first person pronoun: "I shall not say I again, ever again". Along these lines, one could agree with Vivian Mercier's suggestion that, gender aside, Not I is a kind of dramatization of The Unnamable - one should only add that, in Not I, we get the talking partial coupled/supplemented with a minimal figure of the big Other. – Here, then, is the text of this piece in its entirety:

Stage in darkness but for MOUTH, upstage audience right, faintly lit from close-up and below, rest of face in shadow. Invisible microphone. AUDITOR, downstage audience left, tall standing figure, sex undeterminable, enveloped from head to foot in loose black djellaba, with hood, fully faintly lit, standing on invisible podium about 4 feet high shown by attitude alone to be facing diagonally across stage intent on MOUTH, dead still throughout but for four brief movements where indicated. As house lights down MOUTH`S voice unintelligible behind curtain. House lights out. Voice continues unintelligible behind curtain, l0 seconds. With rise of curtain ad-libbing from text as required leading when curtain fully up and attention sufficient into:

Renormalization Automated by Hopf Algebra D.J. Broadhurst1, D. http://dx.doi.org/10.1006/jsco.1999.0283

(1) It was recently shown that the renormalization of quantum field theory **is (e&eb) organized by** the Hopf algebra of decorated rooted trees, whose coproduct identifies the divergences requiring subtraction and whose antipode achieves this.
(2) Authors automate this process in a few lines of recursive symbolic code, which **deliver a** finite renormalized expression for any Feynman diagram.

(3) They thus verify a representation of the operator product expansion, **which (e&eb) generalizes** Chen's Lemma for iterated integrals.
(4) The subset of diagrams whose forest structure entails a unique primitive sub divergence **provides a** representation of the Hopf algebra HRof undecorated rooted trees.
(5) Undecorated Hopf algebra program is designed to process the 24 213 878 BPHZ **contributions (eb) to** the renormalization of 7813 diagrams, with up to 12 loops.
(6) Authors consider 10 models, each in nine renormalization schemes. The two simplest models reveal a notable feature of the subalgebra of Connes and Moscovici, corresponding to the commutative part of the Hopf algebra HTof the diffeomorphism group: it assigns to Feynman diagrams those weights **which (e) remove** zeta values from the counterterms of the minimal subtraction scheme. Authors devise a fast algorithm for these weights, whose squares are summed with a permutation factor, to give rational counterterms.

G_{16} : Category one of subtraction from a symbolic field all the wealth of its differences; Commutative rings; integrally closed domains; principal ideal domains; time; renormalization of quantum field theory; Hopf algebra of decorated rooted trees, whose coproduct identifies the divergences requiring subtraction and whose antipode achieves this; recursive symbolic code; representation of the operator product expansion; subset of diagrams whose forest structure entails a unique primitive sub divergence; zeta values from the counterterms of the minimal subtraction scheme;

G_{17} : Category two of subtraction from a symbolic field all the wealth of its differences; Commutative rings integrally closed domains; principal ideal domains; time; renormalization of quantum field theory; Hopf algebra of decorated rooted trees, whose coproduct identifies the divergences requiring subtraction and whose antipode achieves this; recursive symbolic code; representation of the operator product expansion; subset of diagrams whose forest structure entails a unique primitive sub divergence; zeta values from the counterterms of the minimal subtraction scheme;

G_{18} : Category three of subtraction from a symbolic field all the wealth of its differences; Commutative rings; integrally closed domains; principal ideal domains; time; renormalization of quantum field theory; Hopf algebra of decorated rooted trees, whose coproduct identifies the divergences requiring subtraction and whose antipode achieves this; recursive symbolic code; representation of the operator product expansion; subset of diagrams whose forest structure entails a unique primitive sub divergence; zeta values from the counterterms of the minimal subtraction scheme;

T_{16} : Category one of minimum of antagonism; integral domains; unique factorization domains; Euclidean domains; space; Hopf algebra of decorated rooted trees, whose coproduct identifies the divergences requiring subtraction and whose antipode achieves this; renormalization of quantum field theory; renormalized expression for any Feynman diagram; **generalization of** Chen's Lemma for iterated integrals; representation of the Hopf algebra HRof undecorated rooted trees; subalgebra of Connes and Moscovici, corresponding to the commutative part of the Hopf algebra HTof the diffeomorphism group: it assigns to Feynman diagrams those weights

T_{17} : Category two of minimum of antagonism; integral domains; unique factorization domains; Euclidean domains; space; Hopf algebra of decorated rooted trees, whose coproduct identifies the divergences requiring subtraction and whose antipode achieves this; renormalization of quantum field theory; renormalized expression for any Feynman diagram; **generalization of** Chen's Lemma for iterated integrals; representation of the Hopf algebra HRof undecorated rooted trees; subalgebra of Connes and Moscovici, corresponding to the commutative part of the Hopf algebra HTof the diffeomorphism group: it assigns to Feynman diagrams those weights

T_{18} : Category three of minimum of antagonism; integral domains; unique factorization domains; Euclidean domains; space; Hopf algebra of decorated rooted trees, whose coproduct identifies the divergences requiring subtraction and whose antipode achieves this; renormalization of quantum field theory; renormalized expression for any Feynman diagram; **generalization of** Chen's Lemma for iterated integrals;

representation of the Hopf algebra HRof undecorated rooted trees; subalgebra of Connes and Moscovici, corresponding to the commutative part of the Hopf algebra HTof the diffeomorphism group: it assigns to Feynman diagrams those weights

Module three

Lacan himself is here not beyond reproach, since he gets sometimes **seduced (eb) by** the rhizomatic wealth of language beyond (or, rather, beneath) the formal structure that sustains it.

G_{20} : Category one of subtraction from a symbolic field all the wealth of its differences, **reducing (eb) it to a** minimum of antagonism

G_{21} : Category two of subtraction from a symbolic field all the wealth of its differences, **reducing (eb) it to a** minimum of antagonism

G_{22} : Category three of subtraction from a symbolic field all the wealth of its differences, **reducing (eb) it to a** minimum of antagonism

T_{20} : Category one of rhizomatic wealth of language beyond (or, rather, beneath) the formal structure that sustains it.

T_{21} : Category two of rhizomatic wealth of language beyond (or, rather, beneath) the formal structure that sustains it.

T_{22} : Category three of rhizomatic wealth of language beyond (or, rather, beneath) the formal structure that sustains it.

Module four

It is in this sense that, in the last decade of his teaching, he deployed the notion of lalangue (sometimes simply translated as "language") which stands for language **as the** space of illicit pleasures that defy any normativity the chaotic multitude of homonymies, word-plays, "irregular" metaphoric links and resonances.

G_{24} : Category one of space of illicit pleasures that defy any normativity the chaotic multitude of homonymies, word-plays, "irregular" metaphoric links and resonances.

G_{25} : Category two of space of illicit pleasures that defy any normativity the chaotic multitude of homonymies, word-plays, "irregular" metaphoric links and resonances.

G_{26} : Category three of space of illicit pleasures that defy any normativity the chaotic multitude of homonymies, word-plays, "irregular" metaphoric links and resonances.

T_{24} : Category one of notion of lalangue (sometimes simply translated as "language") which stands for language

T_{25} : Category two of notion of lalangue (sometimes simply translated as "language") which stands for language

T_{26} : Category three of notion of lalangue (sometimes simply translated as "language") which stands for language

Module five

Productive as this notion is, one should be aware of its limitations. Many commentators have noted that Lacan's last great literary reading, that of Joyce to whom his late seminar (XXIII: Le sinthome) is dedicated, is not at the level of his previous great readings (Hamlet, Antigone, Claudel's Coufontaine-trilogy). There is

effectively something fake in Lacan's fascination with late Joyce, with Finnegan's Wake as the latest version of the literary Gesamtkunstwerk with its endless wealth of lalangue in which not only the gap between singular languages, but the very gap between linguistic meaning and jouissance seems **overcome(e)** and the rhizome-like jouis-sense (enjoyment-in-meaning: enjoy-meant) proliferates in all directions.

G_{28} : Category one of rhizome-like jouis-sense (enjoyment-in-meaning: enjoy-meant) proliferates in all directions (See gratification deprivation complex and Bentham for quantification of enjoyment and its manifested feelings). It is important that the individual must feel the enjoyment. Otherwise there are possibilities that the gratification rate differs from individual to individual.

Civil jury instructions are inconsistent in defining what constitutes noneconomic damages, which may include pain, suffering, disability, disfigurement, and loss of enjoyment of life (LEL), among other injury sequelae. This inconsistency has been manifested recently in court decisions that have considered whether LEL should be treated as a separate element of noneconomic damages, distinct from pain and suffering. This paper reviews the case law on this issue and also describes a jury simulation experiment. Mock jurors awarded damages after they received instructions on noneconomic damages in which LEL was (1) not identified as a distinct element of damages; (2) defined an element of damages distinct from pain and suffering, but participants awarded a single amount for noneconomic damages; or (3) defi ned as a distinct element of damages, and participants awarded separate amounts for LEL and pain and suffering. Instructions about LEL resulted in larger awards, but only when mock jurors also made a separate award for that element of damages. **(Measuring Damages for Lost Enjoyment of Life: The View from the Bench and the Jury Box Susan Poser Brian H. Bornstein Erinn Kiernan McGorty)**

The paper argues that the canonical dichotomy between cardinal utility and ordinal utility is inadequate to tell the history of utility theory, and that a third form of utility consistent with the so-called classical understanding of measurement should be added to the traditional picture. According to the classical view, measuring an object consists of assessing the numerical ratio between the object and some other object taken as a unit. In particular, the paper shows that Jevons, Menger and Walras understood measurement in the classical sense, applied this understanding to utility measurement, and therefore were not Cardinalists in the current sense of the term associated with the ranking of utility differences. The paper also analyzes the argumentative strategies adopted by Jevons and Walras to address the conflict between the scientific importance they attributed to measurement, their classical understanding of it, and the apparent immeasurability of the utility featuring in their economic theories. Finally, in order to appreciate the broad intellectual context within which their discussions on utility measurement took place, the paper reviews the understanding of measurement in disciplines that bear some relation to utility theory. This review illustrates that in the years 1870-1910, the period in which Jevons, Menger and Walras were active, the classical understanding of measurement dominated not only utility theory but also all other disciplines surveyed. This circumstance helps to explain why the three marginalists remained committed to the classical understanding even though it did not square with their economic practices. **Were Jevons, Menger and Walras Really Cardinalists? On the Notion of Measurement in Utility Theory, Psychology, Mathematics and Other Disciplines, ca. 1870-1910 Ivan Moscati Bocconi University - Department of Economics JEL Classification: A12, B13, B31, B40**

G_{29} : Category two of rhizome-like jouis-sense (enjoyment-in-meaning: enjoy-meant) proliferates in all directions

G_{30} : Category three of rhizome-like jouis-sense (enjoyment-in-meaning: enjoy-meant) proliferates in all directions

The “art of the smile” lies in the clinician’s ability to recognize the positive elements of beauty in each

patient and then create a strategy to enhance the attributes that fall outside the parameters of the prevailing esthetic concept. New technologies have enhanced our ability to see our patients more dynamically and facilitated the quantification and communication of newer concepts of function and appearance. In a 2-part article, we present a comprehensive methodology for recording, assessing, and planning treatment of the smile in 4 dimensions. In part 1, we discussed the evolution of smile analysis and reviewed the dynamic records needed. In part 2, we review smile analysis and treatment strategies and present a brief case report. (Am J Orthod Dent facial Orthop 2003; 124:116-27 **Dynamic smile visualization and quantification: Part 2. Smile analysis and treatment strategies David M. Sarver, DMD, MS, a and Marc B. Ackerman, DMDb Vestavia Hills, Ala, and Bryn Mawr)**

T_{28} : Category one of endless wealth of lalangue in which not only the gap between singular languages, but the very gap between linguistic meaning and jouissance seems

T_{29} : Category two of endless wealth of lalangue in which not only the gap between singular languages, but the very gap between linguistic meaning and jouissance seems

T_{30} : Category three of endless wealth of lalangue in which not only the gap between singular languages, but the very gap between linguistic meaning and jouissance seems

Module six

The true counterpart to Joyce is, of course, Samuel Becket: after his early period in which he more or less wrote some variations on Joyce, the "true" Becket constituted himself through a true ethical act, a CUT, a **rejection(e)** of the Joycean wealth of enjoy-meant, and the ascetic turn towards a "minimal difference," towards a minimalization, "subtraction," of the narrative content and of language itself (this line is most clearly discernible in his masterpiece, the trilogy Molloy - Malone Dies - L'innomable).

G_{32} : Category one of constitution of human beings through a true ethical act, a CUT

G_{33} : Category two of constitution of human beings through a true ethical act, a CUT

G_{34} : Category three of constitution of human beings through a true ethical act, a CUT

T_{32} : Category one of **rejection(e)** of the Joycean wealth of enjoy-meant, and the ascetic turn towards a "minimal difference," towards a minimalization, "subtraction," of the narrative content and of language itself (this line is most clearly discernible in his masterpiece, the trilogy Molloy - Malone Dies - L'innomable).

T_{33} : Category two of **rejection (e)** of the Joycean wealth of enjoy-meant, and the ascetic turn towards a "minimal difference," towards a minimalization, "subtraction," of the narrative content and of language itself (this line is most clearly discernible in his masterpiece, the trilogy Molloy - Malone Dies - L'innomable).

T_{34} : Category three of **rejection (e)** of the Joycean wealth of enjoy-meant, and the ascetic turn towards a "minimal difference," towards a minimalization, "subtraction," of the narrative content and of language itself (this line is most clearly discernible in his masterpiece, the trilogy Molloy - Malone Dies - L'innomable).

Module seven

So what is the "minimal difference" - the purely parallax gap - that sustains Becket's mature production? One is tempted to propose the thesis that it is the very difference between French and English: as is known, Becket wrote most of his mature works in French (not his mother tongue), and the, desperate at the low

quality of translations, translated them himself into English, and these translations are not mere close translations, but effectively a different text. It is because of this "minimalist" - purely formal and insubstantial - status of the Real that, for Lacan, repetition **precedes (e)** repression - or, as Deleuze put it succinctly: "We do not repeat because we repress, we repress **because** (e) we repeat." (DR-105)

G_{36} : Category one of repression - or, as Deleuze put it succinctly: "We do not repeat because we repress, we repress **because** (e) we repeat." (DR-105)

G_{37} : Category two of repression - or, as Deleuze put it succinctly: "We do not repeat because we repress, we repress **because** (e) we repeat." (DR-105)

G_{38} : Category three of repression - or, as Deleuze put it succinctly: "We do not repeat because we repress, we repress **because** (e) we repeat." (DR-105)

T_{36} : Category one of "minimal difference" - the purely parallax gap - that sustains Becket's mature production? One is tempted to propose the thesis that it is the very difference between French and English: as is known, Becket wrote most of his mature works in French (not his mother tongue), and the, desperate at the low quality of translations, translated them himself into English, and these translations are not mere close translations, but effectively a different text. It is because of this "minimalist" - purely formal and insubstantial - status of the Real that, for Lacan, repetition

T_{37} : Category two of “minimal difference" - the purely parallax gap - that sustains Becket's mature production? One is tempted to propose the thesis that it is the very difference between French and English: as is known, Becket wrote most of his mature works in French (not his mother tongue), and the, desperate at the low quality of translations, translated them himself into English, and these translations are not mere close translations, but effectively a different text. It is because of this "minimalist" - purely formal and insubstantial - status of the Real that, for Lacan, repetition

T_{38} : Category three of "minimal difference" - the purely parallax gap - that sustains Becket's mature production? One is tempted to propose the thesis that it is the very difference between French and English: as is known, Becket wrote most of his mature works in French (not his mother tongue), and the, desperate at the low quality of translations, translated them himself into English, and these translations are not mere close translations, but effectively a different text. It is because of this "minimalist" - purely formal and insubstantial - status of the Real that, for Lacan, repetition

Module eight

It is not that, first, we repress some traumatic content, and then, since we are unable to remember it and thus to clarify **our relationship (e&eb)** to it, this content continues to haunt us, repeating itself in disguised forms. Note repetition Real **is a** minimal difference tion leads to astronomical gratification and deprivation levels

G_{40} : Category one of non repression of some traumatic content, and then, since we are unable to remember it and thus to clarify **our relationship (e&eb);** content continues to haunt us, repeating itself in disguised forms

G_{41} : Category two of non repression of some traumatic content, and then, since we are unable to remember it and thus to clarify **our relationship (e&eb);** content continues to haunt us, repeating itself in disguised forms

G_{42} : Category three of non repression of some traumatic content, and then, since we are unable to remember it and thus to clarify **our relationship (e&eb);** content continues to haunt us, repeating itself in

disguised forms

T_{40} : Category one of content continues to haunt us, repeating itself in disguised forms; non repression of some traumatic content, and then, since we are unable to remember it and thus to clarify **our relationship (e&eb**

T_{41} : Category two of content continues to haunt us, repeating itself in disguised forms; non repression of some traumatic content, and then, since we are unable to remember it and thus to clarify **our relationship (e&eb**

T_{42} : Category three of content continues to haunt us, repeating itself in disguised forms; non repression of some traumatic content, and then, since we are unable to remember it and thus to clarify **our relationship (e&eb**

Module Nine

If the Real **is a** minimal difference, then repetition (that establishes this difference) **is** primordial

G_{44} : Category one of Real **is a** minimal difference. Reality is relativistic

Descriptive versus normative relativism:

The concept of relativism has importance both for philosophers and for anthropologists in another way. In general, anthropologists engage in descriptive relativism, whereas philosophers engage in normative relativism, although there is some overlap (for example, descriptive relativism can pertain to concepts, normative relativism to truth).

Descriptive relativism assumes that certain cultural groups have different modes of thought, standards of reasoning, and so forth, and it is the anthropologist's task to describe, but not to evaluate the validity of these principles and practices of a cultural group. It is possible for an anthropologist in his or her fieldwork to be a descriptive relativist about some things that typically concern the philosopher (e.g., ethical principles) but not about others (e.g., logical principles). However, the descriptive relativist's empirical claims about epistemic principles, moral ideals and the like are often countered by anthropological arguments that such things are universal, and much of the recent literature on these matters is explicitly concerned with the extent of, and evidence for, cultural or moral or linguistic or human universals (see Brown, 1991 for a good discussion).

The fact that the various species of descriptive relativism are empirical claims, may tempt the philosopher to conclude that they are of little philosophical interest, but there are several reasons why this isn't so. First, some philosophers, notably Kant, argue that certain sorts of cognitive differences between human beings (or even all rational beings) are impossible, so such differences could never be found to obtain in fact, an argument that places a priori limits on what empirical inquiry could discover and on what versions of descriptive relativism could be true. Second, claims about actual differences between groups play a central role in some arguments for normative relativism (for example, arguments for normative ethical relativism often begin with claims that different groups in fact have different moral codes or ideals). Finally, the anthropologist's descriptive account of relativism helps to separate the fixed aspects of human nature from those that can vary, and so a descriptive claim that some important aspect of experience or thought does (or does not) vary across groups of human beings tells us something important about human nature and the

human condition.

Normative relativism concerns normative or evaluative claims that modes of thought, standards of reasoning, or the like are only right or wrong relative to a framework. 'Normative' is meant in a general sense, applying to a wide range of views; in the case of beliefs, for example, normative correctness equals truth. This does not mean, of course, that framework-relative correctness or truth is always clear, the first challenge being to explain what it amounts to in any given case (e.g., with respect to concepts, truth, epistemic norms). Normative relativism (say, in regard to normative ethical relativism) therefore implies that things (say, ethical claims) are not simply true in themselves, but only have truth values relative to broader frameworks (say, moral codes). (Many normative ethical relativist arguments run from premises about ethics to conclusions that assert the relativity of truth values, bypassing general claims about the nature of truth, but it is often more illuminating to consider the type of relativism under question directly.)

G_{45} : Category two of Real **is a** minimal difference

G_{46} : Category three of Real **is a** minimal difference

T_{44} : Category one of repetition (that establishes this difference) **is** primordial

T_{45} : Category two of repetition (that establishes this difference) **is** primordial

T_{46} : Category three of repetition (that establishes this difference) **is** primordial

SECTION FIFTEEN

Dark Matter: Nemo Dat Quod Non Habet - Non Est Ad Astra Mollis E Terris Via

INTRODUCTION—VARIABLES USED

Simple theory may explain dark matter 7 hours ago

(1) This is a comparison of an anapole field with common electric and magnetic dipoles. The anapole field, top, is **generated (e) by a** toroidal electrical current.
(2) **As a(eb) result**, the field is confined within the torus, instead of spreading out like the …more
(3) Most of the matter in the universe **may be (e) made out** of particles that possess an unusual, donut-shaped electromagnetic field called an anapole.
(4) This proposal, which endows dark matter particles with a rare form of electromagnetism, **has been (eb) strengthened** by a detailed analysis performed by a pair of theoretical physicists at Vanderbilt University: Professor Robert Scherrer and post-doctoral fellow Chiu Man Ho. An article about the research was published online last month by the journal Physics Letters B.

"There are a great many different theories about the nature of dark matter. What I like about this theory is its simplicity, uniqueness and the fact that it can be tested," said Scherrer.

(5) In the article, titled "Anapole Dark Matter," the physicists propose that dark matter, an invisible form of matter that makes up 85 percent of the all the matter in the universe, **may be (e) made out of a** type of basic particle called the Majorana fermion. The particle's existence was predicted in the 1930's but has stubbornly resisted detection.
(6) A number of physicists have suggested that dark matter **is made(e) from** Majorana particles,
(7) Scherrer and Ho have performed detailed calculations that demonstrate that these particles are uniquely suited **to (e) possess** a rare, donut-shaped type of electromagnetic field called an anapole.
(8) This field gives those properties that differ from those of particles that possess the more common fields possessing two poles (north and south, positive and negative) and **explain**s why they are so difficult to detect.
(9) "Most models for dark matter **assume (eb)** that it interacts through exotic forces that we do not encounter in everyday life.

NOTATION

Module One

This is a comparison of an anapole field with common electric and magnetic dipoles. The anapole field, top, is **generated (e) by a** toroidal electrical current

G_{13} : Category one of toroidal electrical current

G_{14} : Category two of toroidal electrical current

G_{15} : Category three of toroidal electrical current

T_{13} : Category one of anapole field, top

T_{14} : Category two of anapole field, top

T_{15} : Category three of anapole field, top

Module Two

As a(eb) result, the field is confined within the torus, instead of spreading out like the …more

G_{16} : Category one of comparison of an anapole field with common electric and magnetic dipoles. The anapole field, top, is **generated (e) by a** toroidal electrical current

G_{17} : Category two of comparison of an anapole field with common electric and magnetic dipoles. The

anapole field, top, is **generated (e) by a** toroidal electrical current

G_{18} : Category three of comparison of an anapole field with common electric and magnetic dipoles. The anapole field, top, is **generated (e) by a** toroidal electrical current

T_{16} : Category one of field is confined within the torus, instead of spreading out like the …more

T_{17} : Category two of field is confined within the torus, instead of spreading out like the …more

T_{18} : Category three of field is confined within the torus, instead of spreading out like the …more

Module three

Most of the matter in the universe **may be (e) made out** of particles that possess an unusual, donut-shaped electromagnetic field called an anapole

G_{20} : Category one of particles that possess an unusual, donut-shaped electromagnetic field called an anapole

G_{21} : Category two of particles that possess an unusual, donut-shaped electromagnetic field called an anapole

G_{22} : Category three of particles that possess an unusual, donut-shaped electromagnetic field called an anapole

T_{20} : Category one of matter in the universe

T_{21} : Category two of matter in the universe

T_{22} : Category three of matter in the universe

Module four

This proposal, which endows dark matter particles with a rare form of electromagnetism, **has been (eb) strengthened** by a detailed analysis performed by a pair of theoretical physicists at Vanderbilt University: Professor Robert Scherrer and post-doctoral fellow Chiu Man Ho. An article about the research was published online last month by the journal Physics Letters B.

G_{24} : Category one of proposal, which endows dark matter particles with a rare form of electromagnetism

G_{25} : Category two of proposal, which endows dark matter particles with a rare form of electromagnetism

G_{26} : Category three of proposal, which endows dark matter particles with a rare form of electromagnetism

T_{24} : Category one of particles that possess an unusual, donut-shaped electromagnetic field called an anapole.

T_{25} : Category two of particles that possess an unusual, donut-shaped electromagnetic field called an anapole.

T_{26} : Category three of particles that possess an unusual, donut-shaped electromagnetic field called an anapole.

Module five

In the article, titled "Anapole Dark Matter," the physicists propose that dark matter, an invisible form of matter that makes up 85 percent of the all the matter in the universe, **may be (e) made out of a** type of basic particle called the Majorana fermion. The particle's existence was predicted in the 1930's but has stubbornly

resisted detection

G_{28} : Category one of basic particle called the Majorana fermion. The particle's existence was predicted in the 1930's but has stubbornly resisted detection

G_{29} : Category two of basic particle called the Majorana fermion. The particle's existence was predicted in the 1930's but has stubbornly resisted detection

G_{30} : Category three of basic particle called the Majorana fermion. The particle's existence was predicted in the 1930's but has stubbornly resisted detection

T_{28} : Category one of "Anapole Dark Matter," the physicists propose that dark matter, an invisible form of matter that makes up 85 percent of the all the matter in the universe

T_{29} : Category two of "Anapole Dark Matter," the physicists propose that dark matter, an invisible form of matter that makes up 85 percent of the all the matter in the universe

T_{30} : Category three of "Anapole Dark Matter," the physicists propose that dark matter, an invisible form of matter that makes up 85 percent of the all the matter in the universe

Module six

A number of physicists have suggested that dark matter **is made(e) from** Majorana particles

G_{32} : Category one of Majorana particles

G_{33} : Category two of Majorana particles

G_{34} : Category three of Majorana particles

T_{32} : Category one of dark matter

T_{33} : Category two of dark matter

T_{34} : Category three of dark matter

Module seven

Scherrer and Ho have performed detailed calculations that demonstrate that these particles are uniquely suited **to (e) possess** a rare, donut-shaped type of electromagnetic field called an anapole

G_{36} : Category one of rare, donut-shaped type of electromagnetic field called an anapole

G_{37} : Category two of rare, donut-shaped type of electromagnetic field called an anapole

G_{38} : Category three of rare, donut-shaped type of electromagnetic field called an anapole

T_{36} : Category one of detailed calculations that demonstrate that these particles are uniquely suited

T_{37} : Category two of detailed calculations that demonstrate that these particles are uniquely suited

T_{38} : Category three of detailed calculations that demonstrate that these particles are uniquely suited

Module eight

This field gives those properties that differ from those of particles that possess the more common fields possessing two poles (north and south, positive and negative) and **explain**s why they are so difficult to detect

G_{40} : Category one of field gives those properties that differ from those of particles that possess the more common fields possessing two poles (north and south, positive and negative)

G_{41} : Category two of field gives those properties that differ from those of particles that possess the more common fields possessing two poles (north and south, positive and negative)

G_{42} : Category three of field gives those properties that differ from those of particles that possess the more common fields possessing two poles (north and south, positive and negative)

T_{40} : Category one of difficulty in detection concomitant to category one above

T_{41} : Category two of difficulty in detection concomitant to category one above

T_{42} : Category three of difficulty in detection concomitant to category one above

Module Nine

"Most models for dark matter **assume (eb)** that it interacts through exotic forces that we do not encounter in everyday life.

G_{44} : Category one of "Most models for dark matter

G_{45} : Category two of "Most models for dark matter

G_{46} : Category three of "Most models for dark matter

T_{44} : Category one of in interaction through exotic forces that we do not encounter everyday life.

T_{45} : Category two of interaction through exotic forces that we do not encounter everyday life

T_{46} : Category three of interaction through exotic forces that we do not encounter everyday life

SECTION SIXTEEN

Anapole Dark Matter: Nosce Te Ipsum - Omnia Mutantur Nos Et Mutamur In Illis Model

INTRODUCTION —VARIABLES USED

Simple theory may explain dark matter 7 hours ago

(1) Anapole dark matter **makes (e) use of** ordinary electromagnetism that you learned about in school – the same force that makes magnets stick to your refrigerator or makes a balloon rubbed on your hair stick to the ceiling," said Scherrer. "Further, the model makes very specific predictions about the rate at which **it should (eb) show up in** the vast dark matter detectors that are buried underground all over the world. These predictions show that soon the existence of anapole dark matter should either be discovered or ruled out by these experiments."

(2) Fermions are particles like the electron and quark, which are the **building (eb) blocks** of matter.

(3) Their existence was predicted by Paul Dirac in 1928. Ten years later, shortly before he disappeared mysteriously at sea, Italian physicist Ettore Majorana produced a variation of Dirac's formulation that **predicts (eb) the** existence of an electrically neutral fermion. Since then, physicists have been searching for Majorana fermions. The primary candidate has been the neutrino, but scientists have been unable to determine the basic nature of this elusive particle. The existence of dark matter was also first proposed in the 1930's to **explain (e) discrepancies i**n the rotational rate of galactic clusters.

(4) Subsequently, astronomers have discovered that the rate that stars rotate around individual galaxies is similarly out of sync. Detailed observations have shown that stars far from the center of galaxies are moving at much higher velocities than can **be (e) explained by the** amount of visible matter that the galaxies contain. Assuming that they contain a large amount of invisible "dark" matter is the most straightforward way to explain these discrepancies.

(5) Scientists hypothesize that dark matter cannot be seen in telescopes **because (e) i**t does not interact very strongly with light and other electromagnetic radiation. In fact, astronomical observations have basically ruled out the possibility that dark matter particles carry electrical charges.

(6) More recently, though, several physicists have examined dark matter particles that don't carry

electrical charges, but **have (e)** electric or magnetic dipoles. The only problem is that even these more complicated models are ruled out for Majorana particles. That is one of the reasons that Ho and Scherrer took a closer look at dark matter with an anapole magnetic moment.

A. I. Popov et al 2009 EPL 87 67004 doi:10.1209/0295-5075/87/67004 Anapole moment and spin-electric interactions in rare-earth nanoclusters

(1) Spin-electric interactions and magnetoelectric properties of triangular rare-earth molecular nanoclusters are investigated by the example of Dy3. The effective spin-electric Hamiltonian is derived **on the (e) base of** a developed quantum-mechanical model of the cluster spin structure.
(2) It is shown that a toroidal (anapole) moment **is (eb) induced in** such clusters with magnetic field exceeding some threshold value or with crossed electric and magnetic fields (or an electric current).
(3) Due to the considerable relaxation time of the state with non-zero toroidal moment, **it is possible to (e) observe** the non-equilibrium quantum linear magnetoelectric effect.

NOTATION

Module One

Anapole dark matter **makes (e) use of** ordinary electromagnetism that you learned about in school – the same force that makes magnets stick to your refrigerator or makes a balloon rubbed on your hair stick to the ceiling," said Scherrer. "Further, the model makes very specific predictions about the rate at which **it should (eb) show up in** the vast dark matter detectors that are buried underground all over the world. These predictions show that soon the existence of anapole dark matter should either be discovered or ruled out by these experiments."

G_{13} : Category one of predictions about the rate at which **it should (eb) show up**

G_{14} : Category two of predictions about the rate at which **it should (eb) show up**

G_{15} : Category three of predictions about the rate at which **it should (eb) show up**

T_{13} : Category one of vast dark matter detectors that are buried underground all over the world. These predictions show that soon the existence of anapole dark matter should either be discovered or ruled out by these experiments."
T_{14} : Category two of vast dark matter detectors that are buried underground all over the world. These predictions show that soon the existence of anapole dark matter should either be discovered or ruled out by these experiments."
T_{15} : Category three of vast dark matter detectors that are buried underground all over the world. These predictions show that soon the existence of anapole dark matter should either be discovered or ruled out by these experiments."

Module Two

Fermions are particles like the electron and quark, which are the **building (eb) blocks** of matter

G_{16} : Category one of Fermions are particles like the electron and quark,

G_{17} : Category two of Fermions are particles like the electron and quark,

G_{18} : Category three of Fermions are particles like the electron and quark,

T_{16} : Category one of matter

T_{17} : Category two of matter

T_{18} : Category three of matter

Module three

A Liam Fitzpatrick et al JCAP02(2013)004 doi:10.1088/1475-7516/2013/02/004 The effective field theory of dark matter direct detection Liam Fitzpatricka et al

(1) Authors extend and explore the general non-relativistic effective theory of dark matter (DM) direct detection. They describe the basic non-relativistic building blocks of operators and discuss their symmetry properties, writing down all Galilean-invariant operators up to quadratic order in momentum transfer arising from exchange of particles of spin 1 or less. Any DM particle theory **can be (e&eb) translated into** the coefficients of an effective operator and any effective operator can be simply related to most general description of the nuclear response.
(2) They find several operators **which lead to** novel nuclear responses.
(3) These responses differ significantly from the standard minimal WIMP cases in their relative coupling strengths to various elements, **changing (e&eb) how** the results from different experiments should be compared against each other.
(4) Response functions are evaluated for common DM targets — F, Na, Ge, I, and Xe — using standard shell model techniques. They point out that each of the nuclear responses is familiar from past studies of semi-leptonic electroweak interactions, and thus potentially testable in weak interaction studies. We provide tables of the full set of required matrix elements at finite momentum transfer for a range of common elements, making a careful and fully model-independent analysis possible. Finally, they discuss embedding non-relativistic effective theory operators into UV models of dark matter.

G_{20} : Category one of DM particle theories; responses differ significantly from the standard minimal WIMP cases in their relative coupling strengths to various elements

G_{21} : Category two of DM particle theories; responses differ significantly from the standard minimal WIMP cases in their relative coupling strengths to various elements

G_{22} : Category three of DM particle theories; responses differ significantly from the standard minimal WIMP cases in their relative coupling strengths to various elements

T_{20} : Category one of coefficients of an effective operator and any effective operator can be simply related to most general description of the nuclear response; results from different experiments should be compared against each other.

T_{21} : Category two of coefficients of an effective operator and any effective operator can be simply related to most general description of the nuclear response; results from different experiments should be compared against each other.

T_{22} : Category three of coefficients of an effective operator and any effective operator can be simply related to most general description of the nuclear response; results from different experiments should be compared against each other.

Module four

G_{24} : Category one of several operators

G_{25} : Category two of several operators

G_{26} : Category three of several operators

T_{24} : Category one of novel nuclear responses

T_{25} : Category two of novel nuclear responses

T_{26} : Category three of novel nuclear responses

Module five

Their existence was predicted by Paul Dirac in 1928. Ten years later, shortly before he disappeared mysteriously at sea, Italian physicist Ettore Majorana produced a variation of Dirac's formulation that **predicts (eb) the** existence of an electrically neutral fermion. Since then, physicists have been searching for Majorana fermions. The primary candidate has been the neutrino, but scientists have been unable to determine the basic nature of this elusive particle. The existence of dark matter was also first proposed in the 1930's to **explain (e) discrepancies i**n the rotational rate of galactic clusters

G_{28} : Category one of existence of dark matter

G_{29} : Category two of existence of dark matter

G_{30} : Category three of existence of dark matter

T_{28} : Category one of rotational rate of galactic clusters

T_{29} : Category two of rotational rate of galactic clusters

T_{30} : Category three of rotational rate of galactic clusters

Module six

Subsequently, astronomers have discovered that the rate that stars rotate around individual galaxies is similarly out of sync. Detailed observations have shown that stars far from the center of galaxies are moving at much higher velocities than can **be (e) explained by the** amount of visible matter that the galaxies contain. Assuming that they contain a large amount of invisible "dark" matter is the most straightforward way to explain these discrepancies.

G_{32} : Category one of amount of visible matter that the galaxies contain. Assuming that they contain a large amount of invisible "dark" matter is the most straightforward way to explain these discrepancies.

G_{33} : Category two of amount of visible matter that the galaxies contain. Assuming that they contain a large amount of invisible "dark" matter is the most straightforward way to explain these discrepancies.

G_{34} : Category three of amount of visible matter that the galaxies contain. Assuming that they contain a large amount of invisible "dark" matter is the most straightforward way to explain these discrepancies.

T_{32} : Category one of observations have shown that stars far from the center of galaxies are moving at much higher velocities

T_{33} : Category two of observations have shown that stars far from the center of galaxies are moving at much higher velocities

T_{34} : Category three of observations have shown that stars far from the center of galaxies are moving at much higher velocities

Module seven

Scientists hypothesize that dark matter cannot be seen in telescopes **because (e) i**t does not interact very strongly with light and other electromagnetic radiation. In fact, astronomical observations have basically ruled out the possibility that dark matter particles carry electrical charges

G_{36} : Category one of non interaction very strongly with light and other electromagnetic radiation. In fact, astronomical observations have basically ruled out the possibility that dark matter particles carry electrical charges

G_{37} : Category two of non interaction very strongly with light and other electromagnetic radiation. In fact, astronomical observations have basically ruled out the possibility that dark matter particles carry electrical charges

G_{38} : Category three of non interaction very strongly with light and other electromagnetic radiation. In fact, astronomical observations have basically ruled out the possibility that dark matter particles carry electrical charges

T_{36} : Category one of dark matter cannot be seen in telescopes

T_{37} : Category two of dark matter cannot be seen in telescopes

T_{38} : Category three of dark matter cannot be seen in telescopes

Module eight

Spin-electric interactions and magnetoelectric properties of triangular rare-earth molecular nanoclusters are investigated by the example of Dy3. The effective spin-electric Hamiltonian is derived **on the (e) base of** a developed quantum-mechanical model of the cluster spin structure

G_{40} : Category one of developed quantum-mechanical model of the cluster spin structure

G_{41} : Category two of developed quantum-mechanical model of the cluster spin structure

G_{42} : Category three of developed quantum-mechanical model of the cluster spin structure

T_{40} : Category one of Spin-electric interactions and magnetoelectric properties of triangular rare-earth molecular nanoclusters are investigated by the example of Dy3. The effective spin-electric Hamiltonian

T_{41} : Category two of Spin-electric interactions and magnetoelectric properties of triangular rare-earth molecular nanoclusters are investigated by the example of Dy3. The effective spin-electric Hamiltonian

T_{42} : Category three of Spin-electric interactions and magnetoelectric properties of triangular rare-earth molecular nanoclusters are investigated by the example of Dy3. The effective spin-electric Hamiltonian

Module Nine

It is shown that a toroidal (anapole) moment **is (eb) induced in** such clusters with magnetic field exceeding some threshold value or with crossed electric and magnetic fields (or an electric current).

G_{44} : Category one of toroidal (anapole) moment

G_{45} : Category two of toroidal (anapole) moment

G_{46} : Category three of toroidal (anapole) moment

T_{44} : Category one of clusters with magnetic field exceeding some threshold value or with crossed electric and magnetic fields (or an electric current).

T_{45} : Category two of clusters with magnetic field exceeding some threshold value or with crossed electric and magnetic fields (or an electric current).

T_{46} : Category three of clusters with magnetic field exceeding some threshold value or with crossed electric and magnetic fields (or an electric current).

SECTION SEVENTEEN

Majorana Fermions And Anapole: Quod Me Nutrit Me Destruit. Qui Tacet Consentire Model

INTRODUCTION—VARIABLES USED

Simple theory may explain dark matter 7 hours ago

(1) "Although Majorana fermions are electrically neutral, fundamental symmetries of nature **forbid (e) them** from acquiring any electromagnetic properties accept the anapole," Ho said. The existence of a magnetic anapole was predicted by the Soviet physicist Yakov Zel'dovich in 1958. Since then it has been observed in the magnetic structure of the nuclei of cesium-133 and ytterbium-174 atoms.
(2) Particles with familiar electrical and magnetic dipoles **interact (e&eb) with** electromagnetic fields even when they are stationary.
(3) Particles with anapole fields must be moving before they interact and the faster they move **the (eb)** stronger the interaction.
(4) As a result, anapole particles would have been have been much more interactive during the early days of the universe **and** would have **become (=)** less and less interactive as the universe expanded and cooled.
(5) The anapole dark matter particles suggested by Ho and Scherrer would **annihilate** in the early universe just like other proposed dark matter particles
(6) Left-over particles from the process **would (eb) form** the dark matter we see today.
(7) But **because(e)** dark matter is moving so much more slowly at the present day, and **because(e)** the anapole interaction depends on how fast it moves, these particles would have escaped detection so far, but only just barely. Read more at: http://phys.org/news/2013-06-simple-theory-dark.html#jCp
(8) Two proposed forms for dark energy **are(=) the** cosmological constant, a constant energy density filling space homogeneously, and scalar fields such as quintessence or moduli, dynamic quantities whose energy density can vary in time and space. Contributions from scalar fields that are constant in space are usually **also (e) included in** the cosmological constant.
(9) The cosmological constant is physically **equivalent (=) to** vacuum energy. Scalar fields which do change in space can be difficult to distinguish from a cosmological constant because the change may be extremely slow

A Yu Kitaev 2001 Phys.-Usp. 44 131 doi:10.1070/1063-7869/44/10S/S29 Unpaired Majorana fermions in quantum wires

(10) Certain one-dimensional Fermi systems have an energy gap in the bulk spectrum while boundary states are **described (e) by** one Majorana operator per boundary point. A finite system of length L possesses two ground states with an energy difference proportional to exp (-L/l0) and different fermionic parities.
(11) Such systems **can be used as** qubits since they are intrinsically immune to decoherence. The property of a system to have boundary Majorana fermions is expressed as a condition on the bulk electron spectrum.
(12) The condition is satisfied in the presence of an arbitrary small energy gap induced by proximity of a three-dimensional p-wave superconductor, **provided(eb) that** the normal spectrum has an odd number of Fermi points in each half of the Brillouin zone (each spin component counts separately).

NOTATION

Module One

"Although Majorana fermions are electrically neutral, fundamental symmetries of nature **forbid (e) them**

from acquiring any electromagnetic properties accept the anapole," Ho said. The existence of a magnetic anapole was predicted by the Soviet physicist Yakov Zel'dovich in 1958. Since then it has been observed in the magnetic structure of the nuclei of cesium-133 and ytterbium-174 atoms.

G_{13} : Category one of acquisition any electromagnetic properties that are accepted by the anapole

G_{14} : Category two of acquisition any electromagnetic properties that are accepted by the anapole

G_{15} : Category three of acquisition any electromagnetic properties that are accepted by the anapole

T_{13} : Category one of Majorana fermions are electrically neutral, fundamental symmetries of nature

T_{14} : Category two of Majorana fermions are electrically neutral, fundamental symmetries of nature

T_{15} : Category three of Majorana fermions are electrically neutral, fundamental symmetries of nature

Module Two

Particles with familiar electrical and magnetic dipoles **interact (e&eb) with** electromagnetic fields even when they are stationary.

G_{16} : Category one of particles with familiar electrical and magnetic dipoles; electromagnetic fields even when they are stationary.

G_{17} : Category two of particles with familiar electrical and magnetic dipoles; electromagnetic fields even when they are stationary.

G_{18} : Category three of particles with familiar electrical and magnetic dipoles; electromagnetic fields even when they are stationary.

T_{16} : Category one of electromagnetic fields even when they are stationary; particles with familiar electrical and magnetic dipoles

T_{17} : Category two of electromagnetic fields even when they are stationary; particles with familiar electrical and magnetic dipoles

T_{18} : Category three of electromagnetic fields even when they are stationary; particles with familiar electrical and magnetic dipoles

Module three

Particles with anapole fields must be moving before they interact and the faster they move **the (eb)** stronger the interaction

G_{20} : Category one of Particles with anapole fields must be moving before they interact and the faster they move

G_{21} : Category two of Particles with anapole fields must be moving before they interact and the faster they move

G_{22} : Category three of Particles with anapole fields must be moving before they interact and the faster they move

T_{20} : Category one of stronger the interaction

T_{21} : Category two of stronger the interaction

T_{22} : Category three of stronger the interaction

Module four

As a result, anapole particles would have been have been much more interactive during the early days of the universe **and** would have **become (=)** less and less interactive as the universe expanded and cooled.

G_{24} : Category one of less and less interactive as the universe expanded and cooled

G_{25} : Category two of less and less interactive as the universe expanded and cooled

G_{26} : Category three of less and less interactive as the universe expanded and cooled

T_{24} : Category one of anapole particles would have been have been much more interactive during the early days of the universe

T_{25} : Category two of anapole particles would have been have been much more interactive during the early days of the universe

T_{26} : Category three of anapole particles would have been have been much more interactive during the early days of the universe

Module five

anapole dark matter particles suggested by Ho and Scherrer would **annihilate** in the early universe just like other proposed dark matter particles

G_{28} : Category one of early universe just like other proposed dark matter particles

G_{29} : Category two of early universe just like other proposed dark matter particles

G_{30} : Category three of early universe just like other proposed dark matter particles

T_{28} : Category one of anapole dark matter particles suggested by Ho and Scherrer

T_{29} : Category two of anapole dark matter particles suggested by Ho and Scherrer

T_{30} : Category three of anapole dark matter particles suggested by Ho and Scherrer

Module six

left-over particles from the process anapole dark matter particles suggested by Ho and Scherrer would **annihilate** in the early universe just like other proposed dark matter particles **would (eb) form** the dark matter we see today

G_{32} : Category one of left-over particles from the process anapole dark matter particles suggested by Ho and Scherrer would **annihilate** in the early universe just like other proposed dark matter particles

G_{33} : Category two of left-over particles from the process anapole dark matter particles suggested by Ho and Scherrer would **annihilate** in the early universe just like other proposed dark matter particles

G_{34} : Category three of left-over particles from the process anapole dark matter particles suggested by Ho and Scherrer would **annihilate** in the early universe just like other proposed dark matter particles

T_{32} : Category one of dark matter we see today

T_{33} : Category two of dark matter we see today

T_{34} : Category three of dark matter we see today

Module seven

But **because(e)** is dark matter moving so much more slowly at the present day, and **because(e)** the anapole interaction depends on how fast it moves, there are **possibilities(eb)** these particles would have escaped detection so far, but only just barely

G_{36} : Category one of dark matter moving so much more slowly at the present day, and **because(e)** the anapole interaction depends on how fast it moves

G_{37} : Category two of dark matter moving so much more slowly at the present day, and **because(e)** the anapole interaction depends on how fast it moves

G_{38} : Category three of dark matter moving so much more slowly at the present day, and **because(e)** the anapole interaction depends on how fast it moves

T_{36} : Category one of particles would have escaped detection so far, but only just barely

T_{37} : Category two of particles would have escaped detection so far, but only just barely

T_{38} : Category three of particles would have escaped detection so far, but only just barely

Module eight

two proposed forms for dark energy **are(=) the** cosmological constant, a constant energy density filling space homogeneously, and scalar fields such as quintessence or moduli, dynamic quantities whose energy density can vary in time and space. Contributions from scalar fields that are constant in space are usually **also (e) included in** the cosmological constant

G_{40} : Category one of cosmological constant, a constant energy density filling space homogeneously, and scalar fields such as quintessence or moduli, dynamic quantities whose energy density can vary in time and space; cosmological constant

G_{41} : Category two of cosmological constant, a constant energy density filling space homogeneously, and scalar fields such as quintessence or moduli, dynamic quantities whose energy density can vary in time and space; cosmological constant

G_{42} : Category three of cosmological constant, a constant energy density filling space homogeneously, and scalar fields such as quintessence or moduli, dynamic quantities whose energy density can vary in time and space; cosmological constant

T_{40} : Category one of two proposed forms for dark energy; Contributions from scalar fields that are constant in space

T_{41} : Category two of two proposed forms for dark energy; Contributions from scalar fields that are constant in space

T_{42} : Category three of two proposed forms for dark energy; Contributions from scalar fields that are constant in space

Module Nine

The cosmological constant is physically **equivalent (=) to** vacuum energy. Scalar fields which do change in space can be difficult to distinguish from a cosmological constant because the change may be extremely slow
Certain one-dimensional Fermi systems have an energy gap in the bulk spectrum while boundary states are **described (e) by** one Majorana operator per boundary point. A finite system of length L possesses two ground states with an energy difference proportional to exp (-L/l0) and different fermionic parities

Such systems (one-dimensional Fermi systems have an energy gap in the bulk spectrum while boundary states are **described (e) by** one Majorana operator per boundary point. A finite system of length L possesses two ground states with an energy difference proportional to exp (-L/l0) and different fermionic parities) **can be used as** qubits since they are intrinsically immune to decoherence. The property of a system to have boundary Majorana fermions is expressed as a condition on the bulk electron spectrum.

condition is satisfied in the presence of an arbitrary small energy gap induced by proximity of a three-dimensional p-wave superconductor, **provided(eb) that** the normal spectrum has an odd number of Fermi points in each half of the Brillouin zone (each spin component counts separately).
G_{44} : Category one of vacuum energy; Majorana operator per boundary point. A finite system of length L possesses two ground states with an energy difference proportional to exp (-L/l0) and different fermionic parities; qubits since they are intrinsically immune to decoherence. The property of a system to have boundary Majorana fermions is expressed as a condition on the bulk electron spectrum; condition is satisfied in the presence of an arbitrary small energy gap induced by proximity of a three-dimensional p-wave superconductor;

G_{45} : Category two of vacuum energy; Majorana operator per boundary point. A finite system of length L possesses two ground states with an energy difference proportional to exp (-L/l0) and different fermionic parities; qubits since they are intrinsically immune to decoherence. The property of a system to have boundary Majorana fermions is expressed as a condition on the bulk electron spectrum; condition is satisfied in the presence of an arbitrary small energy gap induced by proximity of a three-dimensional p-wave superconductor

G_{46} : Category three of vacuum energy; Majorana operator per boundary point. A finite system of length L possesses two ground states with an energy difference proportional to exp (-L/l0) and different fermionic parities; qubits since they are intrinsically immune to decoherence. The property of a system to have boundary Majorana fermions is expressed as a condition on the bulk electron spectrum; condition is satisfied in the presence of an arbitrary small energy gap induced by proximity of a three-dimensional p-wave superconductor

T_{44} : Category one of cosmological constant change may be extremely slow; certain one-dimensional Fermi systems have an energy gap in the bulk spectrum while boundary states t; Scalar fields which do change in space can be difficult to distinguish from a cosmological constant; one-dimensional Fermi systems have an energy gap in the bulk spectrum while boundary states are **described (e) by** one Majorana operator per boundary point. A finite system of length L possesses two ground states with an energy difference proportional to exp (-L/l0) and different fermionic parities; normal spectrum has an odd number of Fermi points in each half of the Brillouin zone (each spin component counts separately).

T_{45} : Category two of cosmological constant; Scalar fields which do change in space can be difficult to distinguish from a cosmological constant; change may be extremely slow; certain one-dimensional Fermi systems have an energy gap in the bulk spectrum while boundary states; one-dimensional Fermi systems have an energy gap in the bulk spectrum while boundary states are **described (e) by** one Majorana operator

per boundary point. A finite system of length L possesses two ground states with an energy difference proportional to exp (-L/l0) and different fermionic parities; normal spectrum has an odd number of Fermi points in each half of the Brillouin zone (each spin component counts separately).

T_{46} : Category three of cosmological constant; Scalar fields which do change in space can be difficult to distinguish from a cosmological constant; change may be extremely slow; certain one-dimensional Fermi systems have an energy gap in the bulk spectrum while boundary states; one-dimensional Fermi systems have an energy gap in the bulk spectrum while boundary states are **described (e) by** one Majorana operator per boundary point. A finite system of length L possesses two ground states with an energy difference proportional to exp (-L/l0) and different fermionic parities; normal spectrum has an odd number of Fermi points in each half of the Brillouin zone (each spin component counts separately).

SECTION EIGHTEEN

Zeno's Paradox And High-Fidelity Quantum Computation: Non Causa Pro Causa-Non Bis In Idem Model

INTRODUCTION—VARIABLES USED

J. Math. Phys. 18, 756 (1977); http://dx.doi.org/10.1063/1.523304 The Zeno's paradox in quantum theory B. Misra

(1) Authors seek a quantum-theoretic expression **for (e) the** probability that an unstable particle prepared initially in a well defined state ρ will be found to decay sometime during a given interval.
(2) It is argued that probabilities like this **pertain (e&eb) to** continuous monitoring possess operational meaning.
(3) A simple natural approach to this problem **leads to the** conclusion that an unstable particle which is continuously observed to see whether it decays will never be found to decay!
(4) Since recording the track of an unstable particle (which can be distinguished from its decay

products) approximately realizes such continuous observations, the above conclusion seems **to (eb) pose** a paradox which we call Zeno's paradox in quantum theory.

(5) The relation of this result to that of some previous works and its implications and possible resolutions are briefly discussed. The mathematical transcription of the above-mentioned conclusion **is (=) a** structure theorem concerning semigroups.

(6) Although special cases of this theorem are known, the general formulation and the proof given here are believed to be new. Authors also note that the known ''no-go'' theorem concerning the semigroup law for the reduced evolution of any physical system (including decaying systems) **is (eb) subsumed** under the theorem as a direct corollary.

Efficient high-fidelity quantum computation using matter qubits and linear optics Sean D. Barrett et al

(7) Authors propose a practical, scalable, and efficient scheme for quantum computation **using (e)** spatially separated matter qubits and single-photon interference effects.

(8) The qubit systems can be nitrogen-vacancy centers in diamond, Pauli-blockade quantum dots with an excess electron, or trapped ions with optical transitions, which are each, placed in a cavity and subsequently entangled **using (e) a** double-heralded single-photon detection scheme.

(9) The fidelity of the resulting entanglement is extremely robust and remains constant **against (e&eb)** the most important errors such as detector loss, spontaneous emission, and mismatch of cavity parameters. Authors demonstrate how this entangling operation can be used to efficiently generate cluster states of many qubits, which, together with single-qubit operations and readout, can be used to implement universal quantum computation. Existing experimental parameters indicate that high-fidelity clusters can be generated with a moderate constant overhead.URL:http://link.aps.org/doi/10.1103/PhysRevA.71.060310DOI:10.1103/PhysRevA.71.060310

NOTATION

Module One

Authors seek a quantum-theoretic expression **for (e) the** probability that an unstable particle prepared initially in a well defined state ρ will be found to decay sometime during a given interval

G_{13} : Category one of probability that an unstable particle prepared initially in a well defined state ρ will be found to decay sometime during a given interval

G_{14} : Category two of probability that an unstable particle prepared initially in a well defined state ρ will be found to decay sometime during a given interval

G_{15} : Category three of probability that an unstable particle prepared initially in a well defined state ρ will be found to decay sometime during a given interval

T_{13} : Category one of quantum-theoretic expression

T_{14} : Category two of quantum-theoretic expression

T_{15} : Category three of quantum-theoretic expression

Module Two

It is argued that probabilities like this **pertain (e&eb) to** continuous monitoring possess operational meaning.

G_{16} : Category one of probabilities like the one mentioned in the foregoing (probability that an unstable particle prepared initially in a well defined state ρ will be found to decay sometime during a given

interval); continuous monitoring possess operational meaning.

G_{17} : Category two of probabilities like the one mentioned in the foregoing (probability that an unstable particle prepared initially in a well defined state ρ will be found to decay sometime during a given interval); continuous monitoring possess operational meaning.

G_{18} : Category three of probabilities like the one mentioned in the foregoing (probability that an unstable particle prepared initially in a well defined state ρ will be found to decay sometime during a given interval); continuous monitoring possess operational meaning.

T_{16} : Category one of continuous monitoring possess operational meaning; probabilities like the one mentioned in the foregoing (probability that an unstable particle prepared initially in a well defined state ρ will be found to decay sometime during a given interval)

T_{17} : Category two of continuous monitoring possesses operational meaning; probabilities like the one mentioned in the foregoing (probability that an unstable particle prepared initially in a well defined state ρ will be found to decay sometime during a given interval); continuous monitoring possess operational meaning.

T_{18} : Category three of continuous monitoring possess operational meaning; probabilities like the one mentioned in the foregoing (probability that an unstable particle prepared initially in a well defined state ρ will be found to decay sometime during a given interval); continuous monitoring possess operational meaning.

Module three

A simple natural approach to this problem **leads to the** conclusion that an **unstable particle which is continuously observed to see whether it decays will never be found to decay**

G_{20} : Category one of natural approach to this problem

G_{21} : Category two of natural approach to this problem

G_{22} : Category three of natural approach to this problem

T_{20} : Category one of unstable particle which is continuously observed to see whether it decays will never be found to decay

T_{21} : Category two of unstable particle which is continuously observed to see whether it decays will never be found to decay

T_{22} : Category three of unstable particle which is continuously observed to see whether it decays will never be found to decay

Module four

Since recording the track of an unstable particle (which can be distinguished from its decay products) approximately realizes such continuous observations, the above conclusion seems **to (eb) pose** a paradox which we call Zeno's paradox in quantum theory.

G_{24} : Category one of recording the track of an unstable particle (which can be distinguished from its decay products) approximately realizes such continuous observations, the above conclusion

G_{25} : Category two of recording the track of an unstable particle (which can be distinguished from its decay products) approximately realizes such continuous observations, the above conclusion

G_{26} : Category three of recording the track of an unstable particle (which can be distinguished from its

decay products) approximately realizes such continuous observations, the above conclusion

T_{24} : Category one of paradox which we call Zeno's paradox in quantum theory.

T_{25} : Category two of paradox which we call Zeno's paradox in quantum theory.

T_{26} : Category three of paradox which we call Zeno's paradox in quantum theory.

Module five

The mathematical transcription of the above-mentioned conclusion **is a** structure theorem concerning semigroups

In mathematics and computer science, the Krohn–Rhodes theory (or algebraic automata theory) is an approach to the study of finite semigroups and an automaton that seeks to decompose them in terms of elementary components. These components correspond to finite aperiodic semigroups and finite simple groups that are combined together in a feedback-free manner (called a "wreath product" or "cascade"). Krohn and Rhodes found a general decomposition for finite automata. In doing their research, though, the authors discovered and proved an unexpected major result in finite semigroup theory, revealing a deep connection between finite automata and semigroups. The Krohn–Rhodes theorem for finite semigroups states that every finite semigroup S is a divisor of a finite alternating wreath product of finite simple groups, each a divisor of S, and finite aperiodic semigroups (which contain no nontrivial subgroups). Wreath product of group theory is a specialized product of two groups, based on a semidirect product. Wreath products are an important tool in the classification of permutation groups and also provide a way of constructing interesting examples of groups. Semidirect product is a particular way in which a group can be constructed from two subgroups, one of which is a normal subgroup. A semidirect product is a generalization of a direct product. It is a Cartesian product as a set, but with a particular multiplication operation

G_{28} : Category one of structure theorem concerning semigroups. Note we are here expatiating upon the characterstics of various investigating systems that satisfy the foregoing condition. Needless to emphasise that there are many such systems in nature that could be investigated upon which satiate and consummate the structure theorem concerning semigroups. There are structure theorem concerning δ-classes of compact semigroup action(David Stadtlander), arbitrary completely regular semigroups (M Petric), structure theorem for compact simple semigroups(A. D. Wallace), Structure theorems for regular semigroups (Gerard Lallement), Structure theorems for weakly B-abundant semigroups (Yanhui Wang),etc.,

G_{29} : Category two of structure theorem concerning semigroups

G_{30} : Category three of structure theorem concerning semigroups

T_{28} : Category one of mathematical transcription of the above-mentioned conclusion

T_{29} : Category two of mathematical transcription of the above-mentioned conclusion

T_{30} : Category three of mathematical transcription of the above-mentioned conclusion

Module six

Although special cases of this theorem are known, the general formulation and the proof given here are believed to be new. Authors also note that the known ''no-go'' theorem concerning the semigroup law for the reduced evolution of any physical system (including decaying systems) **is (eb) subsumed** under the theorem as a direct corollary

G_{32} : Category one of known ''no-go'' theorem concerning the semigroup law for the reduced evolution of any physical system (including decaying systems)

G_{33} : Category two of known ''no-go'' theorem concerning the semigroup law for the reduced evolution of any physical system (including decaying systems)

G_{34} : Category three of known ''no-go'' theorem concerning the semigroup law for the reduced evolution of any physical system (including decaying systems)

T_{32} : Category one of structure theorem concerning semigroups

T_{33} : Category two of structure theorem concerning semigroups

T_{34} : Category three of structure theorem concerning semigroups

Module seven

Authors propose a practical, scalable, and efficient scheme for quantum computation **using (e)** spatially separated matter qubits and single-photon interference effects.

G_{36} : Category one of spatially separated matter qubits and single-photon interference effects.

G_{37} : Category two of spatially separated matter qubits and single-photon interference effects.

G_{38} : Category three of spatially separated matter qubits and single-photon interference effects.

T_{36} : Category one of scheme for quantum computation

T_{37} : Category two of scheme for quantum computation

T_{38} : Category three of scheme for quantum computation

Module eight

The qubit systems can be nitrogen-vacancy centers in diamond, Pauli-blockade quantum dots with an excess electron, or trapped ions with optical transitions, which are each, placed in a cavity and subsequently entangled **using (e) a** double-heralded single-photon detection scheme.

G_{40} : Category one of double-heralded single-photon detection scheme

G_{41} : Category two of double-heralded single-photon detection scheme

G_{42} : Category three of double-heralded single-photon detection scheme

T_{40} : Category one of the qubit systems can be nitrogen-vacancy centers in diamond, Pauli-blockade quantum dots with an excess electron, or trapped ions with optical transitions, which are each, placed in a cavity and subsequently entangled

T_{41} : Category two of the qubit systems can be nitrogen-vacancy centers in diamond, Pauli-blockade quantum dots with an excess electron, or trapped ions with optical transitions, which are each, placed in a cavity and subsequently entangled

T_{42} : Category three of the qubit systems can be nitrogen-vacancy centers in diamond, Pauli-blockade quantum dots with an excess electron, or trapped ions with optical transitions, which are each, placed in a

cavity and subsequently entangled

Module Nine

Fidelity of the resulting entanglement is extremely robust and remains constant **against (e&eb)** the most important errors such as detector loss, spontaneous emission, and mismatch of cavity parameters.

Authors demonstrate how this entangling operation can be used to efficiently generate cluster states of many qubits, which, together with single-qubit operations and readout, can be used to implement universal quantum computation. Existing experimental parameters indicate that high-fidelity clusters can be generated with a moderate constant overhead.

G_{44} : Category one of fidelity of the resulting entanglement is extremely robust and remains constant; cluster states of many qubits, which, together with single-qubit operations and readout, can be used to implement universal quantum computation

G_{45} : Category two of fidelity of the resulting entanglement is extremely robust and remains constant; cluster states of many qubits, which, together with single-qubit operations and readout, can be used to implement universal quantum computation

G_{46} : Category three of fidelity of the resulting entanglement is extremely robust and remains constant ; cluster states of many qubits, which, together with single-qubit operations and readout, can be used to implement universal quantum computation

T_{44} : Category one of the most important errors such as detector loss, spontaneous emission, and mismatch of cavity parameters; entangling operation;

T_{45} : Category two of the most important errors such as detector loss, spontaneous emission, and mismatch of cavity parameters; entangling operation;

T_{46} : Category three of the most important errors such as detector loss, spontaneous emission, and mismatch of cavity parameters; entangling operation;

SECTION NINETEEN

Negative Cosmological Constant: Omnia Causa Fiunt - Perfer Et Obdura-Dolor Hic Tibi Proderit Olim Model

INTRODUCTION—VARIABLES USED

Inflation with Negative Λ James B. Hartle, S. W. Hawking, Thomas Hertog

(1) The evolution of the universe **is (e) determined by** its quantum state.
(2) The wave function of the universe **obeys (eb) the** constraints of general relativity and in particular the Wheeler-DeWitt equation.
(3) For non-zero \Lambda, authors show that the complexified solutions of the Wheeler-DeWitt equation at large volume **have(e)** two regions in which geometries are asymptotically real.
(4) In one the histories **are (=)** Euclidean asymptotically anti-de Sitter
(5) Other history they **(=) are** Lorentzian asymptotically de Sitter.
(6) Authors illustrate this by an explicit calculation in a homogeneous isotropic minisuperspace model **with** negative \Lambda, and a scalar moving in a negative potential.
(7) It is shown that the wave function in this theory **can(eb) predict an** ensemble of inflationary universes that asymptote to de Sitter space.arXiv:1207.6653 [hep-th]

Accelerated Expansion from Negative \Lambda James B. Hartle, S.W. Hawking, Thomas Hertog e-Print: arXiv: 1205.3807 [hep-th] | PDF

(8) Wave functions specifying a quantum state of the universe **must satisfy the** constraints of general relativity, in particular the Wheeler-DeWitt equation (WDWE). Authors show for a wide class of models with non-zero cosmological constant those solutions of the WDWE **exhibit (eb) a** universal semiclassical asymptotic structure for large spatial volumes.
(9) A consequence of this asymptotic structure **is (eb) that a** wave function in a gravitational theory with a negative cosmological constant **can predict** an ensemble of asymptotically classical histories which expand with a positive effective cosmological constant.

NOTATION

Module One

The evolution of the universe **is (e) determined by** its quantum state

G_{13} : Category one of quantum state

G_{14} : Category two of quantum state

G_{15} : Category three of quantum state

T_{13} : Category one of evolution of the universe

T_{14} : Category two of evolution of the universe

T_{15} : Category three of evolution of the universe

Module Two

The wave function of the universe **obeys (eb) the** constraints of general relativity and in particular the Wheeler-DeWitt equation

G_{16} : Category one of wave function of the universe

G_{17} : Category two of wave function of the universe

G_{18} : Category three of wave function of the universe

T_{16} : Category one of constraints of general relativity and in particular the Wheeler-DeWitt equation

T_{17} : Category two of constraints of general relativity and in particular the Wheeler-DeWitt equation

T_{18} : Category three of constraints of general relativity and in particular the Wheeler-DeWitt equation

Module three

We have here assumed that the general ledger of the potential positions of everything in the universe is the Universal wave function and given a model

G_{20} : Category one of general ledger of all positional ties and states of the particles (including human beings)are in potentially

G_{21} : Category two of general ledger of all positional ties and states of the particles (including human beings)are in potentially

G_{22} : Category three of general ledger of all positional ties and states of the particles (including human beings)are in potentially

T_{20} : Category one of wave function of the universe

T_{21} : Category two of wave function of the universe

T_{22} : Category three of wave function of the universe

Module four

for non-zero Lambda, authors show that the complexified solutions of the Wheeler-DeWitt equation at large volume **have(e)** two regions in which geometries are asymptotically real.

G_{24} : Category one of two regions in which geometries are asymptotically real

G_{25} : Category two of two regions in which geometries are asymptotically real

G_{26} : Category three of two regions in which geometries are asymptotically real

T_{24} : Category one of for non-zero Lambda, authors show that the complexified solutions of the Wheeler-DeWitt equation at large volume

T_{25} : Category two of for non-zero Lambda, authors show that the complexified solutions of the Wheeler-DeWitt equation at large volume

T_{26} : Category three of for non-zero Lambda, authors show that the complexified solutions of the Wheeler-DeWitt equation at large volume

Module five

histories **are** Euclidean asymptotically anti-de Sitter

G_{28} : Category one of Euclidean asymptotically anti-de Sitter. There are many spaces that correspond to

many such systems

G_{29} : Category two of Euclidean asymptotically anti-de Sitter

G_{30} : Category three of Euclidean asymptotically anti-de Sitter

T_{28} : Category one of histories

T_{29} : Category two of histories

T_{30} : Category three of histories

Module six

Other histories they **(=) are** Lorentzian asymptotically de Sitter

G_{32} : Category one of Lorentzian asymptotically de Sitter

G_{33} : Category two of Lorentzian asymptotically de Sitter

G_{34} : Category three of Lorentzian asymptotically de Sitter

T_{32} : Category one of Other histories

T_{33} : Category two of Other histories

T_{34} : Category three of Other histories

Module seven

Authors illustrate this by an explicit calculation in a homogeneous isotropic minisuperspace model **with** negative Lambda, and a scalar moving in a negative potential

G_{36} : Category one of homogeneous isotropic minisuperspace model

G_{37} : Category two of homogeneous isotropic minisuperspace model

G_{38} : Category three of homogeneous isotropic minisuperspace model

T_{36} : Category one of negative Lambda, and a scalar moving in a negative potential; homogeneous isotropic minisuperspace model

T_{37} : Category two of negative Lambda, and a scalar moving in a negative potential; homogeneous isotropic minisuperspace model

T_{38} : Category three of negative Lambda, and a scalar moving in a negative potential; homogeneous isotropic minisuperspace model

Module eight

It is shown that the wave function in this theory **can(eb) predict an** ensemble of inflationary universes that asymptote to de Sitter space

G_{40} : Category one of wave function in this theory

G_{41} : Category two of wave function in this theory

G_{42} : Category three of wave function in this theory

T_{40} : Category one of ensemble of inflationary universes that asymptote to de Sitter space

T_{41} : Category two of ensemble of inflationary universes that asymptote to de Sitter space

T_{42} : Category three of ensemble of inflationary universes that asymptote to de Sitter space

Module Nine

Wave functions specifying a quantum state of the universe **must satisfy the** constraints of general relativity, in particular the Wheeler-DeWitt equation (WDWE). Authors show for a wide class of models with non-zero cosmological constant those solutions of the WDWE **exhibit (eb) a** universal semiclassical asymptotic structure for large spatial volumes

A consequence of this asymptotic structure **is (eb) that a** wave function in a gravitational theory with a negative cosmological constant **can predict** an ensemble of asymptotically classical histories which expand with a positive effective cosmological constant

G_{44} : Category one of Wave functions specifying a quantum state of the universe; wide class of models with non-zero cosmological constant those solutions of the WDWE; asymptotic structure

G_{45} : Category two of Wave functions specifying a quantum state of the universe; wide class of models with non-zero cosmological constant those solutions of the WDWE; asymptotic structure

G_{46} : Category three of Wave functions specifying a quantum state of the universe; wide class of models with non-zero cosmological constant those solutions of the WDWE; asymptotic structure

T_{44} : Category one of constraints of general relativity, in particular the Wheeler-DeWitt equation (WDWE); universal semiclassical asymptotic structure for large spatial volumes; wave function in a gravitational theory with a negative cosmological constant **can predict** an ensemble of asymptotically classical histories which expand with a positive effective cosmological constant

T_{45} : Category two of constraints of general relativity, in particular the Wheeler-DeWitt equation (WDWE); universal semiclassical asymptotic structure for large spatial volumes; wave function in a gravitational theory with a negative cosmological constant **can predict** an ensemble of asymptotically classical histories which expand with a positive effective cosmological constant

T_{46} : Category three of constraints of general relativity, in particular the Wheeler-DeWitt equation (WDWE); universal semiclassical asymptotic structure for large spatial volumes; wave function in a gravitational theory with a negative cosmological constant **can predict** an ensemble of asymptotically classical histories which expand with a positive effective cosmological constant

SECTION TWENTY

Numerous Landscapes Of Eternal Inflation And Lambda: Sed Ipse Spiritus Postulat Pro Nobis, Gemitibus Inenarrabilibus Model

INTRODUCTION—VARIABLES USED

Accelerated Expansion from Negative Lambda James B. Hartle, S.W. Hawking, Thomas Hertog e-Print: arXiv: 1205.3807 [hep-th] | PDF

(1) This (consequence of this asymptotic structure **is (eb) that a** wave function in a gravitational theory
(2) with a negative cosmological constant **can predict** an ensemble of asymptotically classical histories which expand with a positive effective cosmological constant) **raises the possibility** that even fundamental theories with a negative cosmological constant **can be consistent** with our low-energy observations of a classical, accelerating universe.
(3) General framework is illustrated with the specific example of the no-boundary wave function **in (eb) its** holographic form. The implications of these results for model building in string cosmology are discussed.

Local Observation in Eternal Inflation Hartle, James; Hawking, S. W.; Hertog, Thomas Physical Review Letters, vol. 106, Issue 14, id. 141302 (PhRvL Homepage) DOI: 10.1103/PhysRevLett.106.141302 Bibliographic Code: 2011PhRvL.106n1302H

(4) Authors consider landscape models **that (e) admit** several regions where the conditions for eternal inflation hold.
(5) It is shown that the no-boundary wave function **can(e) be used** to calculate small departures from homogeneity within our past light cone despite the possibility of much larger fluctuations on super horizon scales.
(6) The dominant contribution for small departures from homogeneity within our past light cone despite the possibility of much larger fluctuations on super horizon scales comes **from (e) the** history exiting eternal inflation at the lowest value of the potential.

(7) In a class of landscape models **this (eb) predicts** a tensor to scalar ratio of about 10%.

(8) In this way the no-boundary wave function **defines (eb) a** measure for the prediction of local cosmological observations.

No-boundary measure in the regime of eternal inflation Hartle, James; Hawking, S. W.; Hertog, Thomas Physical Review D, vol. 82, Issue 6, id. 063510 (PhRvD Homepage)
DOI: 10.1103/PhysRevD.82.063510

(9) The no-boundary wave function (NBWF) **specifies (eb)** a measure for prediction in cosmology that selects inflationary histories and remains well behaved for spatially large or infinite universes.

(10) This paper explores the predictions of the NBWF for linear scalar fluctuations about homogeneous and isotropic backgrounds in models **with a** single scalar field moving in a quadratic potential.

NOTATION

Module One

This (consequence of this asymptotic structure **is (eb) that a** wave function in a gravitational theory with a negative cosmological constant **can predict** an ensemble of asymptotically classical histories which expand with a positive effective cosmological constant) **raises the possibility** that even fundamental theories with a negative cosmological constant **can be consistent** with our low-energy observations of a classical, accelerating universe

G_{13} : Category one of consequence of this asymptotic structure **is (eb) that a** wave function in a gravitational theory with a negative cosmological constant **can predict** an ensemble of asymptotically classical histories which expand with a positive effective cosmological constant; fundamental theories with a negative cosmological constant

G_{14} : Category two of consequence of this asymptotic structure **is (eb) that a** wave function in a gravitational theory with a negative cosmological constant **can predict** an ensemble of asymptotically classical histories which expand with a positive effective cosmological constant; fundamental theories with a negative cosmological constant

G_{15} : Category three of consequence of this asymptotic structure **is (eb) that a** wave function in a gravitational theory with a negative cosmological constant **can predict** an ensemble of asymptotically classical histories which expand with a positive effective cosmological constant; fundamental theories with a negative cosmological constant

T_{13} : Category one of fundamental theories with a negative cosmological constant **can be consistent** with our low-energy observations of a classical, accelerating universe; low-energy observations of a classical, accelerating universe; fundamental theories with a negative cosmological constant

T_{14} : Category two of fundamental theories with a negative cosmological constant **can be consistent** with our low-energy observations of a classical, accelerating universe; low-energy observations of a classical, accelerating universe ; fundamental theories with a negative cosmological constant

T_{15} : Category three of fundamental theories with a negative cosmological constant **can be consistent** with our low-energy observations of a classical, accelerating universe; low-energy observations of a classical, accelerating universe ; fundamental theories with a negative cosmological constant

Module Two

General framework is illustrated with the specific example of the no-boundary wave function **in (eb) its** holographic form. The implications of these results for model building in string cosmology are discussed.

G_{16} : Category one of no-boundary wave function; holographic form

G_{17} : Category two of no-boundary wave function; holographic form

G_{18} : Category three of no-boundary wave function; holographic form

T_{16} : Category one of holographic form; no-boundary wave function

T_{17} : Category two of holographic form; no-boundary wave function

T_{18} : Category three of holographic form; no-boundary wave function

Module three

Authors consider landscape models **that (e) admit** several regions where the conditions for eternal inflation hold

G_{20} : Category one of several regions where the conditions for eternal inflation hold

G_{21} : Category two of several regions where the conditions for eternal inflation hold

G_{22} : Category three of several regions where the conditions for eternal inflation hold

T_{20} : Category one of landscape models

T_{21} : Category two of landscape models

T_{22} : Category three of landscape models

Module four

It is shown that the no-boundary wave function **can(e) be used** to calculate small departures from homogeneity within our past light cone despite the possibility of much larger fluctuations on super horizon scales

G_{24} : Category one of calculate small departures from homogeneity within our past light cone despite the possibility of much larger fluctuations on super horizon scales

G_{25} : Category two of calculate small departures from homogeneity within our past light cone despite the possibility of much larger fluctuations on super horizon scales

G_{26} : Category three of calculate small departures from homogeneity within our past light cone despite the possibility of much larger fluctuations on super horizon scales

T_{24} : Category one of no-boundary wave function

T_{25} : Category two of no-boundary wave function

T_{26} : Category three of no-boundary wave function

Module five

Hartle–Hawking state **:**Wikipedia

Hartle–Hawking state, named after James Hartle and Stephen Hawking, **is a proposal** concerning the state of the universe prior to the Planck epoch. Hartle–Hawking **is** essentially a no-boundary proposal that the universe **is** infinitely finite:

That there was no time before the Big Bang **because** time did not exist before the formation of spacetime associated with the Big Bang and subsequent expansion of the universe in space and time.

Hartle and Hawking suggest that if we could travel backward in time toward the beginning of the universe, we would note that quite near what might have otherwise been the beginning, time gives way **to** space such that at first there is only space and no time.

Beginnings are entities that have to do with time; **because** time did not exist before the Big Bang, the concept of a beginning of the universe is meaningless.

According to the Hartle–Hawking proposal, the universe has no origin as we would understand it: the universe **was a** singularity in both space and time, pre-Big Bang.

Thus, the Hartle–Hawking state universe has no beginning, but it is not the steady state universe of Hoyle; **it simply has** no initial boundaries in time or space.

Technical explanation

The Hartle–Hawking state **is the** wave function of the Universe–a notion meant to figure out how the Universe started–that is calculated from Feynman's path integral.

More precisely, it is a hypothetical vector in the Hilbert space of a theory of quantum gravity **that describes** this wave functional.

Hartle–Hawking state a functional of the metric tensor **defined at** a (D − 1)-dimensional compact surface, the Universe, where D is the spacetime dimension.

The precise form of the Hartle–Hawking state **is the** path integral over all D-dimensional geometries that have the required induced metric on their boundary

According to the theory time diverged from three state dimensions – as we know the time now – after the universe was at the age of the Planck time. Such a wave function of the Universe **can be shown to satisfy** the Wheeler–DeWitt equation.

G_{28} : Category one of Hartle–Hawking proposal and enumeration of the universe. Note that we are not talking that there are three hawking-Hartle proposals. Classification schema and doxa is based on the characterised based on the idiosyncrasies of the systems that are being investigated. Part of the universe, conceived universe, visible universe, perceived universe; time did not exist before the formation of spacetime associated with the Big Bang and subsequent expansion of the universe in space and time; travel backward in time toward the beginning of the universe, we would note that quite near what might have otherwise been the beginning, time gives way; time did not exist before the Big Bang, the concept of a beginning of the universe is meaningless; singularity in both space and time, pre-Big Bang.; wave function of the Universe–a notion meant to figure out how the Universe started–that is calculated from Feynman's path integral.; Hartle–Hawking state a functional of the metric tensor; path integral over all D-dimensional geometries that have the required induced metric on their boundary; According to the theory time diverged from three state dimensions – as we know the time now – after the universe was at the age of the Planck time. Such a wave function of the Universe;

G_{29} : Category two of Hartle–Hawking proposal and enumeration of the universe; time did not exist before the formation of spacetime associated with the Big Bang and subsequent expansion of the universe in space and time; travel backward in time toward the beginning of the universe, we would note that quite near what

might have otherwise been the beginning, time gives way; time did not exist before the Big Bang, the concept of a beginning of the universe is meaningless; singularity in both space and time, pre-Big Bang.; wave function of the Universe–a notion meant to figure out how the Universe started–that is calculated from Feynman's path integral; Hartle–Hawking state a functional of the metric tensor; path integral over all D-dimensional geometries that have the required induced metric on their boundary; According to the theory time diverged from three state dimensions – as we know the time now – after the universe was at the age of the Planck time. Such a wave function of the Universe;

G_{30} : Category three of Hartle–Hawking proposal and enumeration of the universe; time did not exist before the formation of spacetime associated with the Big Bang and subsequent expansion of the universe in space and time; travel backward in time toward the beginning of the universe, we would note that quite near what might have otherwise been the beginning, time gives way; time did not exist before the Big Bang, the concept of a beginning of the universe is meaningless; singularity in both space and time, pre-Big Bang.; wave function of the Universe–a notion meant to figure out how the Universe started–that is calculated from Feynman's path integral; Hartle–Hawking state a functional of the metric tensor; path integral over all D-dimensional geometries that have the required induced metric on their boundary; According to the theory time diverged from three state dimensions – as we know the time now – after the universe was at the age of the Planck time. Such a wave function of the Universe;

T_{28} : Category one of a no-boundary proposal that the universe **is** infinitely finite; there was no time before the Big Bang; space such that at first there is only space and no time;. Beginnings are entities that have to do with time; universe has no origin as we would understand it: the universe; Hartle–Hawking state9note again that the investigating systems characterstics are taken in to consideration. This does not mean that there are three hawking-Hartle states; (D − 1)-dimensional compact surface, the Universe, where D is the spacetime dimension; Hartle–Hawking state; Wheeler–DeWitt equation;

T_{29} : Category two of a no-boundary proposal that the universe **is** infinitely finite; there was no time before the Big Bang; space such that at first there is only space and no time;. Beginnings are entities that have to do with time; universe has no origin as we would understand it: the universe; Hartle–Hawking state; (D − 1)-dimensional compact surface, the Universe, where D is the spacetime dimension.; Hartle–Hawking state; Wheeler–DeWitt equation;

T_{30} : Category three of a no-boundary proposal that the universe **is** infinitely finite; there was no time before the Big Bang; space such that at first there is only space and no time.; Beginnings are entities that have to do with time; universe has no origin as we would understand it: the universe; Hartle–Hawking state; (D − 1)-dimensional compact surface, the Universe, where D is the spacetime dimension; Hartle–Hawking state; Wheeler–DeWitt equation;

Module six

The dominant contribution for small departures from homogeneity within our past light cone despite the possibility of much larger fluctuations on super horizon scales comes **from (e) the** histories exiting eternal inflation at the lowest value of the potential.
In this way the no-boundary wave function **defines (eb) a** measure for the prediction of local cosmological observations
G_{32} : Category one of histories exiting eternal inflation at the lowest value of the potential; no-boundary wave function;
G_{33} : Category two of histories exiting eternal inflation at the lowest value of the potential; no-boundary wave function;
G_{34} : Category three of histories exiting eternal inflation at the lowest value of the potential; no-boundary

wave function;

T_{32} : Category one of dominant contribution for small departures from homogeneity within our past light cone despite the possibility of much larger fluctuations on super horizon scales comes; measure for the prediction of local cosmological observations;

T_{33} : Category two of dominant contribution for small departures from homogeneity within our past light cone despite the possibility of much larger fluctuations on super horizon scales comes; measure for the prediction of local cosmological observations;

T_{34} : Category three of dominant contribution for small departures from homogeneity within our past light cone despite the possibility of much larger fluctuations on super horizon scales comes; measure for the prediction of local cosmological observations;

Module seven

The no-boundary wave function (NBWF) **specifies (eb)** a measure for prediction in cosmology that selects inflationary histories and remains well behaved for spatially large or infinite universes

G_{36} : Category one of no-boundary wave function (NBWF)

G_{37} : Category two of no-boundary wave function (NBWF)

G_{38} : Category three of no-boundary wave function (NBWF)

T_{36} : Category one of measure for prediction in cosmology that selects inflationary histories and remains well behaved for spatially large or infinite universes

T_{37} : Category two of measure for prediction in cosmology that selects inflationary histories and remains well behaved for spatially large or infinite universes

T_{38} : Category three of measure for prediction in cosmology that selects inflationary histories and remains well behaved for spatially large or infinite universes

Module eight

This paper explores the predictions of the NBWF for linear scalar fluctuations about homogeneous and isotropic backgrounds in models **with a** single scalar field moving in a quadratic potential.

G_{40} : Category one of predictions of the NBWF for linear scalar fluctuations about homogeneous and isotropic backgrounds

G_{41} : Category two of predictions of the NBWF for linear scalar fluctuations about homogeneous and isotropic backgrounds

G_{42} : Category three of predictions of the NBWF for linear scalar fluctuations about homogeneous and isotropic backgrounds

T_{40} : Category one of single scalar field moving in a quadratic potential

T_{41} : Category two of single scalar field moving in a quadratic potential

T_{42} : Category three of single scalar field moving in a quadratic potential

Module Nine(case two)

G_{44} : Category one of single scalar field moving in a quadratic potential

G_{45} : Category two of single scalar field moving in a quadratic potential

G_{46} : Category three of single scalar field moving in a quadratic potential

T_{44} : Category one of predictions of the NBWF for linear scalar fluctuations about homogeneous and isotropic backgrounds

T_{45} : Category two of predictions of the NBWF for linear scalar fluctuations about homogeneous and isotropic backgrounds

T_{46} : Category three of predictions of the NBWF for linear scalar fluctuations about homogeneous and isotropic backgrounds

SECTION TWENTY ONE

Regime Of Eternal Inflation: Non Obstante Veredicto Model

INTRODUCTION—VARIABLES USED

No-boundary measure in the regime of eternal inflation Hartle, James; Hawking, S. W.; Hertog, Thomas Physical Review D, vol. 82, Issue 6, id. 063510 (PhRvD Homepage)
DOI: 10.1103/PhysRevD.82.063510.

(1) Authors treat both the spacetime geometry of the universe **and the** observers inhabiting it quantum

mechanically.

(2) They evaluate top-down probabilities for local observations **that are (e&eb) conditioned on the** NBWF and on part of our data as observers of the universe.

(3) For models where the most probable histories do not have a regime of eternal inflation, the NBWF **predicts** homogeneity on large scales, a spectrum of observable fluctuations with a small non-Gaussian component, and a small amount of inflation in our past.

(4) By contrast, for models where the dominant histories have a regime of eternal inflation, the NBWF **predicts** significant inhomogeneity on scales much larger than the present horizon, a Gaussian spectrum of observable fluctuations, and a long period of inflation in our past.

(5) The absence or presence of non-Gaussianity in our observable universe **therefore provides** information about its global structure, assuming the NBWF.

Quantum effects can render w<-1 on cosmological scales V. K. Onemli et al

(6) Authors report on a revision of their previous computation of the renormalized expectation value of the stress-energy tensor **of (e&eb) a** massless, minimally coupled scalar with a quartic self-interaction on a locally de Sitter background.

(7) This model is important because it demonstrates that quantum effects **can lead** to violations of the weak energy condition on cosmological scales—on average, not just in fluctuations—although the effect in this particular model is far too small to be observed.

(8) The revision consists of modifying the propagator so that dimensional regularization **can (e) be used** when the dimension of the renormalized theory is not four.

(9) Although the finite part of the stress-energy tensor does not change (in D=4) from our previous result, the counterterms do. They also speculate that a certain, finite and separately conserved part of the stress tensor **can be (eb)subsumed** into a natural correction of the initial state from free Bunch-Davies vacuum.URL: http://link.aps.org/doi/10.1103/PhysRevD.70.107301DOI: 10.1103/PhysRevD.70.107301

NOTATION

Module One

Authors treat both the spacetime geometry of the universe **and the** observers inhabiting it quantum mechanically.

G_{13} : Category one of quantum mechanical space time geometry and universe and quantum mechanical observers

G_{14} : Category two of quantum mechanical space time geometry and universe and quantum mechanical observers

G_{15} : Category three ofquantum mechanical space time geometry and universe and quantum mechanical observers

T_{13} : Category one of quantum mechanical observers; quantum mechanical space time geometry and universe

T_{14} : Category two of quantum mechanical observers; quantum mechanical space time geometry and universe

T_{15} : Category three of quantum mechanical observers; quantum mechanical space time geometry and universe

Module Two

They evaluate top-down probabilities for local observations **that are (e&eb) conditioned on the** NBWF and

on part of our data as observers of the universe

G_{16} : Category one of NBWF and on part of our data as observers of the universe

G_{17} : Category two of NBWF and on part of our data as observers of the universe

G_{18} : Category three of NBWF and on part of our data as observers of the universe

T_{16} : Category one of top-down probabilities for local observations

T_{17} : Category two of top-down probabilities for local observations

T_{18} : Category three of top-down probabilities for local observations

Module three

For models where the most probable histories do not have a regime of eternal inflation, the NBWF **predicts** homogeneity on large scales, a spectrum of observable fluctuations with a small non-Gaussian component, and a small amount of inflation in our past.

G_{20} : Category one of models where the most probable histories do not have a regime of eternal inflation, the NBWF

G_{21} : Category two of models where the most probable histories do not have a regime of eternal inflation, the NBWF

G_{22} : Category three of models where the most probable histories do not have a regime of eternal inflation, the NBWF

T_{20} : Category one of homogeneity on large scales, a spectrum of observable fluctuations with a small non-Gaussian component, and a small amount of inflation in our past

T_{21} : Category two of homogeneity on large scales, a spectrum of observable fluctuations with a small non-Gaussian component, and a small amount of inflation in our past

T_{22} : Category three of homogeneity on large scales, a spectrum of observable fluctuations with a small non-Gaussian component, and a small amount of inflation in our past

Module four

By contrast, for models where the dominant histories have a regime of eternal inflation, the NBWF **predicts** significant inhomogeneity on scales much larger than the present horizon, a Gaussian spectrum of observable fluctuations, and a long period of inflation in our past

G_{24} : Category one of models where the dominant histories have a regime of eternal inflation, the NBWF

G_{25} : Category two of models where the dominant histories have a regime of eternal inflation, the NBWF

G_{26} : Category three of models where the dominant histories have a regime of eternal inflation, the NBWF

T_{24} : Category one of significant inhomogeneity on scales much larger than the present horizon, a Gaussian spectrum of observable fluctuations, and a long period of inflation in our past

T_{25} : Category two of significant inhomogeneity on scales much larger than the present horizon, a Gaussian spectrum of observable fluctuations, and a long period of inflation in our past

T_{26} : Category three of significant inhomogeneity on scales much larger than the present horizon, a

Gaussian spectrum of observable fluctuations, and a long period of inflation in our past

Module five

The absence or presence of non-Gaussianity in our observable universe **therefore provides** information about its global structure, assuming the NBWF.

G_{28} : Category one of absence or presence of non-Gaussianity in our observable universe

G_{29} : Category two of absence or presence of non-Gaussianity in our observable universe

G_{30} : Category three of absence or presence of non-Gaussianity in our observable universe

T_{28} : Category one of information about its global structure, assuming the NBWF.

T_{29} : Category two of information about its global structure, assuming the NBWF.

T_{30} : Category three of information about its global structure, assuming the NBWF.

Module six

Authors report on a revision of their previous computation of the renormalized expectation value of the stress-energy tensor **of (e&eb) a** massless, minimally coupled scalar with a quartic self-interaction on a locally de Sitter background

G_{32} : Category one of renormalized expectation value of the stress-energy tensor

G_{33} : Category two of renormalized expectation value of the stress-energy tensor

G_{34} : Category three of renormalized expectation value of the stress-energy tensor

T_{32} : Category one of massless, minimally coupled scalar with a quartic self-interaction on a locally de Sitter background ;renormalized expectation value of the stress-energy tensor

T_{33} : Category two of massless, minimally coupled scalar with a quartic self-interaction on a locally de Sitter background ;renormalized expectation value of the stress-energy tensor

T_{34} : Category three of massless, minimally coupled scalar with a quartic self-interaction on a locally de Sitter background; renormalized expectation value of the stress-energy tensor

Module seven

This model is important because it demonstrates that quantum effects **can lead** to violations of the weak energy condition on cosmological scales—on average, not just in fluctuations—although the effect in this particular model is far too small to be observed

G_{36} : Category one of quantum effects

G_{37} : Category two of quantum effects

G_{38} : Category three of quantum effects

T_{36} : Category one of violations of the weak energy condition on cosmological scales—on average, not just in fluctuations—although the effect in this particular model is far too small to be observed

T_{37} : Category two of violations of the weak energy condition on cosmological scales—on average, not just in fluctuations—although the effect in this particular model is far too small to be observed

T_{38} : Category three of violations of the weak energy condition on cosmological scales—on average, not just in fluctuations—although the effect in this particular model is far too small to be observed

Module eight

The revision consists of modifying the propagator so that dimensional regularization **can (e) be used** when the dimension of the renormalized theory is not four

G_{40} : Category one of dimension of the renormalized theory is not four

G_{41} : Category two of dimension of the renormalized theory is not four

G_{42} : Category three of dimension of the renormalized theory is not four


T_{40} : Category one of modifying the propagator so that dimensional regularization

T_{41} : Category two of modifying the propagator so that dimensional regularization

T_{42} : Category three of modifying the propagator so that dimensional regularization

Module Nine

Although the finite part of the stress-energy tensor does not change (in D=4) from our previous result, the counterterms do. They also speculate that a certain, finite and separately conserved part of the stress tensor **can be (eb)subsumed** into a natural correction of the initial state from free Bunch-Davies vacuum



G_{44} : Category one of finite part of the stress-energy tensor does not change (in D=4) but speculate that a certain, finite and separately conserved part of the stress tensor; classical mechanics; classical mechanics; quantum mechanics; special relativity; quantum field theory; general relativity; Newtonian gravity; electromagnetism

G_{45} : Category two of finite part of the stress-energy tensor does not change (in D=4) but speculate that a certain, finite and separately conserved part of the stress tensor; classical mechanics; classical mechanics; quantum mechanics; special relativity; quantum field theory; general relativity; Newtonian gravity; electromagnetism;

G_{46} : Category three of finite part of the stress-energy tensor does not change (in D=4) but speculate that a certain, finite and separately conserved part of the stress tensor; classical mechanics; classical mechanics; quantum mechanics; special relativity; quantum field theory; general relativity; Newtonian gravity; electromagnetism;

T_{44} : Category one of natural correction of the initial state from free Bunch-Davies vacuum; special relativity; quantum mechanics; quantum field theory ; quantum field theory ;quantum field theory in curved space and time; quantum field theory in curved space and time; general relativity; quantum field theory;

T_{45} : Category two of natural correction of the initial state from free Bunch-Davies vacuum; quantum field theory ; quantum field theory ; quantum field theory in space and time; quantum field theory in curved space and time; general relativity; quantum field theory;

T_{46} : Category three of natural correction of the initial state from free Bunch-Davies vacuum; quantum field theory ; quantum field theory ; quantum field theory in space and time; quantum field theory in curved space and time; general relativity; quantum field theory;

SECTION TWENTY TWO

Quantum Dynamics And Casimir Energy: Eheu Fugaces Labuntur Anni

INTRODUCTION—VARIABLES USED

Quantum dynamics of Kaluza-Klein theories Thomas Appelquist

(1) Some of the quantum properties **of** Kaluza-Klein theories are studied. The classical features of these theories are reviewed, and the quantization of the gravitational field in an arbitrary number of dimensions is described. These results are then applied to a detailed analysis of the five-dimensional Kaluza-Klein model. The fifth dimension **is taken (=) to be** compact and a quantum effective potential, as a function of the five-five component of the metric, is constructed. It is argued that the one-loop computation **is (eb) reliable** as long as the distance around the fifth dimension is large compared to the Planck length.
(2) The effective potential **separates i**nto two pieces: an induced cosmological constant, independent of the size of the fifth dimension, and a distance-dependent "Casimir" energy.
(3) The cosmological term is subtracted, leaving an attractive Casimir potential **which (e) will contract** the fifth dimension to a size on the order of the Planck length. Consequences of this result are discussed and some of the ways in which it can be generalized are outlined. URL:http://link.aps.org/doi/10.1103/PhysRevD.28.772DOI: 10.1103/PhysRevD.28.772

Relaxing the cosmological constant N.C. Tsamis et al http://dx.doi.org/10.1016/0370-2693 (93)91162-G

(4) A non-zero cosmological constant **allows** massless gravitons to self-interact via a coupling of dimension three.
(5) If the cosmological constant is positive **then (eb)** the background geometry will subject gravitons to an enormous redshift.
(6) These two facts are together **responsible (eb) for** the severe infrared divergences which detailed, explicit calculations [N.C. Tsamis and R.P. Woodard, Strong infrared effects in quantum gravity, preprint CRETE-92-17, UFIFT-92-24] reveal in loop corrections to the gravitational force law.
(7) Authors argue that, as a result, the cosmological interaction **has (e) a** finite lifetime.
(8) Furthermore, this lifetime **is consistent** with inflation.

Decoherence and the transition from quantum to classical – REVISITED Wojciech H. Zurek

(9) The environment surrounding a quantum system can, in effect, **monitor** some of the systems observables. As a **result,(eb) the** eigenstates of these observables continuously decohere and can behave like classical states. ArXiv: quant-ph/0306072

NOTATION

Module One

Some of the quantum properties **of** Kaluza-Klein theories are studied. The classical features of these theories are reviewed, and the quantization of the gravitational field in an arbitrary number of dimensions is described. These results are then **applied to** a detailed analysis of the five-dimensional Kaluza-Klein model. The fifth dimension **is taken to be** compact and a quantum effective potential, as a function of the five-five

component of the metric, is constructed. It is argued that the one-loop computation **is (eb) reliable** as long as the distance around the fifth dimension is large compared to the Planck length

G_{13} : Category one of quantization of the gravitational field in an arbitrary number of dimensions and quantum properties **of** Kaluza-Klein theories

G_{14} : Category two of quantization of the gravitational field in an arbitrary number of dimensions and quantum properties **of** Kaluza-Klein theories

G_{15} : Category three of quantization of the gravitational field in an arbitrary number of dimensions and quantum properties **of** Kaluza-Klein theories

T_{13} : Category one of five-dimensional Kaluza-Klein model with fifth dimension **taken to be** compact and a quantum effective potential, as a function of the five-five component of the metric

T_{14} : Category two of five-dimensional Kaluza-Klein model with fifth dimension **taken to be** compact and a quantum effective potential, as a function of the five-five component of the metric

T_{15} : Category three of five-dimensional Kaluza-Klein model with fifth dimension **taken to be** compact and a quantum effective potential, as a function of the five-five component of the metric

Module Two

The effective potential **separates i**nto two pieces: an induced cosmological constant, independent of the size of the fifth dimension, and a distance-dependent "Casimir" energy.

G_{16} : Category one of induced cosmological constant independent of the size of the fifth dimension,

G_{17} : Category two of induced cosmological constant independent of the size of the fifth dimension,

G_{18} : Category three of induced cosmological constant independent of the size of the fifth dimension,

T_{16} : Category one of distance-dependent "Casimir" energy

T_{17} : Category two of distance-dependent "Casimir" energy

T_{18} : Category three of distance-dependent "Casimir" energy

Module three

G_{20} : Category one of effective potential

G_{21} : Category two of effective potential

G_{22} : Category three of effective potential

T_{20} : Category one of induced cosmological constant independent of the size of the fifth dimension and distance-dependent "Casimir" energy

T_{21} : Category two of induced cosmological constant independent of the size of the fifth dimension and distance-dependent "Casimir" energy

T_{22} : Category three of induced cosmological constant independent of the size of the fifth dimension and distance-dependent "Casimir" energy

Module four

The cosmological term is subtracted, leaving an attractive Casimir potential **which (e) will contract** the fifth dimension to a size on the order of the Planck length. Consequences of this result are discussed and some of

the ways in which it can be generalized are outlined

G_{24} : Category one of fifth dimension to a size on the order of the Planck length.

G_{25} : Category two of fifth dimension to a size on the order of the Planck length.

G_{26} : Category three of fifth dimension to a size on the order of the Planck length.

T_{24} : Category one of cosmological term is subtracted, leaving an attractive Casimir potential

T_{25} : Category two of cosmological term is subtracted, leaving an attractive Casimir potential

T_{26} : Category three of cosmological term is subtracted, leaving an attractive Casimir potential

Module five

A non-zero cosmological constant **allows** massless gravitons to self-interact via a coupling of dimension three

G_{28} : Category one of non-zero cosmological constant

G_{29} : Category two of non-zero cosmological constant

G_{30} : Category three of non-zero cosmological constant

T_{28} : Category one of massless gravitons to self-interact via a coupling of dimension three

T_{29} : Category two of massless gravitons to self-interact via a coupling of dimension three

T_{30} : Category three of massless gravitons to self-interact via a coupling of dimension three

Module six

If the cosmological constant is positive **then (eb)** the background geometry will subject gravitons to an enormous redshift

G_{32} : Category one of cosmological constant is positive. Note the systemic classification and the investigating systems and their characterstics form the fundamental basis for the classification schema.

G_{33} : Category two of cosmological constant is positive

G_{34} : Category three of cosmological constant is positive

T_{32} : Category one of background geometry will subject gravitons to an enormous redshift

T_{33} : Category two of background geometry will subject gravitons to an enormous redshift

T_{34} : Category three of background geometry will subject gravitons to an enormous redshift

Module seven

These two facts are together **responsible (eb) for** the severe infrared divergences which detailed, explicit calculations

G_{36} : Category one of non-zero cosmological constant **allows** massless gravitons to self-interact via a coupling of dimension three and cosmological constant positivity **then (eb)** the background geometry will subject gravitons to an enormous redshift

G_{37} : Category two of non-zero cosmological constant **allows** massless gravitons to self-interact via a coupling of dimension three and cosmological constant positivity **then (eb)** the background geometry will subject gravitons to an enormous redshift

G_{38} : Category three of non-zero cosmological constant **allows** massless gravitons to self-interact via a coupling of dimension three and cosmological constant positivity **then (eb)** the background geometry will subject gravitons to an enormous redshift

T_{36} : Category one of severe infrared divergences

T_{37} : Category two of severe infrared divergences

T_{38} : Category three of severe infrared divergences

Module eight

The environment surrounding a quantum system can, in effect, **monitor** some of the systems observables. As a **result,(eb) the** eigenstates of these observables continuously decohere and can behave like classical states

G_{40} : Category one of environment surrounding a quantum system can, in effect, **monitor** some of the systems observables

G_{41} : Category two of environment surrounding a quantum system can, in effect, **monitor** some of the systems observables

G_{42} : Category three of environment surrounding a quantum system can, in effect, **monitor** some of the systems observables

T_{40} : Category one of eigenstates of these observables continuously decohere and can behave like classical states

T_{41} : Category two of eigenstates of these observables continuously decohere and can behave like classical states

T_{42} : Category three of eigenstates of these observables continuously decohere and can behave like classical states

Module Nine

Authors argue that, as a result, the cosmological interaction **has (e) a** finite lifetime. Furthermore, this lifetime **is consistent** with inflation

G_{44} : Category one of finite life time consistent with inflation

G_{45} : Category two of finite life time consistent with inflation

G_{46} : Category three of finite life time consistent with inflation

T_{44} : Category one of non-zero cosmological constant **allows** massless gravitons to self-interact via a coupling of dimension three and cosmological constant positivity **then (eb)** the background geometry will subject gravitons to an enormous redshift, by virtue of which cosmological interaction

T_{45} : Category two of non-zero cosmological constant **allows** massless gravitons to self-interact via a coupling of dimension three and cosmological constant positivity **then (eb)** the background geometry will subject gravitons to an enormous redshift, by virtue of which cosmological interaction

T_{46} : Category three of non-zero cosmological constant **allows** massless gravitons to self-interact via a coupling of dimension three and cosmological constant positivity **then (eb)** the background geometry will subject gravitons to an enormous redshift, by virtue of which cosmological interaction

SECTION TWENTY THREE

Stylobate Problem In Many-Worlds Theories: Saltus In Demonstrando-Salva Veritate Model

INTRODUCTION—VARIABLES USED

The basic problem in many-worlds theories H P Stapp Canadian Journal of Physics, 2002, 80(9): 1043-1052, 10.1139/p02-068

(1) It is emphasized that a many-worlds interpretation of quantum theory **exists (EB) only** to the extent that the associated basis problem is solved. The core basis problem is that the robust enduring states specified by environmental decoherence effects **are (=) essentially** Gaussian wave packets that form continua of non-orthogonal states.

(2) Hence they are not a discrete set of orthogonal basis states to which finite probabilities **can be**

(EB) assigned by the usual rules.

(3) The natural way to get an orthogonal basis without going outside the Schrödinger dynamics **is to (E) use the** eigenstates of the reduced density matrix, and this idea is the basis of some recent attempts by many-worlds proponents to solve the basis problem.

(4) But these eigenstates do not enjoy the locality and quasi-classicality properties of the states **defined (E) by** environmental decoherence effects, and hence are not satisfactory preferred basis states.This core problem **is (EB) obscured** by approaches that treat the universe as a quantum computer, but it needs to be addressed and resolved before a many-worlds-type interpretation can be said to exist.

Feedback control of quantum systems using continuous state estimation. C. Doherty et al

(5) Authors present a formulation of feedback in quantum systems in which the best estimates of the dynamical variables **are (e) obtained** continuously from the measurement record, and fed back to control the system.

(6) They apply this method to the problem of cooling and confining a single quantum degree of freedom, **and compare it** to current schemes in which the measurement signal is fed back directly in the manner usually considered in existing treatments of quantum feedback.


(7) Direct feedback may be combined with feedback by estimation, and the resulting combination, performed on a linear system, is closely **analogou**s(=) to classical linear-quadratic-Gaussian control theory with residual feedback.URL:http://link.aps.org/doi/10.1103/PhysRevA.60.2700DOI:10.1103/PhysRevA.60.2700

Kelemen filter; (Wikipedia)

NOTATION

Module One

Kelemen filter; (Wikipedia):



G_{13} : Category one of prior knowledge of the state ; Next time step; $P_{k-1/k-1;}$; $x_{k-1/k-1}$; $P_{k-1/k-1;}$; $x_{k-1/k-1}$; prediction step

G_{14} : Category two of prior knowledge of the state; Next time step; $P_{k-1/k-1;}$; $x_{k-1/k-1}$; $P_{k-1/k-1;}$; $x_{k-1/k-1}$; prediction step

G_{15} : Category three of prior knowledge of the state; Next time step ; $P_{k-1/k-1;}$; $x_{k-1/k-1}$; $P_{k-1/k-1;}$; $x_{k-1/k-1}$; prediction step

T_{13} : Category one of $P_{k-1/k-1;}$; $x_{k-1/k-1}$; $P_{k-1/k-1;}$; $x_{k-1/k-1}$; prediction step; update step

T_{14} : Category two of $P_{k-1/k-1;}$; $x_{k-1/k-1}$; $P_{k-1/k-1;}$; $x_{k-1/k-1}$; prediction step; update step

T_{15} : Category three of $P_{k-1/k-1;}$; $x_{k-1/k-1}$; $P_{k-1/k-1;}$; $x_{k-1/k-1}$; prediction step; update step

Module Two

It is emphasized that a many-worlds interpretation of quantum theory **exists (EB) only** to the extent that the associated basis problem is solved. The core basis problem is that the robust enduring states specified by environmental decoherence effects **are essentially** Gaussian wave packets that form continua of non-orthogonal states.

G_{16} : Category one of Gaussian wave packets that form continua of non-orthogonal states.

G_{17} : Category two of Gaussian wave packets that form continua of non-orthogonal states.

G_{18} : Category three of Gaussian wave packets that form continua of non-orthogonal states.

T_{16} : Category one of core basis problem is that the robust enduring states specified by environmental decoherence effects

T_{17} : Category two of core basis problem is that the robust enduring states specified by environmental decoherence effects

T_{18} : Category three of core basis problem is that the robust enduring states specified by environmental decoherence effects

Module three

Hence they are not a discrete set of orthogonal basis states to which finite probabilities **can be (EB) assigned** by the usual rules

G_{20} : Category one of assignment of finite probabilities by usual rules and axiomatic predications and postulation alcovishness

G_{21} : Category two of assignment of finite probabilities by usual rules and axiomatic predications and postulation alcovishness

G_{22} : Category three of assignment of finite probabilities by usual rules and axiomatic predications and postulation alcovishness

T_{20} : Category one of non existence of discrete set of orthogonal basis states

T_{21} : Category two of non existence of discrete set of orthogonal basis states

T_{22} : Category three of non existence of discrete set of orthogonal basis states

Module four

The natural way to get an orthogonal basis without going outside the Schrödinger dynamics **is to (E) use the** eigenstates of the reduced density matrix, and this idea is the basis of some recent attempts by many-worlds

proponents to solve the basis problem

G_{24} : Category one of eigenstates of the reduced density matrix, and this idea is the basis of some recent attempts by many-worlds proponents to solve the basis problem

G_{25} : Category two of eigenstates of the reduced density matrix, and this idea is the basis of some recent attempts by many-worlds proponents to solve the basis problem

G_{26} : Category three of eigenstates of the reduced density matrix, and this idea is the basis of some recent attempts by many-worlds proponents to solve the basis problem

T_{24} : Category one of natural way to get an orthogonal basis without going outside the Schrödinger dynamics

T_{25} : Category two of natural way to get an orthogonal basis without going outside the Schrödinger dynamics

T_{26} : Category three of natural way to get an orthogonal basis without going outside the Schrödinger dynamics

Module five

But these eigenstates do not enjoy the locality and quasi-classicality properties of the states **defined (E) by** environmental decoherence effects, and hence are not satisfactory preferred basis states.

G_{28} : Category one of environmental decoherence effects, and hence are not satisfactory preferred basis states

G_{29} : Category two of environmental decoherence effects, and hence are not satisfactory preferred basis states

G_{30} : Category three of environmental decoherence effects, and hence are not satisfactory preferred basis states

T_{28} : Category one of eigenstates do not enjoy the locality and quasi-classicality properties of the states

T_{29} : Category two of eigenstates do not enjoy the locality and quasi-classicality properties of the states

T_{30} : Category three of eigenstates do not enjoy the locality and quasi-classicality properties of the states

Module six

Authors present a formulation of feedback in quantum systems in which the best estimates of the dynamical variables **are (e) obtained** continuously from the measurement record, and fed back to control the system.

G_{32} : Category one of **obtained** continuously from the measurement record, and fed back to control the system

G_{33} : Category two of **obtained** continuously from the measurement record, and fed back to control the system

G_{34} : Category three of **obtained** continuously from the measurement record, and fed back to control the

system

T_{32} : Category one of formulation of feedback in quantum systems in which the best estimates of the dynamical variables

T_{33} : Category two of formulation of feedback in quantum systems in which the best estimates of the dynamical variables

T_{34} : Category three of formulation of feedback in quantum systems in which the best estimates of the dynamical variables

Module seven

They apply this method to the problem of cooling and confining a single quantum degree of freedom, **and compare it** to current schemes in which the measurement signal is fed back directly in the manner usually considered in existing treatments of quantum feedback.

G_{36} : Category one of formulation of feedback in quantum systems in which the best estimates of the dynamical variables **are (e) obtained** continuously from the measurement record, and fed back to control the system

G_{37} : Category two of formulation of feedback in quantum systems in which the best estimates of the dynamical variables **are (e) obtained** continuously from the measurement record, and fed back to control the system

G_{38} : Category three of formulation of feedback in quantum systems in which the best estimates of the dynamical variables **are (e) obtained** continuously from the measurement record, and fed back to control the system

T_{36} : Category one of problem of cooling and confining a single quantum degree of freedom, **and compare it** to current schemes in which the measurement signal is fed back directly in the manner usually considered in existing treatments of quantum feedback

T_{37} : Category two of problem of cooling and confining a single quantum degree of freedom, **and compare it** to current schemes in which the measurement signal is fed back directly in the manner usually considered in existing treatments of quantum feedback

T_{38} : Category three of problem of cooling and confining a single quantum degree of freedom, **and compare it** to current schemes in which the measurement signal is fed back directly in the manner usually considered in existing treatments of quantum feedback

Module eight

Direct feedback may be combined with feedback by estimation, and the resulting combination, performed on a linear system, is closely **analogou**s to classical linear-quadratic-Gaussian control theory with residual feedback.

G_{40} : Category one of classical linear-quadratic-Gaussian control theory with residual feedback.

G_{41} : Category two of classical linear-quadratic-Gaussian control theory with residual feedback.

G_{42} : Category three of classical linear-quadratic-Gaussian control theory with residual feedback.

T_{40} : Category one of Direct feedback may be combined with feedback by estimation, and the resulting combination, performed on a linear system

T_{41} : Category two of Direct feedback may be combined with feedback by estimation, and the resulting combination, performed on a linear system

T_{42} : Category three of Direct feedback may be combined with feedback by estimation, and the resulting combination, performed on a linear system

Module Nine(Unsolved Problem)

Let a Single Input Single Output (SISO) transfer function G(s) be given. Find transfer functions G0(s) and H(s) such that:

1. G (s) = G0 (H (s));

2. H (s) **preserves (eb)** proper stable transfer functions under substitution of the variable s by H (s), and:

3. The degree of the denominator of H(s) **is the** maximum with the properties 1 and 2.

The resultant equation provides such a function with of course the initial value given.

G_{44} : Category one of G0 (H (s)); H (s); maximum with the properties 1 and 2.

G_{45} : Category two of G0 (H (s)); H (s); maximum with the properties 1 and 2.

G_{46} : Category three of G0 (H (s)); H (s); maximum with the properties 1 and 2.

T_{44} : Category one of G (s); proper stable transfer functions under substitution of the variable s by H (s); degree of the denominator of H(s)

T_{45} : Category two of G (s); proper stable transfer functions under substitution of the variable s by H (s); degree of the denominator of H(s)

T_{46} : Category three of G (s); proper stable transfer functions under substitution of the variable s by H (s); degree of the denominator of H(s)

SECTION TWENTY FOUR

Coherent Quantum Control: Non Est Ad Astra Mollis E Terris Via Model

INTRODUCTION—VARIABLES USED

Nature 396, 239-242 (19 November 1998) | doi:10.1038/24329; Coherent quantum control of two-photon transitions by a femtosecond laser pulse Doron Meshulach & Yaron Silberberg

(1) Coherent quantum control1, 2, 3 has attracted interest **as a (e) means** to influence the outcome of a quantum-mechanical interaction.
(2) In principle, the quantum system can be **steered (e&eb) towards** a desired state by its interaction with light.
(3) For example, in photoinduced transitions between atomic energy levels, quantum interference effects **can lead to** enhancement or cancellation of the total transition probability.
(4) The interference depends on the spectral phase distribution of the incident beam; as this phase distribution can be tuned, one can **control(e&eb)** the outcome of the interaction can in principle be controlled.
(5) Here authors demonstrate that a femtosecond laser pulse can be tailored, **using (e)** ultrashort pulse-shaping4, 5, 6, 7 techniques, to control two-photon transitions in caesium.
(6) By varying the spectral phases of the pulse components, observe the predicted cancellation of the transitions **due to** destructive quantum interference; the power spectrum and energy of these 'dark pulses' are unchanged.
(7) They also show that the pulse shape **can be (e&eb) modified** extensively without affecting the two-photon transition probability.

Elegance and Enigma The Frontiers Collection 2011, pp 87-108 Quantum States Maximilian Schlosshauer

(8) ONCE UPON A TIME, there was a quantum state. He considered himself someone who was leading a purposeful life, because he knew that physicists around the world relied on him for doing their fancy calculations and experiments. And he helped them get along as best as he could, and this was something he did indeed extremely well, and he was proud of his role. Yet in his more introspective and philosophical moments, he couldn’t quite shake the feeling that he had been languishing in a sort of perennial **identity crisis.**

Science 9 April 2004: Vol. 304 no. 5668 pp. 270-273 DOI: 10.1126/science.1095374 Real-Time Quantum Feedback Control of Atomic Spin-Squeezing JM Geremia et al

(9) Real-time feedback performed during a quantum nondemolition measurement of atomic spin-angular momentum **allowed (eb) to** influence the quantum statistics of the measurement outcome.

(10) They showed that it is possible **to harness** measurement backaction as a form of actuation in quantum control, and thus describe a valuable tool for quantum information science.Feedback-mediated procedure **generates** spin-squeezing, for which the reduction in quantum uncertainty and resulting atomic entanglement are not conditioned on the measurement outcome.

NOTATION

Module One

Coherent quantum control1, 2, 3 has attracted interest **as a (e) means** to influence the outcome of a quantum-mechanical interaction

G_{13} : Category one of influence the outcome of a quantum-mechanical interaction

G_{14} : Category two of influence the outcome of a quantum-mechanical interaction

G_{15} : Category three of influence the outcome of a quantum-mechanical interaction

T_{13} : Category one of Coherent quantum control1, 2, 3 has attracted interest

T_{14} : Category two of Coherent quantum control1, 2, 3 has attracted interest

T_{15} : Category three of Coherent quantum control1, 2, 3 has attracted interest

Module Two

In principle, the quantum system can be **steered (e&eb) towards** a desired state by its interaction with light.

G_{16} : Category one of desired state by its interaction with light

G_{17} : Category two of desired state by its interaction with light

G_{18} : Category three of desired state by its interaction with light

T_{16} : Category one of quantum system can be **steered**

T_{17} : Category two of quantum system can be **steered**

T_{18} : Category three of quantum system can be **steered**

Module three

For example, in photoinduced transitions between atomic energy levels, quantum interference effects **can lead to** enhancement or cancellation of the total transition probability.

G_{20} : Category one of photoinduced transitions between atomic energy levels, quantum interference effects

G_{21} : Category two of photoinduced transitions between atomic energy levels, quantum interference effects

G_{22} : Category three of photoinduced transitions between atomic energy levels, quantum interference effects

T_{20} : Category one of enhancement or cancellation of the total transition probability

T_{21} : Category two of enhancement or cancellation of the total transition probability

T_{22} : Category three of enhancement or cancellation of the total transition probability

Module four

The interference depends on the spectral phase distribution of the incident beam; as this phase distribution

can be tuned, one can **control(e&eb)** the outcome of the interaction can in principle be controlled.

G_{24} : Category one of interference depends on the spectral phase distribution of the incident beam; as this phase distribution can be tuned,

G_{25} : Category two of interference depends on the spectral phase distribution of the incident beam; as this phase distribution can be tuned,

G_{26} : Category three of interference depends on the spectral phase distribution of the incident beam; as this phase distribution can be tuned,

T_{24} : Category one of outcome of the interaction can in principle be controlled; interference depends on the spectral phase distribution of the incident beam; as this phase distribution can be tuned,

T_{25} : Category two of outcome of the interaction can in principle be controlled; interference depends on the spectral phase distribution of the incident beam; as this phase distribution can be tuned,

T_{26} : Category three of outcome of the interaction can in principle be controlled ;interference depends on the spectral phase distribution of the incident beam; as this phase distribution can be tuned,

Module five

Here authors demonstrate that a femtosecond laser pulse can be tailored, **using (e)** ultrashort pulse-shaping4, 5, 6, 7 techniques, to control two-photon transitions in caesium

G_{28} : Category one of ultrashort pulse-shaping4, 5, 6, 7 techniques, to control two-photon transitions in caesium

G_{29} : Category two of ultrashort pulse-shaping4, 5, 6, 7 techniques, to control two-photon transitions in caesium

G_{30} : Category three of ultrashort pulse-shaping4, 5, 6, 7 techniques, to control two-photon transitions in caesium

T_{28} : Category one of femtosecond laser pulse can be tailored

T_{29} : Category two of femtosecond laser pulse can be tailored

T_{30} : Category three of femtosecond laser pulse can be tailored

Module six

By varying the spectral phases of the pulse components, observe the predicted cancellation of the transitions **due to** destructive quantum interference; the power spectrum and energy of these 'dark pulses' are unchanged

G_{32} : Category one of destructive quantum interference; the power spectrum and energy of these 'dark pulses' are unchanged

G_{33} : Category two of destructive quantum interference; the power spectrum and energy of these 'dark pulses' are unchanged

G_{34} : Category three of destructive quantum interference; the power spectrum and energy of these 'dark

pulses' are unchanged

T_{32} : Category one of variation of the spectral phases of the pulse components and observation of the predicted cancellation of the transitions

T_{33} : Category two of variation of the spectral phases of the pulse components and observation of the predicted cancellation of the transitions

T_{34} : Category three of variation of the spectral phases of the pulse components and observation of the predicted cancellation of the transitions

Module seven

They also show that the pulse shape **can be (e&eb) modified** extensively without affecting the two-photon transition probability

G_{36} : Category one of pulse shape

G_{37} : Category two of pulse shape

G_{38} : Category three of pulse shape

T_{36} : Category one of extensive modification without affecting the two-photon transition probability

T_{37} : Category two of extensive modification without affecting the two-photon transition probability

T_{38} : Category three of extensive modification without affecting the two-photon transition probability

Module eight

Real-time feedback performed during a quantum nondemolition measurement of atomic spin-angular momentum **allowed (eb) to** influence the quantum statistics of the measurement outcome

G_{40} : Category one of Real-time feedback performed during a quantum nondemolition measurement of atomic spin-angular momentum

G_{41} : Category two of Real-time feedback performed during a quantum nondemolition measurement of atomic spin-angular momentum

G_{42} : Category three of Real-time feedback performed during a quantum nondemolition measurement of atomic spin-angular momentum

T_{40} : Category one of influence the quantum statistics of the measurement outcome

T_{41} : Category two of influence the quantum statistics of the measurement outcome

T_{42} : Category three of influence the quantum statistics of the measurement outcome

Module Nine

They showed that it is possible **to harness** measurement backaction as a form of actuation in quantum control, and thus describe a valuable tool for quantum information science .Feedback-mediated procedure **generates** spin-squeezing, for which the reduction in quantum uncertainty and resulting atomic

entanglement are not conditioned on the measurement outcome

G_{44} : Category one of Feedback-mediated procedure

G_{45} : Category two of Feedback-mediated procedure

G_{46} : Category three of Feedback-mediated procedure

T_{44} : Category one of spin-squeezing, for which the reduction in quantum uncertainty and resulting atomic entanglement are not conditioned on the measurement outcome

T_{45} : Category two of spin-squeezing, for which the reduction in quantum uncertainty and resulting atomic entanglement are not conditioned on the measurement outcome

T_{46} : Category three of spin-squeezing, for which the reduction in quantum uncertainty and resulting atomic entanglement are not conditioned on the measurement outcome

SECTION TWENTY FIVE

Quantum Formalism And Holographic Renormalization Group: Non Causa Pro Causa-Non Bis In Idem Model

INTRODUCTION—VARIABLES USED

PHILIPPE GRANGIER, Int. J. Quantum Inform. 03, 17 (2005). DOI: 10.1142/S0219749905000396 CONTEXTUAL OBJECTIVITY AND THE QUANTUM FORMALISM PHILIPPE GRANGIER

(1) The "new orthodoxy" of quantum mechanics (QM) **based on the** decoherence approach **requires** many-worlds as an essential ingredient for logical consistency, and one may wonder what status to give to all these "other worlds."
(2) Here we advocate that it is possible to build a consistent approach to QM where no other worlds are needed, and where the quantum formalism appears **as a (e) consequence of** requiring the enumerability of physical properties.
(3) Such a quantization hypothesis is closely **related to** indistinguishability, and is deeply inconsistent with classical physics.

Quantum (in) stability of dilatonic AdS backgrounds and the holographic renormalization group with gravity Shin'ichi Nojiri

(4) The stability of dilatonic AdS spaces **due (E) to** quantum effects of dilaton coupled conformal matter is considered.
(5) When such spaces do not exist on a classical level, **there shall (eb) be generation** of their

dynamical generation.

(6) Explicit examples corresponding to the quantum creation of a 4D dilatonic AdS universe **and a** 2D dilatonic AdS black hole (BH) are presented.

(7) Motivated by the holographic RG, in a similar approach, the complete 5D effective action is discussed. The intermediate region where it is the sum of two parts, the bulk (classical gravity) and boundary quantum action, is investigated. The effective equation solution representing a 5D AdS universe with a warp scale factor is found. A four-dimensional de Sitter or AdS world **is generated on** the boundary of such a universe as a result of quantum effectsURL:http://link.aps.org/doi/10.1103/PhysRevD.62.064006DOI: 10.1103/PhysRevD.62.064006

Holographic Quantum Liquid A. Karch et al

(8) Quantum liquids **are (e) characterized by** the distinctive properties such as the low-temperature behavior of heat capacity and the spectrum of low-energy quasiparticle excitations. In particular, at low temperature, Fermi liquids **exhibit the** zero sound, predicted by Landau in 1957 and subsequently observed in liquid He-3. In this Letter, we ask whether such characteristic behavior is present in theories with a holographically dual description.

NOTATION

Module One

The "new orthodoxy" of quantum mechanics (QM) **based on the** decoherence approach **requires** many-worlds as an essential ingredient for logical consistency, and one may wonder what status to give to all these "other worlds."

G_{13} : Category one of many-worlds as an essential ingredient for logical consistency, and one may wonder what status to give to all these "other worlds."

G_{14} : Category two of many-worlds as an essential ingredient for logical consistency, and one may wonder what status to give to all these "other worlds."

G_{15} : Category three of many-worlds as an essential ingredient for logical consistency, and one may wonder what status to give to all these "other worlds."

T_{13} : Category one of “new orthodoxy" of quantum mechanics (QM) **based on the** decoherence approach

T_{14} : Category two of "new orthodoxy" of quantum mechanics (QM) **based on the** decoherence approach

T_{15} : Category three of "new orthodoxy" of quantum mechanics (QM) **based on the** decoherence approach

Module Two

Here we advocate that it is possible to build a consistent approach to QM where no other worlds are needed, and where the quantum formalism appears **as a (e) consequence of** requiring the enumerability of physical properties

G_{16} : Category one of requiring the enumerability of physical properties

G_{17} : Category two of requiring the enumerability of physical properties

G_{18} : Category three of requiring the enumerability of physical properties

T_{16} : Category one of consistent approach to QM where no other worlds are needed, and where the quantum

formalism appears

T_{17} : Category two of consistent approach to QM where no other worlds are needed, and where the quantum formalism appears

T_{18} : Category three of consistent approach to QM where no other worlds are needed, and where the quantum formalism appears

Module three

Such a quantization hypothesis is closely **related to** indistinguishability, and is deeply inconsistent with classical physics.

G_{20} : Category one of quantization hypothesis; indistinguishability, and is deeply inconsistent with classical physics.

G_{21} : Category two of quantization hypothesis; indistinguishability, and is deeply inconsistent with classical physics.

G_{22} : Category three of quantization hypothesis; indistinguishability, and is deeply inconsistent with classical physics.

T_{20} : Category one of indistinguishability, and is deeply inconsistent with classical physics; quantization hypothesis

T_{21} : Category two of indistinguishability, and is deeply inconsistent with classical physics; quantization hypothesis

T_{22} : Category three of indistinguishability, and is deeply inconsistent with classical physics; quantization hypothesis

Module four

The stability of dilatonic AdS spaces **due (E) to** quantum effects of dilaton coupled conformal matter is considered

The non-critical string theory describes the relativistic string without enforcing the critical dimension. Although this allows the construction of a string theory in 4 spacetime dimensions, such a theory usually does not describe a Lorentz invariant background. However, there are recent developments which make possible Lorentz invariant quantization of string theory in 4-dimensional Minkowski space-time.

There are several applications of the non-critical string. Through the AdS/CFT correspondence it provides a holographic description of gauge theories which are asymptotically free. It may then have applications to the study of the QCD, the theory of strong interactions between quarks. Another area of much research is two-dimensional string theory which provides simple toy models of string theory. There also exists a duality to the 3-dimensional Ising model

G_{24} : Category one of quantum effects of dilaton coupled conformal matter

G_{25} : Category two of quantum effects of dilaton coupled conformal matter

G_{26} : Category three of quantum effects of dilaton coupled conformal matter

T_{24} : Category one of stability of dilatonic AdS spaces

T_{25} : Category two of stability of dilatonic AdS spaces

T_{26} : Category three of stability of dilatonic AdS spaces

Module five

When such spaces do not exist on a classical level, **there shall (eb) be generation** of their dynamical generation. **Note that we have not taken condition as the one experienced by the conditioned**

G_{28} : Category one of non existence of such spaces

G_{29} : Category two of non existence of such spaces

G_{30} : Category three of non existence of such spaces

T_{28} : Category one of dynamical generation

T_{29} : Category two of dynamical generation

T_{30} : Category three of dynamical generation

Module six

Explicit examples corresponding to the quantum creation of a 4D dilatonic AdS universe **and a** 2D dilatonic AdS black hole (BH) are presented.

G_{32} : Category one of quantum creation of a 4D dilatonic AdS universe

G_{33} : Category two of quantum creation of a 4D dilatonic AdS universe

G_{34} : Category three of quantum creation of a 4D dilatonic AdS universe

T_{32} : Category one of 2D dilatonic AdS black holes (BH).

T_{33} : Category two of2D dilatonic AdS black holes (BH).

T_{34} : Category three of 2D dilatonic AdS black holes (BH).

Module seven

Motivated by the holographic RG, in a similar approach, the complete 5D effective action is discussed. The intermediate region where it is the sum of two parts, the bulk (classical gravity) and boundary quantum action, is investigated. The effective equation solution representing a 5D AdS universe with a warp scale factor is found. A four-dimensional de Sitter or AdS world **is generated on** the boundary of such a universe as a result of quantum effects

G_{36} : Category one of boundary of such a universe as a result of quantum effects

G_{37} : Category two of boundary of such a universe as a result of quantum effects

G_{38} : Category three of boundary of such a universe as a result of quantum effects

T_{36} : Category one of four-dimensional de Sitter or AdS world

T_{37} : Category two of four-dimensional de Sitter or AdS world

T_{38} : Category three of four-dimensional de Sitter or AdS world

Module eight

Anti De Sitter Space And Holography Edward Witten

Recently, it has been proposed by Maldacena that large N limits of certain conformal field theories in d dimensions **can be described** in terms of supergravity (and string theory) on the product of $d+1$-dimensional AdS space with a compact manifold.

Here authors elaborate on this idea and propose a precise correspondence between conformal field theory observables **and** those of supergravity:

Correlation functions in conformal field theory **are given by** the dependence of the supergravity action on the asymptotic behavior at infinity.

In particular, dimensions of operators in conformal field theory **are given by** masses of particles in supergravity.

As quantitative confirmation of this correspondence, they note that the Kaluza-Klein modes of Type IIB supergravity on $AdS_5\times {\bf S} ^5$ **match (=) with the** chiral operators of $\N=4$ super Yang-Mills theory in four dimensions.

With some further assumptions, **one can deduce** a Hamiltonian version of the correspondence and show that the $\N=4$ theory has a large N phase transition related to the thermodynamics of AdS black holes. Adv.Theor.Math.Phys.2:253-291, 1998 arXiv: hep-th/9802150

G_{40} : Category one of large N limits of certain conformal field theories in d dimensions; supergravity; Correlation functions in conformal field theory; dimensions of operators in conformal field theory; chiral operators of $\N=4$ super Yang-Mills theory in four dimensions; Hamiltonian version of the correspondence and show that the $\N=4$ theory has a large N phase transition; thermodynamics of AdS black holes

G_{41} : Category two of large N limits of certain conformal field theories in d dimensions; supergravity; Correlation functions in conformal field theory; dimensions of operators in conformal field theory; chiral operators of $\N=4$ super Yang-Mills theory in four dimensions; Hamiltonian version of the correspondence and show that the $\N=4$ theory has a large N phase transition; thermodynamics of AdS black holes

G_{42} : Category three of large N limits of certain conformal field theories in d dimensions; supergravity; Correlation functions in conformal field theory; dimensions of operators in conformal field theory; chiral operators of $\N=4$ super Yang-Mills theory in four dimensions; Hamiltonian version of the correspondence and show that the $\N=4$ theory has a large N phase transition; thermodynamics of AdS black holes

T_{40} : Category one of supergravity (and string theory) on the product of $d+1$-dimensional AdS space with a compact manifold; conformal field theory observables; masses of particles in supergravity; quantitative confirmation of this correspondence, they note that the Kaluza-Klein modes of Type IIB supergravity on $AdS_5\times {\bf S} ^5$; thermodynamics of AdS black holes ;Hamiltonian version of the correspondence and show that the $\N=4$ theory has a large N phase transition;

T_{41} : Category two of supergravity (and string theory) on the product of $d+1$-dimensional AdS space with a compact manifold; conformal field theory observables; masses of particles in supergravity; quantitative confirmation of this correspondence, they note that the Kaluza-Klein modes of Type IIB supergravity on $AdS_5\times {\bf S} ^5$; thermodynamics of AdS black holes ;Hamiltonian version of the correspondence and show that the $\N=4$ theory has a large N phase transition;

T_{42} : Category three of supergravity (and string theory) on the product of $d+1$-dimensional AdS space with a compact manifold; conformal field theory observables; masses of particles in supergravity; quantitative confirmation of this correspondence, they note that the Kaluza-Klein modes of Type IIB supergravity on $AdS_5\times {\bf S} ^5$; thermodynamics of AdS black holes ;Hamiltonian version of the correspondence and show that the $\N=4$ theory has a large N phase transition

Module Nine

Quantum liquids **are (e) characterized by** the distinctive properties such as the low-temperature behavior of heat capacity and the spectrum of low-energy quasiparticle excitations.,

In particular, at low temperature, Fermi liquids **exhibit the** zero sound, predicted by Landau in 1957 and subsequently observed in liquid He-3. In this Letter, we ask whether such characteristic behavior is present in theories with a holographically dual description

G_{44} : Category one of distinctive properties such as the low-temperature behavior of heat capacity and the spectrum of low-energy quasiparticle excitations; , Fermi liquids; zero sound, predicted by Landau in 1957 and subsequently observed in liquid He-3.

G_{45} : Category two of distinctive properties such as the low-temperature behavior of heat capacity and the spectrum of low-energy quasiparticle excitations; , Fermi liquids; zero sound, predicted by Landau in 1957 and subsequently observed in liquid He-3.

G_{46} : Category three of distinctive properties such as the low-temperature behavior of heat capacity and the spectrum of low-energy quasiparticle excitations; , Fermi liquids; zero sound, predicted by Landau in 1957 and subsequently observed in liquid He-3.

T_{44} : Category one of Quantum liquids; zero sound, predicted by Landau in 1957 and subsequently observed in liquid He-3; theories with a holographically dual description

T_{45} : Category two of Quantum liquids; zero sound, predicted by Landau in 1957 and subsequently observed in liquid He-3; theories with a holographically dual description

T_{46} : Category three of Quantum liquids; zero sound, predicted by Landau in 1957 and subsequently observed in liquid He-3; theories with a holographically dual description

SECTION TWENTY SIX

Holographic Quantum Fields: Naturam Expellas Furca, Tamen Usque Recurret

INTRODUCTION—VARIABLES USED

Predicational anteriorities, ontological consonance, primordial exactitude, substantive limits of Two-point Functions and Quantum Fields in de Sitter Universe was studied by Jacques Bros and Ugo Moschella. They present a theory of general two-point functions and of generalized free fields in d-dimensional de Sitter space-time which **closely parallels** the corresponding Minkowskian theory. The usual spectral condition is **now replaced by a** certain geodesic spectral condition, **equivalent to a** precise thermal characterization of the corresponding "vacuum "states. Their method is **based on the** geometry of the complex de Sitter space-time and on the introduction of a class of holomorphic functions on this manifold, called perikernels, **which reproduce** mutatis mutandis the structural properties of the two-point correlation functions of the Minkowskian quantum field theory. The theory **contains as basic elementary case** the linear massive field models in their "preferred" representation. The latter are described by the introduction of de Sitter plane waves in their tube domains **which lead to** a new integral representation of the two-point functions and to a Fourier Laplace type transformation on the hyperboloid. The Hilbert space structure of these theories is then analysed **by using this** transformation. In particular **authors show the** Reeh-Schlieder property. For general two-point functions, a substitute to the Wick rotation is defined both in complex space-time and in the complex mass variable, and substantial results concerning the derivation of K¨allen-Lehmann type representation are obtained. **Yurtsever U.** shows how the holographic entropy bound can be derived from elementary flat-spacetime quantum field theory when the total energy of Fock states **is constrained** gravitationally. This energy constraint makes the Fock space dimension (whose logarithm is the maximum entropy) finite for both bosons and fermions. Despite the elementary nature of the analysis, **it results in an** upper limit on entropy in remarkable agreement with the holographic bound, and **also provides** a

microscopic deviation of a more general entropy bound recently introduced by Gour. PMID: 12906651 [PubMed] **Phys Rev Lett. 2003 Jul 25; 91(4):041302. Epub 2003 Jul 23.Holographic entropy bound and local quantum field theory. Yurtsever U.**

Holographic Quantum Liquid A. Karch et al

(1) Authors consider a class of gauge theories **with** fundamental matter fields whose holographic dual in the appropriate limit is given in terms of the Dirac-Born-Infeld action in anti–de Sitter space.
(2) They find that these systems **also exhibit a** sound mode at zero temperature despite having a non-Fermi-liquid behavior of the specific heat.
(3) These properties **suggest that** holography identifies a new type of quantum liquid which potentially could be experimentally realized in strongly correlated systems.URL:http://link.aps.org/doi/10.1103/PhysRevLett.102.051602DOI:10.1103/PhysRevLett.102.051602

Holographic Quantum States Tobias J. Osborne

(4) Authors show how continuous matrix product states of quantum fields **can be described** in terms of the dissipative nonequilibrium dynamics of a lower-dimensional auxiliary boundary field by demonstrating that the spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field.
(5) Spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field equivalence illustrates an intimate connection between the theory of continuous quantum measurement **and** quantum field theory
(6) Spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field. **gives an explicit (eb) construction** of the boundary field allowing the extension of real-space renormalization group methods to arbitrary dimensional quantum field theories without the introduction of a lattice parameter
(7) Spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field y**ields(eb) a** novel interpretation of recent cavity QED experiments in terms of quantum field theory, and hence paves the way toward observing genuine quantum phase transitions in such zero-dimensional driven quantum systems.URL:http://link.aps.org/doi/10.1103/PhysRevLett.105.260401DOI:10.1103/PhysRevLett.105.260401

Journal of High Energy Physics January 2010, 2010:114 Holographic quantum liquids in 1+1 dimension sling-Yan Hung, et al

(8) In this paper we initiate the study of holographic quantum liquids **in (eb)** 1+1 dimensions.
(9) Since the Landau Fermi liquid theory **breaks (e) down** in 1+1 dimensions, it is of interest to see what holographic methods have to say about similar models

NOTATION

Module One

Authors consider a class of gauge theories **with** fundamental matter fields whose holographic dual in the appropriate limit is given in terms of the Dirac-Born-Infeld action in anti–de Sitter space

G_{13} : Category one of class of gauge theories

G_{14} : Category two of class of gauge theories

G_{15} : Category three of class of gauge theories

T_{13} : Category one of fundamental matter fields whose holographic dual in the appropriate limit is given in terms of the Dirac-Born-Infeld action in anti–de Sitter space

T_{14} : Category two of fundamental matter fields whose holographic dual in the appropriate limit is given in terms of the Dirac-Born-Infeld action in anti–de Sitter space

T_{15} : Category three of fundamental matter fields whose holographic dual in the appropriate limit is given in terms of the Dirac-Born-Infeld action in anti–de Sitter space

Module Two

They find that these systems **also exhibit a** sound mode at zero temperature despite having a non-Fermi-liquid behavior of the specific heat.

G_{16} : Category one of gauge theories **with** fundamental matter fields whose holographic dual in the appropriate limit is given in terms of the Dirac-Born-Infeld action in anti–de Sitter space

G_{17} : Category two of gauge theories **with** fundamental matter fields whose holographic dual in the appropriate limit is given in terms of the Dirac-Born-Infeld action in anti–de Sitter space

G_{18} : Category three of gauge theories **with** fundamental matter fields whose holographic dual in the appropriate limit is given in terms of the Dirac-Born-Infeld action in anti–de Sitter space

T_{16} : Category one of sound mode at zero temperature despite having a non-Fermi-liquid behavior of the specific heat.

T_{17} : Category two of sound mode at zero temperature despite having a non-Fermi-liquid behavior of the specific heat.

T_{18} : Category three of sound mode at zero temperature despite having a non-Fermi-liquid behavior of the specific heat.

Module three

These properties **suggest that** holography identifies a new type of quantum liquid which potentially could be experimentally realized in strongly correlated systems.

G_{20} : Category one of systems **also exhibit a** sound mode at zero temperature despite having a non-Fermi-liquid behavior of the specific heat

G_{21} : Category two of systems **also exhibit a** sound mode at zero temperature despite having a non-Fermi-liquid behavior of the specific heat

G_{22} : Category three of systems **also exhibit a** sound mode at zero temperature despite having a non-Fermi-liquid behavior of the specific heat

T_{20} : Category one of holography identifies a new type of quantum liquid which potentially could be experimentally realized in strongly correlated systems.

T_{21} : Category two of holography identifies a new type of quantum liquid which potentially could be experimentally realized in strongly correlated systems.

T_{22} : Category three of holography identifies a new type of quantum liquid which potentially could be

experimentally realized in strongly correlated systems.

Module four

Authors show how continuous matrix product states of quantum fields **can be described** in terms of the dissipative nonequilibrium dynamics of a lower-dimensional auxiliary boundary field by demonstrating that the spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field.

G_{24} : Category one of continuous matrix product states of quantum fields **can be described** in terms of the dissipative nonequilibrium dynamics of a lower-dimensional auxiliary boundary field

G_{25} : Category two of continuous matrix product states of quantum fields **can be described** in terms of the dissipative nonequilibrium dynamics of a lower-dimensional auxiliary boundary field

G_{26} : Category three of continuous matrix product states of quantum fields **can be described** in terms of the dissipative nonequilibrium dynamics of a lower-dimensional auxiliary boundary field

T_{24} : Category one of spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field

T_{25} : Category two of spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field

T_{26} : Category three of spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field

Module five

Spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field equivalence illustrates an intimate connection between the theory of continuous quantum measurement **and** quantum field theory

G_{28} : Category one of Spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field equivalence

G_{29} : Category two of Spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field equivalence

G_{30} : Category three of Spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field equivalence

T_{28} : Category one of intimate connection between the theory of continuous quantum measurement **and** quantum field theory

T_{29} : Category two of intimate connection between the theory of continuous quantum measurement **and** quantum field theory

T_{30} : Category three of intimate connection between the theory of continuous quantum measurement **and** quantum field theory

Module six

Spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field. **gives an explicit (eb) construction** of the boundary field allowing the extension of real-space renormalization group methods to arbitrary dimensional quantum field theories without the introduction of a lattice parameter

G_{32} : Category one of Spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field

G_{33} : Category two of Spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field

G_{34} : Category three of Spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field

T_{32} : Category one of boundary field allowing the extension of real-space renormalization group methods to arbitrary dimensional quantum field theories without the introduction of a lattice parameter

T_{33} : Category two of boundary field allowing the extension of real-space renormalization group methods to arbitrary dimensional quantum field theories without the introduction of a lattice parameter

T_{34} : Category three of boundary field allowing the extension of real-space renormalization group methods to arbitrary dimensional quantum field theories without the introduction of a lattice parameter

Module seven

Spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field y**ields(eb) a** novel interpretation of recent cavity QED experiments in terms of quantum field theory, and hence paves the way toward observing genuine quantum phase transitions in such zero-dimensional driven quantum systems

G_{36} : Category one of Spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field

G_{37} : Category two of Spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field

G_{38} : Category three of Spatial correlation functions of the bulk field **correspond to** the temporal statistics of the boundary field

T_{36} : Category one of novel interpretation of recent cavity QED experiments in terms of quantum field theory, and hence paves the way toward observing genuine quantum phase transitions in such zero-dimensional driven quantum systems

T_{37} : Category two of novel interpretation of recent cavity QED experiments in terms of quantum field theory, and hence paves the way toward observing genuine quantum phase transitions in such zero-dimensional driven quantum systems

T_{38} : Category three of novel interpretation of recent cavity QED experiments in terms of quantum field theory, and hence paves the way toward observing genuine quantum phase transitions in such zero-dimensional driven quantum systems

Module eight

In this paper we initiate the study of holographic quantum liquids **in (eb)** 1+1 dimensions

G_{40} : Category one of holographic quantum liquids

G_{41} : Category two of holographic quantum liquids

G_{42} : Category three of holographic quantum liquids

T_{40} : Category one of1+1 dimensions

T_{41} : Category two of 1+1 dimensions

T_{42} : Category three of 1+1 dimensions

Module Nine

Since the Landau Fermi liquid theory **breaks (e) down** in 1+1 dimensions, it is of interest to see what holographic methods have to say about similar models

G_{44} : Category one of1+1 dimensions, it is of interest to see what holographic methods have to say about similar models

G_{45} : Category two of 1+1 dimensions, it is of interest to see what holographic methods have to say about similar models

G_{46} : Category three of 1+1 dimensions, it is of interest to see what holographic methods have to say about similar models

T_{44} : Category one of Landau Fermi liquid theory

T_{45} : Category two of Landau Fermi liquid theory

T_{46} : Category three of Landau Fermi liquid theory

SECTION TWENTY SEVEN

Artificial Intelligence And Uncertainty Modeling: Non In Legendo Sed In Intelligendo Legis Consistent

INTRODUCTION—VARIABLES USED

Journal of High Energy Physics January 2010, 2010:114 Holographic quantum liquids in 1+1 dimension sling-Yan Hung, et al."

(1) For theories with a gapless branch, the Luttinger conjecture **states that there** is an effective description of the physics in terms of a Luttinger liquid which is specified by two parameters.
(2) The theory we consider is the defect CFT arising **due (e) to** a probe D3 brane in the AdS Schwarzschild planar black hole background.
(3) They turn on a fundamental string density on the worldvolume. Unlike higher dimensional defects, a persistent dissipationless zero sound mode is found. The thermodynamic aspects of these models are considered carefully and certain subtleties with boundary terms are explained which are unique to 1+1 dimensions. **Spectral functions of bosonic and fermionic fluctuations** are also considered **and quasinormal modes** are analysed. A prescription is given to compute spectral functions when there are mixing due to the worldvolume gauge field. Authors comment on the **Luttinger conjecture** in the light of our findings.

Annals of Mathematics and Artificial Intelligence August 2001, Volume 32, Issue 1-4, pp 35-66 Possibility Theory, Probability Theory and Multiple-Valued Logics: A Clarification Didier Dubois, Henri Prade

(4) There has been a long-lasting misunderstanding in the literature of artificial intelligence and uncertainty modeling, regarding **the (e&eb) role of** fuzzy set theory and many-valued logics.
(5) The recurring question is that of the mathematical and pragmatic meaningfulness of a compositional calculus **and (e&eb) the** validity of the excluded middle law.
(6) This confusion pervades the early developments of probabilistic logic, despite early warnings of some philosophers of probability. This paper tries to clarify this situation. It emphasizes three main points. First, **it suggests that** the root of the controversies **lies (e) in the** unfortunate confusion

between degrees of belief and what logicians call "degrees of truth". The latter are usually compositional, while the former cannot be so. **This claim is first illustrated** by laying bare the non-compositional belief representation embedded in the standard propositional calculus. It turns out to be an all-or-nothing version of possibility theory. This framework **is then extended** to discuss the case of fuzzy logic versus graded possibility theory.

(7) Next, it is demonstrated that any belief representation where compositionality is taken for granted is bound **to at worst (e) collapse** to a Boolean truth assignment and at best to a poorly expressive tool.

(8) Lastly, some claims pertaining to an alleged compositionality of possibility theory are refuted, thus **clarifying (eb) a** pervasive confusion between possibility theory axioms and fuzzy set basic connectives.

NOTATION

Module One

For theories with a gapless branch, the Luttinger conjecture **states that there** is an effective description of the physics in terms of a Luttinger liquid which is specified by two parameters.

G_{13} : Category one of theories with a gapless branch, the Luttinger conjecture

G_{14} : Category two of theories with a gapless branch, the Luttinger conjecture

G_{15} : Category three of theories with a gapless branch, the Luttinger conjecture

T_{13} : Category one of effective description of the physics in terms of a Luttinger liquid which is specified by two parameters.

T_{14} : Category two of effective description of the physics in terms of a Luttinger liquid which is specified by two parameters.

T_{15} : Category three of effective description of the physics in terms of a Luttinger liquid which is specified by two parameters.

Module Two

The theory we consider is the defect CFT arising **due (e) to** a probe D3 brane in the AdS Schwarzschild planar black hole background.

G_{16} : Category one of probe D3 brane in the AdS Schwarzschild planar black hole background.

G_{17} : Category two of probe D3 brane in the AdS Schwarzschild planar black hole background.

G_{18} : Category three of probe D3 brane in the AdS Schwarzschild planar black hole background.

T_{16} : Category one of theory considered is the defect CFT

T_{17} : Category two of theory considered is the defect CFT

T_{18} : Category three of theory considered is the defect CFT

Module three

Oliver DeWolfe et al studied primary focus and locus of homologous receptiveness of Karch Randall conjecture in **Holography and Defect Conformal Field Theories. They** develop both the gravity and field theory sides of the Karch-Randall conjecture that the near-horizon description of a certain D5-D3 brane

configuration in string theory, **realized (eb) as** AdS_5 x S^5 bisected by an AdS_4 x S^2 "brane", **is dual (=) to** N=4 Super Yang-Mills theory in R^4 coupled to an R^3 defect.

Authors propose a complete Lagrangian for the field theory dual, a novel "defect superconformal field theory" wherein a subset of the fields of N=4 SYM **interacts with** a d=3 SU (N) fundamental hypermultiplet on the defect preserving conformal invariance and 8 supercharges.

The Kaluza-Klein reduction of wrapped D5 modes on AdS_4 x S^2 **leads to** towers of short representations of OSp (4|4), and construct the map to a set of dual gauge-invariant defect operators O_3 possessing integer conformal dimensions.

Gravity calculations of <O_4> and <O_4O_3> are given. Spacetime and N-dependence **matches expectations** from dCFT, while the behavior as functions of lambda = g^2 N at strong and weak coupling is generically different.

Author's comments on a class of correlators for which a non-renormalization theorem **may (eb) still exist**. Partial evidence for the Conformality of the quantum theory is given, including a complete argument for the special case of a U (1) gauge group. Some weak coupling arguments which illuminate the duality are presented.Phys.Rev.D66:025009, 2002DOI: 10.1103/PhysRevD.66.025009ALT-68-2359, CITUSC/01-041, NSF-ITP-01-172, MIT-CTP-3212 arXiv: hep-th/0111135

G_{20} : Category one of gravity and field theory sides of the Karch-Randall conjecture that the near-horizon description of a certain D5-D3 brane configuration in string theory; complete Lagrangian for the field theory dual, a novel "defect superconformal field theory" wherein a subset of the fields of N=4 SYM; Kaluza-Klein reduction of wrapped D5 modes on AdS_4 x S^2; dCFT, while the behavior as functions of lambda = g^2 N at strong and weak coupling is generically different; non-renormalization theorem

G_{21} : Category two of gravity and field theory sides of the Karch-Randall conjecture that the near-horizon description of a certain D5-D3 brane configuration in string theory; complete Lagrangian for the field theory dual, a novel "defect superconformal field theory" wherein a subset of the fields of N=4 SYM; Kaluza-Klein reduction of wrapped D5 modes on AdS_4 x S^2; dCFT, while the behavior as functions of lambda = g^2 N at strong and weak coupling is generically different; non-renormalization theorem

G_{22} : Category three of gravity and field theory sides of the Karch-Randall conjecture that the near-horizon description of a certain D5-D3 brane configuration in string theory; complete Lagrangian for the field theory dual, a novel "defect superconformal field theory" wherein a subset of the fields of N=4 SYM; Kaluza-Klein reduction of wrapped D5 modes on AdS_4 x S^2; dCFT, while the behavior as functions of lambda = g^2 N at strong and weak coupling is generically different; non-renormalization theorem

T_{20} : Category one of AdS_5 x S^5 bisected by an AdS_4 x S^2 "brane", **is dual (=) to** N=4 Super Yang-Mills theory in R^4 coupled to an R^3 defect; d=3 SU (N) fundamental hypermultiplet on the defect preserving conformal invariance and 8 supercharges; complete Lagrangian for the field theory dual, a novel "defect superconformal field theory" wherein a subset of the fields of N=4 SYM; towers of short representations of OSp (4|4), and construct the map to a set of dual gauge-invariant defect operators O_3 possessing integer conformal dimensions; Gravity calculations of <O_4> and <O_4O_3> are given. Spacetime and N-dependence; class of correlators

T_{21} : Category two of AdS_5 x S^5 bisected by an AdS_4 x S^2 "brane", **is dual (=) to** N=4 Super Yang-Mills theory in R^4 coupled to an R^3 defect; d=3 SU (N) fundamental hypermultiplet on the defect preserving conformal invariance and 8 supercharges; complete Lagrangian for the field theory dual, a novel "defect superconformal field theory" wherein a subset of the fields of N=4 SYM; towers of short

representations of OSp (4|4), and construct the map to a set of dual gauge-invariant defect operators O_3 possessing integer conformal dimensions; Gravity calculations of <O_4> and <O_4O_3> are given. Spacetime and N-dependence; class of correlators

T_{22} : Category three of AdS_5 x S^5 bisected by an AdS_4 x S^2 "brane", **is dual (=) to** N=4 Super Yang-Mills theory in R^4 coupled to an R^3 defect; d=3 SU (N) fundamental hypermultiplet on the defect preserving conformal invariance and 8 supercharges; complete Lagrangian for the field theory dual, a novel "defect superconformal field theory" wherein a subset of the fields of N=4 SYM; towers of short representations of OSp (4|4), and construct the map to a set of dual gauge-invariant defect operators O_3 possessing integer conformal dimensions; Gravity calculations of <O_4> and <O_4O_3> are given. Spacetime and N-dependence; class of correlators

Module four

N=4 Super Yang-Mills theory is a mathematical and physical model created to study particles through a simple system, similar to string theory, with conformal symmetry. It was knowingly defined to be false, i.e. to describe properties of the physical world that don't exist in reality. It describes a universe **containing (e)** boson fields and fermion fields which are related by 4 supersymmetries (this means that swapping boson, fermion and scalar fields in a certain way leaves the predictions of the theory invariant). It is one of the simplest (because it has no free parameters) and one of the few finite quantum field theories in 4 dimensions. It can be thought of as the most symmetric field theory that does not involve gravity. This theory is important also in the context of the holographic principle. There is a duality between Type IIB string theory **on** AdS5 × S5 space (a product of 5-dimensional AdS space with a 5-dimensional sphere) and N = 4 Super Yang-Mills on the 4-dimensional boundary of AdS5. It is the most successful realization of the holographic principle, **a speculative idea about** quantum gravity originally proposed by Gerard't Hooft and improved and promoted by Leonard Susskind.

Lagrangian

The Lagrangian for the theory is

$$L = tr\left\{-\frac{1}{2g^2}F_{\mu\nu}F^{\mu\nu} + \frac{\theta_I}{8\pi^2}F_{\mu\nu}\tilde{F}^{\mu\nu} - i\lambda^a\sigma^\mu D_\mu\lambda_a - D_\mu X^i D^\mu X^i + gC_i^{ab}\lambda_a[X^i,\lambda_b] + C_{iab}\lambda^a[X^i,\lambda^b] + \frac{g^2}{2}[X^i,X^j]^2\right\}$$

…… where $F^k_{\mu\nu} = \partial_\mu A^k_\nu - \partial_\nu A^k_\mu + gf^{klm}A^l_\mu A^m_\nu$

and the indices i,j=1..6 and the indices a,b=1..4.

Ten-dimensional Lagrangian

The above Lagrangian can be found by beginning with the simpler ten-dimensional Lagrangian

$$L = tr\left\{\frac{1}{g^2}F_{IJ}F^{IJ} - i\bar{\lambda}\Gamma^I D_I\lambda\right\}$$

where I and J are now run from 0 through 9 and Γ^I are the 32 by 32 gamma matrices. $(32 = 2^{10/2})$ followed by adding the term with θ_I which is a topological term.

The components A_i of the gauge field for i=5 to 9 become scalars upon eliminating the extra dimensions. This also gives an interpretation of the SO (6) R-symmetry as rotations in the extra compact dimensions.

By compactification on a T^6, all the supercharges are preserved, giving N=4 in the 4 dimensional theory. The conformal symmetry manifest in this presentation also survives the compactification giving

a superconformal field theory.

A Type IIB string theory interpretation of the theory is the worldvolume theory of a stack of D3-branes.

The coupling constants θ_I and g naturally pair together in the form: $\tau = \frac{\theta}{2\pi} + \frac{4\pi i}{g^2}.$

Context: Ed. Witten recently wrote a potentially revolutionary paper where he showed that under certain conditions, a Chern-Simons path integral in three dimensions **is equivalent to** an N = 4 path integral in four dimensions (this is the standard d=4, N=4 super Yang Mills theory)

Speculation: Witten had shown that the Chern-Simons topological quantum field theory can be solved in terms of Jones polynomials. A quantum computer can simulate a TQFT, and thereby approximate the Jones polynomial. (Source: Wikipedia and this paper) Now I haven't completed reading Witten's paper and I wouldn't understand much of it anyways. But the idea is that if a quantum computer can simulate a path integral (or a Chern-Simons TQF) and since now Witten has shown in both of them to be dual descriptions in some sense, a quantum computer, atleast theoretically might be able to simulate a QFT. Also by the extension of that, Maldacena proposed that the specific field theory that Witten is using to be dual to type-II B string theory in AdS/CFT so then it may also be possible (only theoretically) to simulate a string theory**.(Quantum computation Blog in Wikipedia)**

G_{24} : Category one of LHS of the equation of the Lagrangian for the theory N=4 Super Yang Mills Theory; an N = 4 path integral in four dimensions (this is the standard d=4, N=4 super Yang Mills theory) ;LHS of the equation $\tau = \frac{\theta}{2\pi} + \frac{4\pi i}{g^2}.$

G_{25} : Category two of LHS of the equation of the Lagrangian for the theory N=4 Super Yang Mills Theory; an N = 4 path integral in four dimensions (this is the standard d=4, N=4 super Yang Mills theory);LHS of the above equation

G_{26} : Category three of LHS of the equation of the Lagrangian for the theory N=4 Super Yang Mills Theory; an N = 4 path integral in four dimensions (this is the standard d=4, N=4 super Yang Mills theory);LHS of the above equation

---

T_{24} : Category one of RHS of the equation of the Lagrangian for the theory N=4 Super Yang Mills Theory; Chern-Simons path integral in three dimensions; RHS of the equation $\tau = \frac{\theta}{2\pi} + \frac{4\pi i}{g^2}.$

T_{25} : Category two of RHS of the equation of the Lagrangian for the theory N=4 Super Yang Mills Theory; Chern-Simons path integral in three dimensions; RHS of the equation above

T_{26} : Category three of RHS of the equation of the Lagrangian for the theory N=4 Super Yang Mills Theory; Chern-Simons path integral in three dimensions ; RHS of the equation above

Module five

There has been a long-lasting misunderstanding in the literature of artificial intelligence and uncertainty modeling, regarding **the (e&eb) role of** fuzzy set theory and many-valued logics

G_{28} : Category one of artificial intelligence and uncertainty modeling; fuzzy set theory and many-valued logics

G_{29} : Category two of artificial intelligence and uncertainty modeling; fuzzy set theory and many-valued logics

G_{30} : Category three of artificial intelligence and uncertainty modeling; fuzzy set theory and many-valued logics

T_{28} : Category one of fuzzy set theory and many-valued logics; artificial intelligence and uncertainty modeling,

T_{29} : Category two of fuzzy set theory and many-valued logics; artificial intelligence and uncertainty modeling,

T_{30} : Category three of fuzzy set theory and many-valued logics ;artificial intelligence and uncertainty modeling,

Module six

This confusion pervades the early developments of probabilistic logic, despite early warnings of some philosophers of probability. This paper tries to clarify this situation. It emphasizes three main points. First, **it suggests that** the root of the controversies **lies (e) in the** unfortunate confusion between degrees of belief and what logicians call "degrees of truth". The latter are usually compositional, while the former cannot be so. **This claim is first illustrated** by laying bare the non-compositional belief representation embedded in the standard propositional calculus. It turns out to be an all-or-nothing version of possibility theory. This framework **is then extended** to discuss the case of fuzzy logic versus graded possibility theory.

G_{32} : Category one of non-compositional belief representation embedded in the standard propositional calculus. It turns out to be an all-or-nothing version of possibility theory. This framework

G_{33} : Category two of non-compositional belief representation embedded in the standard propositional calculus. It turns out to be an all-or-nothing version of possibility theory. This framework

G_{34} : Category three of non-compositional belief representation embedded in the standard propositional calculus. It turns out to be an all-or-nothing version of possibility theory. This framework

T_{32} : Category one of fuzzy logic versus graded possibility theory

T_{33} : Category two of fuzzy logic versus graded possibility theory

T_{34} : Category three of fuzzy logic versus graded possibility theory

Module seven

Next, it is demonstrated that any belief representation where compositionality is taken for granted is bound **to at worst (e) collapse** to a Boolean truth assignment and at best to a poorly expressive tool

G_{36} : Category one of Boolean truth assignment and at best to a poorly expressive tool

G_{37} : Category two of Boolean truth assignment and at best to a poorly expressive tool

G_{38} : Category three of Boolean truth assignment and at best to a poorly expressive tool

T_{36} : Category one of belief representation where compositionality is taken for granted is bound to collapse

T_{37} : Category two of belief representation where compositionality is taken for granted is bound to collapse

T_{38} : Category three of belief representation where compositionality is taken for granted is bound to collapse

Module eight

The Dilatation Operator of N = 4 Super Yang-Mills Theory and Integrability Dissertation Eingereicht an der Humboldt-Universit¨at zu Berlin Uberarbeitete Fassung Niklas Beisert

(1) The dilatation generator **measures(amplitude, analysis, appraisal**, area, assessment, **calibration, capacity, computation**, degree, density, depth, **determination, dimension, distance, estimation**, evaluation, extent, frequency, height, judgment, length, magnitude, mass, measure,**mensuration, metage, pitch**, **quantification, quantity, range, reach, scope, size, survey**) **(EB)** the scaling dimensions of local operators in a conformal field theory.

(2) In this thesis authors consider the example of maximally supersymmetric gauge theory in four dimensions and develop and extend techniques **to derive, investigate and apply** the dilatation operator

(3) These tools are perfectly **suited (e) for** precision tests of the dynamical AdS/CFT conjecture.

(4) In particular, they have been successfully **applied in the context of** strings on plane waves and spinning strings.

(5) They construct the dilatation operator by purely algebraic means: **Relying on** the symmetry algebra and structural properties of Feynman diagrams we are able to bypass involved, higher-loop field theory computations.

(6) In this way obtain the complete one-loop dilatation operator and the planar, three-loop deformation in an interesting subsector. These results **allow us to address** the issue of integrability within a planar four-dimensional gauge theory

(7) Authors prove that the complete dilatation generator **is (=)** integrable at one-loop and present the corresponding Bethe ansatz.

(8) Authors furthermore argue that integrability extends to three-loops and beyond and assuming that it holds indeed, finally construct a novel spin chain model at five-loops and propose a Bethe ansatz which **might be valid** at arbitrary loop-order! They illustrate **the use of technology** in several examples and also present two key applications for the AdS/CFT correspondence:

(9) They derive the plane-waves string field theory Hamiltonian **from the** dilatation operator and compute the energy shift on the torus. Furthermore, **they use** the Bethe ansatz to find scaling dimensions of operators with large quantum numbers. A comparison to the energy of spinning strings shows an intricate functional agreement

Note: Bethe ansatz for the one-dimensional s=1/2 Heisenberg ferromagnet is introduced at an elementary level. A detailed description and a complete classification of all two-magnon scattering states and two-magnon bound states can be given for finite and infinite chains.

G_{40} : Category one of dilatation generator; maximally supersymmetric gauge theory in four dimensions and develop and extend techniques; precision tests of the dynamical AdS/CFT conjecture; symmetry algebra and structural properties of Feynman diagrams we are able to bypass involved, higher-loop field theory computations; complete one-loop dilatation operator and the planar, three-loop deformation in an interesting subsector; issue of integrability within a planar four-dimensional gauge theory; arbitrary loop-order;

G_{41} : Category two of dilatation generator; maximally supersymmetric gauge theory in four dimensions and

develop and extend techniques; precision tests of the dynamical AdS/CFT conjecture; symmetry algebra and structural properties of Feynman diagrams we are able to bypass involved, higher-loop field theory computations; complete one-loop dilatation operator and the planar, three-loop deformation in an interesting subsector; issue of integrability within a planar four-dimensional gauge theory; arbitrary loop-order;

G_{42} : Category three of dilatation generator; maximally supersymmetric gauge theory in four dimensions and develop and extend techniques; precision tests of the dynamical AdS/CFT conjecture; symmetry algebra and structural properties of Feynman diagrams we are able to bypass involved, higher-loop field theory computations; complete one-loop dilatation operator and the planar, three-loop deformation in an interesting subsector; issue of integrability within a planar four-dimensional gauge theory; arbitrary loop-order;

T_{40} : Category one of scaling dimensions of local operators in a conformal field theory; dilatation operator; dilatation generator; construction of the dilatation operator; integrability extends to three-loops and beyond and assuming that it holds indeed, finally construct a novel spin chain model at five-loops and propose a Bethe ansatz;

T_{41} : Category two of scaling dimensions of local operators in a conformal field theory; dilatation operator; dilatation generator; construction of the dilatation operator; integrability extends to three-loops and beyond and assuming that it holds indeed, finally construct a novel spin chain model at five-loops and propose a Bethe ansatz;

T_{42} : Category three of scaling dimensions of local operators in a conformal field theory; dilatation operator; dilatation generator; construction of the dilatation operator; integrability extends to three-loops and beyond and assuming that it holds indeed, finally construct a novel spin chain model at five-loops and propose a Bethe ansatz;

Module Nine

G_{44} : Category one of integrable at one-loop with concomitant bethe -ansatz

G_{45} : Category two of integrable at one-loop with concomitant bethe -ansatz

G_{46} : Category three of integrable at one-loop with concomitant bethe -ansatz

T_{44} : Category one of complete dilatation generator

T_{45} : Category two of complete dilatation generator

T_{46} : Category three of complete dilatation generator

SECTION TWENTY SEVEN

Quantum Weirdness: Mutantur Omnia Nos Et Mutamur In Illis

INTRODUCTION—VARIABLES USED

Web Exclusive: Quantum Weirdness? It's All in Your Mind: Scientific American

A new version of quantum theory sweeps away the bizarre paradoxes of the microscopic world. The cost? Quantum information exists only in your imagination. Caleb Charland. Image is deleted due to spatial constraints.

Flawlessly accounting for the behavior of matter on scales from the subatomic to the astronomical, quantum mechanics is the most successful theory in all the physical sciences. It is also the weirdest.

(1) In the quantum realm, particles seem **to (e) be in two** places at once
(2) Information appears to **travel faster** than the speed of light. Velocity of information**(e)** velocity of light
(3) cats can be dead at a time t(1) and alive at the same time t(1)

Physicists have grappled with the quantum world's apparent paradoxes for nine decades, with little to show for their struggles. Unlike evolution and cosmology, whose truths have been incorporated into the general intellectual landscape, quantum theory is still considered (even by many physicists) to be a bizarre anomaly, a powerful recipe book for building gadgets but good for little else. The deep confusion about the meaning of quantum theory will continue to add fuel to the perception that the deep things it is so urgently trying to tell us about our world are irrelevant to everyday life and too weird to matter.

In 2001 a team of researchers began to develop a model that either eliminates the quantum paradoxes or puts them in a less troubling form. The model, known as Quantum Bayesianism or QBism for short, reimagines the entity that lies at the heart of quantum weirdness—the wave function.

(4) In the conventional view of quantum theory, an object such as an electron is represented by its wave function, a mathematical expression that describes the object's properties. If you want to predict how the electron will behave, you calculate how its wave function **evolves (eb) in** time.
(5) The result of the calculation **gives you the** probability that the electron will have a certain property (like being in one place and not another). But problems arise when physicists assume that a wave function is real.
(6) QBism, combines quantum theory **with** probability theory,
(7) QBism maintains that the wave function **has no_** objective reality.
(8) Instead QBism portrays the wave function as a user's manual, a mathematical tool that an observer **uses(e) to** make wiser decisions about the surrounding world—the quantum world.
(9) Specifically, the observer employs the wave function **to(eb) assign** his or her personal belief that a quantum system will have a specific property, realizing that the individual's own choices and actions affect the system in an inherently uncertain way. Another observer, using a wave function that describes the world as the person sees it, may **come (eb) to a completely** different conclusion about the same quantum system.

NOTATION

Module One

In the quantum realm, particles seem **to (e) be in two** places at once

G_{13} : Category one of seem **to (e) be in two** places at once

G_{14} : Category two of seem **to (e) be in two** places at once

G_{15} : Category three of seem **to (e) be in two** places at once

T_{13} : Category one of quantum realm, particles

T_{14} : Category two of quantum realm, particles

T_{15} : Category three of quantum realm, particles

Module Two

Velocity of information**(e)** velocity of light

G_{16} : Category one of velocity of light

G_{17} : Category two of velocity of light

G_{18} : Category three of velocity of light

T_{16} : Category one of Velocity of information

T_{17} : Category two of Velocity of information

T_{18} : Category three of Velocity of information

Module three

cats can be dead at a time t(1) and alive at the same time t(1)

G_{20} : Category one of cats can be dead at a time t(1)

G_{21} : Category two of cats can be dead at a time t(1)

G_{22} : Category three of cats can be dead at a time t(1)

T_{20} : Category one of cats can be alive at the same time t(1)

T_{21} : Category two of cats can be alive at the same time t(1)

T_{22} : Category three of cats can be alive at the same time t(1)

Module four

In the conventional view of quantum theory, an object such as an electron is represented by its wave function, a mathematical expression that describes the object's properties. If you want to predict how the electron will behave, you calculate how its wave function **evolves (eb) in** time.

G_{24} : Category one of evolution of wave function in time

G_{25} : Category two of evolution of wave function in time

G_{26} : Category three of evolution of wave function in time

T_{24} : Category one of behaviour of electrons

T_{25} : Category two of behaviour of electrons

T_{26} : Category three of behaviour of electrons

Module five

The result of the calculation **gives you the** probability that the electron will have a certain property (like being in one place and not another). But problems arise when physicists assume that a wave function is real

G_{28} : Category one of result of the calculation

G_{29} : Category two of result of the calculation

G_{30} : Category three of result of the calculation

T_{28} : Category one of probability that the electron will have a certain property (like being in one place and not another).

T_{29} : Category two of probability that the electron will have a certain property (like being in one place and not another).

T_{30} : Category three of probability that the electron will have a certain property (like being in one place and not another).

Module six

QBism, combines quantum theory **with** probability theory

G_{32} : Category one of quantum theory. We donot mean there are three quantum theories. Classification is based on the characterstics of millions of quantum systems under investigation.

G_{33} : Category two of quantum theory

G_{34} : Category three of quantum theory

T_{32} : Category one of probability theory

T_{33} : Category two of probability theory

T_{34} : Category three of probability theory

Module seven

QBism maintains that the wave function **has no_** objective reality.

G_{36} : Category one of objective reality

G_{37} : Category two of objective reality

G_{38} : Category three of objective reality

T_{36} : Category one of wave function

T_{37} : Category two of wave function

T_{38} : Category three of wave function

Module eight

Instead QBism portrays the wave function as a user's manual, a mathematical tool that an observer **uses(e) to** make wiser decisions about the surrounding world—the quantum world.

G_{40} : Category one of wiser decisions about the surrounding world—the quantum world

G_{41} : Category two of wiser decisions about the surrounding world—the quantum world

G_{42} : Category three of wiser decisions about the surrounding world—the quantum world

T_{40} : Category one of wave function as a user's manual, a mathematical tool that an observer **uses**

T_{41} : Category two of wave function as a user's manual, a mathematical tool that an observer **uses**

T_{42} : Category three of wave function as a user's manual, a mathematical tool that an observer **uses**

Module Nine

Specifically, the observer employs the wave function **to(eb) assign** his or her personal belief that a quantum system will have a specific property, realizing that the individual's own choices and actions affect the system in an inherently uncertain way. Another observer, using a wave function that describes the world as the person sees it, may **come (eb) to a completely** different conclusion about the same quantum system.

G_{44} : Category one of observer employs the wave function

G_{45} : Category two of observer employs the wave function

G_{46} : Category three of observer employs the wave function

T_{44} : Category one of **assign** his or her personal belief that a quantum system will have a specific property, realizing that the individual's own choices and actions affect the system in an inherently uncertain way. Another observer, using a wave function that describes the world as the person sees it, may **come (eb) to a completely** different conclusion about the same quantum system.

T_{45} : Category two of **assign** his or her personal belief that a quantum system will have a specific property, realizing that the individual's own choices and actions affect the system in an inherently uncertain way. Another observer, using a wave function that describes the world as the person sees it, may **come (eb) to a completely** different conclusion about the same quantum system.

T_{46} : Category three of **assign** his or her personal belief that a quantum system will have a specific property, realizing that the individual's own choices and actions affect the system in an inherently uncertain way. Another observer, using a wave function that describes the world as the person sees it, may **come (eb) to a**

completely different conclusion about the same quantum system.

SECTION TWENTY EIGHT

QBism: Veritas Vos Liberabit - Vincit Qui Se Vincit

INTRODUCTION—VARIABLES USED

Web Exclusive: Quantum Weirdness? It's All in Your Mind: Scientific American

(1) One system—one event—**can (e) have** as many different wave functions as there are observers.
(2) After observers have communicated with one another and **modified (e&eb)** their private wave functions to account for the newly acquired knowledge, a coherent worldview emerges.

Seen this way, the wave function "may well be the most powerful abstraction we have ever found," says theoretical physicist N. David Mermin of Cornell University, a recent convert to QBism.

(3) The notion that the wave function isn't real dates back to the 1930s and the writings of Niels Bohr, one of the founding fathers of quantum mechanics. He considered it **part (e&eb) of** quantum

theory's "purely symbolic" formalism—a computational tool, no more.

(4) QBism is the first model to give mathematical backbone to Bohr's assertion. It melds quantum theory with Bayesian statistics, a 200-year-old discipline that defines "probability" **as (=) something like** "subjective belief."

(5) Bayesian statistics **also (eb) gives** formal mathematical rules for how to update one's subjective beliefs in light of new information.

(6) By interpreting the wave function as a subjective belief and subject to revision by the rules of Bayesian statistics, **there shall be (e) obliteration of** mysterious paradoxes of quantum mechanics, QBism's proponents say.

(7) We know that each time we detect an electron; we find **it (e) in** one particular location. But when we're not looking, the electron's wave function can spread out, **representing (eb) the** possibility that the electron is in many different places at once.

(8) Now make a measurement again. You'll find the electron back **in a** particular location. According to the standard way of thinking, the observation c**auses (eb) the** wave function to instantaneously "collapse" back to a single particular value.

NOTATION

Module One

Anagrammatic knowledge(reductionism) kills truth(My model)

Reductionism is a philosophical position which holds that a complex system is nothing but the sum of its parts, and that an account of it can be reduced to accounts of individual constituents. This can be said of objects, phenomena, explanation, theories, and meanings. Fragmentalism is an alternative term for reductionism, although fragmentalism is frequently used in a pejorative sense.

Reductionism strongly reflects a certain perspective on causality. In a reductionist framework, phenomena that can be explained completely in terms of relations between other more fundamental phenomena are called epiphenomena. Often there is an implication that the epiphenomenon exerts no causal agency on the fundamental phenomena that explain it.

Philosophy employs what Plato calls 'the eye of the mind' to go beyond the immediacy revealed by sense experience so as to access the true reality revealed by the intellect. In this view of philosophy, there are levels of cognition which ascend from instincts and desires at the level of immediacy up to reason and intellect at the highest level. The path to truth, knowledge and freedom leads at the summit, the opening at the top of the bottle. This is the way of philosophy. The intellect shows the true reality. This is reality as seen through the eye of the mind. The phrase 'the eye of the mind' comes from Plato. Wittgenstein's fly in the bottle is really a simplified version of Plato's allegory of the prisoners in the cave who, in mistaking the shadows on the wall for the true reality, remain chained by their senses to empirical necessity. Philosophy is the path to enlightenment, to see by the light of the Sun in the real world outside of the cave of illusion. Most people dwell in the blissful ignorance of partial knowledge. For philosophers, however, ignorance is never bliss; it is a passive contentment that leaves human beings living unfulfilled lives which fall far short of the potentialities individuals are born with. Partial knowledge, at best, leads only to a partial freedom and a partial life. But if ignorance isn't bliss, then neither too is knowledge in itself. It depends on the extent to which knowledge **is used to** transform existence in such a way as to enable human beings to live full and flourishing lives. Short of that end, knowledge can be the way of pain, misery and suffering. If we knew everything about the inner workings of the systems, governments, and corporations that run our world, all those forces which cast their shadows on the wall and beguile us with illusion, **then the result might** well be a perpetual state of anxiety. Rather than stand up and be counted, many might well prefer the tranquillity and

contentment of the world of illusion, the cozy world in which reality is as the puppet masters say it is and individuals prefer not to question. The problem is that these troubling questions do not go away and most people, at some point in their lives, wonder whether reality really is as they constantly told it is. The questions are not expressed in a public way but instead go underground, are sublimated, and create some very uneasy consciences. The ability of the human mind continuously exceeds its conceptual grasp, giving human beings an enduring tendency to transcendence. Human beings have an innate predisposition to push their ideas and beliefs to the extreme and beyond. Ultimately, there is nowhere else to go but the transcendent. This is something we need to be aware of and learn to understand. It is an **ineliminable aspect** of human nature. This does not mean that those ideas and beliefs are right. On the contrary, we need to be ever sceptical, alert to the dangers of wishful thinking and the mess that human beings can get them into by investing their existence with transcendental meaning. Human beings develop their culture, their institutions and their practices over time, as they attempt to increase their control over their environments. Human beings are meaning-seeking, rule-following animals by nature. This is reason to be cautious and sceptical when it comes to the world of value. Human beings have a will to believe. Sigmund Freud was resigned to the fact that even after psychology has achieved all it can achieve; human beings will tend to believe what they want to believe. If human beings have a natural tendency to conform to the rules, institutions and practices of the society in which they live, so are they inclined to entrench those social arrangements with transcendent meaning and value. This will to believe reveals nothing of the veracity of the transcendent meaning and value human beings find in the world. Philosophy is not about any old meaning and philosophers have frequently fallen foul for having challenged prevailing social norms and values. Human beings are meaning seeking creatures, and a viable social order needs to speak to that aspect of human nature. But – and this is where philosophy comes in – social norms and rules are not right simply on account of being invested with transcendent meaning and value. The value of philosophy lies in weighing the standards of good, right, truth and value rather than simply accepting those norms and values that prevail in particular times and places. That said, philosophy amounts to more than the policing of belief, language and meaning. **Alfred North Whitehead** was surely correct when he argued: "But if men cannot live on bread alone, still less can they do so on disinfectants."To think that this aspiration to the transcendent **can be undermined by** reason, logic and evidence is a crude eighteenth century error likely to rebound unspectacular fashion, bringing about the opposite of the rational world sought. Many scientists nevertheless commit this error, exhibiting a coldness that is frankly inhumane and unlikely to issue in a civilization that is at peace. Take these blunt words from Victor Stenger: I do not think science has to make any apologies. It looks at the world and tell it like it is. And we all live longer, better lives because of this dispassionate view.

The question whether objective truth can be attributed to human thinking is not a question of theory but is a practical question. Man must prove the truth, i.e. the reality and power, the **this-sidedness** of his thinking in practice. The dispute over the reality or non-reality of thinking that is isolated from practice is a purely scholastic question. This is a call for human beings to live up to their reason, to realise the philosophical ideal without by exercising the rational capacity that is innate within them. Marx's praxis is not the same thing as pragmatism. Marx is not simply arguing the pragmatic case that something is true because it works. His praxis is the unity of theory and practice, the incorporation of the philosophical ideal into practice and the realisation of this ideal in and through practice. In this sense, Marx's revolutionary-critical praxis **is the combination of** both logos and mythos. It emphasizes the human species as the maker of its own reality. Put into practice, the philosophical ideal is both logos and mythos in **revealing something** profoundly true about our essential humanity. There is a need to be clear here. In arguing for philosophy as ethos, as praxis, as a way of life, **one has to be careful** to distinguish philosophy from religion. Religion is not primarily a body of beliefs, a theology, an intellectual appreciation of all that exists but a practice, something that people does rather than think. Religion may therefore be defined as a way of life. Atheists like Richard Dawkins spectacularly miss the point when they bring the weapons of scientific reason to bear upon religious beliefs, as though religion rests on the quality of its intellectual propositions. These beliefs, even the highest

achievements of theology – and these have been high, much higher than Dawkins is capable of appreciating - rationalizations of truths acquired by practical action. Criticism of these beliefs donot dents the meaning that this action brings to believers. It may be a controversial point, but many believers would be hard pressed to state, let alone explain, their beliefs in anything like a coherent intellectual fashion. It can be done, and theologians have done it. But in an important sense, it doesn't matter. I like what Kenneth Clark argues with respect to the remarkable success of the Catholic response to the challenge of the Reformation (**Dr Peter Critchley Philosophizing Through the Eye of the Mind PHILOSOPHISING THROUGH THE EYE OF THE MIND Philosophy as Ethos and Praxis2010B)**

Reductionism does not preclude the existence of what might be called emergent phenomena, but it does imply the ability to understand those phenomena completely in terms of the processes from which they are composed. This reductionist understanding is very different from that usually implied by the term 'emergence', which typically intends that what emerges is more than the sum of the processes from which it emerges.

Religious reductionism generally attempts to explain religion by boiling it down to certain nonreligious causes. A few examples of reductionistic explanations for the presence of religion are: that religion can be reduced to humanity's conceptions of right and wrong, that religion is fundamentally a primitive attempt at controlling our environments, that religion is a way to explain the existence of a physical world, and that religion confers an enhanced survivability for members of a group and so is reinforced by natural selection. Anthropologists Edward Burnett Tylor and James George Frazer employed some religious reductionist arguments. Sigmund Freud's idea that religion is nothing more than an illusion, or even mental illnesses, and the Marxist view that religion is "the sigh of the oppressed," providing only "the illusory happiness of the people," are two other influential reductionist explanations of religion.

More you think about a theory or your theory more and more aggrandizement and exacerbated the conviction of the righteousness of the Theory. There is a song here wherein a lover (a scezophreniac) starts attributing motives to every action of lover. He starts imagining that when she is looking at the horizon, she is mulling and introspecting about him, when she bows her head down, there is she is immersed in his thoughts. I have often wondered why people defend their theories in the most exigent and contingent of circumstances. And it is nothing but the repetitive thinking about their own theories. Now this is a story. Mafia, crime syndicates, hoodlum mugger aggregates, and raucous ribaldry congregates, loutish jeers elements have all joined against you. To others it is just a traffic jam or one hundred people in the park. You know you are being watched. You know you could be attacked. Unless you have the consciousness that you are being followed, to be attacked you never identify the slew of cars and vehicles as meant for your distraction. People can gesticulate, talk in sign language to disparage and denigrate you, and the car moving well within city limits can crush you. Colours are the natural weapon of the vested interests. Because you identify red with danger. They will again and again expose you to the repetitive stances which only go to increase your gratification or deprivation complex. This is distortion of nature. Prakruti Kshobha and Prakruti Siddhhi. By ingeniously using the heritage, the halcyon days and history and hoary past, such "Prakruti Kshobha could be created among large cross section of people. This is more so in a country like India where individual consciousness is that expanded. Effective creation of such states in US may be less due to expansive individual consciousness. In Brahman –Anti Brahman Model I had mentioned, there is virtually no difference between Brahman and Anti Brahman. Both follow the same path. Verily in the subterranean realm and ceratoid dualism of the Brahman-Anti Brahman, there is no dualities' (either you are Chairman or a Chaprasi- a janitor) and there is absolute non polarisation of good and evil.

Dairy Notes: 19th March 2013 night 2.15 PM

"Mind is essentially anagrammatic!"

"So what?"

"Hatred for some and love for some. Everything is subjective"

"Creation?"

"Yes! You are both the observer and the actor but not the script writer?"

"Who does that writing"

"Brahman- Anti Brahman Agency"

"You said they are same"

"Yes! He is both warm, affectionate, protecting and on the other a thunderstorm Thermistor and verkrampte reactionary for obliteration and cosmolysis. Like it is the same energy that gives light and creates nuclear arsenal"

"Nature?"

"Nature pretends. She does show her real self"

"Why?"

" Because she is simulated, Maya"

"Mind is somewhere!"

"What do you mean?"

"You see! It is association in space time that is responsible for all the problems. Moment you are associating there shall be million axiomatic predications, postulation alcovishness and primordial exactitudes arise. Hicks in fact said: Structuralists were seeking subjective noumenal categories, and the Phenomenologists were content with describing the phenomena without asking what connection to an external reality those experiences might have. Association leads to gratification and deprivation complex. It brings you repetitive thoughts. Conscious actions produce sense perceptions which produce association which produce concatenation which leads to Prakruti Kshobha. Neuron DNA stores all your actions and passions. It is your general ledger. That is true in pure consciousness and impure consciousness which separates Brahman and Anti Brahman atleast by action types. "

"You are a Dr. Watson"

"I am yourself!"

G_{13} : Category one of **truth**

G_{14} : Category two of **truth**

G_{15} : Category three of **truth**

T_{13} : Category one of **Anagrammatic knowledge(reductionism)**

T_{14} : Category two of **Anagrammatic knowledge(reductionism)**

T_{15} : Category three of **Anagrammatic knowledge(reductionism)**

Module Two

One system—one event—**can (e) have** as many different wave functions as there are observers

G_{16} : Category one of many different wave functions as there are observers

G_{17} : Category two of many different wave functions as there are observers

G_{18} : Category three of many different wave functions as there are observers

T_{16} : Category one of One system—one event

T_{17} : Category two of One system—one event

T_{18} : Category three of One system—one event

Module three

After observers have communicated with one another and **modified (e&eb)** their private wave functions to account for the newly acquired knowledge, a coherent worldview emerges

G_{20} : Category one of coherent worldview emerges

G_{21} : Category two of coherent worldview emerges

G_{22} : Category three of coherent worldview emerges

T_{20} : Category one of observers have communicated with one another and **modified (e&eb)** their private wave functions to account for the newly acquired knowledge

T_{21} : Category two of observers have communicated with one another and **modified (e&eb)** their private wave functions to account for the newly acquired knowledge

T_{22} : Category three of observers have communicated with one another and **modified (e&eb)** their private wave functions to account for the newly acquired knowledge

Module four

The notion that the wave function isn't real dates back to the 1930s and the writings of Niels Bohr, one of the founding fathers of quantum mechanics. He considered it **part (e&eb) of** quantum theory's "purely symbolic" formalism—a computational tool, no more.

G_{24} : Category one of wave function

G_{25} : Category two of wave function

G_{26} : Category three of wave function

T_{24} : Category one of quantum theory's "purely symbolic" formalism—a computational tool, no more.

T_{25} : Category two of quantum theory's "purely symbolic" formalism—a computational tool, no more.

T_{26} : Category three of quantum theory's "purely symbolic" formalism—a computational tool, no more.

Module five

QBism is the first model to give mathematical backbone to Bohr's assertion. It melds quantum theory with Bayesian statistics, a 200-year-old discipline that defines "probability" **as (=) something like** "subjective belief."

G_{28} : Category one of "subjective belief."

G_{29} : Category two of "subjective belief."

G_{30} : Category three of "subjective belief."

T_{28} : Category one of probability

T_{29} : Category two of probability

T_{30} : Category three of probability

Module six

Bayesian statistics **also (eb) gives** formal mathematical rules for how to update one's subjective beliefs in light of new information.

G_{32} : Category one of Bayesian statistics

G_{33} : Category two of Bayesian statistics

G_{34} : Category three of Bayesian statistics

T_{32} : Category one of formal mathematical rules for how to update one's subjective beliefs in light of new information.

T_{33} : Category two of formal mathematical rules for how to update one's subjective beliefs in light of new information.

T_{34} : Category three of formal mathematical rules for how to update one's subjective beliefs in light of new information.

Module seven

By interpreting the wave function as a subjective belief and subject to revision by the rules of Bayesian statistics, **there shall be (e) obliteration of** mysterious paradoxes of quantum mechanics, QBism's proponents say.

G_{36} : Category one of mysterious paradoxes of quantum mechanics, QBism's proponents say.

G_{37} : Category two of mysterious paradoxes of quantum mechanics, QBism's proponents say.

G_{38} : Category three of mysterious paradoxes of quantum mechanics, QBism's proponents say.

T_{36} : Category one of wave function as a subjective belief and subject to revision by the rules of Bayesian statistics,

T_{37} : Category two of wave function as a subjective belief and subject to revision by the rules of Bayesian statistics,

T_{38} : Category three of wave function as a subjective belief and subject to revision by the rules of Bayesian statistics,

Module eight

We know that each time we detect an electron; we find **it (e) in** one particular location. But when we're not looking, the electron's wave function can spread out, **representing (eb) the** possibility that the electron is in many different places at once.

G_{40} : Category one of electron's wave function can spread out when not observed

G_{41} : Category two of electron's wave function can spread out when not observed

G_{42} : Category three of electron's wave function can spread out when not observed

T_{40} : Category one of possibility that the electron is in many different places at once.

T_{41} : Category two of possibility that the electron is in many different places at once.

T_{42} : Category three of possibility that the electron is in many different places at once.

Module Nine

Now make a measurement again. You'll find the electron back **in a** particular location. According to the standard way of thinking, the observation c**auses (eb) the** wave function to instantaneously “collapse” back to a single particular value

G_{44} : Category one of observation

G_{45} : Category two of observation

G_{46} : Category three of observation

T_{44} : Category one of wave function to instantaneously “collapse” back to a single particular value

T_{45} : Category two of wave function to instantaneously “collapse” back to a single particular value

T_{46} : Category three of wave function to instantaneously “collapse” back to a single particular value

SECTION TWENTY NINE

God Does Not Play Dice: Quod Me Nutrit Me Destruit Qui Tacet Consentire

INTRODUCTION—VARIABLES USED

Concerning Dice and Divinity D.M.Appleby

(1) Einstein initially objected to the probabilistic aspect of quantum mechanics - the idea that God is playing at dice. Later he changed his ground, and focussed instead on the point that the Copenhagen Interpretation **leads to what** Einstein saw as the abandonment of physical realism.
(2) Authors argue here that Einstein's initial intuition was perfectly sound, and that it is precisely the fact that quantum mechanics is a fundamentally probabilistic theory **which is (e) at the root** of all the controversies regarding its interpretation.
(3) Probability is an intrinsically logical concept. **This means that** the quantum state has an essentially logical significance.
(4) It is extremely difficult to reconcile that fact with Einstein's belief, that it is the task of physics **to give us a vision of** the world apprehended sub specie aeternitatis.
(5) Quantum mechanics thus presents us with a simple choice: either to follow Einstein in looking for a theory which is not probabilistic at the fundamental level, **or implies not** to accept that physics does not in fact put us in the position of God looking down on things from above.
(6) There is a widespread fear that the latter alternative must inevitably **lead to a** greatly impoverished, positivistic view of physical theory. It appears to us, however, that the truth is just the opposite. The Einsteinian vision is much less attractive than it seems at first sight. In particular, **it is closely connected with** philosophical reductionism. Contribution to proceedings of Foundations of Probability and Physics, Vaxjo, 2006 arXiv: quant-ph/0611261

Foundations of Physics April 2003, Volume 33, Issue 4, pp 665-676The Information Interpretation and the Conceptual Problems of Quantum Mechanics Miguel Ferrero

(7) It has been traditionally considered that Quantum Mechanics has two conceptual kinds of problems, namely, those related with local-realism and the so-called measurement problem. That is, the uniqueness of the result when we make a measurement. With the development of what is called generically Quantum Information Theory, a new form of the Copenhagen interpretation of the formalism has taken shape. In this paper, we will analyse if this information interpretation **is able (e) to clarify** these old problems. Although this interpretation seems to be the most promising approach we have, we have reached the conclusion that the answer cannot be given in a positive and clear way yet.

High energy astrophysical neutrinos: The upper bound is robust John Bahcall et al

(8) Authors elucidate the physical basis for the upper bound on high energy neutrino fluxes **implied (e) by** the observed cosmic ray flux.

(9) They stress that the bound **is (e)valid** for neutrinos produced either by p, γ reactions or by p-p(n) reactions in sources which are optically thin for high energy protons to photo-meson and nucleon-meson interactions.

(10) Authors show that the upper bound is robust and conservative. The Waxman-Bahcall bound overestimates the most likely neutrino flux by a factor ∼5/τ, for small optical depths τ. The upper limit cannot be plausibly **evaded (e) by** invoking magnetic fields, optically thick active galactic nuclei (AGNs), or large hidden fluxes of extragalactic protons.

(11) They describe the implications of the bound for future experiments including the AMANDA, ANTARES, Auger, ICECUBE, NESTOR, and OWL/AIRWATCH detectors.© 2001 The American Physical societyURLhttp://link.aps.org/doi/10.1103/PhysRevD.64.023002DOI:10.1103/PhysRevD.64.023002

NOTATION

Module One

Einstein initially objected to the probabilistic aspect of quantum mechanics - the idea that God is playing at dice. Later he changed his ground, and focussed instead on the point that the Copenhagen Interpretation **leads to what** Einstein saw as the abandonment of physical realism

G_{13} : Category one of Copenhagen Interpretation

G_{14} : Category two of Copenhagen Interpretation

G_{15} : Category three of Copenhagen Interpretation

T_{13} : Category one of abandonment of physical realism

T_{14} : Category two of abandonment of physical realism

T_{15} : Category three of abandonment of physical realism

Module Two

Authors argue here that Einstein's initial intuition was perfectly sound, and that it is precisely the fact that quantum mechanics is a fundamentally probabilistic theory **which is (e) at the root** of all the controversies regarding its interpretation.

G_{16} : Category one of controversies regarding its interpretation.

G_{17} : Category two of controversies regarding its interpretation.

G_{18} : Category three of controversies regarding its interpretation.

T_{16} : Category one of Einstein's initial intuition was perfectly sound, and that it is precisely the fact that quantum mechanics is a fundamentally probabilistic theory

T_{17} : Category two of Einstein's initial intuition was perfectly sound, and that it is precisely the fact that quantum mechanics is a fundamentally probabilistic theory

T_{18} : Category three of Einstein's initial intuition was perfectly sound, and that it is precisely the fact that quantum mechanics is a fundamentally probabilistic theory

Module three

Probability is an intrinsically logical concept. **This means that** the quantum state has an essentially logical significance

G_{20} : Category one of Probability is an intrinsically logical concept

G_{21} : Category two of Probability is an intrinsically logical concept

G_{22} : Category three of Probability is an intrinsically logical concept

T_{20} : Category one of quantum state has an essentially logical significance

T_{21} : Category two of quantum state has an essentially logical significance

T_{22} : Category three of quantum state has an essentially logical significance

Module four

It is extremely difficult to reconcile that fact with Einstein's belief, that it is the task of physics **to give us a vision of** the world apprehended sub specie aeternitatis

G_{24} : Category one of task of physics

G_{25} : Category two of task of physics

G_{26} : Category three of task of physics

T_{24} : Category one of **vision of** the world apprehended sub specie aeternitatis

T_{25} : Category two of **vision of** the world apprehended sub specie aeternitatis

T_{26} : Category three of **vision of** the world apprehended sub specie aeternitatis

Module five

Quantum mechanics thus presents us with a simple choice: either to follow Einstein in looking for a theory which is not probabilistic at the fundamental level, **or implies not** to accept that physics does not in fact put us in the position of God looking down on things from above.

G_{28} : Category one of follow Einstein in looking for a theory which is not probabilistic at the fundamental level. Classification is done based on the characterstics of the investigating systems.

G_{29} : Category two of Einstein in looking for a theory which is not probabilistic at the fundamental level. Classification is done based on the characterstics of the investigating systems

G_{30} : Category three of Einstein in looking for a theory which is not probabilistic at the fundamental level. Classification is done based on the characterstics of the investigating systems

T_{28} : Category one of physics does not in fact put us in the position of God looking down on things from above.

T_{29} : Category two of physics does not in fact put us in the position of God looking down on things from

above.

T_{30} : Category three of physics does not in fact put us in the position of God looking down on things from above.

Module six

There is a widespread fear that the latter alternative must inevitably **lead to a** greatly impoverished, positivistic view of physical theory. It appears to us, however, that the truth is just the opposite. The Einsteinian vision is much less attractive than it seems at first sight. In particular, **it is closely connected with** philosophical reductionism. Contribution to proceedings of Foundations of Probability and Physics, Vaxjo, 2006 arXiv: quant-ph/0611261

G_{32} : Category one of Einsteinian vision is much less attractive than it seems at first sight; philosophical reductionism

G_{33} : Category two of Einsteinian vision is much less attractive than it seems at first sight; philosophical reductionism

G_{34} : Category three of Einsteinian vision is much less attractive than it seems at first sight; philosophical reductionism

T_{32} : Category one of philosophical reductionism; Einsteinian vision is much less attractive than it seems at first sight

T_{33} : Category two of philosophical reductionism; Einsteinian vision is much less attractive than it seems at first sight

T_{34} : Category three of philosophical reductionism; Einsteinian vision is much less attractive than it seems at first sight

Module seven

Authors elucidate the physical basis for the upper bound on high energy neutrino fluxes **implied (e) by** the observed cosmic ray flux

G_{36} : Category one of observed cosmic ray flux

G_{37} : Category two of observed cosmic ray flux

G_{38} : Category three of observed cosmic ray flux

T_{36} : Category one of physical basis for the upper bound on high energy neutrino fluxes

T_{37} : Category two of physical basis for the upper bound on high energy neutrino fluxes

T_{38} : Category three of physical basis for the upper bound on high energy neutrino fluxes

Module eight

They stress that the bound **is (e)valid** for neutrinos produced either by p, γ reactions or by p-p(n) reactions in sources which are optically thin for high energy protons to photo-meson and nucleon-meson interactions.

G_{40} : Category one of neutrinos produced either by p, γ reactions or by p-p(n) reactions in sources which are optically thin for high energy protons to photo-meson and nucleon-meson interactions

G_{41} : Category two of neutrinos produced either by p, γ reactions or by p-p(n) reactions in sources which are optically thin for high energy protons to photo-meson and nucleon-meson interactions

G_{42} : Category three of neutrinos produced either by p, γ reactions or by p-p(n) reactions in sources which are optically thin for high energy protons to photo-meson and nucleon-meson interactions

T_{40} : Category one of bound mentioned in the foregoing

T_{41} : Category two of bound mentioned in the foregoing

T_{42} : Category three of bound mentioned in the foregoing

Module Nine

Authors show that the upper bound is robust and conservative; the Waxman-Bahcall bound overestimates the most likely neutrino flux by a factor ∼5/τ, for small optical depths τ; the upper limit cannot be plausibly **evaded (e) by** invoking magnetic fields, optically thick active galactic nuclei (AGNs), or large hidden fluxes of extragalactic protons

G_{44} : Category one of invoking magnetic fields, optically thick active galactic nuclei (AGNs), or large hidden fluxes of extragalactic protons

G_{45} : Category two of invoking magnetic fields, optically thick active galactic nuclei (AGNs), or large hidden fluxes of extragalactic protons

G_{46} : Category three of invoking magnetic fields, optically thick active galactic nuclei (AGNs), or large hidden fluxes of extragalactic protons

T_{44} : Category one of upper bound is robust and conservative; the Waxman-Bahcall bound overestimates the most likely neutrino flux by a factor ∼5/τ, for small optical depths τ; the upper limit cannot be plausibly evaded

T_{45} : Category two of upper bound is robust and conservative; the Waxman-Bahcall bound overestimates the most likely neutrino flux by a factor ∼5/τ, for small optical depths τ; the upper limit cannot be plausibly evaded

T_{46} : Category three of upper bound is robust and conservative; the Waxman-Bahcall bound overestimates the most likely neutrino flux by a factor ∼5/τ, for small optical depths τ; the upper limit cannot be plausibly evaded

SECTION THIRTY

Nagarjuna's Tetra Lemma : Legum Servi Sumus Ut Liberi Esse Possimus - Mutantur Omnia Nos Et Mutamur In Illis

INTRODUCTION— Logics of tensor strain in co-Tangent bundles over a general Manifold for Musculpt by VirFut Q-Pro μTm Scenarios, T4(M) Some paragraphs are deleted due to spatial constraints (normativity the chaotic multitude of homonymies, word-plays, "irregular" metaphoric links and resonances. in quantum computation)

(1) As noted in MOON, I read Christopher Isherwood's translation of Adi Shankara's Crest-Jewel of Discrimination sometime in the late-'70s and lost interest altogether when it became apparent that Shankara regarded the appearance of infinite regress as being **equivalent to** what modern logicians call a fallacy.

(2) How could anyone seriously engaged with concentration in self-observation possibly entertain such a notion? Sakyamuni Buddha after 1000 incarnations appearing discretely in 3-space at a particular memetime as Trikaya Gautama, Trinitarian, i.e., 3-bodied -- the 3-valued being a numinous **violation (e) of the** prevalent, socially-condoned bi-valent logic of the period, East and Middle East).

(3) Authenticity deals in **are (=)** indisputably transfinite, and the residue of understanding of this is much more elaborately on display in Bon Tzog-chen than in Madhyamika.

(4) We would like to further clarify Nagarjuna's tetra lemma. Scholar Edward Conze presents the four lemmas as: (i) x exists (ii) x does not exist, (iii) x both exists and does not exist, (iv) x neither exists nor does not exist."

(5) The modern Buddhist philosopher T. R. V. Murti characterizes it thusly: "These four alternatives represent all the possible standpoints from which every problem can be viewed; they **also provide** a schema under which all systems of philosophy can be classified." Another way of looking at this is provided by Richard Robinson, a contemporary translator of Eastern texts, who views each lemma as representing an "existential quantification," each differing in quantity of its constituent terms, so that (1) all x is A, (2) no x is A, (3) some x is A and some is not A, and (4) no x is A and no x is not A.

(6) What about the problem of the constitution of the phenomenological object, that problem which Edmund Husserl addressed in developing "eidetic reduction" (a practice an actual practitioner of walking meditation might engage in -- even if Husserl clearly didn't)? Does not each such object have a transfinite collection (mdo) of (Platonic) "horizons" -- **each of which (e&eb) corresponds** to a "standpoint" from which the object (i.e., the "problem", the problem of how an object with a transfinite set of distinct and different horizons is cognitively constituted as a simply-identified single-valued whatnot) "can be viewed"?

(7) This problem does have something to do with questions relevant to the notions of maya and samsara which Buddhism does concern itself with. If I am not mistaken, a set of four elements, both as regards cardinality and ordinality, is considerably **less (e) than all** of the elements of a transfinite set.

(8) Which takes us to Robinson's lemmas? None of his four lemmas **violate (eb)** the standard Aristotelian-Baconian-Boolean rules of 1T2 logic.

(9) The forth lemma is the most likely to be mistaken as representing **such a(e) violation**, but then there is always B, C, D… to consider, along with their identity relations to A.

NOTATION

Module One

As noted in MOON, I read Christopher Isherwood's translation of Adi Shankara's Crest-Jewel of Discrimination sometime in the late-'70s and lost interest altogether when it became apparent that Shankara regarded the appearance of infinite regress as being **equivalent to** what modern logicians call a fallacy.

G_{13} : Category one of what modern logicians call a fallacy

G_{14} : Category two of what modern logicians call a fallacy

G_{15} : Category three of what modern logicians call a fallacy

T_{13} : Category one of appearance of infinite regress

T_{14} : Category two of appearance of infinite regress

T_{15} : Category three of appearance of infinite regress

Module Two

How could anyone seriously engaged with concentration in self-observation possibly entertain such a notion? Sakyamuni Buddha after 1000 incarnations appearing discretely in 3-space at a particular memetime as Trikaya Gautama, Trinitarian, i.e., 3-bodied -- the 3-valued being a numinous **violation (e) of the** prevalent, socially-condoned bi-valent logic of the period, East and Middle East).

G_{16} : Category one of prevalent, socially-condoned bi-valent logic of the period, East and Middle East).

G_{17} : Category two of prevalent, socially-condoned bi-valent logic of the period, East and Middle East).

G_{18} : Category three of prevalent, socially-condoned bi-valent logic of the period, East and Middle East).

T_{16} : Category one of Trinitarian, i.e., 3-bodied -- the 3-valued being a numinous

T_{17} : Category two of Trinitarian, i.e., 3-bodied -- the 3-valued being a numinous

T_{18} : Category three of Trinitarian, i.e., 3-bodied -- the 3-valued being a numinous

Module three

authenticity deals in **are (=)** indisputably transfinite, and the residue of understanding of this is much more elaborately on display in Bon Tzog-chen than in Madhyamika

G_{20} : Category one of indisputably transfinite, and the residue of understanding of this is much more elaborately on display in Bon Tzog-chen than in Madhyamika

G_{21} : Category two of indisputably transfinite, and the residue of understanding of this is much more elaborately on display in Bon Tzog-chen than in Madhyamika

G_{22} : Category three of indisputably transfinite, and the residue of understanding of this is much more elaborately on display in Bon Tzog-chen than in Madhyamika

T_{20} : Category one of authenticity expatiation ,enucleation and elucidation

T_{21} : Category two of authenticity expatiation ,enucleation and elucidation

T_{22} : Category three of authenticity expatiation ,enucleation and elucidation

Module four

We would like to further clarify Nagarjuna's tetra lemma. Scholar Edward Conze presents the four lemmas as: (i) x exists (ii) x does not exist, (iii) x both exists and does not exist, (iv) x neither exists nor does not exist."

G_{24} : Category one of x exists; x both exists and does not exist

G_{25} : Category two of x exists; x both exists and does not exist

G_{26} : Category three of x exists; x both exists and does not exist

T_{24} : Category one of x does not exist; "x neither exists nor does not exist."

T_{25} : Category two of x does not exist; "x neither exists nor does not exist."

T_{26} : Category three of x does not exist; "x neither exists nor does not exist."

Module five

The modern Buddhist philosopher T. R. V. Murti characterizes it thusly: "These four alternatives represent all the possible standpoints from which every problem can be viewed; they **also provide** a schema under which all systems of philosophy can be classified." Another way of looking at this is provided by Richard Robinson, a contemporary translator of Eastern texts, who views each lemma as representing an "existential quantification," each differing in quantity of its constituent terms, so that (1) all x is A, (2) no x is A, (3) some x is A and some is not A, and (4) no x is A and no x is not A.

G_{28} : Category one of all x is A; some x is A and some is not A

G_{29} : Category two of all x is A; some x is A and some is not A

G_{30} : Category three of all x is A; some x is A and some is not A

T_{28} : Category one of no x is A; no x is A and no x is not A.

T_{29} : Category two of no x is A; no x is A and no x is not A.

T_{30} : Category three of no x is A; no x is A and no x is not A.

Module six

What about the problem of the constitution of the phenomenological object, that problem which Edmund Husserl addressed in developing "eidetic reduction" (a practice an actual practitioner of walking meditation might engage in -- even if Husserl clearly didn't)? Does not each such object have a transfinite collection (mdo) of (Platonic) "horizons" -- **each of which (e&eb) corresponds** to a "standpoint" from which the object (i.e., the "problem", the problem of how an object with a transfinite set of distinct and different horizons is cognitively constituted as a simply-identified single-valued whatnot) "can be viewed"?

Standpoint theory is a postmodern method **for analyzing** inter-subjective discourses. This theory concerns the authority **generated by** people's knowledge and the power such authority has to shape people's opinions in daily life. Standpoint theory's most important concept is that an individual's own perspectives are shaped by his or her experiences in social locations and social groups. Standpoints always involve more than one factor. For example, if you look at different Hispanic women, their standpoints may resemble one another in terms of race and biological sex categories; however, if their socioeconomic status is different, their standpoints are not completely the same. These perspectives are the core point of view for individuals to see

the world. Standpoint theory focuses especially on gender perspectives to see how feminine viewpoints shape women's communication with themselves, others, and the world. Standpoint theory has a huge effect on how people's perceptions change from one thing to another. A standpoint is a place from which one views and sees the world, which determines both what one, focuses on as well as what is obscured. Depending on one's situation, one's standpoint may vary from that of another individual who may be of a similar status.

Standpoint theories are said to remind people why a naturalistic conception of knowing is important. Knowledge helps people understand part of the world that they normally tend to not understand. Gaining knowledge occurs only in specific circumstances and has real consequences. These consequences can have an effect on how a person can live his or her life. It matters politically as well as epistemically which concepts are intelligible, which claims are heard and understood by whom, which features of the world are perceptually salient, and which reasons are understood to be relevant and forceful, as well as which conclusions credible.

Standpoint theory supports what feminist theorist Sandra Harding calls strong objectivity, or the notion that the perspectives of marginalized and/or oppressed individuals can help to create more objective accounts of the world. Through the outsider-within phenomenon, these individuals are placed in a unique position to point to patterns of behavior that those immersed in the dominant group culture are unable to recognize Standpoint theory gives voice to the marginalized groups by allowing them to challenge the status quo as the outsider within. The status quo representing the dominant white male position of privilege.

The predominant culture in which all groups exist is not experienced in the same way by all persons or groups. The views of those who belong to groups with more social power are validated more than those in marginalized groups. Those in marginalized groups must learn to be bicultural, or to "pass" in the dominant culture to survive, even though that perspective is not their own. For persons of color, in an effort to help organizations achieve their diversity initiatives, there is an expectation that they will check their color at the door in order to assimilate into the existing culture and discursive practices.(Wikipedia)

G_{32} : Category one of constitution of the phenomenological object, that problem which Edmund Husserl addressed in developing “eidetic reduction” (a practice an actual practitioner of walking meditation might engage in -- even if Husserl clearly didn’t)? Does not each such object have a transfinite collection (mdo) of (Platonic) “horizons” --

G_{33} : Category two of constitution of the phenomenological object, that problem which Edmund Husserl addressed in developing “eidetic reduction” (a practice an actual practitioner of walking meditation might engage in -- even if Husserl clearly didn’t)? Does not each such object have a transfinite collection (mdo) of (Platonic) “horizons” --

G_{34} : Category three of constitution of the phenomenological object, that problem which Edmund Husserl addressed in developing “eidetic reduction” (a practice an actual practitioner of walking meditation might engage in -- even if Husserl clearly didn’t)? Does not each such object have a transfinite collection (mdo) of (Platonic) “horizons” --

T_{32} : Category one of “standpoint” from which the object (i.e., the “problem”, the problem of how an object with a transfinite set of distinct and different horizons is cognitively constituted as a simply-identified single-valued whatnot) “can be viewed”?

T_{33} : Category two of “standpoint” from which the object (i.e., the “problem”, the problem of how an object with a transfinite set of distinct and different horizons is cognitively constituted as a simply-identified single-valued whatnot) “can be viewed”?

T_{34} : Category three of “standpoint” from which the object (i.e., the “problem”, the problem of how an object with a transfinite set of distinct and different horizons is cognitively constituted as a simply-identified

single-valued whatnot) "can be viewed"?

Module seven

Which takes us to Robinson's lemmas? None of Robinson's lemmas four lemmas **violate (eb)** the standard Aristotelian-Baconian-Boolean rules of 1T2 logic.

G_{36} : Category one of None of Robinson's lemmas four lemmas

G_{37} : Category two of None of Robinson's lemmas four lemmas

G_{38} : Category three of None of Robinson's lemmas four lemmas

T_{36} : Category one of standard Aristotelian-Baconian-Boolean rules of 1T2 logic.

T_{37} : Category two of standard Aristotelian-Baconian-Boolean rules of 1T2 logic.

T_{38} : Category three of standard Aristotelian-Baconian-Boolean rules of 1T2 logic.

Module eight

This problem does have something to do with questions relevant to the notions of maya and samsara which Buddhism does concern itself with. If I am not mistaken, a set of four elements, both as regards cardinality and ordinality, is considerably **less (e) than all** of the elements of a transfinite set.

G_{40} : Category one of elements of a transfinite set

G_{41} : Category two of elements of a transfinite set

G_{42} : Category three of elements of a transfinite set

T_{40} : Category one of set of four elements, both as regards cardinality and ordinality

T_{41} : Category two of set of four elements, both as regards cardinality and ordinality

T_{42} : Category three of set of four elements, both as regards cardinality and ordinality

Module Nine

The fourth lemma is the most likely to be mistaken as representing **such a(e) violation**, but then there is always B, C, D… to consider, along with their identity relations to A.

G_{44} : Category one of there is always B, C, D… to consider, along with their identity relations to A.

G_{45} : Category two of there is always B, C, D… to consider, along with their identity relations to A.

G_{46} : Category three of there is always B, C, D… to consider, along with their identity relations to A.

T_{44} : Category one of fourth lemma (is the most likely to be mistaken as representing **such a(e) violation)**

T_{45} : Category two of fourth lemma (is the most likely to be mistaken as representing **such a(e) violation)**

T_{46} : Category three of fourth lemma(is the most likely to be mistaken as representing **such a(e) violation)**

SECTION THIRTY ONE

Cyclotomic Quinary: Fiat Lux Model

INTRODUCTION—VARIABLES USED

Crowley described the encounter in detail in The Equinox of the Gods, saying Wikipedia)

The Voice of Aiwass came apparently from over my left shoulder, from the furthest corner of the room. It seemed to echo itself in my physical heart in a very strange manner, hard to describe. I have noticed a similar phenomenon when I have been waiting for a message fraught with great hope or dread. The voice was passionately poured, as if Aiwass were alert about the time- limit. [...] The voice was of deep timbre, musical and expressive, its tones solemn, voluptuous, tender, fierce or aught else as suited the moods of the message. Not bass --perhaps a rich tenor or baritone. The English was free of either native or foreign accent, perfectly pure of local or caste mannerisma, thus startling and even uncanny at first hearing. I had a strong impression that the speaker was actually in the corner where he seemed to be, in a body of "fine matter," transparent as a veil of gauze, or a cloud of incense-smoke. He seemed to be a tall, dark man in his thirties, well-knit, active and strong, with the face of a savage king, and eyes veiled lest their gaze should destroy what they saw. The dress was not Arab; it suggested Assyria or Persia, but very vaguely. I took little note of it, for to me at that time Aiwass and an "angel" such as I had often seen in visions, a being purely astral.

In the later-written Liber 418, the voice of the 8th Aethyr says "my name is called Aiwass," and "in The Book of the Law did I write the secrets of truth that are like unto a star and a snake and a sword." Crowley says this later manifestation took the form of a pyramid of light.

Identity

Crowley went to great pains to argue that Aiwass was an objectively separate being from himself, possessing far more knowledge than he or any other human could possibly have. He wrote "...no forger could have prepared so complex a set of numerical and literal puzzles [...]" As Crowley writes in his Confessions: "I was bound to admit that Aiwass had shown a knowledge of the Cabbala immeasurably superior to my own" and "We are forced to conclude that the author of The Book of the Law is an intelligence both alien and superior to myself, yet acquainted with my inmost secrets; and, most important point of all, that this intelligence is discarnate." Finally, this excerpt (also from Confessions, ch.49):

The existence of true religion presupposes that of some discarnate intelligence, whether we call him God or anything else. And this is exactly what no religion had ever proved scientifically. And this is what The Book

of the Law does prove by internal evidence, altogether independent of any statement of mine. This proof is evidently the most important step in science that could possibly be made: for it opens up an entirely new avenue to knowledge. The immense superiority of this particular intelligence, AIWASS, to any other with which mankind has yet been in conscious communication is shown not merely by the character of the book itself, but by the fact of his comprehending perfectly the nature of the proof necessary to demonstrate the fact of his own existence and the conditions of that existence. And, further, having provided the proof required.

However, Crowley also spoke of Aiwass in symbolic terms. In The Law is for All, he goes on at length in comparison to various other deities and spiritual concepts, but most especially to The Fool. For example, he writes of Aiwass: "In his absolute innocence and ignorance he is The Fool; he is the Saviour, being the Son who shall trample on the crocodiles and tigers, and avenge his father Osiris. Thus we see him as the Great Fool of Celtic legend, the Pure Fool of Act I of Parsifal, and, generally speaking, the insane person whose words have always been taken for oracles."

Perhaps more importantly, Crowley later identified Aiwass as his own personal Holy Guardian Angel and more. Again from Equinox of the Gods: "I now incline to believe that Aiwass is not only the God once held holy in Sumer, and mine own Guardian Angel, but also a man as I am, insofar as He uses a human body to make His magical link with Mankind, whom He loves, and that He is thus an Ipsissimus, the Head of the A.'.A.'."

Logics of tensor strain in co-Tangent bundles over a general Manifold for Musculpt by VirFut Q-Pro µTm Scenarios, T4(M) Some paragraphs are deleted due to spatial constraints (normativity the chaotic multitude of homonymies, word-plays, "irregular" metaphoric links and resonances. in quantum computation)

(1) I would note here that this general treatment of classic Madhyamika (in its several varieties) slides in the direction **of(e) removing** truth-value as the foundational notion of logic without explicitly developing order-type/index of identity-transparency (quantal non-simple identity, no mere matter of "existential quantification") as a replacement, **for that would(E) require** that the good, the true, and the beautiful be recognized as context-dependent values, not numinous irreducible universals -- a notion anathema to any sort of fundamentalism, which is always generative and hence necessarily recursive and consequently dependent upon elaboration of precedence relations, personified or not

(2) Such multiple linear scenarios, generated by iterations of quadripolar scenario logics, are regarded alternative possible futures, only one of which could become actual; a notion differing considerably from how such scenarios would be conceived under Vir-Fut Q-Pro. Targ and Hurtak in The End of Suffering propose a kind of logical set of Cartesian coordinates, or possibly a logical Feynman diagram, by analogy only, of course. The first two Conze-Nagarjuna lemmas **are placed (eb) at** the ends of an indefinitely extended line; the last two, at the ends of another such line.

(3) The two lines are placed rectilinearly to form a kind of logical plane. Where the two lemma-lines intersect is the origin, and the origin **represents (eb) t**he 4-valued proposition required to establish presence of a 4-valued logic: actual statement of this origin-"statement" is never actually stated. Notice, as Targ and Hurtak apparently did not, that Conze-Nagarjuna's third and forth lemmas can be "true" **iff (eb)** x is transfinitely m-polyvalent (one class of all such x being any member of the set of all phenomenological objects considered under Husserl's eidetic reduction, i.e., objects with a transfinite collection [mdo] of [Platonic] "horizons").

(4) The class of non-simple identity referred to by lemma three is precisely that which Cantor **demonstrated(eb) with** his notorious "diagonal proof"; the class of non-simple identity referred to by lemma four is precisely that which Cantor referred to with his famous "continuum hypothesis".

(5) But a 4-valued logic is only adequate to raise a few of the issues involved, not to provide an exhaustive account thereof. Our present understanding of this is very much due to the efforts of Niels Henrik Abel and the number 5, not the number 4. His theorem on the quintic, an equation with a fifth-power-exponent variable, particularly as explicated by Eric Temple Bell in 1912, The Cyclotomic Quinary Quintic, (non-generative, non-recursive; the cyclotomic cascade as Samantabhadra **being(=)** decomposition of the already always there/here by mathematical involutes, the decomposition of the already always there/here in no way violating absence of precedence relations, logical, ontological, and temporal, for where is precedence in the already always there/here, regardless of what is done with it?) showed not only the necessity of transfinite process algebras (i.e., transcendental to "al-jabr proper"), but also of the multiformity of variables, variables lacking simple-identity by virtue of being m-valued, variables, that is, that are m-polyvalent (p-adic rather than monadic: untruncated p-adicity being a prerequisite to unbounded wholeness).

(6) Thelema of Aiwass: Each and every number is infinite (Liber AL vel Legis, sub figura CCXX, or rather CTC-xk-1+……xn-k). Multi-valued variables **are not (e&eb) the same as** multi-valued logics, but the former inevitably leads to the latter, and the path between the two passes through transfinite sets.

(7) After Abel, after Cantor, quasi-after Lukasiewicz, there was Analytical Cubism, which Jean Gebser in The Ever-Present Origin (which Targ and Hurtak argue is 4-valued) **involved (e) a** "concretion of time"

(8) Whereupon Husserl's transfinite (Platonic) horizons of the phenomenological object are **superposed (e&eb)** in just such a fashion as to constitute a Ponderable object with Piaget's property of object constancy (Picasso and Braque brought the subliminal superposition into relief on the canvas by painting signifying referents thereto).

(9) Phenomenological constitution of the discrete Ponderable macroscopic object **requires (e)** menhir, i.e., Tantric Schrödinger superposition, not a collapse of the wave-function (such a collapse would have been a matter of considerable embarrassment for Schrödinger in midst of the type of double-slit experiment he was conducting at the time he wrote his time-independent wave equation: ménage à trios).

NOTATION

Module One

I would note here that this general treatment of classic Madhyamika (in its several varieties) slides in the direction **of(e) removing** truth-value as the foundational notion of logic without explicitly developing order-type/index of identity-transparency (quantal non-simple identity, no mere matter of "existential quantification") as a replacement, **for that would(E) require** that the good, the true, and the beautiful be recognized as context-dependent values, not numinous irreducible universals -- a notion anathema to any sort of fundamentalism, which is always generative and hence necessarily recursive and consequently dependent upon elaboration of precedence relations, personified or not

G_{13} : Category one of good, the true, and the beautiful be recognized as context-dependent values, not numinous irreducible universals -- a notion anathema to any sort of fundamentalism, which is always generative and hence necessarily recursive and consequently dependent upon elaboration of precedence relations, personified or not

G_{14} : Category two of good, the true, and the beautiful be recognized as context-dependent values, not numinous irreducible universals -- a notion anathema to any sort of fundamentalism, which is always generative and hence necessarily recursive and consequently dependent upon elaboration of precedence relations, personified or not

G_{15} : Category three of good, the true, and the beautiful be recognized as context-dependent values, not numinous irreducible universals -- a notion anathema to any sort of fundamentalism, which is always generative and hence necessarily recursive and consequently dependent upon elaboration of precedence

relations, personified or not

T_{13} : Category one of Madhyamika (in its several varieties) slides in the direction **of(e) removing** truth-value as the foundational notion of logic without explicitly developing order-type/index of identity-transparency (quantal non-simple identity, no mere matter of "existential quantification") as a replacement,

T_{14} : Category two of Madhyamika (in its several varieties) slides in the direction **of(e) removing** truth-value as the foundational notion of logic without explicitly developing order-type/index of identity-transparency (quantal non-simple identity, no mere matter of "existential quantification") as a replacement,

T_{15} : Category three of Madhyamika (in its several varieties) slides in the direction **of(e) removing** truth-value as the foundational notion of logic without explicitly developing order-type/index of identity-transparency (quantal non-simple identity, no mere matter of "existential quantification") as a replacement,

Module Two

Such multiple linear scenarios, **generated by** iterations of quadripolar scenario logics, are regarded alternative possible futures, only one of which could become actual; a notion differing considerably from how such scenarios would be conceived under Vir-Fut Q-Pro. Targ and Hurtak in The End of Suffering propose a kind of logical set of Cartesian coordinates, or possibly a logical Feynman diagram, by analogy only, of course. The first two Conze-Nagarjuna lemmas **are placed (eb) at** the ends of an indefinitely extended line; the last two, at the ends of another such line.

G_{16} : Category one of iterations of quadripolar scenario logics, are regarded alternative possible futures, only one of which could become actual; a notion differing considerably from how such scenarios would be conceived under Vir-Fut Q-Pro. Targ and Hurtak in The End of Suffering propose a kind of logical set of Cartesian coordinates, or possibly a logical Feynman diagram, by analogy

G_{17} : Category two of iterations of quadripolar scenario logics, are regarded alternative possible futures, only one of which could become actual; a notion differing considerably from how such scenarios would be conceived under Vir-Fut Q-Pro. Targ and Hurtak in The End of Suffering propose a kind of logical set of Cartesian coordinates, or possibly a logical Feynman diagram, by analogy

G_{18} : Category three of iterations of quadripolar scenario logics, are regarded alternative possible futures, only one of which could become actual; a notion differing considerably from how such scenarios would be conceived under Vir-Fut Q-Pro. Targ and Hurtak in The End of Suffering propose a kind of logical set of Cartesian coordinates, or possibly a logical Feynman diagram, by analogy

T_{16} : Category one of multiple linear scenarios

T_{17} : Category two of multiple linear scenarios

T_{18} : Category three of multiple linear scenarios

Module three

The two lines are placed rectilinearly to form a kind of logical plane. Where the two lemma-lines intersect is the origin, and the origin **represents (eb) t**he 4-valued proposition required to establish presence of a 4-valued logic: actual statement of this origin-"statement" is never actually stated. Notice, as Targ and Hurtak apparently did not, that Conze-Nagarjuna's third and forth lemmas can be "true" **iff (eb)** x is transfinitely m-polyvalent (one class of all such x being any member of the set of all phenomenological objects considered under Husserl's eidetic reduction, i.e., objects with a transfinite collection [mdo] of [Platonic] "horizons").

G_{20} : Category one of The two lines are placed rectilinearly to form a kind of logical plane. Where the two lemma-lines intersect is the origin, and the origin; Conze-Nagarjuna's third and forth lemmas can be "true";

G_{21} : Category two of The two lines is placed rectilinearly to form a kind of logical plane. Where the two lemma-lines intersect is the origin, and the origin; Conze-Nagarjuna's third and forth lemmas can be “true”;

G_{22} : Category three of The two lines is placed rectilinearly to form a kind of logical plane. Where the two lemma-lines intersect is the origin, and the origin; Conze-Nagarjuna's third and forth lemmas can be “true”;

T_{20} : Category one of4-valued proposition required to establish presence of a 4-valued logic: actual statement of this origin-“statement”; x is transfinitely m-polyvalent (one class of all such x being any member of the set of all phenomenological objects considered under Husserl's eidetic reduction, i.e., objects with a transfinite collection [mdo] of [Platonic] “horizons”).

T_{21} : Category two of4-valued proposition required to establish presence of a 4-valued logic: actual statement of this origin-“statement”; x is transfinitely m-polyvalent (one class of all such x being any member of the set of all phenomenological objects considered under Husserl's eidetic reduction, i.e., objects with a transfinite collection [mdo] of [Platonic] “horizons”).

T_{22} : Category three of4-valued proposition required to establish presence of a 4-valued logic: actual statement of this origin-“statement”; x is transfinitely m-polyvalent (one class of all such x being any member of the set of all phenomenological objects considered under Husserl's eidetic reduction, i.e., objects with a transfinite collection [mdo] of [Platonic] “horizons”).

Module four;

The class of non-simple identity referred to by lemma three is precisely that which Cantor **demonstrated(eb) with** his notorious “diagonal proof”; the class of non-simple identity referred to by lemma four is precisely that which Cantor referred to with his famous “continuum hypothesis”.

G_{24} : Category one of class of non-simple identity referred to by lemma three is precisely that which Cantor

G_{25} : Category two of class of non-simple identity referred to by lemma three is precisely that which Cantor

G_{26} : Category three of class of non-simple identity referred to by lemma three is precisely that which Cantor

T_{24} : Category one of “diagonal proof”; the class of non-simple identity referred to by lemma four is precisely that which Cantor referred to with his famous “continuum hypothesis”.

T_{25} : Category two of “diagonal proof”; the class of non-simple identity referred to by lemma four is precisely that which Cantor referred to with his famous “continuum hypothesis”.

T_{26} : Category three of “diagonal proof”; the class of non-simple identity referred to by lemma four is precisely that which Cantor referred to with his famous “continuum hypothesis”.

Module five

But a 4-valued logic is only adequate to raise a few of the issues involved, not to provide an exhaustive account thereof. Our present understanding of this is very much due to the efforts of Niels Henrik Abel and the number 5, not the number 4. His theorem on the quintic, an equation with a fifth-power-exponent variable, particularly as explicated by Eric Temple Bell in 1912, The Cyclotomic Quinary Quintic, (non-generative, non-recursive; the cyclotomic cascade as Samantabhadra **being(=)** decomposition of the already always there/here by mathematical involutes, the decomposition of the already always there/here in no way violating absence of precedence relations, logical, ontological, and temporal, for where is precedence in the already always there/here, regardless of what is done with it?) showed not only the necessity of transfinite process algebras (i.e., transcendental to “al-jabr proper”), but also of the multiformity of variables, variables

lacking simple-identity by virtue of being m-valued, variables, that is, that are m-polyvalent (p-adic rather than monadic: untruncated p-adicity being a prerequisite to unbounded wholeness).

G_{28} : Category one of decomposition of the already always there/here by mathematical involutes, the decomposition of the already always there/here in no way violating absence of precedence relations, logical, ontological, and temporal, for where is precedence in the already always there/here, regardless of what is done with it?) showed not only the necessity of transfinite process algebras (i.e., transcendental to "al-jabr proper"), but also of the multiformity of variables, variables lacking simple-identity by virtue of being m-valued, variables, that is, that are m-polyvalent (p-adic rather than monadic: untruncated p-adicity being a prerequisite to unbounded wholeness).

G_{29} : Category two of decomposition of the already always there/here by mathematical involutes, the decomposition of the already always there/here in no way violating absence of precedence relations, logical, ontological, and temporal, for where is precedence in the already always there/here, regardless of what is done with it?) showed not only the necessity of transfinite process algebras (i.e., transcendental to "al-jabr proper"), but also of the multiformity of variables, variables lacking simple-identity by virtue of being m-valued, variables, that is, that are m-polyvalent (p-adic rather than monadic: untruncated p-adicity being a prerequisite to unbounded wholeness).

G_{30} : Category three of decomposition of the already always there/here by mathematical involutes, the decomposition of the already always there/here in no way violating absence of precedence relations, logical, ontological, and temporal, for where is precedence in the already always there/here, regardless of what is done with it?) showed not only the necessity of transfinite process algebras (i.e., transcendental to "al-jabr proper"), but also of the multiformity of variables, variables lacking simple-identity by virtue of being m-valued, variables, that is, that are m-polyvalent (p-adic rather than monadic: untruncated p-adicity being a prerequisite to unbounded wholeness).

T_{28} : Category one of Cyclotomic Quinary Quintic, (non-generative, non-recursive; the cyclotomic cascade as Samantabhadra

T_{29} : Category two of Cyclotomic Quinary Quintic, (non-generative, non-recursive; the cyclotomic cascade as Samantabhadra

T_{30} : Category three of Cyclotomic Quinary Quintic, (non-generative, non-recursive; the cyclotomic cascade as Samantabhadra

Module six

Thelema of Aiwass: Each and every number is infinite (Liber AL vel Legis, sub figura CCXX, or rather CTC-xk-1+……xn-k). Multi-valued variables **are not (e&eb) the same as** multi-valued logics, but the former inevitably leads to the latter, and the path between the two passes through transfinite sets.

G_{32} : Category one of Multi-valued variables; multi-valued logics, but the former inevitably leads to the latter, and the path between the two passes through transfinite sets.

G_{33} : Category two of Multi-valued variables; multi-valued logics, but the former inevitably leads to the latter, and the path between the two passes through transfinite sets.

G_{34} : Category three of Multi-valued variables; multi-valued logics, but the former inevitably leads to the latter, and the path between the two passes through transfinite sets.

T_{32} : Category one of multi-valued logics, but the former inevitably leads to the latter, and the path between the two passes through transfinite sets; Multi-valued variables

T_{33} : Category two of multi-valued logics, but the former inevitably leads to the latter, and the path between the two passes through transfinite sets.; Multi-valued variables

T_{34} : Category three of multi-valued logics, but the former inevitably leads to the latter, and the path between the two passes through transfinite sets.; Multi-valued variables

Module seven

After Abel, after Cantor, quasi-after Lukasiewicz, there was Analytical Cubism, which Jean Gebser in The Ever-Present Origin (which Targ and Hurtak argue is 4-valued) **involved (e) a** "concretion of time"

G_{36} : Category one of "concretion of time"

G_{37} : Category two of "concretion of time"

G_{38} : Category three of "concretion of time"

T_{36} : Category one of Ever-Present Origin (which Targ and Hurtak argue is 4-valued)

T_{37} : Category two of Ever-Present Origin (which Targ and Hurtak argue is 4-valued)

T_{38} : Category three of Ever-Present Origin (which Targ and Hurtak argue is 4-valued)

Module eight

Whereupon Husserl's transfinite (Platonic) horizons of the phenomenological object are **superposed (e&eb)** in just such a fashion as to constitute a Ponderable object with Piaget's property of object constancy (Picasso and Braque brought the subliminal superposition into relief on the canvas by painting signifying referents thereto).

Badiou makes repeated use of several concepts throughout his philosophy. One of the aims of his thought is to show that his categories of truth are useful for any type of philosophical critique. Therefore, he uses them to interrogate art and history as well as ontology and scientific discovery. Johannes Thumfart argues that Badiou's philosophy can be regarded as a contemporary reinterpretation of Platonism.

Conditions

According to Badiou, philosophy is suspended from four conditions (art, love, politics, and science), each of them fully independent "truth procedures." (For Badiou's notion of truth procedures, see below.) Badiou consistently maintains throughout his work (but most systematically in Manifesto for Philosophy) that philosophy must avoid the temptation to suture itself (that is, to hand over its entire intellectual effort) to any of these independent truth procedures. When philosophy does suture itself to one of its conditions (and Badiou argues that the history of philosophy during the nineteenth and twentieth centuries is primarily a history of sutures), what results is a philosophical "disaster." Consequently, philosophy is, according to Badiou, a thinking of the compossibility of the several truth procedures, whether this is undertaken through the investigation of the intersections between distinct truth procedures (the intersection of art and love in the novel, for instance), or whether this is undertaken through the more traditionally philosophical work of addressing categories like truth or the subject (concepts that are, as concepts, external to the individual truth procedures, though they are functionally operative in the truth procedures themselves). For Badiou, when

philosophy addresses the four truth procedures in a genuinely philosophical manner, rather than through a suturing abandonment of philosophy as such, it speaks of them with a theoretical terminology that marks its philosophical character: "inaesthetics" rather than art; metapolitics rather than politics; ontology rather than science; etc.

Truth, for Badiou, is a specifically philosophical category. While philosophy's several conditions are, on their own terms, "truth procedures" (i.e., they produce truths as they are pursued), it is only philosophy that can speak of the several truth procedures as truth procedures. (The lover, for instance, does not think of her love as a question of truth, but simply and rightly as a question of love. Only the philosopher sees in the true lover's love the unfolding of a truth.) Badiou has a very rigorous notion of truth, one that is strongly against the grain of much of contemporary European thought. Badiou at once embraces the traditional modernist notion that truths are genuinely invariant (always and everywhere the case, eternal and unchanging) and the incisively postmodernist notion that truths are constructed through processes. Badiou's theory of truth, exposited throughout his work, accomplishes this strange mixture by uncoupling invariance and self-evidence (such that invariance does not imply self-evidence), as well as by uncoupling constructedness from relativity (such that constructedness does not lead to relativism).

The idea, here, is that a truth's invariance makes it genuinely indiscernible: because a truth is everywhere and always the case, it passes unnoticed unless there is a rupture in the laws of being and appearance, during which the truth in question becomes, but only for a passing moment, discernible. Such a rupture is what Badiou calls an event, according to a theory originally worked out in Being and Event and fleshed out in important ways in Logics of Worlds. The individual, who chances to witness such an event, if he is faithful to what he has glimpsed, can then introduce the truth by naming it into worldly situations. For Badiou, it is by positioning oneself to the truth of an event that a human animal becomes a subject; subjectivity is not an inherent human trait. According to a process or procedure that subsequently unfolds only if those who subject themselves to the glimpsed truth continue faithful in the work of announcing the truth in question, genuine knowledge is produced (knowledge often appears in Badiou's work under the title of the "veridical"). While such knowledge **is produced in** the process of being faithful to a truth event, it should be noted that, for Badiou, knowledge, in the figure of the encyclopedia, **always remains fragile**, subject to what may yet be produced as faithful subjects of the event produce further knowledge. According to Badiou, truth procedures proceed to infinity, such that faith (fidelity) outstrips knowledge. (Badiou, following both Lacan and Heidegger, distances truth from knowledge.) The dominating ideology of the day, which Badiou terms "democratic materialism," denies the existence of truth and only recognize "bodies" and "languages." Badiou proposes a turn towards the "materialist dialectic," which recognizes that there are only bodies and languages, except there are also truths.

Inaesthetic

In Handbook of Inaesthetics Badiou both draws on the original Greek meaning and later Kantian concept of "aesthesis" as "material perception" and coins the phrase "inaesthetic" to refer to a concept of artistic creation that denies "the reflection/object relation" yet, at the same time, in reaction against the bourgeois idea of mimesis, or poetic reflection of "nature", **he affirms that** art is "immanent" and "singular". Art is immanent in the sense that its truth is given in its immediacy in a given work of art, and singular in that its truth is found in art and art alone—hence reviving the ancient materialist concept of "aesthesis". His view of the link between philosophy and art is tied into the motif of pedagogy, which he claims functions so as to "arrange the forms of knowledge in a way that some truth may come to pierce a hole in them". He develops these ideas with examples from the prose of Samuel Beckett and the poetry of Stéphane Mallarmé and Fernando Pessoa (who he argues has developed a body of work that philosophy is currently incapable of incorporating), among others.

Introduction to Being and Event

The major propositions of Badiou's philosophy all find their basis in Being and Event, in which he continues his attempt (which he began in Théorie du sujet) to reconcile a notion of the subject with ontology, and in particular post-structuralist and constructivist ontologies. A frequent criticism of post-structuralist work is that it prohibits, through its fixation on semiotics and language, any notion of a subject. Badiou's work is, by his own admission, an attempt to break out of contemporary philosophy's fixation upon language, which he sees almost as a straitjacket. This effort leads him, in Being and Event, to combine rigorous mathematical formulae with his readings of poets such as Mallarmé and Hölderlin and religious thinkers such as Pascal. His philosophy draws upon both 'analytical' and 'continental' traditions. In Badiou's own opinion, this combination places him awkwardly relative to his contemporaries, meaning that his work had been only slowly taken up. Being and Event offers an example of this slow uptake, in fact: it was translated into English only in 2005, a full seventeen years after its French publication.

As is implied in the title of the book, two elements mark the thesis of Being and Event: the place of ontology, or 'the science of being qua being' (being in itself), and the place of the event – which is seen as a rupture in being – through which the subject finds realization and reconciliation with truth. This situation of being and the rupture which characterizes the event are thought in terms of set theory, and specifically Zermelo–Fraenkel set theory (with the axiom of choice), to which Badiou accords a fundamental role in a manner quite distinct from the majority of either mathematicians or philosophers.

Mathematics as ontology (Set theory portal: Mathematics portal)

For Badiou the problem which the Greek tradition of philosophy has faced and never satisfactorily dealt with is that while beings themselves are plural, and thought in terms of multiplicity, being **itself is thought** to be singular; that is, it is thought in terms of the one. He proposes as the solution to this impasse the following declaration: that the one is not. This is why Badiou accords set theory (the axioms of which he refers to as the Ideas of the multiple) such stature, and refers to mathematics as the very place of ontology: Only set theory allows one to conceive a 'pure doctrine of the multiple'. Set theory does not operate in terms of definite individual elements in groupings but only functions insofar as what belongs to a set is of the same relation as that set (that is, another set too). What separates sets out therefore is not an existential positive proposition, but other multiples whose properties validate its presentation; which is to say their structural relation. The structure of being thus secures the regime of the count-as-one. So if one is to think of a set – for instance, the set of people, or humanity – as counting as one the elements which belong to that set, it can then secure the multiple (the multiplicities of humans) as one consistent concept (humanity), but only in terms of what does not belong to that set. What is, in following, crucial for Badiou is that the structural form of the count-as-one, which makes multiplicities thinkable, implies that the proper name of being does not belong to an element as such (an original 'one'), but rather the void set (written Ø), the set to which nothing (not even the void set itself) belongs. It may help to understand the concept 'count-as-one' if it is associated with the concept of 'terming': a multiple is not one, but it is referred to with 'multiple': one word. To count a set as one is to mention that set. How the being of terms such as 'multiple' does not contradict the non-being of the one can be understood by considering the multiple nature of terminology: for there to be a term without there also being a system of terminology, within which the difference between terms gives context and meaning to any one term, does not coincide with what is understood by 'terminology', which is precisely difference (thus multiplicity) conditioning meaning. Since the idea of a term without meaning is incoherent, the count-as-one is a structural effect or a situational operation and not an event of truth. Multiples which are 'composed' or 'consistent' are count-effects; inconsistent multiplicity is the presentation of presentation.

Badiou's use of set theory in this manner is not just illustrative or heuristic. Badiou uses the axioms of Zermelo–Fraenkel set theory to identify the relationship of being to history, Nature, the State, and God. Most

significantly this use means that (as with set theory) there is a strict prohibition on self-belonging; a set cannot contain or belong to itself. Russell's paradox famously ruled that possibility out of formal logic. So too does the axiom of foundation – or to give an alternative name the axiom of regularity – enact such a prohibition (cf. p. 190 in Being and Event) (cf. p. 190 in Being and Event). (This axiom states that all sets contain an element for which only the void [empty] set names what is common to both the set and its element.) Badiou's philosophy draws two major implications from this prohibition. Firstly, it secures the inexistence of the 'one': there cannot be a grand overarching set, and thus it is fallacious to conceive of a grand cosmos, a whole Nature, or a Being of God. Badiou is therefore – against Cantor, from whom he draws heavily – staunchly atheist. However, secondly, this prohibition prompts him to introduce the event. Because, according to Badiou, the axiom of foundation 'founds' all sets in the void, it ties all being to the historico-social situation of the multiplicities of de-centred sets – thereby effacing the positivity of subjective action, or an entirely 'new' occurrence. And whilst this is acceptable ontologically, it is unacceptable, Badiou holds, philosophically. Set theory mathematics has consequently 'pragmatically abandoned' an area which philosophy cannot. And so, Badiou argues, there is therefore only one possibility remaining: that ontology can say nothing about the event.

Roger Scruton has questioned Badiou's grasp of the foundation of mathematics. Scruton wrote in 2012, "There is no evidence that I can find in Being and Event that the author really understands what he is talking about when he invokes (as he constantly does) Georg Cantor's theory of transfinite cardinals, the axioms of set theory, Gödel's incompleteness proof or Paul Cohen's proof of the independence of the continuum hypothesis. When these things appear in Badiou's texts it is always allusively, with fragments of symbolism detached from the context that endows them with sense, and often with free variables and bound variables colliding randomly. No proof is clearly stated or examined, and the jargon of set theory is waved like a magician's wand, to give authority to bursts of all but unintelligible metaphysics".[13]

The event and the subject

The principle of the event is where Badiou diverges from the majority of late twentieth century philosophy and social thought, and in particular the likes of Foucault, Butler, Lacan and Deleuze, among others. In short, it represents that which is outside of ontology. Badiou's problem here is, unsurprisingly, the question of how to 'make use' of that which cannot be discerned. But it is a problem he views as vital, because if one constructs the world only from that which can be discerned and therefore given a name, it results in either the destitution of subjectivity and the removal of the subject from ontology (the criticism continually leveled at Foucault's discursive universe), or the Panglossian solution of Leibniz: that God **is** language in its supposed completion.

Badiou again turns here to mathematics and set theory – Badiou's language of ontology – to study the possibility of an indiscernible element existing extrinsically to the situation of ontology. He employs the strategy of the mathematician Paul J. Cohen, using what are called the conditions of sets. These conditions are thought of in terms of domination, a domination being that which defines a set. (If one takes, in binary language, the set with the condition 'items marked only with ones', any item marked with zero negates the property of the set. The condition which has only ones is thus dominated by any condition which has zeros in it [cf. p. 367-71 in Being and Event].) Badiou reasons using these conditions that every discernible (nameable or constructible) set is dominated by the conditions which don't possess the property that makes it discernible as a set. (The property 'one' is always dominated by 'not one'.) These sets are, in line with constructible ontology, relative to one's being-in-the-world and one's being in language (where sets and concepts, such as the concept 'humanity', get their names). However, he continues, the dominations themselves are, whilst being relative concepts, not necessarily intrinsic to language and constructible thought; rather one can axiomatically define a domination – in the terms of mathematical ontology – as a set

of conditions such that any condition outside the domination is dominated by at least one term inside the domination. One does not necessarily need to refer to constructible language to conceive of a 'set of dominations', which he refers to as the indiscernible set, or the generic set. It is therefore, he continues, possible to think beyond the strictures of the relativistic constructible universe of language, by a process Cohen calls forcing. And he concludes in following that while ontology can mark out a space for an inhabitant of the constructible situation to decide upon the indiscernible, it falls to the subject – about which the ontological situation cannot comment – to nominate this indiscernible, this generic point; and thus nominate, and give name to, the undecidable event. Badiou thereby marks out a philosophy by which to refute the apparent relativism or apoliticism in post-structuralist thought.

Badiou's ultimate ethical maxim is therefore one of: 'decide upon the undecidable'. It is to name the indiscernible; the generic set, and thus name the event that re-casts ontology in a new light. He identifies four domains in which a subject (who, it is important to note, becomes a subject through this process) can potentially witness an event: love, science, politics and art. By enacting fidelity to the event within these four domains one performs a 'generic procedure', which in its undecidability is necessarily experimental, and one potentially recasts the situation in which being takes place. Through this maintenance of fidelity, truth has the potentiality to emerge.

In line with his concept of the event, Badiou maintains, politics is not about politicians, but activism based on the present situation and the evental [sic] (his translators' neologism) rupture. So too does love have this characteristic of becoming anew. Even in science the guesswork that marks the event is prominent. He vigorously rejects the tag of 'decisionist' (the idea that once something is decided it 'becomes true'), but rather argues that the recasting of a truth comes prior to its veracity or verifiability. As he says of Galileo (p. 401):

When Galileo announced the principle of inertia, he was still separated from the truth of the new physics by all the chance encounters that are named in subjects such as Descartes or Newton. How could he, with the names he fabricated and displaced (because they were at hand – ‘movement’, ‘equal proportion’, etc.), have supposed the veracity of his principle for the situation to-come that was the establishment of modern science; that is, the supplementation of his situation with the indiscernible and unfinishable part that one has to name ‘rational physics’?

While Badiou is keen to reject equivalence between politics and philosophy, he correlates nonetheless his political activism and skepticism toward the parliamentary-democratic process with his philosophy, based around singular, situated truths, and potential revolutions.

G_{40} : Category one of transfinite (Platonic) horizons of the phenomenological object; in just such a fashion as to constitute a Ponderable object with Piaget's property of object constancy (Picasso and Braque brought the subliminal superposition into relief on the canvas by painting signifying referents thereto)

G_{41} : Category two of transfinite (Platonic) horizons of the phenomenological object; in just such a fashion as to constitute a Ponderable object with Piaget's property of object constancy (Picasso and Braque brought the subliminal superposition into relief on the canvas by painting signifying referents thereto).

G_{42} : Category three of transfinite (Platonic) horizons of the phenomenological object; in just such a fashion as to constitute a Ponderable object with Piaget's property of object constancy (Picasso and Braque brought the subliminal superposition into relief on the canvas by painting signifying referents thereto).

T_{40} : Category one of in just such a fashion as to constitute a Ponderable object with Piaget's property of object constancy (Picasso and Braque brought the subliminal superposition into relief on the canvas by painting signifying referents thereto); transfinite (Platonic) horizons of the phenomenological object

T_{41} : Category two of in just such a fashion as to constitute a Ponderable object with Piaget's property of object constancy (Picasso and Braque brought the subliminal superposition into relief on the canvas by painting signifying referents thereto); transfinite (Platonic) horizons of the phenomenological object

T_{42} : Category three of in just such a fashion as to constitute a Ponderable object with Piaget's property of object constancy (Picasso and Braque brought the subliminal superposition into relief on the canvas by painting signifying referents thereto); transfinite (Platonic) horizons of the phenomenological object

Module Nine

Phenomenological constitution of the discrete Ponderable macroscopic object **requires (e)** menhir, i.e., Tantric Schrödinger superposition, not a collapse of the wave-function (such a collapse would have been a matter of considerable embarrassment for Schrödinger in midst of the type of double-slit experiment he was conducting at the time he wrote his time-independent wave equation: ménage à trios).

Schrödinger's Carousel (Tantric Schrõdinger superposition): Quantum Blog (Wikipedia: emphasis mine some sections removed due to spatial constraints)

Every club has its mascot and among my tribe of physicists no mascot is more fitting than Schrödinger's Cat.

Erwin Schrödinger was a sophisticated Austrian, a reluctant participant in the Quantum Revolution sweeping through Europe in the late 1920's. Schrödinger expressed his displeasure at the new quantum world he was helping to construct by focusing attention **on its strange nesses**. He was the first to notice that quantum entanglement--an odd kind of instant voodoo action--was an inescapable consequence of his own quantum equations when applied to two or more particles of matter.

And he was particularly troubled by the **curious notion of quantum superposition**. In theory at least, an atom could not only be in state A or in state B but could exist also in a superposition state 1/2 A and 1/2 B with some "quantum phase" between them. Today this superposition principle forms the basis for quantum computers which instead of manipulating binary bits 1 and 0, compute using "qubits", quantum superpositions of binary bits.

Schrödinger tried **to demonstrate** the absurdity of the New Physics by taking the notion of quantum superposition to extremes. If one could superpose any number of atoms, as the equations allowed, then why could not one superpose a cat in two different states, say, alive and dead? Schrödinger showed, in a famous thought experiment, exactly how this might be accomplished and left us with an image of not just an atom, but a macroscopic object (covered with fur) existing at the same time in two extremely different states of being.

The quantum **rules describe the** world as waves (of possibility) when not observed and particles (of actuality) when observed. So Schrödinger's Cat maintains her twofold existence only as long as she is wavelike, that is only until she is observed, whereupon she departs her superposition state (and in a manner complete mysterious to physicists) becomes definitively either dead or alive but not both.

Schrödinger's Cat has been the subject of much theoretical discussion and a kind of informal contest among experimentalists to see how big a system they can coax into a quantum superposition. Superpositions of "big" systems have come to be called "cat states" as in the case of a recent experiment at Oxford University

in which researchers placed **13 nuclear spins in phased-linked superposition--Magnetic Field Sensors Using Large Cat States**.

Thirteen spins may seem a long way from a 10-pound pussy cat, but the recent production of Bose-Einstein Condensates has now opened the way to manipulate not just a few spins but a few million rubidium atoms into a single state and then coax that state into a quantum superposition. The mass of a few million rubidium atoms is comparable to the mass of a virus so we are still a very long way from being able to quantum-superpose cats.

A Bose-Einstein Condensate (BEC) is a dilute gas of atoms confined by electromagnetic fields and cooled to a super-low temperature (a few nanodegrees above Absolute Zero) until the atoms collapse spontaneously into the same quantum state.

A new way to turn a BEC into a cat state was recently proposed by three **theorists (Thanvanthri, Kapale and Dowling) a**t Louisiana State University. Instead of confining the BEC inside a simple potential well, TKD propose to use a "**sombrero potential**" for confinement. In a sombrero potential the particles are trapped in the brim of an electromagnetic Mexican hat to form a kind **of "moat of quantum waviness**" around the central peak of the hat.

Now that you have an unmoving "ring of bright water" made of quantum stuff, the next thing to do is to get that ring to rotate by shining light that contains angular momentum on the ring. When the BEC absorbs the spinning light, it too begins to spin, forming a **Quantized Vortex State**, a million atoms all rotating CW, say, around the central sombrero peak. You can also get the Vortex to rotate CCW by illuminating it with light spinning in the opposite direction.

Now comes the Schrödinger Cat part. Compared to getting matter to superpose, light is easy. So now you shine a superposition of CW and CCW light on the BEC and a million Rubidium atoms begin to swirl in two directions at once--**an odd state of excitation one might call "Schrödinger's Carousel".**

Make no mistake, Schrödinger's Carousel is not a simple classical situation where half of the atoms are rotating in one direction and half in the other, but a truly quantum state in which ALL OF THE ATOMS are rotating halfly CW and halfly CCW around the sombrero peak. Cool quantum cat, man.

But wait, there's more. Because this experiment is carried out on the Earth which is rotating once a day, one of the cat halves finds itself spinning slightly faster than the other which **gives rise to** a phase shift between CW and CCW vortex matter waves that can be easily measured. The Schrödinger Carousel then behaves as an ultrasensitive gyrocompass that can accurately pinpoint the direction of True North. Nice Kitty.

G_{44} : Category one of an odd state of excitation one might call "Schrödinger's Carousel".

G_{45} : Category two of an odd state of excitation one might call "Schrödinger's Carousel".

G_{46} : Category three of an odd state of excitation one might call "Schrödinger's Carousel".

T_{44} : Category one of Phenomenological constitution of the discrete Ponderable macroscopic object

T_{45} : Category two of Phenomenological constitution of the discrete Ponderable macroscopic object

T_{46} : Category three of Phenomenological constitution of the discrete Ponderable macroscopic object

SECTION THIRTY TWO

Pan-Hylism And Pan-Psychism: A Uncomfortable Menage A Trios

INTRODUCTION—VARIABLES USED

Logics of tensor strain in co-Tangent bundles over a general Manifold for Musculpt by VirFut Q-Pro μTm Scenarios, T4(M) Some paragraphs are deleted due to spatial constraints (normativity the chaotic multitude of homonymies, word-plays, "irregular" metaphoric links and resonances. in quantum computation)

(1) The first statement is legitimate in 1T2 logic: exemplified in the rule that "no A **is** not-A".
(2) The second statement, however, **violates the** 1T2-logical rule that "the case is either A or not-A"; in so doing, it is a statement "typical of theosophical thinking".
(3) In both statements a distinction is drawn, establishing two possible logical values to the given proposition; this is indicated by the 2 to the right-hand side of the T. According to the first statement, only one or the other will be true (indicated by the 1 to the left-hand side of the T), not both simultaneously: no A (inner qualities) is not-A (outward qualities). The definition of "qualia" offered in the second statement, however, doesn't fit this formal-logic mode. It **fits (eb) the mode of** Conze-Nagarjuna's third lemma (earlier discussed): "x both exists and does not exist".

(4) The distinctive property of "qualia" as given by Wilberg is resident in the fact that "x exists" (qualities of awareness) and simultaneously "does not exist" (qualities we are not aware of). The formal logic of this statement is 2T2: there are two possible mutually exclusive distinct logical values, both of which can simultaneously be **"true" (eb)** (the notion of truth-value is not actually applicable). In shift from the either/or 1T2 logic to the both/and 2T2 logic, the objectively real linear-time resident in iterations of either/or is lost to the simultaneity of both/and.

(5) What is the formal logic mode of Conze-Nagarjuna's forth lemma, "x neither exists nor does not exist"? This also is 2T2 logic, but one would have to invert the T so that it is upside down. In terms of the notation employed for the "calculus of indications" described in G. Spencer Brown's Laws of Form, the "tokens" or "marks" or "crosses" indicating a "crossing" (here denoted by T) are inverted (see The Moon of Hoa Binh, Vol. 2, pp. 334-44, for a discussion of this). The inverted cross or T denotes logical operations in numerical counterspace, a reference space of counter-hypernumbers beyond counter-quaternions. It is important to recognize that Brown's calculus is exclusively confined to 1T2 logic, but in "proving" Sheffer's postulates for Boolean algebras he simultaneously **demonstrated that** the notion of "distinction" is more fundamental than that of "truth-value", and this opened up understanding of Lukasiewicz µTm-valued logics relative to states of non-simple identity-transparency.

(6) Interpretation of Lukasiewicz logics as indicating the infinitude of states of gray or fuzziness over the real-number line between 0 and 1 -- whatever the pragmatic utility – **is(=) a** dissimulation derivative of the mistaken retention of truth-value as the foundational notion of logic.

(7) Arbitrarily cutting the line, the plane, the solid, the polytope -- whatever referent one may choose -- at two poles, 0 and 1, positive and negative, in space or counter-space -- however the representation is made -- is to illegitimately **orient (e&eb) that** which is nonorientable to the mth-degree of hypertwisting: the "logical walk" over the reference space/counter-space is uninterruptedly without 1T2-logical boundary.

(8) When, in "crossing" from 1T2 logic to 2T2 logic, objectively real linear-time and truth-value **are (e) lost** (this loss of time being quasi-equivalent to Gebser's "concretion of time"), just try to imagine all we do not know spread over transfinitely µTm-valued Lukasiewicz logics! If x is particle and not-x is wave, and the double-slit experiment is equivalent to the "cross" or the T, then we begin to see how sophomoric contemporary science is. Karl Pribram, reviewing, in the Journal of Nervous and Mental Diseases, Ignacio Matte Blanco's The Unconscious as Infinite Sets (Duckworth, 1975), said: "Its charts of terra incognita are as good as any we have as yet available." This is surely an enormous understatement, and, that being the case, we can understand something of why Gödel ran from Lukasiewicz into nervous and mental disorders late in life -- and why the human species has been collectively following Gödel.

(9) Where I have difficulty with Rudolf Steiner, Rupert Sheldrake, Peter Wilberg, the material channeled by Jane Roberts through her Seth, and a good portion of phenomenology, particularly of the descriptive variety, is the inability **to(e) overcome** incredulity that the 19th century notions of "field" and "resonance", which are not general enough to unify theories even of matter, are somehow general enough to explain both quantitative material-quanta and qualitative awareness-qualia, i.e., to unify pan-hylism and pan-psychism on some more fundamental, non-binary level -- and that this can be done, not only largely in the frame established by 1T2 logic, with just a few minor variances there from, but also in a fashion substantively communicable in speakable languages and notatable between the covers of books, even books written on a commercially viable, popular level.

NOTATION

Module One

The first statement is legitimate in 1T2 logic: exemplified in the rule that "no A **is** not-A".

G_{13} : Category one of not A

G_{14} : Category two of not A

G_{15} : Category three of not A

T_{13} : Category one of no A

T_{14} : Category two of no A

T_{15} : Category three of no A

Module Two

The second statement, however, **violates the** 1T2-logical rule that "the case is either A or not-A"; in so doing, it is a statement "typical of theosophical thinking".

G_{16} : Category one of 1T2-logical rule that "the case is either A or not-A"; in so doing, it is a statement "typical of theosophical thinking".

G_{17} : Category two of 1T2-logical rule that "the case is either A or not-A"; in so doing, it is a statement "typical of theosophical thinking".

G_{18} : Category three of 1T2-logical rule that "the case is either A or not-A"; in so doing, it is a statement "typical of theosophical thinking".

T_{16} : Category one of second statement. Many systems are being investigated and the value of x changes. Characterstics of these systems form the bastion, stylobate, sentinel **bulwark, citadel, defense, fortification** of the classification from the angle, attitude, judgment, opinion, outlook of the stratification doxa based on set axiomatic predications with locus and focus of essence sense and perception.

T_{17} : Category two of second statement

T_{18} : Category three of second statement

Module three

In both statements a distinction is drawn, establishing two possible logical values to the given proposition; this is indicated by the 2 to the right-hand side of the T. According to the first statement, only one or the other will be true (indicated by the 1 to the left-hand side of the T), not both simultaneously: no A (inner qualities) is not-A (outward qualities). The definition of "qualia" offered in the second statement, however, doesn't fit this formal-logic mode. It **fits (eb) the mode of** Conze-Nagarjuna's third lemma (earlier discussed): "x both exists and does not exist".

G_{20} : Category one of definition of "qualia" offered in the second statement

G_{21} : Category two of definition of "qualia" offered in the second statement

G_{22} : Category three of definition of "qualia" offered in the second statement

T_{20} : Category one of **mode of** Conze-Nagarjuna's third lemma (earlier discussed): "x both exists and does not exist".

T_{21} : Category two of **mode of** Conze-Nagarjuna's third lemma (earlier discussed): "x both exists and does not exist".

T_{22} : Category three of **mode of** Conze-Nagarjuna's third lemma (earlier discussed): "x both exists and does

not exist".

Module four

The distinctive property of "qualia" as given by Wilberg is resident in the fact that "x exists" (qualities of awareness) and simultaneously "does not exist" (qualities we are not aware of). The formal logic of this statement is 2T2: there are two possible mutually exclusive distinct logical values, both of which can simultaneously be **"true" (eb)** (the notion of truth-value is not actually applicable). In shift from the either/or 1T2 logic to the both/and 2T2 logic, the objectively real linear-time resident in iterations of either/or is lost to the simultaneity of both/and

G_{24} : Category one of two possible mutually exclusive distinct logical values

G_{25} : Category two of two possible mutually exclusive distinct logical values

G_{26} : Category three of two possible mutually exclusive distinct logical values

T_{24} : Category one of both of which can simultaneously be true. Note we are dealing with different investigative systems with different propensities and properties.

T_{25} : Category two of both of which can simultaneously be true

T_{26} : Category three of both of which can simultaneously be true

Module five

What is the formal logic mode of Conze-Nagarjuna's forth lemma, "x neither exists nor does not exist"? This also is 2T2 logic, but one would have to invert the T so that it is upside down. In terms of the notation employed for the "calculus of indications" described in G. Spencer Brown's Laws of Form, the "tokens" or "marks" or "crosses" indicating a "crossing" (here denoted by T) are inverted (see The Moon of Hoa Binh, Vol. 2, pp. 334-44, for a discussion of this). The inverted cross or T **denotes** logical operations in numerical counterspace, a reference space of counter-hypernumbers beyond counter-quaternions. It is important to recognize that Brown's calculus is exclusively confined to 1T2 logic, but in "proving" Sheffer's postulates for Boolean algebras he simultaneously **demonstrated that** the notion of "distinction" is more fundamental than that of "truth-value", and this opened up understanding of Lukasiewicz µTm-valued logics relative to states of non-simple identity-transparency

G_{28} : Category one of 2T2 logic; Brown's calculus is exclusively confined to 1T2 logic, but in "proving" Sheffer's postulates for Boolean algebras he simultaneously;

G_{29} : Category two of 2T2 logic; Brown's calculus is exclusively confined to 1T2 logic, but in "proving" Sheffer's postulates for Boolean algebras he simultaneously;

G_{30} : Category three of2T2 logic; Brown's calculus is exclusively confined to 1T2 logic, but in "proving" Sheffer's postulates for Boolean algebras he simultaneously;

T_{28} : Category one of "x neither exists nor does not exist"; notion of "distinction" is more fundamental than that of "truth-value", and this opened up understanding of Lukasiewicz µTm-valued logics relative to states of non-simple identity-transparency

T_{29} : Category two of "x neither exists nor does not exist"; notion of "distinction" is more fundamental than that of "truth-value", and this opened up understanding of Lukasiewicz µTm-valued logics relative to states

of non-simple identity-transparency

T_{30} : Category three of "x neither exists nor does not exist"; notion of "distinction" is more fundamental than that of "truth-value", and this opened up understanding of Lukasiewicz μTm-valued logics relative to states of non-simple identity-transparency

Module six

Interpretation of Lukasiewicz logics as indicating the infinitude of states of gray or fuzziness over the real-number line between 0 and 1 -- whatever the pragmatic utility – **is(=) a** dissimulation derivative of the mistaken retention of truth-value as the foundational notion of logic.

G_{32} : Category one of dissimulation derivative of the mistaken retention of truth-value as the foundational notion of logic.

G_{33} : Category two of dissimulation derivative of the mistaken retention of truth-value as the foundational notion of logic.

G_{34} : Category three of dissimulation derivative of the mistaken retention of truth-value as the foundational notion of logic.

T_{32} : Category one of Interpretation of Lukasiewicz logics as indicating the infinitude of states of gray or fuzziness over the real-number line between 0 and 1 -- whatever the pragmatic utility

T_{33} : Category two of Interpretation of Lukasiewicz logics as indicating the infinitude of states of gray or fuzziness over the real-number line between 0 and 1 -- whatever the pragmatic utility

T_{34} : Category three of Interpretation of Lukasiewicz logics as indicating the infinitude of states of gray or fuzziness over the real-number line between 0 and 1 -- whatever the pragmatic utility

Module seven(Both cases are discussed in a single module)

arbitrarily cutting the line, the plane, the solid, the polytope -- whatever referent one may choose -- at two poles, 0 and 1, positive and negative, in space or counter-space -- however the representation is made -- is to illegitimately **orient (e&eb) that** line which is nonorientable to the mth-degree of hypertwisting: the "logical walk" over the reference space/counter-space is uninterruptedly without 1T2-logical boundary.

G_{36} : Category one of arbitrarily cutting the line, the plane, the solid, the polytope -- whatever referent one may choose -- at two poles, 0 and 1, positive and negative, in space or counter-space -- however the representation is made -- is to illegitimately; line which is nonorientable to the mth-degree of hypertwisting: the "logical walk" over the reference space/counter-space is uninterruptedly without 1T2-logical boundary

G_{37} : Category two of arbitrarily cutting the line, the plane, the solid, the polytope -- whatever referent one may choose -- at two poles, 0 and 1, positive and negative, in space or counter-space -- however the representation is made -- is to illegitimately; line which is nonorientable to the mth-degree of hypertwisting: the "logical walk" over the reference space/counter-space is uninterruptedly without 1T2-logical boundary

G_{38} : Category three of arbitrarily cutting the line, the plane, the solid, the polytope -- whatever referent one may choose -- at two poles, 0 and 1, positive and negative, in space or counter-space -- however the representation is made -- is to illegitimately; line which is nonorientable to the mth-degree of hypertwisting: the "logical walk" over the reference space/counter-space is uninterruptedly without 1T2-logical boundary

T_{36} : Category one of line which is nonorientable to the mth-degree of hypertwisting: the "logical walk" over the reference space/counter-space is uninterruptedly without 1T2-logical boundary; arbitrarily cutting the line, the plane, the solid, the polytope -- whatever referent one may choose -- at two poles, 0 and 1, positive and negative, in space or counter-space -- however the representation is made -- is to illegitimately

T_{37} : Category two of line which is nonorientable to the mth-degree of hypertwisting: the "logical walk" over the reference space/counter-space is uninterruptedly without 1T2-logical boundary; arbitrarily cutting the line, the plane, the solid, the polytope -- whatever referent one may choose -- at two poles, 0 and 1, positive and negative, in space or counter-space -- however the representation is made -- is to illegitimately

T_{38} : Category three of line which is nonorientable to the mth-degree of hypertwisting: the "logical walk" over the reference space/counter-space is uninterruptedly without 1T2-logical boundary; arbitrarily cutting the line, the plane, the solid, the polytope -- whatever referent one may choose -- at two poles, 0 and 1, positive and negative, in space or counter-space -- however the representation is made -- is to illegitimately

Module eight

When, in "crossing" from 1T2 logic to 2T2 logic, objectively real linear-time and truth-value **are (e) lost** (this loss of time being quasi-equivalent to Gebser's "concretion of time"), just try to imagine all we do not know spread over transfinitely µTm-valued Lukasiewicz logics! If x is particle and not-x is wave, and the double-slit experiment **is equivalent to** the "cross" or the T, then we begin to see how sophomoric contemporary science is. Karl Pribram, reviewing, in the Journal of Nervous and Mental Diseases, Ignacio Matte Blanco's The Unconscious as Infinite Sets (Duckworth, 1975), said: "Its charts of terra incognita are as good as any we have as yet available." This is surely an enormous understatement, and, that being the case, we can understand something of why Gödel ran from Lukasiewicz into nervous and mental disorders late in life -- and why the human species has been collectively following Gödel

G_{40} : Category one of objectively real linear-time and truth-value(this loss of time being quasi-equivalent to Gebser's "concretion of time"),

G_{41} : Category two of objectively real linear-time and truth-value(this loss of time being quasi-equivalent to Gebser's "concretion of time"),

G_{42} : Category three of objectively real linear-time and truth-value(this loss of time being quasi-equivalent to Gebser's "concretion of time"),

T_{40} : Category one of "crossing" from 1T2 logic to 2T2 logic

T_{41} : Category two of "crossing" from 1T2 logic to 2T2 logic

T_{42} : Category three of "crossing" from 1T2 logic to 2T2 logic

Module Nine

Where I have difficulty with Rudolf Steiner, Rupert Sheldrake, Peter Wilberg, the material channeled by Jane Roberts through her Seth, and a good portion of phenomenology, particularly of the descriptive variety, is the inability **to(e) overcome** incredulity that the 19th century notions of "field" and "resonance", which are not general enough to unify theories even of matter, **are somehow general enough to explain** both quantitative material-quanta and qualitative awareness-qualia, i.e., to unify pan-hylism and pan-psychism on some more fundamental, non-binary level -- and that this can be done, not only largely in the frame established by 1T2 logic, with just a few minor variances there from, but also in a fashion substantively

communicable in speakable languages and notatable between the covers of books, even books written on a commercially viable, popular level.

G_{44} : Category one of phenomenology, particularly of the descriptive variety

G_{45} : Category two of phenomenology, particularly of the descriptive variety

G_{46} : Category three of phenomenology, particularly of the descriptive variety

T_{44} : Category one of quantitative material-quanta and qualitative awareness-qualia, i.e., to unify pan-hylism and pan-psychism on some more fundamental, non-binary level -- and that this can be done, not only largely in the frame established by 1T2 logic, with just a few minor variances there from, but also in a fashion substantively communicable in speakable languages and notatable between the covers of books, even books written on a commercially viable, popular level.

T_{45} : Category two of quantitative material-quanta and qualitative awareness-qualia, i.e., to unify pan-hylism and pan-psychism on some more fundamental, non-binary level -- and that this can be done, not only largely in the frame established by 1T2 logic, with just a few minor variances there from, but also in a fashion substantively communicable in speakable languages and notatable between the covers of books, even books written on a commercially viable, popular level.

T_{46} : Category three of quantitative material-quanta and qualitative awareness-qualia, i.e., to unify pan-hylism and pan-psychism on some more fundamental, non-binary level -- and that this can be done, not only largely in the frame established by 1T2 logic, with just a few minor variances there from, but also in a fashion substantively communicable in speakable languages and notatable between the covers of books, even books written on a commercially viable, popular level.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$
$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$
$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**
$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)}\ \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{array}\right] G_{18} \quad 63$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for*

category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category $1, 2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & \boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} & \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} & \end{bmatrix} G_{30} \quad 81$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & \boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} & \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} & \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & \boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} & \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} & \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & \boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} & \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} & \end{bmatrix} T_{30} \quad 84$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ *are second detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for

category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a_{32}')^{(6)} & +(a_{32}'')^{(6)}(T_{33},t) & +(a_{28}'')^{(5,5,5)}(T_{29},t) & +(a_{24}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a_{33}')^{(6)} & +(a_{33}'')^{(6)}(T_{33},t) & +(a_{29}'')^{(5,5,5)}(T_{29},t) & +(a_{25}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a_{34}')^{(6)} & +(a_{34}'')^{(6)}(T_{33},t) & +(a_{30}'')^{(5,5,5)}(T_{29},t) & +(a_{26}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a_{32}'')^{(6)}(T_{33},t)$, $+(a_{33}'')^{(6)}(T_{33},t)$, $+(a_{34}'')^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a_{28}'')^{(5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a_{24}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) & \\ -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) & \\ -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) & \\ -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{34} \quad 90$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37},t) & +(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{c} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{c} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T_{21}'| e^{-(\hat{M}_{20})^{(3)} t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} ||G_{23} - G_{23}'|| e^{-(\hat{M}_{20})^{(3)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(G) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(H) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(I) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(J)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(K) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(L) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$
satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\leq\ \left(\hat{P}_{13}\right)^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\leq\ (\hat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\leq\ (\hat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\leq\ (\hat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 156

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(b) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)} t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

208

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)}$$

209

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)}$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

214

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40{,}41{,}42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(8)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0, (\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0, (\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}\left((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)}\right)}\left[e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}\left((S_1)^{(3)} - (a'_{22})^{(3)}\right)}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}\left((R_1)^{(3)} - (b'_{22})^{(3)}\right)}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}\left((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)}\right)}\left[e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right) - (b''_{25})^{(4)}\left((G_{27}), t\right) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(\nu_1)^{(6)}, (\nu_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, \nu^{(6)}, u^{(6)}$: 350

By $(\nu_1)^{(6)} > 0$, $(\nu_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)},, (\bar{\nu}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, $\boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad (354)$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad (355)$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad (356)$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^{0}e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^{0}e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^{0}}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t}-e^{-(b'_{34})^{(6)}t}\right]+T_{34}^{0}e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^{0}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T_{34}^{0}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)}+(C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(b) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^{*} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^{*} for which $f(T_{33}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

Definition and uniqueness of T_{37}^{*} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^{*} for which $f(T_{37}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523

A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTY THREE

Consciousness And Quantum Computing: Veritas Vos Liberabit Model

INTRODUCTION—VARIABLES USED

Consciousness and Logic in a Quantum-Computing Universe Paola Zizzi

(1) The early inflationary universe **can be described (e&eb) in terms** of quantum information.
(2) More specifically, the inflationary universe **can be (=) viewed** as a superposed state of quantum registers.
(3) Actually, during inflation, **one (eb) can speak of** a quantum superposition of universes.
(4) At the end of inflation, only one universe is selected, **by (e) a mechanism** called self-reduction, which is consistent with Penrose's objective reduction (OR) model.
(5) The quantum gravity threshold of (OR) is reached at the end of inflation, **and corresponds** to a superposed state of 109 quantum registers.
(6) This is also the number of superposed tubulins — qubits in our brain, which undergo the Penrose-Hameroff orchestrated objective reduction, (Orch OR), **leading to a** conscious event.
(7) Then, an analogy naturally arises between the very early quantum-computing universe**, and** our mind.
(8) In fact, we argue that at the end of in- flation, the universe underwent a cosmic conscious event, the so-called "Big Wow", which acted as an **imprinting (e&eb) for** the future minds to come, with future modes of computation, consciousness and logic.
(9) The post inflationary universe organized itself as a cellular automaton (CA) with two computational modes: quantum and classical, like the two conformations assumed by the cellular automaton of tubulins in our brain, as in Hameroff's model. In the quantum configuration, the universe quantum-evaluates recursive functions, **which are** the laws of physics in their most abstract form.

NOTATION

Module One

The early inflationary universe **can be described (e&eb) in terms** of quantum information

G_{13} : Category one of early inflationary universe

G_{14} : Category two of early inflationary universe

G_{15} : Category three of early inflationary universe

T_{13} : Category one of quantum information

T_{14} : Category two of quantum information

T_{15} : Category three of quantum information

Module Two

More specifically, the inflationary universe **can be (=) viewed** as a superposed state of quantum registers.

G_{16} : Category one of superposed state of quantum registers

G_{17} : Category two of superposed state of quantum registers

G_{18} : Category three of superposed state of quantum registers

T_{16} : Category one of inflationary universe

T_{17} : Category two of inflationary universe

T_{18} : Category three of inflationary universe

Module three

Conservation Law can be seem as tallied quantum registers from everything in the universe(individual general ledger becomes nature's general ledger: Like general ledgers from all branches are tallied at the central office: my statement)

G_{20} : Category one of tallied general ledger at the Central Office of various branches'

G_{21} : Category two of tallied general ledger at the Central Office of various branches'

G_{22} : Category three of tallied general ledger at the Central Office of various branches'

T_{20} : Category one of conservation laws

T_{21} : Category two of conservation laws

T_{22} : Category three of conservation laws

Module four

Actually, during inflation, **one (eb) can speak of** a quantum superposition of universes

G_{24} : Category one of during inflation (assume various parts of the universe)

G_{25} : Category two of during inflation (assume various parts of the universe)

G_{26} : Category three of during inflation (assume various parts of the universe)

T_{24} : Category one of quantum superposition of universes

T_{25} : Category two of quantum superposition of universes

T_{26} : Category three of quantum superposition of universes

Module five

At the end of inflation, only one universe is selected, **by (e) a mechanism** called self-reduction, which is consistent with Penrose's objective reduction (OR) model

G_{28} : Category one of **mechanism** called self-reduction, which is consistent with Penrose's objective reduction (OR) model

G_{29} : Category two of **mechanism** called self-reduction, which is consistent with Penrose's objective reduction (OR) model

G_{30} : Category three of **mechanism** called self-reduction, which is consistent with Penrose's objective reduction (OR) model

T_{28} : Category one of one universe is selected at the end of the inflation

T_{29} : Category two of one universe is selected at the end of the inflation

T_{30} : Category three of one universe is selected at the end of the inflation

Module six

The quantum gravity threshold of (OR) is reached at the end of inflation, **and corresponds** to a superposed state of 109 quantum registers

G_{32} : Category one of quantum gravity threshold of (OR) is reached at the end of inflation; superposed state of 109 quantum registers

G_{33} : Category two of quantum gravity threshold of (OR) is reached at the end of inflation; superposed state of 109 quantum registers

G_{34} : Category three of quantum gravity threshold of (OR) is reached at the end of inflation; superposed state of 109 quantum registers

T_{32} : Category one of superposed state of 109 quantum registers ;quantum gravity threshold of (OR) is reached at the end of inflation

T_{33} : Category two of superposed state of 109 quantum registers; quantum gravity threshold of (OR) is reached at the end of inflation

T_{34} : Category three of superposed state of 109 quantum registers ;quantum gravity threshold of (OR) is reached at the end of inflation

Module seven

This is also the number of superposed tubulins — qubits in our brain, which undergo the Penrose-Hameroff orchestrated objective reduction, (Orch OR), **leading to a** conscious event

Then, an analogy naturally arises between the very early quantum-computing universe**, and** our mind.

G_{36} : Category one of number of superposed tubulins — qubits in our brain, which undergo the Penrose-Hameroff orchestrated objective reduction, (Orch OR); mind.

G_{37} : Category two of number of superposed tubulins — qubits in our brain, which undergo the Penrose-Hameroff, orchestrated objective reduction, (Orch OR); mind.

G_{38} : Category three of number of superposed tubulins — qubits in our brain, which undergo the Penrose-Hameroff, orchestrated objective reduction, (Orch OR); mind.

T_{36} : Category one of conscious event; very early quantum-computing universe

T_{37} : Category two of conscious event; very early quantum-computing universe

T_{38} : Category three of conscious event; very early quantum-computing universe

Module eight

In fact, we argue that at the end of in- flation, the universe underwent a cosmic conscious event, the so-called "Big Wow", which acted as an **imprinting (e&eb) for** the future minds to come, with future modes of computation, consciousness and logic.

G_{40} : Category one of universe underwent a cosmic conscious event, the so-called "Big Wow", which acted as an **imprinting;** future minds to come, with future modes of computation, consciousness and logic.

G_{41} : Category two of universe underwent a cosmic conscious event, the so-called "Big Wow", which acted as an **imprinting;** future minds to come, with future modes of computation, consciousness and logic.

G_{42} : Category three of universe underwent a cosmic conscious event, the so-called "Big Wow", which acted as an **imprinting;** future minds to come, with future modes of computation, consciousness and logic.

T_{40} : Category one of future minds to come, with future modes of computation, consciousness and logic; universe underwent a cosmic conscious event, the so-called "Big Wow", which acted as an **imprinting**

T_{41} : Category two of future minds to come, with future modes of computation, consciousness and logic; universe underwent a cosmic conscious event, the so-called "Big Wow", which acted as an **imprinting**

T_{42} : Category three of future minds to come, with future modes of computation, consciousness and logic; universe underwent a cosmic conscious event, the so-called "Big Wow", which acted as an **imprinting**

Module Nine

The post inflationary universe organized itself as a cellular automaton (CA) with two computational modes: quantum and classical, like the two conformations assumed by the cellular automaton of tubulins in our brain, as in Hameroff's model. In the quantum configuration, the universe quantum-evaluates recursive functions, **which are** the laws of physics in their most abstract form

G_{44} : Category one of laws of physics in their most abstract form

G_{45} : Category two of laws of physics in their most abstract form

G_{46} : Category three of laws of physics in their most abstract form

T_{44} : Category one of post inflationary universe organized itself as a cellular automaton (CA) with two computational modes: quantum and classical, like the two conformations assumed by the cellular automaton of tubulins in our brain, as in Hameroff's model. In the quantum configuration, the universe quantum-evaluates recursive functions,

T_{45} : Category two of post inflationary universe organized itself as a cellular automaton (CA) with two computational modes: quantum and classical, like the two conformations assumed by the cellular automaton of tubulins in our brain, as in Hameroff's model. In the quantum configuration, the universe quantum-evaluates recursive functions,

T_{46} : Category three of post inflationary universe organized itself as a cellular automaton (CA) with two computational modes: quantum and classical, like the two conformations assumed by the cellular automaton of tubulins in our brain, as in Hameroff's model. In the quantum configuration, the universe quantum-evaluates recursive functions,

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$

$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$
$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,

2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad 73$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad 74$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad 76$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)} & \boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ are first augmentation coefficients for category 1, 2 and 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32} \quad 85$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b_{36}')^{(7)}\boxed{-(b_{36}'')^{(7)}(G_{39},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b_{37}')^{(7)}\boxed{-(b_{37}'')^{(7)}(G_{39},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b_{38}')^{(7)}\boxed{-(b_{38}'')^{(7)}(G_{39},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)}$ 103

$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)}$ 104

$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)}$ 105

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t}$ 107

$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t}$ 108

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1$ 110

$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1$ 111

Where we suppose

$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22$ 112

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with (116)
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \quad (117)$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(M) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(N) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$

132

(O) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(P)

$\lim_{G\to\infty} (b_i'')^{(7)}((G_{39}),t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}((G_{39}),t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(Q) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(R) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad 146\ A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{20})^{(3)}, T_i^0 \le (\hat{Q}_{20})^{(3)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad (169)$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that (170)

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that (172)

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that (174)

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(c) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $\left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$sup_i\{max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}ds_{(16)} +$

$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_{0}^{t}(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_{0}^{t}\{(a_{20}')^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a_{20}'')^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a_{20}'')^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a_{20}'')^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a_{20}')^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a_{20}'')^{(3)}$ and $(b_{20}'')^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_{+}$. 215

If instead of proving the existence of the solution on $\mathbb{R}_{+}$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a_{22}')^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ (228)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$
$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}\}$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t} \; and \; (\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2 \; and \; \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37} \; and \; G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)},, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \qquad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \qquad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \qquad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

406

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case,

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

407

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(c) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)} G_{14}^{*}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)} G_{14}^{*}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)} T_{14}^{*}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)} T_{14}^{*}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)} G_{17}^{*}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)} G_{17}^{*}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)} T_{17}^{*}}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)} T_{17}^{*}}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)} G_{21}^{*}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)} G_{21}^{*}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)} T_{21}^{*}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)} T_{21}^{*}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)} T_{45}^{*}}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)} T_{45}^{*}}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTY FOUR

Pastry Chef And Contemporary Computer Science: Sacrosuchus Imperator-Bonvivavt Antelope Model

INTRODUCTION—VARIABLES USED

Consciousness and Logic in a Quantum-Computing Universe Paola Zizzi

(1) To do so (universe quantum-evaluates recursive functions, **which are** the laws of physics in their most abstract form) in a very efficient way, the universe **uses(e) as** subroutines, black holes - quantum computers and quantum minds, which operate in parallel
(2) The outcomes of the overall quantum computation **are the** universals, the attributes of things in themselves.
(3) These universals are partially **obtained(e) also by** the quantum minds,
(4) Universals **endowed (e)** subjective meaning.
(5) The units of the subjective universals **are** qualia, which are strictly related to the (virtual) existence of Planckian black holes.
(6) Further, authors consider two aspects of the quantum mind, which are not algorithmic in the usual sense: the self **and** mathematical intuition.
(7) The self **is (e) due to a** reversible self-measurement of a quantum state of superposed tubulins.
(8) Mathematical intuition **is due to the** paraconsistent logic of the internal observer in a quantum-computing universe.

Minds and Machines May 2002, Volume 12, Issue 2, Toward a Formal Philosophy of Hypercomputation Selmer Bringsjord, Michael Zenzen

(9) Does what guides a pastry chef stand on par, from the standpoint of contemporary computer science, with what guides a supercomputer? Did Betty Crocker, when telling us how to bake a cake, provide an effective procedure, in the sense of `effective' used in computer science? According to Cleland, the answer in both cases is ``Yes''. One consequence of Cleland's affirmative answer is supposed to be that hypercomputation is, to use her phrase, ``theoretically viable''. Unfortunately, though we applaud Cleland's ``gadfly philosophizing'' (as, in fact, seminal), we believe that unless such a modus operandi **is (e&eb) married to** formal philosophy, nothing conclusive will be produced (as evidenced by the problems plaguing Cleland's work that we uncover). Herein, we attempt to pull off not the complete marriage for hypercomputation, but perhaps at least the

beginning of a courtship that others can subsequently help along.

NOTATION

Module One

To do so (universe quantum-evaluates recursive functions, **which are** the laws of physics in their most abstract form) in a very efficient way, the universe **uses(e) as** subroutines, black holes - quantum computers and quantum minds, which operate in parallel

G_{13} : Category one of subroutines, black holes - quantum computers and quantum minds, which operate in parallel

G_{14} : Category two of subroutines, black holes - quantum computers and quantum minds, which operate in parallel

G_{15} : Category three of subroutines, black holes - quantum computers and quantum minds, which operate in parallel

T_{13} : Category one of universe quantum-evaluates recursive functions, **which are** the laws of physics in their most abstract form in a very efficient way, the universe

T_{14} : Category two of universe quantum-evaluates recursive functions, **which are** the laws of physics in their most abstract form in a very efficient way, the universe

T_{15} : Category three of universe quantum-evaluates recursive functions, **which are** the laws of physics in their most abstract form in a very efficient way, the universe

Module Two

The outcomes of the overall quantum computation **are the** universals, the attributes of things in themselves.

G_{16} : Category one of universals, the attributes of things in themselves

G_{17} : Category two of universals, the attributes of things in themselves

G_{18} : Category three of universals, the attributes of things in themselves

T_{16} : Category one of outcomes of the overall quantum computation

T_{17} : Category two of outcomes of the overall quantum computation

T_{18} : Category three of outcomes of the overall quantum computation

Module three

These universals are partially **obtained(e) also by** the quantum minds

Universals **endowed (e)** subjective meaning

G_{20} : Category one of quantum minds; subjective meaning

G_{21} : Category two of quantum minds; subjective meaning

G_{22} : Category three of quantum minds ; subjective meaning

T_{20} : Category one of universals are partially; Universals

T_{21} : Category two of universals are partially; Universals

T_{22} : Category three of universals are partially; Universals

Module four

The units of the subjective universals **are** qualia, which are strictly related to the (virtual) existence of Planckian black holes

G_{24} : Category one of qualia, which are strictly related to the (virtual) existence of Planckian black holes

G_{25} : Category two of qualia, which are strictly related to the (virtual) existence of Planckian black holes

G_{26} : Category three of qualia, which are strictly related to the (virtual) existence of Planckian black holes

T_{24} : Category one of units of the subjective universals

T_{25} : Category two of units of the subjective universals

T_{26} : Category three of units of the subjective universals

Module five

Further, authors consider two aspects of the quantum mind, which are not algorithmic in the usual sense: the self **and** mathematical intuition.

G_{28} : Category one of self

G_{29} : Category two of self

G_{30} : Category three of self

T_{28} : Category one of mathematical intuition.

T_{29} : Category two of mathematical intuition.

T_{30} : Category three of mathematical intuition.

Module six

The self **is (e) due to a** reversible self-measurement of a quantum state of superposed tubulins

G_{32} : Category one of reversible self-measurement of a quantum state of superposed tubulins

G_{33} : Category two of reversible self-measurement of a quantum state of superposed tubulins

G_{34} : Category three of reversible self-measurement of a quantum state of superposed tubulins

T_{32} : Category one of self

T_{33} : Category two of self

T_{34} : Category three of self

Module seven

Mathematical intuition **is due to the** paraconsistent logic of the internal observer in a quantum-computing universe.

G_{36} : Category one of paraconsistent logic of the internal observer in a quantum-computing universe.

G_{37} : Category two of paraconsistent logic of the internal observer in a quantum-computing universe.

G_{38} : Category three of paraconsistent logic of the internal observer in a quantum-computing universe.

T_{36} : Category one of Mathematical intuition

T_{37} : Category two of Mathematical intuition

T_{38} : Category three of Mathematical intuition

Module eight

Does what guides a pastry chef stand on par, from the standpoint of contemporary computer science, with what guides a supercomputer? Did Betty Crocker, when telling us how to bake a cake, provide an effective procedure, in the sense of `effective' used in computer science? According to Cleland, the answer in both cases is ``Yes". One consequence of Cleland's affirmative answer is supposed to be that hypercomputation is, to use her phrase, ``theoretically viable". Unfortunately, though we applaud Cleland's ``gadfly philosophizing" (as, in fact, seminal), authors believe that unless such a modus operandi **is (e&eb) married to** formal philosophy, nothing conclusive will be produced (as evidenced by the problems plaguing Cleland's work that we uncover). Herein, we attempt to pull off not the complete marriage for hypercomputation, but perhaps at least the beginning of a courtship that others can subsequently help along.

G_{40} : Category one of hypercomputation

G_{41} : Category two of hypercomputation

G_{42} : Category three of hypercomputation

T_{40} : Category one of ``theoretically viable"

T_{41} : Category two of ``theoretically viable"

T_{42} : Category three of ``theoretically viable"

Module Nine(Both cases discussed)

G_{44} : Category one of hypercomputation; formal philosophy

G_{45} : Category two of hypercomputation; formal philosophy

G_{46} : Category three of hypercomputation; formal philosophy

T_{44} : Category one of formal philosophy; hypercomputation

T_{45} : Category two of formal philosophy ;hypercomputation

T_{46} : Category three of formal philosophy ;hypercomputation

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \qquad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**
$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**
Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$
$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$
$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$
$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$
$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$
$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**
$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**
Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**
Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**
Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition

coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{20})^{(3)}\left[+(a''_{20})^{(3)}(T_{21},t)\right]\left[+(a''_{16})^{(2,2,2)}(T_{17},t)\right]\left[+(a''_{13})^{(1,1,1,)}(T_{14},t)\right] \\ \left[+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)\right]\left[+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)\right]\left[+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{36})^{(7,7,7,7)}(T_{37},t)\right]\left[+(a''_{40})^{(8,8,8,8)}(T_{41},t)\right]\left[+(a''_{44})^{(9,9,9)}(T_{45},t)\right] \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c} (a'_{21})^{(3)}\left[+(a''_{21})^{(3)}(T_{21},t)\right]\left[+(a''_{17})^{(2,2,2)}(T_{17},t)\right]\left[+(a''_{14})^{(1,1,1,)}(T_{14},t)\right] \\ \left[+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)\right]\left[+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)\right]\left[+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{37})^{(7,7,7,7)}(T_{37},t)\right]\left[+(a''_{41})^{(8,8,8,8)}(T_{41},t)\right]\left[+(a''_{45})^{(9,9,9)}(T_{45},t)\right] \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{22})^{(3)}\left[+(a''_{22})^{(3)}(T_{21},t)\right]\left[+(a''_{18})^{(2,2,2)}(T_{17},t)\right]\left[+(a''_{15})^{(1,1,1,)}(T_{14},t)\right] \\ \left[+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)\right]\left[+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)\right]\left[+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{38})^{(7,7,7,7)}(T_{37},t)\right]\left[+(a''_{42})^{(8,8,8,8)}(T_{41},t)\right]\left[+(a''_{46})^{(9,9,9)}(T_{45},t)\right] \end{array}\right] G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for

category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad 78$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

are fourth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{16}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{17}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{18}'')^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{20}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{21}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{22}'')^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{46}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{45}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{44}'')^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{28}')^{(5)}\boxed{+(a_{28}'')^{(5)}(T_{29},t)} & \boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)} & \boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a_{29}')^{(5)}\boxed{+(a_{29}'')^{(5)}(T_{29},t)} & \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)} & \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{30}')^{(5)}\boxed{+(a_{30}'')^{(5)}(T_{29},t)} & \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)} & \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad 81$$

Where $\boxed{+(a_{28}'')^{(5)}(T_{29},t)}, \boxed{+(a_{29}'')^{(5)}(T_{29},t)}, \boxed{+(a_{30}'')^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3
And $\boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3
$\boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3
$\boxed{+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3
$\boxed{+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3
$\boxed{+(a_{40}'')^{(8,8\,,8,8,8,8)}(T_{41},t)}, \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3
$\boxed{+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} & +(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) \quad +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{array}\right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) \quad -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{array}\right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) \quad -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{array}\right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) \quad -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{array}\right] T_{34} \quad 90$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)} & +(a_{36}'')^{(7)}(T_{37},t) & +(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)} & +(a_{37}'')^{(7)}(T_{37},t) & +(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)} & +(a_{38}'')^{(7)}(T_{37},t) & +(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a_{36}'')^{(7)}(T_{37},t)$, $(a_{37}'')^{(7)}(T_{37},t)$, $(a_{38}'')^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\,-(b''_{36})^{(7)}(G_{39},t) & -(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\,-(b''_{37})^{(7)}(G_{39},t) & -(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\,-(b''_{38})^{(7)}(G_{39},t) & -(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)} G_{44}$$
$$- \left[\begin{array}{ccc} (a'_{45})^{(9)} & +(a''_{45})^{(9)}(T_{45},t) & +(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t) \quad +(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)} G_{45}$$
$$- \left[\begin{array}{ccc} (a'_{46})^{(9)} & +(a''_{46})^{(9)}(T_{37},t) & +(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t) \quad +(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t) \end{array} \right] G_{15}$$

Where $+(a''_{44})^{(9)}(T_{45},t)$, $+(a''_{45})^{(9)}(T_{45},t)$, $+(a''_{46})^{(9)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)} T_{45} -$$
$$\left[\begin{array}{ccc} (b'_{44})^{(9)} & -(b''_{44})^{(9)}(G_{47},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t) \quad -(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t) \end{array} \right] T_{13}$$
$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ \ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(S) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(T) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(U) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(V)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(W) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(X) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^{0} + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^{0} + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^{0} + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad (169)$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that (170)

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that (172)

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that (174)

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(d) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+\ (a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right) d\left(\left((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\tilde{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\,((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \le \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \le \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \ge (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \ge \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \ge \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$
Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},\ \boldsymbol{if}\ (v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t)\right) - (a_{41}'')^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(d) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)} T_{45}^{*}}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)} T_{45}^{*}}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}$, $\frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTY FIVE

Organizational Invariance And Conscious Experience Aeì Koloiòs Parà Koloiôi Hizánei Model

INTRODUCTION—VARIABLES USED

Absent Qualia, Fading Qualia, Dancing Qualia David J. Chalmers

The principle of organizational invariance

(1) It is widely accepted that conscious experience has a physical basis. That is, the properties of experience (phenomenal properties, or qualia) systematically **depend on** physical properties according to some lawful relation.

(2) There are two key questions about this relation. The first concerns the strength of the laws: are they logically or metaphysically necessary, or are they merely nomologically necessary? This question about the strength of the psychophysical link **is the basis for** debates over physicalism and property dualism.

(3) The second question concerns the nature of the laws: precisely how do phenomenal properties **depend on** physical properties? What sort of physical properties **enter into** the laws' antecedents, for instance; consequently, what sort of physical systems **can give rise** to conscious experience? It is this second question that I will address in this paper. To put the issue differently, even once it is accepted that experience **arises from** physical systems, the question remains open: in virtue of what sort of physical properties does conscious experience arise? Some property that brains can possess, presumably, but it is far from obvious what sort of property this is. Some have suggested biochemical properties; some have suggested quantum-mechanical properties; many have professed uncertainty. A natural suggestion is that when experience arises from a physical system, **it does (E) so in virtue** of the system's functional organization. On this view, the chemical and indeed the quantum substrate of the brain is not directly relevant to the existence of consciousness. What counts is rather the brain's abstract causal organization, an organization that might be realized in many different physical substrates.

(4) In this paper I defend this view. Specifically, I defend a principle of organizational invariance, holding that experience **is i**nvariant across systems with the same fine-grained functional organization. More precisely, the principle states that given any system that has conscious experiences, **then any system** that has the same functional organization at a fine enough grain will have qualitatively identical conscious experiences. A full specification of a system's fine-grained functional organization **will fully determine** any conscious experiences that arise.

(5) To clarify this, we must first clarify the notion of functional organization. This is best understood as the abstract pattern of causal interaction between the components of a system, and perhaps between these components and external inputs and outputs. A functional organization **is determined (E) by**

specifying (1) a number of abstract components, (2) for each component, a number of different possible states, and (3) a system of dependency relations, specifying how the states of each component depends on the previous states of all components and on inputs to the system, and how outputs from the system depend on previous component states. Beyond specifying their number and their dependency relations, the nature of the components and the states is left unspecified.

(6) A physical system realizes a given functional organization when the system can be divided into an appropriate number of physical components each with the appropriate number of possible states, such that the causal dependency relations between the components of the system, inputs, and outputs precisely **reflect (EB)** the dependency relations given in the specification of the functional organization. A given functional organization can be realized by diverse physical systems. For example, the organization realized by the brain at the neural level might in principle be realized by a silicon system.

NOTATION

Module One

It is widely accepted that conscious experience **has a** physical basis. That is, the properties of experience (phenomenal properties, or qualia) systematically **depend on** physical properties according to some lawful relation.

G_{13} : Category one of properties of experience (phenomenal properties, or qualia) systematically; physical properties according to some lawful relation.

G_{14} : Category two of properties of experience (phenomenal properties, or qualia) systematically; physical properties according to some lawful relation.

G_{15} : Category three of properties of experience (phenomenal properties, or qualia) systematically; physical properties according to some lawful relation.

T_{13} : Category one of physical properties according to some lawful relation; properties of experience (phenomenal properties, or qualia) systematically

T_{14} : Category two of physical properties according to some lawful relation; properties of experience (phenomenal properties, or qualia) systematically

T_{15} : Category three of physical properties according to some lawful relation; properties of experience (phenomenal properties, or qualia) systematically

Module Two

There are two key questions about this relation. The first concerns the strength of the laws: are they logically or metaphysically necessary, or are they merely nomologically necessary? This question about the strength of the psychophysical link **is the basis for** debates over physicalism and property dualism.

G_{16} : Category one of physicalism and property dualism

G_{17} : Category two of physicalism and property dualism

G_{18} : Category three of physicalism and property dualism

T_{16} : Category one of strength of the psychophysical link

T_{17} : Category two of strength of the psychophysical link

T_{18} : Category three of strength of the psychophysical link

Module three

The second question concerns the nature of the laws: precisely how do phenomenal properties **depend on** physical properties? What sort of physical properties **enter into** the laws' antecedents, for instance; consequently, what sort of physical systems **can give rise** to conscious experience? It is this second question that I will address in this paper. To put the issue differently, even once it is accepted that experience **arises from** physical systems, the question remains open: in virtue of what sort of physical properties does conscious experience arise? Some property that brains can possess, presumably, but it is far from obvious what sort of property this is. Some have suggested biochemical properties; some have suggested quantum-mechanical properties; many have professed uncertainty. A natural suggestion is that when experience arises from a physical system, **it does (E) so in virtue** of the system's functional organization. On this view, the chemical and indeed the quantum substrate of the brain is not directly relevant to the existence of consciousness. What counts is rather the brain's abstract causal organization, an organization that might be realized in many different physical substrates.

G_{20} : Category one of physical properties; phenomenal properties; physical systems; physical systems with physical properties; system's functional organization

G_{21} : Category two of physical properties; phenomenal properties; physical systems; physical systems with physical properties; system's functional organization

G_{22} : Category three of physical properties; phenomenal properties; physical systems; physical systems with physical properties; system's functional organization

T_{20} : Category one of phenomenal properties; physical properties; conscious experience; experience; experience arises from a physical system;

T_{21} : Category two of phenomenal properties; physical properties; conscious experience; experience; experience arises from a physical system;

T_{22} : Category three of phenomenal properties; physical properties; conscious experience; experience; experience arises from a physical system;

Module four

In this paper I defend this view. Specifically, I defend a principle of organizational invariance, holding that experience **is i**nvariant across systems with the same fine-grained functional organization. More precisely, the principle states that given any system that has conscious experiences, **then any system** that has the same functional organization at a fine enough grain will have qualitatively identical conscious experiences. A full specification of a system's fine-grained functional organization **will fully determine** any conscious experiences that arise

G_{24} : Category one of **i**nvariant across systems with the same fine-grained functional organization; given any system that has conscious experiences; specification of a system's fine-grained functional organization;

G_{25} : Category two of **i**nvariant across systems with the same fine-grained functional organization; given any system that has conscious experiences; specification of a system's fine-grained functional organization;

G_{26} : Category three of **i**nvariant across systems with the same fine-grained functional organization; given any system that has conscious experiences; specification of a system's fine-grained functional organization;

T_{24} : Category one of experience; **system** that has the same functional organization at a fine enough grain will have qualitatively identical conscious experiences; any conscious experiences that arise;

T_{25} : Category two of experience; **system** that has the same functional organization at a fine enough grain will have qualitatively identical conscious experiences; any conscious experiences that arise;

T_{26} : Category three of experience; **system** that has the same functional organization at a fine enough grain will have qualitatively identical conscious experiences; any conscious experiences that arise;

Module five

To clarify this, we must first clarify the notion of functional organization. This is best understood as the abstract pattern of causal interaction between the components of a system, and perhaps between these components and external inputs and outputs. A functional organization **is determined (E) by** specifying (1) a number of abstract components, (2) for each component, a number of different possible states, and (3) a system of dependency relations, specifying how the states of each component depends on the previous states of all components and on inputs to the system, and how outputs from the system depend on previous component states. Beyond specifying their number and their dependency relations, the nature of the components and the states is left unspecified

G_{28} : Category one of number of abstract components

G_{29} : Category two of number of abstract components

G_{30} : Category three of number of abstract components

T_{28} : Category one of functional organization

T_{29} : Category two of functional organization

T_{30} : Category three of functional organization

Module six

G_{32} : Category one of for each component, a number of different possible states

G_{33} : Category two of for each component, a number of different possible states

G_{34} : Category three of for each component, a number of different possible states

T_{32} : Category one of functional organization

T_{33} : Category two of functional organization

T_{34} : Category three of functional organization

Module seven

G_{36} : Category one of system of dependency relations, specifying how the states of each component depends on the previous states of all components and on inputs to the system, and how outputs from the system depend on previous component states. Beyond specifying their number and their dependency relations, the nature of the components and the states is left unspecified

G_{37} : Category two of system of dependency relations, specifying how the states of each component depend on the previous states of all components and on inputs to the system, and how outputs from the system depend on previous component states. Beyond specifying their number and their dependency relations, the nature of the components and the states is left unspecified

G_{38} : Category three of system of dependency relations, specifying how the states of each component depend on the previous states of all components and on inputs to the system, and how outputs from the system depend on previous component states. Beyond specifying their number and their dependency relations, the nature of the components and the states is left unspecified

T_{36} : Category one of functional organization

T_{37} : Category two of functional organization

T_{38} : Category three of functional organization

Module eight

A physical system realizes a given functional organization when the system can be divided into an appropriate number of physical components each with the appropriate number of possible states, such that the causal dependency relations between the components of the system, inputs, and outputs precisely **reflect (EB)** the dependency relations given in the specification of the functional organization. A given functional organization can be realized by diverse physical systems. For example, the organization realized by the brain at the neural level might in principle be realized by a silicon system.

G_{40} : Category one of physical system realizes a given functional organization when the system can be divided into an appropriate number of physical components each with the appropriate number of possible states, such that the causal dependency relations between the components of the system, inputs, and outputs precisely

G_{41} : Category two of physical system realizes a given functional organization when the system can be divided into an appropriate number of physical components each with the appropriate number of possible states, such that the causal dependency relations between the components of the system, inputs, and outputs precisely

G_{42} : Category three of physical system realizes a given functional organization when the system can be divided into an appropriate number of physical components each with the appropriate number of possible states, such that the causal dependency relations between the components of the system, inputs, and outputs precisely

T_{40} : Category one of dependency relations given in the specification of the functional organization. A given functional organization can be realized by diverse physical systems. For example, the organization realized by the brain at the neural level might in principle be realized by a silicon system

T_{41} : Category two of dependency relations given in the specification of the functional organization. A given functional organization can be realized by diverse physical systems. For example, the organization realized by the brain at the neural level might in principle be realized by a silicon system

T_{42} : Category three of dependency relations given in the specification of the functional organization. A given functional organization can be realized by diverse physical systems. For example, the organization

realized by the brain at the neural level might in principle be realized by a silicon system

Module Nine (Unsolved Problem in Neuro Science)

THE GENERAL LEDGER (UNSOLVED PROBLEM)

The binding problem is when you look at what's happening in the brain, you find there's a division of labor. You have some parts of your brain that care about **vision**, some about **hearing,** and some about **touch.** And even within a system, like vision, you have parts that care about **colors, parts that care about orientations**, parts that care **about angles**. And **how this all comes together** so that you have a unified perception of the world is one of the unsolved mysteries in neuroscience.

We're not aware of that division of labor. Everything seems like it's perfectly unified to us. So this is still something we're all working on.

One thing that's very clear to us now, though, is that vision is not like a camera. It's not like light signals hit your eye and work their way up to the top and they move up some hierarchy and then they get seen. Instead, vision is all about internal activity that's already happening in your head and there's a little bit of data that comes up these cables and modifies or modulates that activity. But, essentially, all you're ever seeing is your internal model of what you believe you're seeing out there.

And this process of balancing shall go on,. And finally the various parameters are stored. As to how when one sees some acquaintance how the link pops up is given by sundry deposit- account identified problem solution discussed in one of the papers. In essence the holistic feelings of vision, hear, touch, sound, angles, orientations are all stored and the liabilities are those of people whose characterstics are dissipated away which come to the fore by Sundry deposit-account identification syndrome, when met and "memory is refreshed and recollected". Model makes allowance for new additions to the extant and existential recordial evidence and also allows for the dissipation, which might be stored in the subconscious. All the three states can be studied using the model. Each of the profile classified has inward additions and outward dissipation.

G_{44} : Category one of assets pertaining to accounts in individual general ledger about vision, hearing, touch, colors, parts, orientations, angles. There are different individuals and we are talking of the idiosyncrasies that form the terra firma and locus standi of classification.

G_{45} : Category two of assets pertaining to accounts in individual general ledger about vision, hearing, touch, colors, parts, orientations, angles. There are different individuals and we are talking of the idiosyncrasies that form the terra firma and locus standi of classification

G_{46} : Category three of assets pertaining to accounts in individual general ledger about vision, hearing, touch, colors, parts, orientations, angles. There are different individuals and we are talking of the idiosyncrasies that form the terra firma and locus standi of classification

T_{44} : Category one of liabilities pertaining to accounts in individual general ledger about vision, hearing, touch, colors, parts, orientations, angles. There are different individuals and we are talking of the idiosyncrasies that form the terra firma and locus standi of classification

T_{45} : Category two of liabilities pertaining to accounts in individual general ledger about vision, hearing, touch, colors, parts, orientations, angles. There are different individuals and we are talking of the

idiosyncrasies that form the terra firma and locus standi of classification

T_{46} : Category three of liabilities pertaining to accounts in individual general ledger about vision, hearing, touch, colors, parts, orientations, angles. There are different individuals and we are talking of the idiosyncrasies that form the terra firma and locus standi of classification

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**
$-(b''_{28})^{(5)}\big((G_{31}),t\big) =$ **First detritions factor**
Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \big[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\big]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \big[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\big]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \big[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\big]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \big[(b'_{32})^{(6)} - (b''_{32})^{(6)}\big((G_{35}),t\big)\big]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \big[(b'_{33})^{(6)} - (b''_{33})^{(6)}\big((G_{35}),t\big)\big]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \big[(b'_{34})^{(6)} - (b''_{34})^{(6)}\big((G_{35}),t\big)\big]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**
Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \big[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\big]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \big[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\big]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \big[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\big]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \big[(b'_{36})^{(7)} - (b''_{36})^{(7)}\big((G_{39}),t\big)\big]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \big[(b'_{37})^{(7)} - (b''_{37})^{(7)}\big((G_{39}),t\big)\big]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \big[(b'_{38})^{(7)} - (b''_{38})^{(7)}\big((G_{39}),t\big)\big]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**
Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \big[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\big]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \big[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\big]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \big[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\big]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \big[(b'_{40})^{(8)} - (b''_{40})^{(8)}\big((G_{43}),t\big)\big]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \big[(b'_{41})^{(8)} - (b''_{41})^{(8)}\big((G_{43}),t\big)\big]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \big[(b'_{42})^{(8)} - (b''_{42})^{(8)}\big((G_{43}),t\big)\big]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \big[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\big]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \big[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\big]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \big[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\big]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \big[(b'_{44})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big)\big]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \big[(b'_{45})^{(9)} - (b''_{45})^{(9)}\big((G_{47}),t\big)\big]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)}\ \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\ \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\ \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{c} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\ \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\ \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\ \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{array}\right] T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{c} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\ \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\ \boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\ \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{array}\right] T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$

$are\ fourth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
$are\ fifth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
$are\ sixth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
$are\ seventh\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
$are\ eighth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

$Where\ -(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ $are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ $are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ $are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad (84)$$

$where\ \boxed{-(b''_{28})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{29})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{30})^{(5)}(G_{31},t)}\ \ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}\ ,\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}\ ,\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt}=(a_{32})^{(6)}G_{33} \qquad 85$$

$$-\left[\begin{matrix}(a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)}\ \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}\ \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{32}$$

$$\frac{dG_{33}}{dt}=(a_{33})^{(6)}G_{32}-\left[\begin{matrix}(a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)}\ \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\ \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{33} \qquad 86$$

$$\frac{dG_{34}}{dt}=(a_{34})^{(6)}G_{33}-\left[\begin{matrix}(a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\ \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\ \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{34} \qquad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}\ ,\boxed{+(a''_{33})^{(6)}(T_{33},t)}\ ,\boxed{+(a''_{34})^{(6)}(T_{33},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}\ ,\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\ ,\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\ are\ second\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}\ \ are\ third\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation

coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34} \quad (90)$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{c} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{c} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(Y) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(Z) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(AA) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(BB)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(CC) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(DD) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\leq\ \left(\hat{P}_{13}\right)^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\leq\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\leq\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\leq\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{20})^{(3)}, T_i^0 \le (\hat{Q}_{20})^{(3)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(e) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)} t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$

In the same way , one can obtain

$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25}\ \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25}\ \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}\ , G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\ , \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40{,}41{,}42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(8)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(\mathrm{T}_{17},t)+(a''_{17})^{(2)}(\mathrm{T}_{17},t)\le-(\sigma_1)^{(2)}$ (293)

$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)}$ (294)

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(\nu_1)^{(6)}, (\nu_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, \nu^{(6)}, u^{(6)}$: 350

By $(\nu_1)^{(6)} > 0$, $(\nu_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)},, (\bar{\nu}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, $\boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad (354)$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad (355)$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad (356)$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^{0}e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^{0}e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^{0}}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t}-e^{-(b'_{34})^{(6)}t}\right]+T_{34}^{0}e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^{0}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T_{34}^{0}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)}+(C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)}=\frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)}(T_{45}, t)\right) - (a_{45}'')^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(e) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^{*} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^{*} for which $f(T_{33}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

Definition and uniqueness of T_{37}^{*} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^{*} for which $f(T_{37}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523

A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^{*}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^{*}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}((G_{31})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b_i'')^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTY SIX

Holographic And Anthropic Multiverse: Aei Ho Theos Geōmetreî

INTRODUCTION—VARIABLES USED

The Geometry of Interaction (GoI) was introduced by Jean-Yves Girard shortly after his work on Linear Logic. In linear logic, proofs can be seen as some kind of networks instead of the flat tree structure of sequent calculus. To distinguish the real proof nets from all the possible networks, Girard devised a criterium involving trips in the network. Trips can in fact be seen as some kind of operator acting on the proof. Drawing from this observation, Girard described directly this operator from the proof and has given a formula, the so-called execution formula, encoding the process of cut elimination at the level of operators. One of the first significant applications of GoI was a better analysis of Lamping's algorithm for optimal

reduction for the lambda calculus. GoI had a strong influence on game semantics for linear logic and PCF. GoI has been applied to deep compiler optimization for lambda calculi. A bounded version of GoI dubbed the Geometry of Synthesis has been used to compile higher-order programming languages directly into static circuits. (Wikipedia)

Physical Traces: Quantum vs. Classical Information Processing Samson Abramskya http://dx.doi.org/10.1016/S1571-0661 (04)80556-5

(1) Within the Geometry of Interaction (GoI) paradigm, authors present a setting that **enables** qualitative differences between classical and quantum processes to be explored.
(2) The key construction is the physical interpretation/realization of the traced monoidal categories of finite dimensional vector spaces with tensor product as monoidal structure and of finite sets and relations with Cartesian product as monoidal structure, **both of them providing** a so-called wave-style GoI.
(3) The developments in this paper reveal that envisioning state update due to quantum measurement as a process powerful tool **for (eb) developing** high-level approaches to quantum information processing.

The Holographic provides a Anthropic Multiverse Formalizing the Complex Geometry of Reality Richard L Amoroso and Elizabeth A Rauscher

(4) Every hundred years or so, a unique groundbreaking Copernican class volume arises unexpectedly. From ashes long thought cold of Einstein's static universe model, for the first time technically viable alternative interpretations to all pillars of Big Bang cosmology are presented in the context of a profound new "continuous-state" cosmological paradigm able to elucidate many contemporary problems plaguing the standard model of particle physics. The cosmology provides an alternative derivation of the string/brane tension formalism derived from large-scale additional dimensions **that leads** to a putative unique background-independent string vacuum without requiring the Higgs mechanism or SUSY superpartners.
(5) Breakthroughs presented, arising from the broad spectrum of explanatory power, include an empirical protocol for violation of the quantum uncertainty principle **that provides** a viable model for the implementation of universal bulk quantum computing.
(6) Other developments naturally arising from utility of the new regime **include a** design for constructing de Broglie-type matter-wave projectile defense shields with far-reaching consequences.

Atomic nuclei intimately entangled by a quantum measurement (figures deleted due to spatial constraints)

(7) Scientists from the Netherlands (Delft University of Technology and the FOM Foundation) and the UK (Element Six) have brought two atomic nuclei in a diamond into a quantum entangled state. This exotic relation was **created** by subjecting the nuclei to a new type of quantum measurement. These experiments mark an important step towards the realization of a quantum computer. The results were published on 14 October 2012 online in Nature Physics.

Electron microscope image of the synthetic diamond sample used. The green hemisphere is a lens that was etched into the synthetic diamond with a focussed ion beam, so that it is placed right on top of the nuclei to be entangled. The blue and yellow structures are metal electrodes that are created on the synthetic diamond using nanolithography. Scale: the total width of the image is 50 micrometer.

Quantum technologies and entanglement

(8) Quantum entanglement is one of the most intriguing phenomena in physics. When two particles are entangled their properties are so strongly connected that **they (e) lose** their own identity.

(9) Measuring both particles **yields** fully correlated outcomes, even when the particles are very far apart. Einstein famously called this feature 'spooky action at a distance'.

NOTATION

Module One

Within the Geometry of Interaction (GoI) paradigm, authors present a setting that **enables** qualitative differences between classical and quantum processes to be explored

G_{13} : Category one of paradigmatic statement within the Geometry of interaction schema and doxa. We are talking of systems that have characterstics which are responsible for the classification of the scheme of presentation.

G_{14} : Category two of paradigmatic statement within the Geometry of interaction schema and doxa

G_{15} : Category three of paradigmatic statement within the Geometry of interaction schema and doxa

T_{13} : Category one of qualitative differences between classical and quantum processes to be explored

T_{14} : Category two of qualitative differences between classical and quantum processes to be explored

T_{15} : Category three of qualitative differences between classical and quantum processes to be explored

Module Two

The key construction is the physical interpretation/realization **of (eb) the** traced monoidal categories of finite dimensional vector spaces with tensor product as monoidal structure and of finite sets and relations with Cartesian product as monoidal structure, **both of them providing** a so-called wave-style GoI.

G_{16} : Category one of key construction is the physical interpretation/realization

G_{17} : Category two of key construction is the physical interpretation/realization

G_{18} : Category three of key construction is the physical interpretation/realization

T_{16} : Category one of traced monoidal categories of finite dimensional vector spaces with tensor product as monoidal structure and of finite sets and relations with Cartesian product as monoidal structure, **both of them providing** a so-called wave-style GoI.

T_{17} : Category two of traced monoidal categories of finite dimensional vector spaces with tensor product as monoidal structure and of finite sets and relations with Cartesian product as monoidal structure, **both of them providing** a so-called wave-style GoI.

T_{18} : Category three of traced monoidal categories of finite dimensional vector spaces with tensor product as monoidal structure and of finite sets and relations with Cartesian product as monoidal structure, **both of them providing** a so-called wave-style GoI.

Module three

The developments in this paper reveal that envisioning state update due to quantum measurement as a process powerful tool **for (eb) developing** high-level approaches to quantum information processing.

G_{20} : Category one of envisioning state update due to quantum measurement as a process powerful tool

G_{21} : Category two of envisioning state update due to quantum measurement as a process powerful tool

G_{22} : Category three of envisioning state update due to quantum measurement as a process powerful tool

T_{20} : Category one of high-level approaches to quantum information processing

T_{21} : Category two of high-level approaches to quantum information processing

T_{22} : Category three of high-level approaches to quantum information processing

Module four

Every hundred years or so, a unique groundbreaking Copernican class volume arises unexpectedly. From ashes long thought cold of Einstein's static universe model, for the first time technically viable alternative interpretations to all pillars of Big Bang cosmology are presented in the context of a profound new "continuous-state" cosmological paradigm able to elucidate many contemporary problems plaguing the standard model of particle physics. The cosmology provides an alternative derivation of the string/brane tension formalism derived from large-scale additional dimensions **that leads** to a putative unique background-independent string vacuum without requiring the Higgs mechanism or SUSY superpartners.

G_{24} : Category one of cosmology provides an alternative derivation of the string/brane tension formalism derived from large-scale additional dimensions

G_{25} : Category two of cosmology provides an alternative derivation of the string/brane tension formalism derived from large-scale additional dimensions

G_{26} : Category three of cosmology provides an alternative derivation of the string/brane tension formalism derived from large-scale additional dimensions

T_{24} : Category one of putative unique background-independent string vacuum without requiring the Higgs mechanism or SUSY superpartners

T_{25} : Category two of putative unique background-independent string vacuum without requiring the Higgs mechanism or SUSY superpartners

T_{26} : Category three of putative unique background-independent string vacuum without requiring the Higgs mechanism or SUSY superpartners

Module five

Breakthroughs presented, arising from the broad spectrum of explanatory power, include an empirical protocol for violation of the quantum uncertainty principle **that provides** a viable model for the implementation of universal bulk quantum computing

G_{28} : Category one of empirical protocol for violation of the quantum uncertainty principle

G_{29} : Category two of empirical protocol for violation of the quantum uncertainty principle

G_{30} : Category three of empirical protocol for violation of the quantum uncertainty principle

T_{28} : Category one of viable model for the implementation of universal bulk quantum computing

T_{29} : Category two of viable model for the implementation of universal bulk quantum computing

T_{30} : Category three of viable model for the implementation of universal bulk quantum computing

Module six

Other developments naturally arising from utility of the new regime **include a** design for constructing de Broglie-type matter-wave projectile defense shields with far-reaching consequences

G_{32} : Category one of design for constructing de Broglie-type matter-wave projectile defense shields with far-reaching consequences

G_{33} : Category two of design for constructing de Broglie-type matter-wave projectile defense shields with far-reaching consequences

G_{34} : Category three of design for constructing de Broglie-type matter-wave projectile defense shields with far-reaching consequences

T_{32} : Category one of developments naturally arising from utility of the new regime

T_{33} : Category two of developments naturally arising from utility of the new regime

T_{34} : Category three of developments naturally arising from utility of the new regime

Module seven

Scientists from the Netherlands (Delft University of Technology and the FOM Foundation) and the UK (Element Six) have brought two atomic nuclei in a diamond into a quantum entangled state. This exotic relation was **created** by subjecting the nuclei to a new type of quantum measurement. These experiments mark an important step towards the realization of a quantum computer. The results were published on 14 October 2012 online in Nature Physics.

G_{36} : Category one of two atomic nuclei in a diamond into a quantum entangled state an exotic relation

G_{37} : Category two of two atomic nuclei in a diamond into a quantum entangled state an exotic relation

G_{38} : Category three of two atomic nuclei in a diamond into a quantum entangled state an exotic relation

T_{36} : Category one of by subjecting the nuclei to a new type of quantum measurement. These experiments mark an important step towards the realization of a quantum computer

T_{37} : Category two of by subjecting the nuclei to a new type of quantum measurement. These experiments mark an important step towards the realization of a quantum computer

T_{38} : Category three of by subjecting the nuclei to a new type of quantum measurement. These experiments mark an important step towards the realization of a quantum computer

Module eight

Quantum entanglement is one of the most intriguing phenomena in physics. When two particles are entangled their properties are so strongly connected that **they (e) lose** their own identity.

G_{40} : Category one of their own identity

G_{41} : Category two of their own identity

G_{42} : Category three of their own identity

T_{40} : Category one of two particles are entangled their properties are so strongly connected

T_{41} : Category two of two particles are entangled their properties are so strongly connected

T_{42} : Category three of two particles are entangled their properties are so strongly connected

Module Nine

Measuring both particles **yields** fully correlated outcomes, even when the particles are very far apart. Einstein famously called this feature 'spooky action at a distance'.

G_{44} : Category one of Measuring both particles

G_{45} : Category two of Measuring both particles

G_{46} : Category three of Measuring both particles

T_{44} : Category one of correlated outcomes, even when the particles are very far apart. Einstein famously called this feature 'spooky action at a distance'.

T_{45} : Category two of correlated outcomes, even when the particles are very far apart. Einstein famously called this feature 'spooky action at a distance'.

T_{46} : Category three of correlated outcomes, even when the particles are very far apart. Einstein famously called this feature 'spooky action at a distance'.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$
$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$
$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}),t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) \quad +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{37}'')^{(7,7,)}(G_{39},t)$, $-(b_{36}'')^{(7,7,)}(G_{39},t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8)}(G_{43},t)$ $-(b_{41}'')^{(8,8)}(G_{43},t)$ $-(b_{42}'')^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17},t) + (a_{13}'')^{(1,1,)}(T_{14},t) + (a_{20}'')^{(3,3,3)}(T_{21},t) \\ + (a_{24}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{28}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{32}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{36}'')^{(7,7,7)}(T_{37},t) + (a_{40}'')^{(8,8,8)}(T_{41},t) + (a_{44}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17},t) + (a_{14}'')^{(1,1,)}(T_{14},t) + (a_{21}'')^{(3,3,3)}(T_{21},t) \\ + (a_{25}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{29}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{33}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{37}'')^{(7,7,7)}(T_{37},t) + (a_{41}'')^{(8,8,8)}(T_{41},t) + (a_{45}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17},t) + (a_{15}'')^{(1,1,)}(T_{14},t) + (a_{22}'')^{(3,3,3)}(T_{21},t) \\ + (a_{26}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{30}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{34}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{38}'')^{(7,7,7)}(T_{37},t) + (a_{42}'')^{(8,8,8)}(T_{41},t) + (a_{46}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad 63$$

Where $+(a_{16}'')^{(2)}(T_{17},t)$, $+(a_{17}'')^{(2)}(T_{17},t)$, $+(a_{18}'')^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for*

category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category $1, 2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a_{32}')^{(6)} & +(a_{32}'')^{(6)}(T_{33},t) & +(a_{28}'')^{(5,5,5)}(T_{29},t) & +(a_{24}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a_{33}')^{(6)} & +(a_{33}'')^{(6)}(T_{33},t) & +(a_{29}'')^{(5,5,5)}(T_{29},t) & +(a_{25}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a_{34}')^{(6)} & +(a_{34}'')^{(6)}(T_{33},t) & +(a_{30}'')^{(5,5,5)}(T_{29},t) & +(a_{26}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a_{32}'')^{(6)}(T_{33},t)$, $+(a_{33}'')^{(6)}(T_{33},t)$, $+(a_{34}'')^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a_{28}'')^{(5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a_{24}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad (90)$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad (91)$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{ccc} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(EE) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(FF) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(GG) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(HH)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(II) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(JJ) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\le\ \left(\hat{P}_{13}\right)^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\le\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\le\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 156

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^{0} + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^{0} + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{20})^{(3)}, T_i^0 \le (\hat{Q}_{20})^{(3)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By 161

$$\bar{G}_{20}(t) = G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^{0} + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^{0} + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(f) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}\leq$$
$$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{\mathrm{A}}_{16})^{(2)}+(\widehat{\mathrm{P}}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)\mathrm{d}\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

208

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)}$$

209

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)}$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),\left((G_{23})^{(2)},(T_{23})^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

214

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\,((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\leq G_{15}(t)\leq\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\leq T_{13}(t)\leq T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\leq T_{13}(t)\leq\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\leq T_{15}(t)\leq \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\leq-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(\mathrm{T}_{17},t)+(a''_{17})^{(2)}(\mathrm{T}_{17},t)\leq-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\leq-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\leq-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$

$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t} \right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)} t} \right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t} \right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(\nu_1)^{(6)}, (\nu_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, \nu^{(6)}, u^{(6)}$: 350

By $(\nu_1)^{(6)} > 0$, $(\nu_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)},, (\bar{\nu}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

$$\Big(\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t}\Big)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{\left((R_1)^{(9)}+(r_{44})^{(9)}\right)t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{\left((R_1)^{(9)}+(r_{44})^{(9)}\right)t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}\left((R_1)^{(9)}-(b'_{46})^{(9)}\right)}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}\left((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)}\right)}\left[e^{\left((R_1)^{(9)}+(r_{44})^{(9)}\right)t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $v^{(4)}$:- $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

For $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le v^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $v^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)},(v_0)^{(7)}$:-

For $0<\boxed{(v_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(v_1)^{(7)}<(\bar{v}_1)^{(7)}$

$$v^{(7)}(t)\geq\frac{(v_1)^{(7)}+(C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(v_1)^{(7)}-(v_0)^{(7)}}{(v_0)^{(7)}-(v_2)^{(7)}}}$$

it follows $(v_0)^{(7)}\leq v^{(7)}(t)\leq(v_1)^{(7)}$

In the same manner , we get 417

$$v^{(7)}(t)\leq\frac{(\bar{v}_1)^{(7)}+(\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{v}_1)^{(7)}-(v_0)^{(7)}}{(v_0)^{(7)}-(\bar{v}_2)^{(7)}}}$$

From which we deduce $(v_0)^{(7)}\leq v^{(7)}(t)\leq(\bar{v}_1)^{(7)}$

If $0<(v_1)^{(7)}<(v_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)}\leq\frac{(v_1)^{(7)}+(C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_2)^{(7)}\right)t\right]}}\leq v^{(7)}(t)\leq$$

$$\frac{(\bar{v}_1)^{(7)}+(\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}\leq(\bar{v}_1)^{(7)}$$

If $0<(v_1)^{(7)}\leq(\bar{v}_1)^{(7)}\leq\boxed{(v_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)}\leq v^{(7)}(t)\leq\frac{(\bar{v}_1)^{(7)}+(\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}\leq(v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)}\leq v^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{v^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(f) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^{*} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^{*} for which $f(T_{33}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

Definition and uniqueness of T_{37}^{*} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^{*} for which $f(T_{37}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523

A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b_i'')^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTY SEVEN

Quantum Transport: Ut Sementem Feceris Ita Metes

INTRODUCTION—VARIABLES USED

Chinmay Belthangady, N. Bar-Gill, L. M. Pham, K. Arai, D. Le Sage, P. Cappellaro, R. L. Walsworth "Dressed-State Resonant Coupling between Bright and Dark Spins in Diamond "Phys. Rev. Lett. 110, 157601(2013)

(1) Under ambient conditions, spin impurities in solid-state systems **are (EB) found in** thermally-mixed states and are optically "dark", i.e., the spin states cannot be optically controlled.\
(2) Nitrogen-vacancy (NV) centers in diamond are an exception in that the electronic spin states are "bright", i.e., they can be polarized by optical pumping, coherently **manipulated(E&EB) w**ith spin-resonance techniques, and read out optically, all at room temperature.
(3) Here authors demonstrate a dressed-state, double-resonance scheme to transfer polarization from bright NV electronic spins **to** dark substitutional-Nitrogen (P1) electronic spins in diamond.

(4) This polarization-transfer mechanism **could be used** to cool a mesoscopic bath of dark spins to near-zero temperature, thus providing a resource for quantum information and sensing, and aiding studies of quantum effects in many-body spin systems.

A. Ajoy and P. Cappellaro "Perfect quantum transport in arbitrary spin networks "Phys. Rev. B 87, 064303 (2013)

(5) Spin chains have been proposed **as (=)** wires to transport information between distributed registers in a quantum information processor.

(6) Unfortunately, the challenges in manufacturing linear chains with engineered couplings have **hindered** experimental implementations. Here authors present strategies to achieve perfect quantum information transport in arbitrary spin networks. Proposal is **based on the** weak coupling limit for pure state transport, where information is transferred between two end spins that are only weakly coupled to the rest of the network.

(7) This regime **allows** ignoring the complex, internal dynamics of the bulk network and relying on virtual transitions or on the coupling to a single bulk eigenmodes.

(8) They further introduce control methods capable **of tuning the** transport process and achieve perfect fidelity with limited resources, involving only **manipulation (e&eb) of** the end qubits.

(9) These strategies control methods capable **of tuning the** transport process and achieve perfect fidelity with limited resources, involving only **manipulation (e&eb) of** the end qubits could **be thus applied** not only to engineered systems with relaxed fabrication precision, but also to naturally occurring networks; specifically, we discuss the practical implementation of quantum state transfer between two separated nitrogen vacancy (NV) centers through a network of nitrogen substitutional impurities.

NOTATION

Module One

Under ambient conditions, spin impurities in solid-state systems **are (EB) found in** thermally-mixed states and are optically "dark", i.e., the spin states cannot be optically controlled

G_{13} : Category one of spin impurities in solid-state systems

G_{14} : Category two of spin impurities in solid-state systems

G_{15} : Category three of spin impurities in solid-state systems

T_{13} : Category one of thermally-mixed states and are optically "dark", i.e., the spin states cannot be optically controlled

T_{14} : Category two of thermally-mixed states and are optically "dark", i.e., the spin states cannot be optically controlled

T_{15} : Category three of thermally-mixed states and are optically "dark", i.e., the spin states cannot be optically controlled

Module Two

Nitrogen-vacancy (NV) centers in diamond are an exception in that the electronic spin states are "bright", i.e., they can be polarized by optical pumping, coherently **manipulated(E&EB) w**ith spin-resonance techniques, and read out optically, all at room temperature

G_{16} : Category one of Nitrogen-vacancy (NV) centers in diamond are an exception in that the electronic spin states are "bright", i.e., they can be polarized by optical pumping, coherently

G_{17} : Category two of Nitrogen-vacancy (NV) centers in diamond are an exception in that the electronic

spin states are "bright", i.e., they can be polarized by optical pumping, coherently

G_{18} : Category three of Nitrogen-vacancy (NV) centers in diamond are an exception in that the electronic spin states are "bright", i.e., they can be polarized by optical pumping, coherently

T_{16} : Category one of spin-resonance techniques, and read out optically, all at room temperature ;Nitrogen-vacancy (NV) centers in diamond are an exception in that the electronic spin states are "bright", i.e., they can be polarized by optical pumping, coherently

T_{17} : Category two of spin-resonance techniques, and read out optically, all at room temperature; Nitrogen-vacancy (NV) centers in diamond are an exception in that the electronic spin states are "bright", i.e., they can be polarized by optical pumping, coherently

T_{18} : Category three of spin-resonance techniques, and read out optically, all at room temperature ;Nitrogen-vacancy (NV) centers in diamond are an exception in that the electronic spin states are "bright", i.e., they can be polarized by optical pumping, coherently

Module three

Here authors demonstrate a dressed-state, double-resonance scheme to transfer polarization from bright NV electronic spins **to** dark substitutional-Nitrogen (P1) electronic spins in diamond.

G_{20} : Category one of dressed-state, double-resonance scheme to transfer polarization from bright NV electronic spins

G_{21} : Category two of dressed-state, double-resonance scheme to transfer polarization from bright NV electronic spins

G_{22} : Category three of dressed-state, double-resonance scheme to transfer polarization from bright NV electronic spins

T_{20} : Category one of dark substitutional-Nitrogen (P1) electronic spins in diamond

T_{21} : Category two of dark substitutional-Nitrogen (P1) electronic spins in diamond

T_{22} : Category three of dark substitutional-Nitrogen (P1) electronic spins in diamond

Module four

This polarization-transfer mechanism **could be used** to cool a mesoscopic bath of dark spins to near-zero temperature, thus providing a resource for quantum information and sensing, and aiding studies of quantum effects in many-body spin systems.

G_{24} : Category one of cool a mesoscopic bath of dark spins to near-zero temperature, thus providing a resource for quantum information and sensing, and aiding studies of quantum effects in many-body spin systems.

G_{25} : Category two of cool a mesoscopic bath of dark spins to near-zero temperature, thus providing a resource for quantum information and sensing, and aiding studies of quantum effects in many-body spin systems.

G_{26} : Category three of cool a mesoscopic bath of dark spins to near-zero temperature, thus providing a resource for quantum information and sensing, and aiding studies of quantum effects in many-body spin systems.

T_{24} : Category one of polarization-transfer mechanism

T_{25} : Category two of polarization-transfer mechanism

T_{26} : Category three of polarization-transfer mechanism

Module five

Spin chains have been proposed **as** wires to transport information between distributed registers in a quantum information processor

G_{28} : Category one of wires to transport information between distributed registers in a quantum information processor

G_{29} : Category two of wires to transport information between distributed registers in a quantum information processor

G_{30} : Category three of wires to transport information between distributed registers in a quantum information processor

T_{28} : Category one of Spin chains

T_{29} : Category two of Spin chains

T_{30} : Category three of Spin chains

. Module six

Unfortunately, the challenges in manufacturing linear chains with engineered couplings have **hindered** experimental implementations. Here authors present strategies to achieve perfect quantum information transport in arbitrary spin networks. Proposal is **based on the** weak coupling limit for pure state transport, where information is transferred between two end spins that are only weakly coupled to the rest of the network.

G_{32} : Category one of weak coupling limit for pure state transport, where information is transferred between two end spins that are only weakly coupled to the rest of the network.

G_{33} : Category two of weak coupling limit for pure state transport, where information is transferred between two end spins that are only weakly coupled to the rest of the network.

G_{34} : Category three of weak coupling limit for pure state transport, where information is transferred between two end spins that are only weakly coupled to the rest of the network.

T_{32} : Category one of strategies to achieve perfect quantum information transport in arbitrary spin networks.

T_{33} : Category two of strategies to achieve perfect quantum information transport in arbitrary spin networks.

T_{34} : Category three of strategies to achieve perfect quantum information transport in arbitrary spin networks.

Module seven

This regime **allows** ignoring the complex, internal dynamics of the bulk network and relying on virtual

transitions or on the coupling to a single bulk eigenmodes

G_{36} : Category one of weak coupling limit for pure state transport, where information is transferred between two end spins that are only weakly coupled to the rest of the network regime

G_{37} : Category two of weak coupling limit for pure state transport, where information is transferred between two end spins that are only weakly coupled to the rest of the network regime

G_{38} : Category three of weak coupling limit for pure state transport, where information is transferred between two end spins that are only weakly coupled to the rest of the network regime

T_{36} : Category one of ignoring the complex, internal dynamics of the bulk network and relying on virtual transitions or on the coupling to a single bulk eigenmodes

T_{37} : Category two of ignoring the complex, internal dynamics of the bulk network and relying on virtual transitions or on the coupling to a single bulk eigenmodes

T_{38} : Category three of ignoring the complex, internal dynamics of the bulk network and relying on virtual transitions or on the coupling to a single bulk eigenmodes

Module eight

They further introduce control methods capable **of tuning the** transport process and achieve perfect fidelity with limited resources, involving only **manipulation (e&eb) of** the end qubits.

G_{40} : Category one of control methods capable **of tuning the** transport process and achieve perfect fidelity with limited resources; end qubits

G_{41} : Category two of control methods capable **of tuning the** transport process and achieve perfect fidelity with limited resources; end qubits

G_{42} : Category three of control methods capable **of tuning the** transport process and achieve perfect fidelity with limited resources; end qubits

T_{40} : Category one of end qubits ;control methods capable **of tuning the** transport process and achieve perfect fidelity with limited resources,

T_{41} : Category two of end qubits ;control methods capable **of tuning the** transport process and achieve perfect fidelity with limited resources,

T_{42} : Category three of end qubits ;control methods capable **of tuning the** transport process and achieve perfect fidelity with limited resources,

Module Nine

These strategies control methods capable **of tuning the** transport process and achieve perfect fidelity with limited resources, involving only **manipulation (e&eb) of** the end qubits could **be thus applied** not only to engineered systems with relaxed fabrication precision, but also to naturally occurring networks specifically, practical implementation of quantum state transfer between two separated nitrogen vacancy (NV) centers through a network of nitrogen substitutional impurities

G_{44} : Category one of control methods capable **of tuning the** transport process and achieve perfect fidelity

with limited resources, involving only **manipulation (e&eb) of** the end qubits; engineered systems with relaxed fabrication precision, but also to naturally occurring networks specifically, practical implementation of quantum state transfer between two separated nitrogen vacancy (NV) centers through a network of nitrogen substitutional impurities

G_{45} : Category two of control methods capable **of tuning the** transport process and achieve perfect fidelity with limited resources, involving only **manipulation (e&eb) of** the end qubits; engineered systems with relaxed fabrication precision, but also to naturally occurring networks specifically, practical implementation of quantum state transfer between two separated nitrogen vacancy (NV) centers through a network of nitrogen substitutional impurities

G_{46} : Category three of control methods capable **of tuning the** transport process and achieve perfect fidelity with limited resources, involving only **manipulation (e&eb) of** the end qubits; engineered systems with relaxed fabrication precision, but also to naturally occurring networks specifically, practical implementation of quantum state transfer between two separated nitrogen vacancy (NV) centers through a network of nitrogen substitutional impurities

T_{44} : Category one of engineered systems with relaxed fabrication precision, but also to naturally occurring networks specifically, practical implementation of quantum state transfer between two separated nitrogen vacancy (NV) centers through a network of nitrogen substitutional impurities; control methods capable **of tuning the** transport process and achieve perfect fidelity with limited resources, involving only **manipulation (e&eb) of** the end qubits

T_{45} : Category two of engineered systems with relaxed fabrication precision, but also to naturally occurring networks specifically, practical implementation of quantum state transfer between two separated nitrogen vacancy (NV) centers through a network of nitrogen substitutional impurities; control methods capable **of tuning the** transport process and achieve perfect fidelity with limited resources, involving only **manipulation (e&eb) of** the end qubits

T_{46} : Category three of engineered systems with relaxed fabrication precision, but also to naturally occurring networks specifically, practical implementation of quantum state transfer between two separated nitrogen vacancy (NV) centers through a network of nitrogen substitutional impurities; control methods capable **of tuning the** transport process and achieve perfect fidelity with limited resources, involving only **manipulation (e&eb) of** the end qubits

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad (1)$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad (2)$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad (3)$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad (4)$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad (5)$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad (6)$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad (7)$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad (8)$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad (9)$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad (10)$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad (11)$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad (12)$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad (13)$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad (14)$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad (15)$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad (16)$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad (17)$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad (18)$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$
$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$
$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$
$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$
$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$
$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**
$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$
$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$
$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$
$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$
$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$
$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**
$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$
$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for

category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8)}(G_{43},t)$ $-(b_{41}'')^{(8,8)}(G_{43},t)$ $-(b_{42}'')^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt}=(a_{16})^{(2)}G_{17}-\begin{bmatrix}(a_{16}')^{(2)} & +(a_{16}'')^{(2)}(T_{17},t) & +(a_{13}'')^{(1,1,)}(T_{14},t) & +(a_{20}'')^{(3,3,3)}(T_{21},t)\\ +(a_{24}'')^{(4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)\\ +(a_{36}'')^{(7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9)}(T_{45},t)\end{bmatrix}G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt}=(a_{17})^{(2)}G_{16}-\begin{bmatrix}(a_{17}')^{(2)} & +(a_{17}'')^{(2)}(T_{17},t) & +(a_{14}'')^{(1,1,)}(T_{14},t) & +(a_{21}'')^{(3,3,3)}(T_{21},t)\\ +(a_{25}'')^{(4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)\\ +(a_{37}'')^{(7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9)}(T_{45},t)\end{bmatrix}G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt}=(a_{18})^{(2)}G_{17}-\begin{bmatrix}(a_{18}')^{(2)} & +(a_{18}'')^{(2)}(T_{17},t) & +(a_{15}'')^{(1,1,)}(T_{14},t) & +(a_{22}'')^{(3,3,3)}(T_{21},t)\\ +(a_{26}'')^{(4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)\\ +(a_{38}'')^{(7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9)}(T_{45},t)\end{bmatrix}G_{18} \quad (63)$$

Where $+(a_{16}'')^{(2)}(T_{17},t)$, $+(a_{17}'')^{(2)}(T_{17},t)$, $+(a_{18}'')^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a_{44}'')^{(9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt}=(b_{16})^{(2)}T_{17}-\begin{bmatrix}(b_{16}')^{(2)} & -(b_{16}'')^{(2)}(G_{19},t) & -(b_{13}'')^{(1,1,)}(G,t) & -(b_{20}'')^{(3,3,3,)}(G_{23},t)\\ -(b_{24}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{28}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{32}'')^{(6,6,6,6,6)}(G_{35},t)\\ -(b_{36}'')^{(7,7,7)}(G_{39},t) & -(b_{40}'')^{(8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9)}(G_{47},t)\end{bmatrix}T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients

for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category $1, 2$ *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)}\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32} \quad 85$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)}\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b_{36}')^{(7)}\boxed{-(b_{36}'')^{(7)}(G_{39},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b_{37}')^{(7)}\boxed{-(b_{37}'')^{(7)}(G_{39},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b_{38}')^{(7)}\boxed{-(b_{38}'')^{(7)}(G_{39},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{matrix} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{matrix} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{matrix} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\,-(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\,-(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\,-(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(KK) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(LL) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$ 132

(MM) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(NN)

$\lim_{G\to\infty}(b_i'')^{(7)}((G_{39}),t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}((G_{39}),t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(OO) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(PP) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ *and* $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(g) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}ds_{(16)} +$

$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right) - (a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)};\ (G_{23})^{(2)}, (T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(3)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)},, (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \qquad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq\frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \qquad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \qquad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$$

$$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$$

$$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)}\leq-(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq-(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)}\leq-(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq-(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \le \nu^{(2)}(t) \le (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \le u^{(2)}(t) \le (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

406

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case,

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

407

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t)\right) - (a_{41}'')^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(g) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)} G_{14}^{*}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)} G_{14}^{*}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)} T_{14}^{*}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)} T_{14}^{*}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)} G_{17}^{*}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)} G_{17}^{*}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)} T_{17}^{*}}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)} T_{17}^{*}}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)} G_{21}^{*}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)} G_{21}^{*}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)} T_{21}^{*}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)} T_{21}^{*}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTY SEVEN

Hamiltonian Engineering: Lucus A Non Lucendo-Lux Et Veritas--Lux Hominum Vita

INTRODUCTION—VARIABLES USED

A. Ajoy and P. Cappellaro "Quantum simulation via filtered Hamiltonian engineering: application to perfect quantum transport in spin networks" Phys. Rev. Lett. 110, 220503 (2013)

(1) Authors propose a method for Hamiltonian engineering that requires no local control **but only (e) relies on** collective qubit rotations and field gradients.
(2) The technique achieves a spatial modulation of the coupling strengths **via a** dynamical construction of a weighting function combined with a Bragg grating.
(3) As an example of the technique achieves a spatial modulation of the coupling strengths **via a** dynamical construction of a weighting function combined with a Bragg grating authors **demonstrate (eb)** how to generate the ideal Hamiltonian for perfect quantum information transport between two separated nodes of a large spin network.
(4) They engineer a spin chain with optimal couplings starting from a large spin network, such as one naturally occurring in crystals, while **decoupling (e)** all unwanted interactions.
(5) For realistic experimental parameters, method can be used **to (eb) drive** almost perfect quantum information transport at room temperature.
(6) The Hamiltonian engineering method can be made more robust under decoherence and coupling disorder **by (e) a** novel apodization scheme.
(7) Thus, the method is quite general and **can be (e) used to** engineer the Hamiltonian of many complex spin lattices with different topologies and interactions.

Decay of spin coherences in one-dimensional spin systems G. Kaur et al

(8) Strategies to protect multi-qubit states against decoherence are difficult to **formulate because (e) of** their complex many-body dynamics.
(9) A better knowledge of the decay dynamics would help in the construction of decoupling control schemes. Here we use solid-state nuclear magnetic resonance techniques to experimentally **investigate (e&eb)** the decay of coherent multi-spin states in linear spin chains. Leveraging on the quasi-one-dimensional geometry of Fluorapatite crystal spin systems, can gain a deeper insight on the multi-spin states **created (e) by** the coherent evolution, and their subsequent decay, than it is

possible in 3D systems.

NOTATION

Module One

Authors propose a method for Hamiltonian engineering that requires no local control **but only (e) relies on** collective qubit rotations and field gradients

G_{13} : Category one of collective qubit rotations and field gradients

G_{14} : Category two of collective qubit rotations and field gradients

G_{15} : Category three of collective qubit rotations and field gradients

T_{13} : Category one of method for Hamiltonian engineering that requires no local control

T_{14} : Category two of method for Hamiltonian engineering that requires no local control

T_{15} : Category three of method for Hamiltonian engineering that requires no local control

Module Two

The technique achieves a spatial modulation of the coupling strengths **via a** dynamical construction of a weighting function combined with a Bragg grating.

G_{16} : Category one of technique achieves a spatial modulation of the coupling strengths; dynamical construction of a weighting function combined with a Bragg grating.

G_{17} : Category two of technique achieves a spatial modulation of the coupling strengths; dynamical construction of a weighting function combined with a Bragg grating.

G_{18} : Category three of technique achieves a spatial modulation of the coupling strengths; dynamical construction of a weighting function combined with a Bragg grating.

T_{16} : Category one of dynamical construction of a weighting function combined with a Bragg grating.; technique achieves a spatial modulation of the coupling strengths

T_{17} : Category two of dynamical construction of a weighting function combined with a Bragg grating.; technique achieves a spatial modulation of the coupling strengths

T_{18} : Category three of dynamical construction of a weighting function combined with a Bragg grating.; technique achieves a spatial modulation of the coupling strengths

Module three

As an example of the technique achieves a spatial modulation of the coupling strengths **via a** dynamical construction of a weighting function combined with a Bragg grating authors **demonstrate (eb)** how to generate the ideal Hamiltonian for perfect quantum information transport between two separated nodes of a large spin network

G_{20} : Category one of example of the technique achieves a spatial modulation of the coupling strengths **via a** dynamical construction of a weighting function combined with a Bragg grating

G_{21} : Category two of example of the technique achieves a spatial modulation of the coupling strengths **via a** dynamical construction of a weighting function combined with a Bragg grating

G_{22} : Category three of example of the technique achieves a spatial modulation of the coupling strengths **via**

a dynamical construction of a weighting function combined with a Bragg grating

T_{20} : Category one of generate the ideal Hamiltonian for perfect quantum information transport between two separated nodes of a large spin network

T_{21} : Category two of generate the ideal Hamiltonian for perfect quantum information transport between two separated nodes of a large spin network

T_{22} : Category three of generate the ideal Hamiltonian for perfect quantum information transport between two separated nodes of a large spin network

Module four

They engineer a spin chain with optimal couplings starting from a large spin network, such as one naturally occurring in crystals, while **decoupling (e)** all unwanted interactions

G_{24} : Category one of unwanted interactions

G_{25} : Category two of unwanted interactions

G_{26} : Category three of unwanted interactions

T_{24} : Category one of spin chains with optimal couplings starting from a large spin network, such as one naturally occurring in crystals

T_{25} : Category two of spin chains with optimal couplings starting from a large spin network, such as one naturally occurring in crystals

T_{26} : Category three of spin chains with optimal couplings starting from a large spin network, such as one naturally occurring in crystals

Module five

For realistic experimental parameters, method can be used **to (eb) drive** almost perfect quantum information transport at room temperature.

G_{28} : Category one of spin chains with optimal couplings starting from a large spin network, such as one naturally occurring in crystals, while **decoupling (e)** all unwanted interactions can be used

G_{29} : Category two of spin chains with optimal couplings starting from a large spin network, such as one naturally occurring in crystals, while **decoupling (e)** all unwanted interactions can be used

G_{30} : Category three of spin chains with optimal couplings starting from a large spin network, such as one naturally occurring in crystals, while **decoupling (e)** all unwanted interactions can be used

T_{28} : Category one of almost perfect quantum information transport at room temperature.

T_{29} : Category two of almost perfect quantum information transport at room temperature.

T_{30} : Category three of almost perfect quantum information transport at room temperature.

Module six

The Hamiltonian engineering method can be made more robust under decoherence and coupling disorder **by**

(e) a novel apodization scheme.

G_{32} : Category one of apodization scheme

G_{33} : Category two of apodization scheme

G_{34} : Category three of apodization scheme

T_{32} : Category one of Hamiltonian engineering method can be made more robust under decoherence and coupling disorder

T_{33} : Category two of Hamiltonian engineering method can be made more robust under decoherence and coupling disorder

T_{34} : Category three of Hamiltonian engineering method can be made more robust under decoherence and coupling disorder

Module seven

The method is quite general and **can be (e) used to** engineer the Hamiltonian of many complex spin lattices with different topologies and interactions.

G_{36} : Category one of engineer the Hamiltonian of many complex spin lattices with different topologies and interactions.

G_{37} : Category two of engineer the Hamiltonian of many complex spin lattices with different topologies and interactions.

G_{38} : Category three of engineer the Hamiltonian of many complex spin lattices with different topologies and interactions.

T_{36} : Category one of method is quite general

T_{37} : Category two of method is quite general

T_{38} : Category three of method is quite general

Module eight

Strategies to protect multi-qubit states against decoherence are difficult to **formulate because (e) of** their complex many-body dynamics

G_{40} : Category one of complex many-body dynamics

G_{41} : Category two of complex many-body dynamics

G_{42} : Category three of complex many-body dynamics

T_{40} : Category one of Strategies to protect multi-qubit states against decoherence are difficult to **formulate**

T_{41} : Category two of Strategies to protect multi-qubit states against decoherence are difficult to **formulate**

T_{42} : Category three of Strategies to protect multi-qubit states against decoherence are difficult to **formulate**

Module Nine

A better knowledge of the decay dynamics would help in the construction of decoupling control schemes. Here authors use solid-state nuclear magnetic resonance techniques to experimentally **investigate (e&eb)** the decay of coherent multi-spin states in linear spin chains. Leveraging on the quasi-one-dimensional geometry of Fluorapatite crystal spin systems, can gain a deeper insight on the multi-spin states **created (e) by** the coherent evolution, and their subsequent decay, than it is possible in 3D systems.

G_{44} : Category one of coherent evolution, and their subsequent decay, than it is possible in 3D systems

G_{45} : Category two of coherent evolution, and their subsequent decay, than it is possible in 3D systems

G_{46} : Category three of coherent evolution, and their subsequent decay, than it is possible in 3D systems

T_{44} : Category one of better knowledge of the decay dynamics would help in the construction of decoupling control schemes. Here authors use solid-state nuclear magnetic resonance techniques to experimentally **investigate (e&eb)** the decay of coherent multi-spin states in linear spin chains. Leveraging on the quasi-one-dimensional geometry of Fluorapatite crystal spin systems, can gain a deeper insight on the multi-spin states

T_{45} : Category two of better knowledge of the decay dynamics would help in the construction of decoupling control schemes. Here authors use solid-state nuclear magnetic resonance techniques to experimentally **investigate (e&eb)** the decay of coherent multi-spin states in linear spin chains. Leveraging on the quasi-one-dimensional geometry of Fluorapatite crystal spin systems, can gain a deeper insight on the multi-spin states

T_{46} : Category three of better knowledge of the decay dynamics would help in the construction of decoupling control schemes. Here authors use solid-state nuclear magnetic resonance techniques to experimentally **investigate (e&eb)** the decay of coherent multi-spin states in linear spin chains. Leveraging on the quasi-one-dimensional geometry of Fluorapatite crystal spin systems, can gain a deeper insight on the multi-spin states

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$

$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$
$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$
$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$
$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}),t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}),t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}),t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$
$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\left[-(b''_{13})^{(1)}(G,t)\right] & \left[-(b''_{16})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\left[-(b''_{14})^{(1)}(G,t)\right] & \left[-(b''_{17})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\left[-(b''_{15})^{(1)}(G,t)\right] & \left[-(b''_{18})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,

2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt}=(a_{16})^{(2)}G_{17}-\begin{bmatrix}(a'_{16})^{(2)}&+(a''_{16})^{(2)}(T_{17},t)&+(a''_{13})^{(1,1,)}(T_{14},t)&+(a''_{20})^{(3,3,3)}(T_{21},t)\\+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)&+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)&+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)\\+(a''_{36})^{(7,7,7)}(T_{37},t)&+(a''_{40})^{(8,8,8)}(T_{41},t)&+(a''_{44})^{(9,9)}(T_{45},t)\end{bmatrix}G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt}=(a_{17})^{(2)}G_{16}-\begin{bmatrix}(a'_{17})^{(2)}&+(a''_{17})^{(2)}(T_{17},t)&+(a''_{14})^{(1,1,)}(T_{14},t)&+(a''_{21})^{(3,3,3)}(T_{21},t)\\+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)&+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)&+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)\\+(a''_{37})^{(7,7,7)}(T_{37},t)&+(a''_{41})^{(8,8,8)}(T_{41},t)&+(a''_{45})^{(9,9)}(T_{45},t)\end{bmatrix}G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt}=(a_{18})^{(2)}G_{17}-\begin{bmatrix}(a'_{18})^{(2)}&+(a''_{18})^{(2)}(T_{17},t)&+(a''_{15})^{(1,1,)}(T_{14},t)&+(a''_{22})^{(3,3,3)}(T_{21},t)\\+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)&+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)&+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)\\+(a''_{38})^{(7,7,7)}(T_{37},t)&+(a''_{42})^{(8,8,8)}(T_{41},t)&+(a''_{46})^{(9,9)}(T_{45},t)\end{bmatrix}G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt}=(b_{16})^{(2)}T_{17}-\begin{bmatrix}(b'_{16})^{(2)}&-(b''_{16})^{(2)}(G_{19},t)&-(b''_{13})^{(1,1,)}(G,t)&-(b''_{20})^{(3,3,3,)}(G_{23},t)\\-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)&-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)&-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)\\-(b''_{36})^{(7,7,7)}(G_{39},t)&-(b''_{40})^{(8,8,8)}(G_{43},t)&-(b''_{44})^{(9,9)}(G_{47},t)\end{bmatrix}T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt}=(b_{17})^{(2)}T_{16}-\begin{bmatrix}(b'_{17})^{(2)}&-(b''_{17})^{(2)}(G_{19},t)&-(b''_{14})^{(1,1,)}(G,t)&-(b''_{21})^{(3,3,3,)}(G_{23},t)\\-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)&-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)&-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)\\-(b''_{37})^{(7,7,7)}(G_{39},t)&-(b''_{41})^{(8,8,8)}(G_{43},t)&-(b''_{45})^{(9,9)}(G_{47},t)\end{bmatrix}T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad 73$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad 74$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad 76$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)} & \boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category $1, 2$ *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8\ ,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)}$ 103

$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)}$ 104

$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)}$ 105

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t}$ 107

$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t}$ 108

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1$ 110

$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1$ 111

Where we suppose

$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22$ 112

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - \ T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(QQ) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(RR) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$ 132

(SS) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(TT)

$\lim_{G\to\infty}(b_i'')^{(7)}((G_{39}),t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}((G_{39}),t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(UU) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(VV) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40,41,42$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\tilde{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\tilde{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad (169)$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that (170)

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that (172)

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that (174)

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(h) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ and\ (\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t} \; and \; (\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2 \; and \; \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37} \; and \; G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\frac{2}{\varepsilon_7}$$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)}$$ 267

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)}$$ 268

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}),(\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}),(\widetilde{T_{43}})\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(h) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)} G_{14}^{*}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)} G_{14}^{*}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)} T_{14}^{*}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)} T_{14}^{*}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)} G_{17}^{*}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)} G_{17}^{*}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)} T_{17}^{*}}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)} T_{17}^{*}}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)} G_{21}^{*}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)} G_{21}^{*}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)} T_{21}^{*}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)} T_{21}^{*}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

(586 A)

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTY EIGHT

Quantum Control Systems: Labor Omnia Vincit-Leges Humanae Nascuntur, Vivunt, Et Moriuntur

INTRODUCTION— Decay of spin coherences in one-dimensional spin systems G. Kaur et al

(1) Authors are then able to formulate an analytical model that **captures (e) the** key features of the decay.
(2) compare the decoherence behavior for different initial states of the spin chain and link their decay rate **to the** state characteristics, in particular their coherence and long-range correlation among spins.
(3) Experimental and theoretical study **shows that** the spin chains undergo a rich dynamics, with a slower decay rate than for the 3D case, and thus might be more **amenable(e&eb)** to decoupling techniques.
(4) While some recent works have looked at their decoherence and control schemes via dynamical-decoupling, the decay was usually **assumed to be(e) induced by** an uncorrelated bath, acting independently on each qubit, although this is often not the case in nature, especially for spatially close spins.
(5) In this paper, we experimentally and theoretically study the decay of such multi-qubit states **under the action** of a correlated spin bath.
(6) In particular, we are interested in investigating the decay rate **dependence on the** correlations in a multi-qubit spin state.
(7) We leverage the low dimensionality of the system studied – the linear coupling geometry **provided (e) by** nuclear spins in apatite crystals [8, 9] – to gain insight into both the many-body states created by the coherent Hamiltonian dynamics and their subsequent decay.
(8) We present a simple analytical model that captures the essential features of the multiqubit decays, and compares well with the experimental data. These results **will be (eb) helpful** in paving the way for the future design of schemes to mitigate the decay

ArXiv: quant-ph/0602014 v2 21 May 2006 HAMILTONIAN ENGINEERING FOR QUANTUM SYSTEMS Sonia G Schirmer

(9) Extending control to the quantum domain, i.e., to physical systems whose behavior is not governed by classical laws but dominated by quantum effects, has become an important area of research recently. It is also an essential prerequisite for the development of novel technologies such as

quantum information processing, as well as new applications in quantum optics, quantum electronics, or quantum chemistry. Choreographing the behavior of interacting quantum particles is a rather difficult task in general, for a variety of reasons, including the destructive effects of uncontrollable interactions with the environment and measurement backaction, both of which **lead to** loss of coherence, for instance.

(10) Yet, many problems in quantum optics, quantum electronics, atomic physics, molecular chemistry, and quantum computing can be **reduced (eb) to** quantum state or process engineering, and optimization of observables, which can be solved using Hamiltonian engineering techniques. In this paper we outline and compare different strategies for accomplishing this in various quantum settings.

CONTROL SYSTEM MODEL

A control system must obviously consist of a system to be controlled and a controller. In quantum control the former is always quantum-mechanical. The latter can be either quantum or classical, and it may seem natural to choose another quantum system as the controller. Indeed, this is useful for some applications, for instance in quantum optics [Yanagisawa and Kimura (2003)]. The main challenge in general though, and the focus of this paper, is control at the interface between the quantum world and the classical world we experience, i.e., control of systems that obey the laws of quantum physics using semi-classical sensors and actuators that interact with the quantum system but accept classical input (such as different settings of the classical control switches of the laboratory equipment) and return classical information (See Fig. 1).

This model is applicable to many quantum control settings. For example, the quantum system could be an ensemble of molecules involved in a chemical reaction, subject to laser pulses **produced by an** actuator consisting of a laser source and pulse shaping equipment, and a detector that might consist of a mass spectrometer to identify the reaction products. It could be a solid-state system such as an ensemble of quantum dots representing qubits, with actuators and sensors consisting of control electrodes and single-electron transistors, respectively.

It could be an ensemble of molecules with nuclear spins subject to actuators **generating** magnetic and radio-frequency fields, and sensors that detect the magnetization of the sample, etc.

SYSTEM DYNAMICS

In the setting defined above the state of the system can be represented either by a Hilbert space vector |Ψ(t)i, or more generally, a density operator ρ(t), i.e., a positive trace-one **operator acting on** the system's Hilbert space H. Although the full Hilbert spaces of most quantum systems are hardly finite or separable—even the Hilbert space of the hydrogen atom is not separable if the full continuum of ionized states is included, we are usually only interested **in controlling the** dynamics of a finite-dimensional subspace of the system's full Hilbert space (e.g., a subset of bound states of an atom), and full control of an infinite dimensional system is generally not possible in feedback controller

NOTATION

Module One

Authors are then able to formulate an analytical model that **captures (e) the** key features of the decay.

G_{13} : Category one of key features of the decay

G_{14} : Category two of key features of the decay

G_{15} : Category three of key features of the decay

T_{13} : Category one of analytical models

T_{14} : Category two of analytical models

T_{15} : Category three of analytical models

Module Two

compare the decoherence behavior for different initial states of the spin chain and link their decay rate **to the** state characteristics, in particular their coherence and long-range correlation among spins.

G_{16} : Category one of state characteristics, in particular their coherence and long-range correlation among spins

G_{17} : Category two of state characteristics, in particular their coherence and long-range correlation among spins

G_{18} : Category three of state characteristics, in particular their coherence and long-range correlation among spins

T_{16} : Category one of compare the decoherence behavior for different initial states of the spin chain and link their decay rate

T_{17} : Category two of compare the decoherence behavior for different initial states of the spin chain and link their decay rate

T_{18} : Category three of compare the decoherence behavior for different initial states of the spin chain and link their decay rate

Module three

Experimental and theoretical study **shows that** the spin chains undergo a rich dynamics, with a slower decay rate than for the 3D case, and thus might be more **amenable(e&eb)** to decoupling techniques.

G_{20} : Category one of Experimental and theoretical study **shows that** the spin chains undergo a rich dynamics, with a slower decay rate than for the 3D case, and thus might be more **amenable;** decoupling techniques

G_{21} : Category two of Experimental and theoretical study **shows that** the spin chains undergo a rich dynamics, with a slower decay rate than for the 3D case, and thus might be more **amenable;** decoupling techniques

G_{22} : Category three of Experimental and theoretical study **shows that** the spin chains undergo a rich dynamics, with a slower decay rate than for the 3D case, and thus might be more **amenable;** decoupling techniques

T_{20} : Category one of decoupling techniques ;Experimental and theoretical study **shows that** the spin chains undergo a rich dynamics, with a slower decay rate than for the 3D case, and thus might be more **amenable**

T_{21} : Category two of decoupling techniques; Experimental and theoretical study **shows that** the spin chains undergo a rich dynamics, with a slower decay rate than for the 3D case, and thus might be more **amenable**

T_{22} : Category three of decoupling techniques ;Experimental and theoretical study **shows that** the spin chains undergo a rich dynamics, with a slower decay rate than for the 3D case, and thus might be more

amenable

Module four

While some recent works have looked at their decoherence and control schemes via dynamical-decoupling, the decay was usually **assumed to be(e) induced by** an uncorrelated bath, acting independently on each qubit, although this is often not the case in nature, especially for spatially close spins.

G_{24} : Category one of uncorrelated bath, acting independently on each qubit, although this is often not the case in nature, especially for spatially close spins.

G_{25} : Category two of uncorrelated bath, acting independently on each qubit, although this is often not the case in nature, especially for spatially close spins.

G_{26} : Category three of uncorrelated bath, acting independently on each qubit, although this is often not the case in nature, especially for spatially close spins.

T_{24} : Category one of recent works have looked at their decoherence and control schemes via dynamical-decoupling, the decay was usually **assumed**

T_{25} : Category two of recent works have looked at their decoherence and control schemes via dynamical-decoupling, the decay was usually **assumed**

T_{26} : Category three of recent works have looked at their decoherence and control schemes via dynamical-decoupling, the decay was usually **assumed**

Module five

In this paper, we experimentally and theoretically study the decay of such multi-qubit states **under the action** of a correlated spin bath

G_{28} : Category one of decay of such multi-qubit states

G_{29} : Category two of decay of such multi-qubit states

G_{30} : Category three of decay of such multi-qubit states

T_{28} : Category one of correlated spin bath

T_{29} : Category two of correlated spin bath

T_{30} : Category three of correlated spin bath

Module six

In particular, we are interested in investigating the decay rate **dependence on the** correlations in a multi-qubit spin state.

G_{32} : Category one of decay rate; correlations in a multi-qubit spin state.

G_{33} : Category two of decay rate; correlations in a multi-qubit spin state.

G_{34} : Category three of decay rate; correlations in a multi-qubit spin state.

T_{32} : Category one of correlations in a multi-qubit spin state; decay rate

T_{33} : Category two of correlations in a multi-qubit spin state; decay rate

T_{34} : Category three of correlations in a multi-qubit spins state. ;decay rate

Module seven

We leverage the low dimensionality of the system studied – the linear coupling geometry **provided (e) by** nuclear spins in apatite crystals [8, 9] – to gain insight into both the many-body states created by the coherent Hamiltonian dynamics and their subsequent decay.

G_{36} : Category one of nuclear spins in apatite crystals [8, 9] – to gain insight into both the many-body states created by the coherent Hamiltonian dynamics and their subsequent decay

G_{37} : Category two of nuclear spins in apatite crystals [8, 9] – to gain insight into both the many-body states created by the coherent Hamiltonian dynamics and their subsequent decay

G_{38} : Category three of nuclear spins in apatite crystals [8, 9] – to gain insight into both the many-body states created by the coherent Hamiltonian dynamics and their subsequent decay

T_{36} : Category one of low dimensionality of the system studied – the linear coupling geometry

T_{37} : Category two of low dimensionality of the system studied – the linear coupling geometry

T_{38} : Category three of low dimensionality of the system studied – the linear coupling geometry

Module eight

We present a simple analytical model that captures the essential features of the multiqubit decays, and compares well with the experimental data. These results **will be (eb) helpful** in paving the way for the future design of schemes to mitigate the decay

G_{40} : Category one of model that captures the essential features of the multiqubit decays, and compares well with the experimental data.

G_{41} : Category two of model that captures the essential features of the multiqubit decays, and compares well with the experimental data.

G_{42} : Category three of model that captures the essential features of the multiqubit decays, and compares well with the experimental data.

T_{40} : Category one of paving the way for the future design of schemes to mitigate the decay

T_{41} : Category two of paving the way for the future design of schemes to mitigate the decay

T_{42} : Category three of paving the way for the future design of schemes to mitigate the decay

Module Nine

Extending control to the quantum domain, i.e., to physical systems whose behavior is not governed by classical laws but dominated by quantum effects, has become an important area of research recently. It is also an essential prerequisite for the development of novel technologies such as quantum information

processing, as well as new applications in quantum optics, quantum electronics, or quantum chemistry. Choreographing the behavior of interacting quantum particles is a rather difficult task in general, for a variety of reasons, including the destructive effects of uncontrollable interactions with the environment and measurement backaction, both of which **lead to** loss of coherence, for instance.

many problems in quantum optics, quantum electronics, atomic physics, molecular chemistry, and quantum computing can be **reduced (eb) to** quantum state or process engineering, and optimization of observables, which can be solved using Hamiltonian engineering techniques. In this paper we outline and compare different strategies for accomplishing this in various quantum settings.

G_{44} : Category one of control to the quantum domain, i.e., to physical systems whose behavior is not governed by classical laws but dominated by quantum effects; Choreographing the behavior of interacting quantum particles is a rather difficult task in general, for a variety of reasons, including the destructive effects of uncontrollable interactions with the environment and measurement backaction; many problems in quantum optics, quantum electronics, atomic physics, molecular chemistry, and quantum computing

G_{45} : Category two of control to the quantum domain, i.e., to physical systems whose behavior is not governed by classical laws but dominated by quantum effects; Choreographing the behavior of interacting quantum particles is a rather difficult task in general, for a variety of reasons, including the destructive effects of uncontrollable interactions with the environment and measurement backaction; many problems in quantum optics, quantum electronics, atomic physics, molecular chemistry, and quantum computing

G_{46} : Category three of control to the quantum domain, i.e., to physical systems whose behavior is not governed by classical laws but dominated by quantum effects; Choreographing the behavior of interacting quantum particles is a rather difficult task in general, for a variety of reasons, including the destructive effects of uncontrollable interactions with the environment and measurement backaction; many problems in quantum optics, quantum electronics, atomic physics, molecular chemistry, and quantum computing

T_{44} : Category one of quantum information processing, as well as new applications in quantum optics, quantum electronics, or quantum chemistry; loss of coherence, for instance; quantum state or process engineering, and optimization of observables, which can be solved using Hamiltonian engineering techniques

T_{45} : Category two of quantum information processing, as well as new applications in quantum optics, quantum electronics, or quantum chemistry; loss of coherence, for instance; quantum state or process engineering, and optimization of observables, which can be solved using Hamiltonian engineering techniques

T_{46} : Category three of quantum information processing, as well as new applications in quantum optics, quantum electronics, or quantum chemistry; loss of coherence, for instance; quantum state or process engineering, and optimization of observables, which can be solved using Hamiltonian engineering techniques

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{16})^{(2)} \; +(a''_{16})^{(2)}(T_{17},t) \; +(a''_{13})^{(1,1,)}(T_{14},t) \; +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) \; +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) \; +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) \; +(a''_{40})^{(8,8,8)}(T_{41},t) \; +(a''_{44})^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)} \; +(a''_{17})^{(2)}(T_{17},t) \; +(a''_{14})^{(1,1,)}(T_{14},t) \; +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) \; +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) \; +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) \; +(a''_{41})^{(8,8,8)}(T_{41},t) \; +(a''_{45})^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{18})^{(2)} \; +(a''_{18})^{(2)}(T_{17},t) \; +(a''_{15})^{(1,1,)}(T_{14},t) \; +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) \; +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) \; +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7)}(T_{37},t) \; +(a''_{42})^{(8,8,8)}(T_{41},t) \; +(a''_{46})^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for*

category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category $1, 2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a_{32}')^{(6)} & +(a_{32}'')^{(6)}(T_{33},t) & +(a_{28}'')^{(5,5,5)}(T_{29},t) & +(a_{24}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a_{33}')^{(6)} & +(a_{33}'')^{(6)}(T_{33},t) & +(a_{29}'')^{(5,5,5)}(T_{29},t) & +(a_{25}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a_{34}')^{(6)} & +(a_{34}'')^{(6)}(T_{33},t) & +(a_{30}'')^{(5,5,5)}(T_{29},t) & +(a_{26}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a_{32}'')^{(6)}(T_{33},t)$, $+(a_{33}'')^{(6)}(T_{33},t)$, $+(a_{34}'')^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a_{28}'')^{(5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a_{24}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a'_{42})^{(8)} & +(a''_{42})^{(8)}(T_{41},t) & +(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41},t)$, $+(a''_{41})^{(8)}(T_{41},t)$, $+(a''_{42})^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(WW) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(XX) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(YY) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(ZZ)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(AAA) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(BBB) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+\ (\hat{P}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+\ (\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\le\ \left(\hat{P}_{13}\right)^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\le\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\le\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 156

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^{0} + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^{0} + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^{0} + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^{0} + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(i) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\,((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)}=\frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)},(v_0)^{(7)}$:-

$$\text{For } 0<\boxed{(v_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(v_1)^{(7)}<(\bar{v}_1)^{(7)}$$

$$v^{(7)}(t)\geq\frac{(v_1)^{(7)}+(C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(v_1)^{(7)}-(v_0)^{(7)}}{(v_0)^{(7)}-(v_2)^{(7)}}}$$

it follows $(v_0)^{(7)}\leq v^{(7)}(t)\leq(v_1)^{(7)}$

In the same manner , we get 417

$$v^{(7)}(t)\leq\frac{(\bar{v}_1)^{(7)}+(\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{v}_1)^{(7)}-(v_0)^{(7)}}{(v_0)^{(7)}-(\bar{v}_2)^{(7)}}}$$

From which we deduce $(v_0)^{(7)}\leq v^{(7)}(t)\leq(\bar{v}_1)^{(7)}$

If $0<(v_1)^{(7)}<(v_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)}\leq\frac{(v_1)^{(7)}+(C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_2)^{(7)}\right)t\right]}}\leq v^{(7)}(t)\leq$$

$$\frac{(\bar{v}_1)^{(7)}+(\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}\leq(\bar{v}_1)^{(7)}$$

If $0<(v_1)^{(7)}\leq(\bar{v}_1)^{(7)}\leq\boxed{(v_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)}\leq v^{(7)}(t)\leq\frac{(\bar{v}_1)^{(7)}+(\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}\leq(v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)}\leq v^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{v^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(i) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523

A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTY NINE

Hamiltonian Quantum Systems: Nanos Gigantum Humeris Insidentes

INTRODUCTION—VARIABLES USED

Complete Control of Hamiltonian Quantum Systems: Engineering of Floquet Evolution G. Harel etal

(1) Authors propose a method for controlling the unitary evolution of quantum systems **by (e)** switching on and off alternatively two distinct constant perturbations.
(2) They show how to find appropriate switching times **in order (eb) to attain** any desired evolution. We suggest an experimental realization of our method in controlling the translational motion of cold atoms. © 1999 The American Physical SocietyURL: http://link.aps.org/doi/10.1103/PhysRevLett.82.1DOI: 10.1103/PhysRevLett.82.1

Lecture Notes in Physics Volume 575, 2001Decoherence, Pointer Engineering and Quantum State Protection A.R.R. Carvalho et al

(3) Authors show that motional states of trapped ions can be **protected(e)** against decoherence by generating, through transitions involving unstable electronic states, artificial reservoirs for the vibrational motion, which have the states to be protected as their pointer states.
(4) Exemplification of procedure with qubits, squeezed states, approximate phase eigenstates **and** superpositions of coherent states.
(5) Authors show that these states may be efficiently **protected (e)** against decoherence from thermal reservoirs and random fields.

Control Theory & Applications 2000-06 Energy-Based Stabilization of Forced Hamiltonian Systems and Its Application to Power Systems CHENG Daizhan et al

(6) The purpose of this paper is twofold. First of all, the stabilization of excitation control of power systems is considered. The system has been formulated **as a** forced Hamiltonian system with dissipation.
(7) The energy-based Lyapunov function is constructed **to (e&eb) investigate** the stability of the forced system.
(8) Secondly, the model of the generalized Hamiltonian systems is proposed, **which consists of** externally supplied energy, dissipation and internal energy source.
(9) The state space is described **as a** manifold equipped with a quadratic tensor field. Global coordinate free model of the generalized Hamiltonian systems is obtained. TM76

NOTATION

Module One

Authors propose a method for controlling the unitary evolution of quantum systems **by (e)** switching on and off alternatively two distinct constant perturbations

G_{13} : Category one of switching on and off alternatively two distinct constant perturbations
G_{14} : Category two of switching on and off alternatively two distinct constant perturbations
G_{15} : Category three of switching on and off alternatively two distinct constant perturbations
T_{13} : Category one of method for controlling the unitary evolution of quantum systems

T_{14} : Category two of method for controlling the unitary evolution of quantum systems

T_{15} : Category three of method for controlling the unitary evolution of quantum systems

Module Two

They show how to find appropriate switching times **in order (eb) to attain** any desired evolution. We suggest an experimental realization of our method in controlling the translational motion of cold atoms.

G_{16} : Category one of appropriate switching times

G_{17} : Category two of appropriate switching times

G_{18} : Category three of appropriate switching times

T_{16} : Category one of any desired evolution. We suggest an experimental realization of our method in controlling the translational motion of cold atoms

T_{17} : Category two of any desired evolution. We suggest an experimental realization of our method in controlling the translational motion of cold atoms

T_{18} : Category three of any desired evolution. We suggest an experimental realization of our method in controlling the translational motion of cold atoms

Module three

Authors show that motional states of trapped ions can be **protected(e)** against decoherence by generating, through transitions involving unstable electronic states, artificial reservoirs for the vibrational motion, which have the states to be protected as their pointer states

G_{20} : Category one of decoherence by generating, through transitions involving unstable electronic states, artificial reservoirs for the vibrational motion, which have the states to be protected as their pointer states

G_{21} : Category two of decoherence by generating, through transitions involving unstable electronic states, artificial reservoirs for the vibrational motion, which have the states to be protected as their pointer states

G_{22} : Category three of decoherence by generating, through transitions involving unstable electronic states, artificial reservoirs for the vibrational motion, which have the states to be protected as their pointer states

T_{20} : Category one of motional states of trapped ions

T_{21} : Category two of motional states of trapped ions

T_{22} : Category three of motional states of trapped ions

Module four

Exemplification of procedure with qubits, squeezed states, approximate phase eigenstates **and** superpositions of coherent states

G_{24} : Category one of qubits, squeezed states, approximate phase eigenstates

G_{25} : Category two of qubits, squeezed states, approximate phase eigenstates

G_{26} : Category three of qubits, squeezed states, approximate phase eigenstates

T_{24} : Category one of superpositions of coherent states

T_{25} : Category two of superpositions of coherent states

T_{26} : Category three of superpositions of coherent states

Module five

Authors show that these states may be efficiently **protected (e)** against decoherence from thermal reservoirs and random fields.

G_{28} : Category one of decoherence from thermal reservoirs and random fields.

G_{29} : Category two of decoherence from thermal reservoirs and random fields.

G_{30} : Category three of decoherence from thermal reservoirs and random fields.

T_{28} : Category one of qubits, squeezed states, approximate phase eigenstates **and** superpositions of coherent states

T_{29} : Category two of qubits, squeezed states, approximate phase eigenstates **and** superpositions of coherent states

T_{30} : Category three of qubits, squeezed states, approximate phase eigenstates **and** superpositions of coherent states

Module six

The purpose of this paper is twofold. First of all, the stabilization of excitation control of power systems is considered. The system has been formulated **as a** forced Hamiltonian system with dissipation.

G_{32} : Category one of forced Hamiltonian system with dissipation

G_{33} : Category two of forced Hamiltonian systems with dissipation

G_{34} : Category three of forced Hamiltonian systems with dissipation

T_{32} : Category one of stabilization of excitation control of power systems

T_{33} : Category two of stabilization of excitation control of power systems

T_{34} : Category three of stabilization of excitation control of power systems

Module seven

The energy-based Lyapunov function is constructed **to (e&eb) investigate** the stability of the forced system.

G_{36} : Category one of energy-based Lyapunov function is constructed

G_{37} : Category two of energy-based Lyapunov function is constructed

G_{38} : Category three of energy-based Lyapunov function is constructed

T_{36} : Category one of stability of the forced system

T_{37} : Category two of stability of the forced system

T_{38} : Category three of stability of the forced system

Module eight

Secondly, the model of the generalized Hamiltonian systems is proposed, **which consists of** externally supplied energy, dissipation and internal energy source.

G_{40} : Category one of externally supplied energy, dissipation and internal energy source.

G_{41} : Category two of externally supplied energy, dissipation and internal energy source.

G_{42} : Category three of externally supplied energy, dissipation and internal energy source.

T_{40} : Category one of generalized Hamiltonian systems

T_{41} : Category two of generalized Hamiltonian systems

T_{42} : Category three of generalized Hamiltonian systems

Module Nine

The state space is described **as a** manifold equipped with a quadratic tensor field. Global coordinate free model of the generalized Hamiltonian systems is obtained. TM76

G_{44} : Category one of manifold equipped with a quadratic tensor field. Global coordinate free model of the generalized Hamiltonian systems is obtained. TM76

G_{45} : Category two of manifold equipped with a quadratic tensor field. Global coordinate free model of the generalized Hamiltonian systems is obtained. TM76

G_{46} : Category three of manifold equipped with a quadratic tensor field. Global coordinate free model of the generalized Hamiltonian systems is obtained. TM76

T_{44} : Category one of state spaces

T_{45} : Category two of state spaces

T_{46} : Category three of state spaces

\

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$
$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$
$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}),t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)}\ \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{array}\right] G_{18} \quad 63$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for*

category 1, 2 and 3

$-(b_{13}'')^{(1,1,1,)}(G,t)$, $-(b_{14}'')^{(1,1,1,)}(G,t)$, $-(b_{15}'')^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b_{36}'')^{(7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7)}(G_{39},t)$ $-(b_{38}'')^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{46}'')^{(9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a_{24}')^{(4)} & +(a_{24}'')^{(4)}(T_{25},t) & +(a_{28}'')^{(5,5,)}(T_{29},t) & +(a_{32}'')^{(6,6,)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1)}(T_{14},t) & +(a_{16}'')^{(2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3)}(T_{21},t) \\ & +(a_{36}'')^{(7,7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a_{25}')^{(4)} & +(a_{25}'')^{(4)}(T_{25},t) & +(a_{29}'')^{(5,5,)}(T_{29},t) & +(a_{33}'')^{(6,6)}(T_{33},t) \\ & +(a_{14}'')^{(1,1,1,1)}(T_{14},t) & +(a_{17}'')^{(2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3)}(T_{21},t) \\ & +(a_{37}'')^{(7,7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a_{26}')^{(4)} & +(a_{26}'')^{(4)}(T_{25},t) & +(a_{30}'')^{(5,5,)}(T_{29},t) & +(a_{34}'')^{(6,6,)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1)}(T_{14},t) & +(a_{18}'')^{(2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3)}(T_{21},t) \\ & +(a_{38}'')^{(7,7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a_{24}'')^{(4)}(T_{25},t)$, $(a_{25}'')^{(4)}(T_{25},t)$, $(a_{26}'')^{(4)}(T_{25},t)$ *are first augmentation coefficients category* $1,2\ 3$

$+(a_{28}'')^{(5,5,)}(T_{29},t)$, $+(a_{29}'')^{(5,5,)}(T_{29},t)$, $+(a_{30}'')^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* $1,2$ *and* 3

$+(a_{32}'')^{(6,6,)}(T_{33},t)$, $+(a_{33}'')^{(6,6,)}(T_{33},t)$, $+(a_{34}'')^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* $1,2$ *and* 3

$+(a_{13}'')^{(1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category $1,2$ *and* 3

$+(a_{16}'')^{(2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category $1,2$ *and* 3

$+(a_{20}'')^{(3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category $1,2$ *and* 3

$+(a_{36}'')^{(7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category $1,2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a_{32}')^{(6)} & +(a_{32}'')^{(6)}(T_{33},t) & +(a_{28}'')^{(5,5,5)}(T_{29},t) & +(a_{24}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) & \\ +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a_{33}')^{(6)} & +(a_{33}'')^{(6)}(T_{33},t) & +(a_{29}'')^{(5,5,5)}(T_{29},t) & +(a_{25}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) & \\ +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a_{34}')^{(6)} & +(a_{34}'')^{(6)}(T_{33},t) & +(a_{30}'')^{(5,5,5)}(T_{29},t) & +(a_{26}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) & \\ +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{34} \quad (87)$$

$+(a_{32}'')^{(6)}(T_{33},t)$, $+(a_{33}'')^{(6)}(T_{33},t)$, $+(a_{34}'')^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a_{28}'')^{(5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a_{24}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{ccc} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\boxed{-(b_{44}'')^{(9)}(G_{47},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\boxed{-(b_{45}'')^{(9)}(G_{47},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\boxed{-(b_{46}'')^{(9)}(G_{47},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}} \ , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G \to \infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(CCC) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(DDD) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(EEE) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(FFF)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(GGG) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(HHH) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\leq\ \left(\hat{P}_{13}\right)^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\leq\ (\hat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\leq\ (\hat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\leq\ (\hat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^{0} + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^{0} + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^{0} + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^{0} + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(j) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18}. indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16}, G_{18} and G_{16}, G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18}. The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2:

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}\ and\ (\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40{,}41{,}42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 275

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(8)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\,((G_{43})(t),t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}\,(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}$, $if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)}$,

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}$, $if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}$, $if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}$,

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(\mathrm{T}_{17},t)+(a''_{17})^{(2)}(\mathrm{T}_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}\left((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)}\right)}\left[e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}\left((S_1)^{(3)} - (a'_{22})^{(3)}\right)}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}\left((R_1)^{(3)} - (b'_{22})^{(3)}\right)}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}\left((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)}\right)}\left[e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right) - (b''_{25})^{(4)}\left((G_{27}), t\right) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(\nu_1)^{(6)}, (\nu_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, \nu^{(6)}, u^{(6)}$: 350

By $(\nu_1)^{(6)} > 0$, $(\nu_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)},, (\bar{\nu}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, $\boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, $\boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, $\boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, $\boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $v^{(4)}$:- $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

For $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)},(v_0)^{(7)}$:-

$$\text{For } 0<\boxed{(v_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(v_1)^{(7)}<(\bar{v}_1)^{(7)}$$

$$v^{(7)}(t)\geq\frac{(v_1)^{(7)}+(C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(v_1)^{(7)}-(v_0)^{(7)}}{(v_0)^{(7)}-(v_2)^{(7)}}}$$

it follows $(v_0)^{(7)}\leq v^{(7)}(t)\leq(v_1)^{(7)}$

In the same manner , we get 417

$$v^{(7)}(t)\leq\frac{(\bar{v}_1)^{(7)}+(\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{v}_1)^{(7)}-(v_0)^{(7)}}{(v_0)^{(7)}-(\bar{v}_2)^{(7)}}}$$

From which we deduce $(v_0)^{(7)}\leq v^{(7)}(t)\leq(\bar{v}_1)^{(7)}$

If $0<(v_1)^{(7)}<(v_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)}\leq\frac{(v_1)^{(7)}+(C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_2)^{(7)}\right)t\right]}}\leq v^{(7)}(t)\leq$$

$$\frac{(\bar{v}_1)^{(7)}+(\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}\leq(\bar{v}_1)^{(7)}$$

If $0<(v_1)^{(7)}\leq(\bar{v}_1)^{(7)}\leq\boxed{(v_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)}\leq v^{(7)}(t)\leq\frac{(\bar{v}_1)^{(7)}+(\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}\leq(v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)}\leq v^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{v^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(j) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523

A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FORTY

Atomic Engineering: Cato Esse Quam Videri Bonus Male Bat

INTRODUCTION—VARIABLES USED

Report from the university:

(1) Researchers at the University of New South Wales have proposed a **new way to (e&eb) distinguish** between quantum bits that are placed only a few nanometres apart in a silicon chip, taking them a step closer to the construction of a large-scale quantum computer. Qubits are the basic building blocks of quantum computers - ultra-powerful devices that will offer enormous advantages for solving complex problems.

(2) Professor Michelle Simmons, leader of the research team, said a qubit based on the spin of an individual electron bound to a phosphorus atom within a silicon chip is one of the most promising systems **for (eb) building a** practical quantum computer, due to silicon's widespread use in the microelectronics industry.

(3) "However, to be able to couple electron-spins on single atom qubits, **it is necessary (e) that** the qubits need to be placed with atomic precision, within just a few tens of nanometres of each other," she says.

(4) This poses a technical problem in how to make them, and an operational problem in how to **control them** independently when they are so close together." In other words, the placement of qubits with atomic precision would **create(eb)** a problem of control when they are so close to each other

The UNSW team, in collaboration with theorists at Sandia National Laboratories in New Mexico, has found a solution to both these problems. Their study is published in the journal Nature Communications.

(5) In a significant feat of atomic engineering, they were able to read-out the spins of individual electrons **on (eb) a** cluster of phosphorus atoms that had been placed precisely in silicon. They also propose a new method for distinguishing between neighboring qubits that are only a few nanometres apart.

"It is a daunting challenge to rotate the spin of each qubit individually," says Holger Büch, lead author of the

new study.

(6) "If each electron spin-qubit is hosted by a single phosphorus atom, every time you try to rotate one qubit, then there **shall be (eb) production of** all the neighbouring qubits will rotate at the same time – and quantum computation will not work."
(7) "But if each electron is hosted by a different number of phosphorus atoms, then the qubits **will(e&eb) respond** to different electromagnetic fields – and each qubit can be distinguished from the others around it," he says.

The UNSW team is part of the Australian Centre of Excellence for Quantum Computation and Communication Technology, a world-leading research centre headquartered in Sydney, Australia.

(8) This is an elegant and satisfying piece of work," says Professor Simmons, centre director and Mr Büch's PhD supervisor. "This first demonstration that **we can (eb) maintains** long spin lifetimes of electrons on multi-donor systems are very powerful. It offers a new method for addressing individual qubits, putting us one step closer to realising a practical, large-scale quantum computer."
(9) To make the tiny device, the researchers deposited a layer of hydrogen on a silicon wafer and used a scanning tunnelling microscope **to create** a pattern on the surface in an ultra-high vacuum.

NOTATION

Module One

Researchers at the University of New South Wales have proposed a **new way to (e&eb) distinguish** between quantum bits that are placed only a few nanometres apart in a silicon chip, taking them a step closer to the construction of a large-scale quantum computer. Qubits are the basic building blocks of quantum computers - ultra-powerful devices that will offer enormous advantages for solving complex problems.

G_{13} : Category one of **new way to (e&eb) distinguish** between quantum bits that are placed only a few nanometres apart in a silicon chip, taking them a step closer to the construction of a large-scale quantum computer. Qubits

G_{14} : Category two of **new way to (e&eb) distinguishes** between quantum bits that are placed only a few nanometres apart in a silicon chip, taking them a step closer to the construction of a large-scale quantum computer. Qubits

G_{15} : Category three of **new way to (e&eb) distinguishes** between quantum bits that are placed only a few nanometres apart in a silicon chip, taking them a step closer to the construction of a large-scale quantum computer. Qubits

T_{13} : Category one of quantum computers - ultra-powerful devices that will offer enormous advantages for solving complex problems.

T_{14} : Category two of quantum computers - ultra-powerful devices that will offer enormous advantages for solving complex problems.

T_{15} : Category three of quantum computers - ultra-powerful devices that will offer enormous advantages for solving complex problems.

Module Two

"However, to be able to couple electron-spins on single atom qubits, **it is necessary (e) that** the qubits need to be placed with atomic precision, within just a few tens of nanometres of each other," she says.

G_{16} : Category one of qubits need to be placed with atomic precision, within just a few tens of nanometres of each other,

G_{17} : Category two of qubits need to be placed with atomic precision, within just a few tens of nanometres of each other,

G_{18} : Category three of qubits need to be placed with atomic precision, within just a few tens of nanometres of each other,

T_{16} : Category one of couplings of electron-spins on single atom qubits

T_{17} : Category two of couplings of electron-spins on single atom qubits

T_{18} : Category three of couplings of electron-spins on single atom qubits

Module three

This poses a technical problem in how to make them, and an operational problem in how to **control them** independently when they are so close together." In other words, the placement of qubits with atomic precision would **create(eb)** a problem of control when they are so close to each other

G_{20} : Category one of placement of qubits with atomic precision

G_{21} : Category two of placement of qubits with atomic precision

G_{22} : Category three of placement of qubits with atomic precision

T_{20} : Category one of problem of control when they are so close to each other

T_{21} : Category two of problem of control when they are so close to each other

T_{22} : Category three of problem of control when they are so close to each other

Module four

In a significant feat of atomic engineering, they were able to read-out the spins of individual electrons **on (eb) a** cluster of phosphorus atoms that had been placed precisely in silicon. They also propose a new method for distinguishing between neighboring qubits that are only a few nanometres apart.

G_{24} : Category one of read-out the spins of individual electrons

G_{25} : Category two of read-out the spins of individual electrons

G_{26} : Category three of read-out the spins of individual electrons

T_{24} : Category one of cluster of phosphorus atoms that had been placed precisely in silicon. They also propose a new method for distinguishing between neighboring qubits that are only a few nanometres apart.

T_{25} : Category two of cluster of phosphorus atoms that had been placed precisely in silicon. They also propose a new method for distinguishing between neighboring qubits that are only a few nanometres apart.

T_{26} : Category three of cluster of phosphorus atoms that had been placed precisely in silicon. They also propose a new method for distinguishing between neighboring qubits that are only a few nanometres apart.

Module five

Professor Michelle Simmons, leader of the research team, said a qubit based on the spin of an individual electron bound to a phosphorus atom within a silicon chip is one of the most promising systems **for (eb) building a** practical quantum computer, due to silicon's widespread use in the microelectronics industry.

G_{28} : Category one of qubit based on the spin of an individual electron bound to a phosphorus atom within a silicon chip is one of the most promising systems

G_{29} : Category two of qubit based on the spin of an individual electron bound to a phosphorus atom within a silicon chip is one of the most promising systems

G_{30} : Category three of qubit based on the spin of an individual electron bound to a phosphorus atom within a silicon chip is one of the most promising systems

T_{28} : Category one of practical quantum computer, due to silicon's widespread use in the microelectronics industry

T_{29} : Category two of practical quantum computer, due to silicon's widespread use in the microelectronics industry

T_{30} : Category three of practical quantum computer, due to silicon's widespread use in the microelectronics industry

Module six

"If each electron spin-qubit is hosted by a single phosphorus atom, every time you try to rotate one qubit, then there **shall be (eb) production of** all the neighbouring qubits will rotate at the same time – and quantum computation will not work."

G_{32} : Category one of each electron spin-qubit is hosted by a single phosphorus atom, every time you try to rotate one qubit

G_{33} : Category two of each electron spin-qubit is hosted by a single phosphorus atom, every time you try to rotate one qubit

G_{34} : Category three of each electron spin-qubit is hosted by a single phosphorus atom, every time you try to rotate one qubit

T_{32} : Category one of neighbouring qubits will rotate at the same time – and quantum computation will not work."

T_{33} : Category two of neighbouring qubits will rotate at the same time – and quantum computation will not work."

T_{34} : Category three of neighbouring qubits will rotate at the same time – and quantum computation will not work."

Module seven

"But if each electron is hosted by a different number of phosphorus atoms, then the qubits **will(e&eb) respond** to different electromagnetic fields – and each qubit can be distinguished from the others around it," he says.

G_{36} : Category one of electrons are hosted by a different number of phosphorus atoms, then the qubits; different electromagnetic fields – and each qubit can be distinguished from the others around it

G_{37} : Category two of electrons are hosted by a different number of phosphorus atoms, then the qubits; different electromagnetic fields – and each qubit can be distinguished from the others around it

G_{38} : Category three of electrons are hosted by a different number of phosphorus atoms, then the qubits ; different electromagnetic fields – and each qubit can be distinguished from the others around it

T_{36} : Category one of different electromagnetic fields – and each qubit can be distinguished from the others around it; electrons are hosted by a different number of phosphorus atoms, then the qubits

T_{37} : Category two of different electromagnetic fields – and each qubit can be distinguished from the others around it; electrons are hosted by a different number of phosphorus atoms, then the qubits

T_{38} : Category three of different electromagnetic fields – and each qubit can be distinguished from the others around it; electrons are hosted by a different number of phosphorus atoms, then the qubits

Module eight

This is an elegant and satisfying piece of work," says Professor Simmons, centre director and Mr Büch's PhD supervisor. "This first demonstrations that **we can (eb) maintains** long spin lifetimes of electrons on multi-donor systems are very powerful and offers a new method for addressing individual qubits, putting us one step closer to realising a practical, large-scale quantum computer."

G_{40} : Category one of demonstrations that **we can (eb) maintains** long spin lifetimes of electrons on multi-donor systems are very powerful and offers a new method for addressing individual qubits

G_{41} : Category two of demonstrations that **we can (eb) maintains** long spin lifetimes of electrons on multi-donor systems are very powerful and offers a new method for addressing individual qubits

G_{42} : Category three of demonstrations that **we can (eb) maintains** long spin lifetimes of electrons on multi-donor systems are very powerful and offers a new method for addressing individual qubits

T_{40} : Category one of realising a practical, large-scale quantum computer

T_{41} : Category two of realising a practical, large-scale quantum computer

T_{42} : Category three of realising a practical, large-scale quantum computer

Module Nine

To make the tiny device, the researchers deposited a layer of hydrogen on a silicon wafer and used a scanning tunnelling microscope **to create** a pattern on the surface in an ultra-high vacuum

G_{44} : Category one of researchers deposited a layer of hydrogen on a silicon wafer and used a scanning tunnelling microscope

G_{45} : Category two of researchers deposited a layer of hydrogen on a silicon wafer and used a scanning tunnelling microscope

G_{46} : Category three of researchers deposited a layer of hydrogen on a silicon wafer and used a scanning

tunnelling microscope

T_{44} : Category one of pattern on the surface in an ultra-high vacuum

T_{45} : Category two of pattern on the surface in an ultra-high vacuum

T_{46} : Category three of pattern on the surface in an ultra-high vacuum

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}),t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}), t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}), t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

59

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

60

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

61

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16}$$

62

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17}$$

63

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18}$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a_{44}'')^{(9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b_{16}')^{(2)} & -(b_{16}'')^{(2)}(G_{19},t) & -(b_{13}'')^{(1,1,)}(G,t) & -(b_{20}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{24}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{28}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{32}'')^{(6,6,6,6,6)}(G_{35},t) \\ -(b_{36}'')^{(7,7,7)}(G_{39},t) & -(b_{40}'')^{(8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b_{17}')^{(2)} & -(b_{17}'')^{(2)}(G_{19},t) & -(b_{14}'')^{(1,1,)}(G,t) & -(b_{21}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{25}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{29}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{33}'')^{(6,6,6,6,6)}(G_{35},t) \\ -(b_{37}'')^{(7,7,7)}(G_{39},t) & -(b_{41}'')^{(8,8,8)}(G_{43},t) & -(b_{45}'')^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b_{18}')^{(2)} & -(b_{18}'')^{(2)}(G_{19},t) & -(b_{15}'')^{(1,1,)}(G,t) & -(b_{22}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{26}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{30}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{34}'')^{(6,6,6,6,6)}(G_{35},t) \\ -(b_{38}'')^{(7,7,7)}(G_{39},t) & -(b_{42}'')^{(8,8,8)}(G_{43},t) & -(b_{46}'')^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad 66$$

where $-(b_{16}'')^{(2)}(G_{19},t)$, $-(b_{17}'')^{(2)}(G_{19},t)$, $-(b_{18}'')^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b_{13}'')^{(1,1,)}(G,t)$, $-(b_{14}'')^{(1,1,)}(G,t)$, $-(b_{15}'')^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b_{20}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b_{24}'')^{(4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b_{28}'')^{(5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b_{32}'')^{(6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt}=(a_{20})^{(3)}G_{21}-\left[\begin{array}{ccc}(a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}\\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}\end{array}\right]G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt}=(a_{21})^{(3)}G_{20}-\left[\begin{array}{ccc}(a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}\\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}\end{array}\right]G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt}=(a_{22})^{(3)}G_{21}-\left[\begin{array}{ccc}(a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}\\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}\end{array}\right]G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt}=(b_{20})^{(3)}T_{21}-\left[\begin{array}{ccc}(b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,1,)}(G,t)}\\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}\end{array}\right]T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1,2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1,2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1,2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category $1,2$ *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category $1,2$ *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$

are sixth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$

are seventh detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$

are eighth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt}=(a_{28})^{(5)}G_{29}-\left[\begin{array}{ccc}(a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt}=(a_{29})^{(5)}G_{28}-\left[\begin{array}{ccc}(a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt}=(a_{30})^{(5)}G_{29}-\left[\begin{array}{ccc}(a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt}=(b_{28})^{(5)}T_{29}-\left[\begin{array}{ccc}(b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}\end{array}\right]T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)} & +(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition

coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37},t) & +(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37},t) & +(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37},t) & +(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a''_{36})^{(7)}(T_{37},t)$, $(a''_{37})^{(7)}(T_{37},t)$, $(a''_{38})^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation

coefficient for 1,2,3

$\frac{dT_{36}}{dt} =$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$\frac{dT_{37}}{dt} =$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$\frac{dT_{38}}{dt} =$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

96

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \begin{bmatrix} (a'_{45})^{(9)} \boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \begin{bmatrix} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - \begin{bmatrix} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(\mathrm{p_i})^{(2)}$, $(\mathrm{r_i})^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}\left((G_{19}), t\right) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ \ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ \ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(III) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(JJJ) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(KKK) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(LLL)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(MMM) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(NNN) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and \ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{16}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\Big] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{20}(s_{(20)})\Big] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\Big] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\Big] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(k) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : (\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying 193
Equations into itself

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$sup_i\{max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}ds_{(16)} +$

$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0$ for $\mathrm{t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$:$\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t} \; and \; (\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2 \; and \; \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37} \; and \; G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \ \boldsymbol{if}\ (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \boldsymbol{if}\ (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

and $\boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \qquad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \qquad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \qquad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(k) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$


$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$


And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.


SECTION FORTY ONE

Nature Nanotechnology: Perfer Et Obdura; Dolor Hic Tibi Proderit Olim

INTRODUCTION—VARIABLES USED







See enlarged images

(1) This was then exposed to phosphine gas and annealed at 350 degrees so phosphorus atoms **became (=)** incorporated precisely into the silicon. The device was **then (eb) buried in** another layer of silicon.

(2) In a quantum computer information is stored in the spin, or magnetic orientation, of an electron. This spin can not only be in two states – up and down – just as in a classical computer. It can also **be (eb) i**n a combination of both states at the same time, allowing exponentially larger amounts of information to be stored and processed in parallel. Read more at: http://phys.org/news/2013-06-method-distinguish-neighboring-quantum-bits.html#jCp

(3) A controllable transistor engineered from a single phosphorus atom has been developed by researchers at the University of New South Wales, Purdue University and the University of Melbourne. The atom, shown here in the center of an image from a computer model, **sits (eb) in a** channel in a silicon crystal.

(4) The atomic-sized transistor and wires **might (eb) allow** researchers to control gated qubits of information in future quantum computers. (Purdue University image) video. Please see enlarged images. Constricted because of spatial constraints.

(5) Although single atoms serving as transistors have been observed before, this is the first time a single-atom transistor has been controllably engineered with atomic precision. The structure even has markers **that allow** researchers to attach contacts and apply a voltage, says Martin Fuechsle, a researcher at the University of New South Wales and lead author on the journal paper.

(6) "The thing that is unique about what we have done is that we have, with atomic precision, positioned this individual atom within our device," Fuechsle says. Simmons says this control is the key step **in making** a single-atom device.

(7) "By achieving the placement of a single atom, we have, at the same time, developed a technique that will **allow us to** be able to place several of these single-atom devices towards the goal of a developing a scalable system."

(8) The single-atom transistor **could lead the way** to building a quantum computer that works by controlling the electrons and thereby the quantum information, or qubits. Some scientists, however,

have doubts that such a device can ever be built. At the start of 2012. Professor Michelle Simmons' lab announced it had created 1-atom tall and 4-atom-wide nanowires.

(9) They embedded a wire **in** crystalline silicon **to (eb)** isolate the dopant atoms from surfaces and interfaces that caused this deactivation leading to highly conductive wires, and this is what happened.

NOTATION

Module One

Deposited a layer of hydrogen on a silicon wafer and used a scanning tunnelling microscope **to create** a pattern on the surface in an ultra-high vacuum exposed to phosphine gas and annealed at 350 degrees so phosphorus atoms **became incorporated** precisely into the silicon. The device was **then (eb) buried in** another layer of silicon.

G_{13} : Category one of **incorporated** precisely into the silicon. The device was **then (eb) buried in** another layer of silicon.

G_{14} : Category two of incorporated precisely into the silicon. The device was **then (eb) buried in** another layer of silicon.

G_{15} : Category three of incorporated precisely into the silicon. The device was **then (eb) buried in** another layer of silicon.

T_{13} : Category one of deposited a layer of hydrogen on a silicon wafer and used a scanning tunnelling microscope **to create** a pattern on the surface in an ultra-high vacuum exposed to phosphine gas and annealed at 350 degrees so phosphorus atoms

T_{14} : Category two of deposited a layer of hydrogen on a silicon wafer and used a scanning tunnelling microscope **to create** a pattern on the surface in an ultra-high vacuum exposed to phosphine gas and annealed at 350 degrees so phosphorus atoms

T_{15} : Category three of deposited a layer of hydrogen on a silicon wafer and used a scanning tunnelling microscope **to create** a pattern on the surface in an ultra-high vacuum exposed to phosphine gas and annealed at 350 degrees so phosphorus atoms

Module Two

Quantum computer information is stored in the spin, or magnetic orientation, of an electron. This spin can not only be in two states – up and down – just as in a classical computer which can also **be (eb) i**n a combination of both states at the same time, allowing exponentially larger amounts of information to be stored and processed in parallel. Read more at: http://phys.org/news/2013-06-method-distinguish-neighboring-quantum-bits.html#jCp

G_{16} : Category one of quantum computer information is stored in the spin, or magnetic orientation, of an electron. This spin can not only be in two states – up and down – just as in a classical computer

G_{17} : Category two of quantum computer information is stored in the spin, or magnetic orientation, of an electron. This spin can not only be in two states – up and down – just as in a classical computer

G_{18} : Category three of quantum computer information is stored in the spin, or magnetic orientation, of an electron. This spin can not only be in two states – up and down – just as in a classical computer

T_{16} : Category one of combination of both states at the same time, allowing exponentially larger amounts of information to be stored and processed in parallel

T_{17} : Category two of combination of both states at the same time, allowing exponentially larger amounts of information to be stored and processed in parallel

T_{18} : Category three of combination of both states at the same time, allowing exponentially larger amounts of information to be stored and processed in parallel

Module three

A controllable transistor engineered from a single phosphorus atom has been developed by researchers at the University of New South Wales, Purdue University and the University of Melbourne. The atom, in the from a computer model, **sits (eb) in a** channel in a silicon crystal.

G_{20} : Category one of controllable transistor engineered from a single phosphorus atom

G_{21} : Category two of controllable transistor engineered from a single phosphorus atom

G_{22} : Category three of controllable transistor engineered from a single phosphorus atom

T_{20} : Category one of channel in a silicon crystal.

T_{21} : Category two of channel in a silicon crystal.

T_{22} : Category three of channel in a silicon crystal.

Module four

The atomic-sized transistor and wires **might (eb) allow** researchers to control gated qubits of information in future quantum computers. (Purdue University image) video. Please see enlarged images. Constricted because of spatial constraints.

G_{24} : Category one of atomic-sized transistor and wires

G_{25} : Category two of atomic-sized transistor and wires

G_{26} : Category three of atomic-sized transistor and wires

T_{24} : Category one of researchers to control gated qubits of information in future quantum computers

T_{25} : Category two of researchers to control gated qubits of information in future quantum computers

T_{26} : Category three of researchers to control gated qubits of information in future quantum computers

Module five

Although single atoms serving as transistors have been observed before, this is the first time a single-atom transistor has been controllably engineered with atomic precision. The structure even has markers **that allow** researchers to attach contacts and apply a voltage, says Martin Fuechsle, a researcher at the University of New South Wales and lead author on the journal paper

G_{28} : Category one of single atoms serving as transistors have been observed before, this is the first time a single-atom transistor has been controllably engineered with atomic precision, the structure even has markers

G_{29} : Category two of single atoms serving as transistors have been observed before, this is the first time a single-atom transistor has been controllably engineered with atomic precision, the structure even has

markers

G_{30} : Category three of single atoms serving as transistors have been observed before, this is the first time a single-atom transistor has been controllably engineered with atomic precision, the structure even has markers

T_{28} : Category one of researchers to attach contacts and apply a voltage

T_{29} : Category two of researchers to attach contacts and apply a voltage

T_{30} : Category three of researchers to attach contacts and apply a voltage

Module six

"The thing that is unique about what we have done is that we have, with atomic precision, positioned this individual atom within our device," Fuechsle says. Simmons says this quantum control is the key step **in making** a single-atom device.

G_{32} : Category one of quantum control

G_{33} : Category two of quantum control

G_{34} : Category three of quantum control

T_{32} : Category one of single-atom device.

T_{33} : Category two of single-atom device.

T_{34} : Category three of single-atom device.

Module seven

"By achieving the placement of a single atom, we have, at the same time, developed a technique that will **allow us to** be able to place several of these single-atom devices towards the goal of a developing a scalable system."

G_{36} : Category one of achieving the placement of a single atom, at the same time has developed a technique

G_{37} : Category two of achieving the placement of a single atom, at the same time has developed a technique

G_{38} : Category three of achieving the placement of a single atom, at the same time has developed a technique

T_{36} : Category one of abilities to place several of these single-atom devices towards the goal of a developing a scalable system.

T_{37} : Category two of abilities to place several of these single-atom devices towards the goal of a developing a scalable system.

T_{38} : Category three of abilities to place several of these single-atom devices towards the goal of a

developing a scalable system.

Module eight

The single-atom transistor **could lead the way** to building a quantum computer that works by controlling the electrons and thereby the quantum information, or qubits. Some scientists, however, have doubts that such a device can ever be built. At the start of 2012. Professor Michelle Simmons' lab announced it had created 1-atom tall and 4-atom-wide nanowires.

G_{40} : Category one of single-atom transistor

G_{41} : Category two of single-atom transistor

G_{42} : Category three of single-atom transistor

T_{40} : Category one of building a quantum computer that works by controlling the electrons and thereby the quantum information, or qubits

T_{41} : Category two of building a quantum computer that works by controlling the electrons and thereby the quantum information, or qubits

T_{42} : Category three of building a quantum computer that works by controlling the electrons and thereby the quantum information, or qubits

Module Nine

They embedded a wire **in** crystalline silicon **to (eb)** isolate the dopant atoms from surfaces and interfaces that caused this deactivation leading to highly conductive wires, and this is what happened.

G_{44} : Category one of embedded a wire **in** crystalline silicon

G_{45} : Category two of embedded a wire **in** crystalline silicon

G_{46} : Category three of embedded a wire **in** crystalline silicon

T_{44} : Category one of isolate the dopant atoms from surfaces and interfaces that caused this deactivation leading to highly conductive wires, and this is what happened

T_{45} : Category two of isolate the dopant atoms from surfaces and interfaces that caused this deactivation leading to highly conductive wires, and this is what happened

T_{46} : Category three of isolate the dopant atoms from surfaces and interfaces that caused this deactivation leading to highly conductive wires, and this is what happened

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$
$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$
$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**
$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)}\ \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{array}\right] G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{array}\right] G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{array}\right] G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for*

category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category $1, 2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a_{32}')^{(6)} & +(a_{32}'')^{(6)}(T_{33},t) & +(a_{28}'')^{(5,5,5)}(T_{29},t) & +(a_{24}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a_{33}')^{(6)} & +(a_{33}'')^{(6)}(T_{33},t) & +(a_{29}'')^{(5,5,5)}(T_{29},t) & +(a_{25}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a_{34}')^{(6)} & +(a_{34}'')^{(6)}(T_{33},t) & +(a_{30}'')^{(5,5,5)}(T_{29},t) & +(a_{26}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a_{32}'')^{(6)}(T_{33},t)$, $+(a_{33}'')^{(6)}(T_{33},t)$, $+(a_{34}'')^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a_{28}'')^{(5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a_{24}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{c} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{c} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(OOO) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(PPP) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(QQQ) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(RRR)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(SSS) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(TTT) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\le\ (\hat{P}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\le\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\le\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(l) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18}. indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16}, G_{18} and G_{16}, G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18}. The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2:

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

208

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)}$$

209

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)}$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),\left((G_{23})^{(2)},(T_{23})^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

214

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\;(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\,((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})},\widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})},\widetilde{(T_{31})}\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$\left|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{36})^{(7)}\left|G_{37}^{(1)}-G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a''_{36})^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\leq \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}$, $if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)}$,

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}$, $if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}$, $if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}$,

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, ***if*** $(u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(\nu_1)^{(6)}, (\nu_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, \nu^{(6)}, u^{(6)}$: 350

By $(\nu_1)^{(6)} > 0$, $(\nu_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)},, (\bar{\nu}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21}, t)\right) - (a_{21}'')^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $v^{(4)}$:- $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

For $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)},(v_0)^{(7)}$:-

$$\text{For } 0<\boxed{(v_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(v_1)^{(7)}<(\bar{v}_1)^{(7)}$$

$$v^{(7)}(t)\geq\frac{(v_1)^{(7)}+(C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(v_1)^{(7)}-(v_0)^{(7)}}{(v_0)^{(7)}-(v_2)^{(7)}}}$$

it follows $(v_0)^{(7)}\leq v^{(7)}(t)\leq(v_1)^{(7)}$

In the same manner , we get 417

$$v^{(7)}(t)\leq\frac{(\bar{v}_1)^{(7)}+(\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{v}_1)^{(7)}-(v_0)^{(7)}}{(v_0)^{(7)}-(\bar{v}_2)^{(7)}}}$$

From which we deduce $(v_0)^{(7)}\leq v^{(7)}(t)\leq(\bar{v}_1)^{(7)}$

If $0<(v_1)^{(7)}<(v_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)}\leq\frac{(v_1)^{(7)}+(C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_2)^{(7)}\right)t\right]}}\leq v^{(7)}(t)\leq$$

$$\frac{(\bar{v}_1)^{(7)}+(\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}\leq(\bar{v}_1)^{(7)}$$

If $0<(v_1)^{(7)}\leq(\bar{v}_1)^{(7)}\leq\boxed{(v_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)}\leq v^{(7)}(t)\leq\frac{(\bar{v}_1)^{(7)}+(\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}\leq(v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)}\leq v^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{v^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(l) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523

A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^{*}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^{*}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}((G_{31})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FORTY TWO

A Single-Atom Transistor: Exitus Acta Probat Errare Humanum Est

INTRODUCTION—VARIABLES USED

POSTED BY BRIAN WANG AT 1/11/2013

(1) Perspective STM image of the device, in which the hydrogen-desorbed regions defining source (S) and drain (D) leads and two gates (G1, G2) appear raised **due (e) to the** increased tunnelling current through the silicon dangling bond states that were created.

(2) Upon subsequent dosing with phosphine, these regions **form** highly phosphorus-doped co-planar transport electrodes of monatomic height, which are registered to a single phosphorus atom in the centre of the device.

(3) Several atomic steps running across the Si (100) surface are also visible. b, Close-up of the inner device area (dashed box in a), where the central bright protrusion is the silicon atom, **which is ejected** when a single phosphorus atom incorporates into the surface.

(4) Schematic of the chemical reaction to deterministically **incorporate a** single phosphorus atom into the surface

(5) Saturation dosing of a three-dimer patch (I) at room temperature (RT) followed by annealing to 350 °C **allows** successive dissociation of PH3 (II–IV) and subsequent incorporation of a single phosphorus atom in the surface layer, ejecting a silicon adatom in the process (V). **POSTED BY BRIAN WANG AT 1/11/2013**

(6) The key to making these wires was combining scanning tunneling microscopy, a technique to image and manipulate individual atoms, **with (e&eb)** molecular beam epitaxy, a way of growing

perfect crystals.

(7) It gave us great precision in all three dimensions, and when combined with a high density of the dopant atoms, **allowed us to create** these highly conductive nanowires.

(8) The Kane quantum computer is a proposal for a scalable quantum computer proposed by Bruce Kane in 1998, then at the University of New South Wales. Often thought of as a hybrid between quantum dot **and** NMR quantum computers, the Kane computer **is based (e) on** an array of individual phosphorus donor atoms embedded in a pure silicon lattice. Both the nuclear spins of the donors and the spins of the donor electrons participate in the computation. The original proposal calls for phosphorus donors to be placed in an array with a spacing of 20 nm, approximately 20 nm below the surface. An insulating oxide layer is grown on top of the silicon. Metal A gates are deposited on the oxide above each donor, and J gates between adjacent donors.

(9) The phosphorus donors are isotropically pure 31P, which have a nuclear spin of 1/2. The silicon substrate is isotropically pure 28Si which has nuclear spin 0. Using the nuclear spin of the P donors as a method to encode qubits has two major advantages. Firstly, the state **has (e) an** extremely long decoherence time, perhaps on the order of 1018 seconds at millikelvin temperatures.

NOTATION

Module One

Perspective STM image of the device, in which the hydrogen-desorbed regions defining source (S) and drain (D) leads and two gates (G1, G2) appear raised **due (e) to the** increased tunnelling current through the silicon dangling bond states that were created.

G_{13} : Category one of increased tunnelling current through the silicon dangling bond states that were created.

G_{14} : Category two of increased tunnelling current through the silicon dangling bond states that were created.

G_{15} : Category three of increased tunnelling current through the silicon dangling bond states that were created.

T_{13} : Category one of Perspective STM image of the device, in which the hydrogen-desorbed regions defining source (S) and drain (D) leads and two gates (G1, G2) appear raised

T_{14} : Category two of Perspective STM image of the device, in which the hydrogen-desorbed regions defining source (S) and drain (D) leads and two gates (G1, G2) appear raised

T_{15} : Category three of Perspective STM image of the device, in which the hydrogen-desorbed regions defining source (S) and drain (D) leads and two gates (G1, G2) appear raised

Module Two

Upon subsequent dosing with phosphine, these regions **form** highly phosphorus-doped co-planar transport electrodes of monatomic height, which are registered to a single phosphorus atom in the centre of the device.

G_{16} : Category one of subsequent dosing with phosphine, these regions

G_{17} : Category two of subsequent dosing with phosphine, these regions

G_{18} : Category three of subsequent dosing with phosphine, these regions

T_{16} : Category one of highly phosphorus-doped co-planar transport electrodes of monatomic height, which

are registered to a single phosphorus atom in the centre of the device

T_{17} : Category two of highly phosphorus-doped co-planar transport electrodes of monatomic height, which are registered to a single phosphorus atom in the centre of the device

T_{18} : Category three of highly phosphorus-doped co-planar transport electrodes of monatomic height, which are registered to a single phosphorus atom in the centre of the device

Module three

Several atomic steps running across the Si (100) surface are also visible. b, Close-up of the inner device area (dashed box in a), where the central bright protrusion is the silicon atom, **which is ejected** when a single phosphorus atom incorporates into the surface.

G_{20} : Category one of Several atomic steps running across the Si (100) surface are also visible. b, Close-up of the inner device area (dashed box in a), where the central bright protrusion is the silicon atom

G_{21} : Category two of Several atomic steps running across the Si (100) surface are also visible. b, Close-up of the inner device area (dashed box in a), where the central bright protrusion is the silicon atom

G_{22} : Category three of Several atomic steps running across the Si (100) surface are also visible. b, Close-up of the inner device area (dashed box in a), where the central bright protrusion is the silicon atom

T_{20} : Category one of when a single phosphorus atom incorporates into the surface.. We Are talking of the characterstics of the various investigating systems.

T_{21} : Category two of when a single phosphorus atom incorporates into the surface.

T_{22} : Category three of when a single phosphorus atom incorporates into the surface.

Module four

Schematic of the chemical reaction to deterministically **incorporate a** single phosphorus atom into the surface

G_{24} : Category one of Schematic representation of the chemical reaction to deterministically; single phosphorus atom into the surface

G_{25} : Category two of Schematic representation of the chemical reaction to deterministically; single phosphorus atom into the surface

G_{26} : Category three of Schematic representation of the chemical reaction to deterministically; single phosphorus atom into the surface

T_{24} : Category one of single phosphorus atom into the surface; Schematic representation of the chemical reaction to deterministically

T_{25} : Category two of single phosphorus atom into the surface; Schematic representation of the chemical reaction to deterministically

T_{26} : Category three of Schematic representation of the chemical reaction to deterministically; single phosphorus atom into the surface

Module five

Saturation dosing of a three-dimer patch (I) at room temperature (RT) followed by annealing to 350 °C **allows** successive dissociation of PH3 (II–IV) and subsequent incorporation of a single phosphorus atom in the surface layer, ejecting a silicon adatom in the process (V). **POSTED BY BRIAN WANG AT 1/11/2013**

G_{28} : Category one of Saturation dosing of a three-dimer patch (I) at room temperature (RT) followed by annealing to 350 °C

G_{29} : Category two of Saturation dosing of a three-dimer patch (I) at room temperature (RT) followed by annealing to 350 °C

G_{30} : Category three of Saturation dosing of a three-dimer patch (I) at room temperature (RT) followed by annealing to 350 °C

T_{28} : Category one of successive dissociation of PH3 (II–IV) and subsequent incorporation of a single phosphorus atom in the surface layer, ejecting a silicon adatom in the process (V). **POSTED BY BRIAN WANG AT 1/11/2013**

T_{29} : Category two of successive dissociation of PH3 (II–IV) and subsequent incorporation of a single phosphorus atom in the surface layer, ejecting a silicon adatom in the process (V). **POSTED BY BRIAN WANG AT 1/11/2013**

T_{30} : Category three of successive dissociation of PH3 (II–IV) and subsequent incorporation of a single phosphorus atom in the surface layer, ejecting a silicon adatom in the process (V). **POSTED BY BRIAN WANG AT 1/11/2013**

Module six

The key to making these wires was combining scanning tunneling microscopy, a technique to image and manipulate individual atoms, **with (e&eb)** molecular beam epitaxy, a way of growing perfect crystals.

G_{32} : Category one of key to making these wires was combining scanning tunneling microscopy, a technique to image and manipulate individual atoms

G_{33} : Category two of key to making these wires was combining scanning tunneling microscopy, a technique to image and manipulate individual atoms

G_{34} : Category three of key to making these wires was combining scanning tunneling microscopy, a technique to image and manipulate individual atoms

T_{32} : Category one of molecular beam epitaxy, a way of growing perfect crystals ;key to making these wires was combining scanning tunneling microscopy, a technique to image and manipulate individual atoms

T_{33} : Category two of molecular beam epitaxy, a way of growing perfect crystals; key to making these wires was combining scanning tunneling microscopy, a technique to image and manipulate individual atoms

T_{34} : Category three of molecular beam epitaxy, a way of growing perfect crystals; key to making these wires was combining scanning tunneling microscopy, a technique to image and manipulate individual atoms

Module seven

It gave us great precision in all three dimensions, and when combined with a high density of the dopant

atoms, **allowed us to create** these highly conductive nanowires

G_{36} : Category one of great precision in all three dimensions, and when combined with a high density of the dopant atoms,

G_{37} : Category two of great precision in all three dimensions, and when combined with a high density of the dopant atoms,

G_{38} : Category three of great precision in all three dimensions, and when combined with a high density of the dopant atoms,

T_{36} : Category one of highly conductive nanowires

T_{37} : Category two of highly conductive nanowires

T_{38} : Category three of highly conductive nanowires

Module eight

The Kane quantum computer is a proposal for a scalable quantum computer proposed by Bruce Kane in 1998, then at the University of New South Wales. Often thought of as a hybrid between quantum dot **and** NMR quantum computers, the Kane computer **is based (e) on** an array of individual phosphorus donor atoms embedded in a pure silicon lattice. Both the nuclear spins of the donors and the spins of the donor electrons participate in the computation. The original proposal calls for phosphorus donors to be placed in an array with a spacing of 20 nm, approximately 20 nm below the surface. An insulating oxide layer is grown on top of the silicon. Metal A gates are deposited on the oxide above each donor, and J gates between adjacent donors

G_{40} : Category one of array of individual phosphorus donor atoms embedded in a pure silicon lattice. Both the nuclear spins of the donors and the spins of the donor electrons participate in the computation. The original proposal calls for phosphorus donors to be placed in an array with a spacing of 20 nm, approximately 20 nm below the surface. An insulating oxide layer is grown on top of the silicon. Metal A gates are deposited on the oxide above each donor, and J gates between adjacent donors

G_{41} : Category two of array of individual phosphorus donor atoms embedded in a pure silicon lattice. Both the nuclear spins of the donors and the spins of the donor electrons participate in the computation. The original proposal calls for phosphorus donors to be placed in an array with a spacing of 20 nm, approximately 20 nm below the surface. An insulating oxide layer is grown on top of the silicon. Metal A gates are deposited on the oxide above each donor, and J gates between adjacent donors

G_{42} : Category three of array of individual phosphorus donor atoms embedded in a pure silicon lattice. Both the nuclear spins of the donors and the spins of the donor electrons participate in the computation. The original proposal calls for phosphorus donors to be placed in an array with a spacing of 20 nm, approximately 20 nm below the surface. An insulating oxide layer is grown on top of the silicon. Metal A gates are deposited on the oxide above each donor, and J gates between adjacent donors

T_{40} : Category one of hybrid between quantum dot **and** NMR quantum computers, the Kane computer

T_{41} : Category two of hybrid between quantum dot **and** NMR quantum computers, the Kane computer

T_{42} : Category three of hybrid between quantum dot **and** NMR quantum computers, the Kane computer

Module Nine

The phosphorus donors are isotropically pure 31P, which have a nuclear spin of ½; the silicon substrate is isotropically pure 28Si which has nuclear spin 0. Using the nuclear spin of the P donors as a method to encode qubits has two major advantages; firstly, the state **has (e) an** extremely long decoherence time, perhaps on the order of 1018 seconds at millikelvin temperatures.

G_{44} : Category one of extremely long decoherence time, perhaps on the order of 1018 seconds at millikelvin temperatures.

G_{45} : Category two of extremely long decoherence time, perhaps on the order of 1018 seconds at millikelvin temperatures.

G_{46} : Category three of extremely long decoherence time, perhaps on the order of 1018 seconds at millikelvin temperatures.

T_{44} : Category one of phosphorus donors are isotropically pure 31P, which have a nuclear spin of ½; the silicon substrate is isotropically pure 28Si which has nuclear spin 0. Using the nuclear spin of the P donors as a method to encode qubits has two major advantages; firstly, the state

T_{45} : Category two of phosphorus donors is isotropically pure 31P, which have a nuclear spin of ½; the silicon substrate is isotropically pure 28Si which has nuclear spin 0. Using the nuclear spin of the P donors as a method to encode qubits has two major advantages; firstly, the state

T_{46} : Category three of phosphorus donors is isotropically pure 31P, which have a nuclear spin of ½; the silicon substrate is isotropically pure 28Si which has nuclear spin 0. Using the nuclear spin of the P donors as a method to encode qubits has two major advantages; firstly, the state

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$

$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$
$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$
$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$
$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) \quad +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{ccc} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) \quad +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right]G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) \quad +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right]G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation

coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\left[-(b''_{13})^{(1)}(G,t)\right] & \left[-(b''_{16})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\left[-(b''_{14})^{(1)}(G,t)\right] & \left[-(b''_{17})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\left[-(b''_{15})^{(1)}(G,t)\right] & \left[-(b''_{18})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad 73$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad 74$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad 76$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)} & \boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad 78$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad 81$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32} \quad 85$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b_{36}')^{(7)}\boxed{-(b_{36}'')^{(7)}(G_{39},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b_{37}')^{(7)}\boxed{-(b_{37}'')^{(7)}(G_{39},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b_{38}')^{(7)}\boxed{-(b_{38}'')^{(7)}(G_{39},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\,-(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\,-(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\,-(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(UUU) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(VVV) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$

132

(WWW) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(XXX)

$\lim_{G\to\infty} (b_i'')^{(7)}((G_{39}),t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}((G_{39}),t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(YYY) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(ZZZ) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad (169)$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that (170)

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that (172)

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that (174)

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(m) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+\ (a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right) d\left(\left((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\tilde{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$sup_i\{max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\tilde{M}_{32})^{(6)}t}, max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\tilde{M}_{32})^{(6)}t}\}$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right) \quad (248)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(6)}$ and $(b''_i)^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(6)} - (a''_i)^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)},, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},, (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad (333)$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad (334)$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad (335)$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad (336)$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \qquad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le \frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \qquad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \qquad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le -(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le -(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(m) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\right)\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\right)\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FORTY TWO A

Kane Computer: Una Hirundo Non Facit Ver- Usus Est Magister Optimus

INTRODUCTION—VARIABLES USED

Kane quantum computer (Wikipedia)

(1) Secondly, the qubits **may be manipulated** by applying an oscillating magnetic field, as in typical NMR proposals. By altering the voltage on the A gates, it should be possible to alter the Larmor frequency of individual donors. **This allows them** to be addressed individually, by bringing specific donors into resonance with the applied oscillating magnetic field.
(2) Nuclear spins alone will not interact significantly with other nuclear spins 20 nm away. Nuclear spin is **useful (e) to** perform single-qubit operations, but to make a quantum computer; two-qubit operations are also required.
(3) This is the role of electron spin in this design. Under A-gate control, the spin is transferred from the nucleus **to (e&eb) the** donor electron.
(4) Then, a potential is applied to the J gate, drawing adjacent donor electrons into a common region, greatly **enhancing the** interaction between the neighbouring spins. By controlling the J gate voltage, two-qubit operations are possible.
(5) Kane's proposal for readout was to apply an electric field **to encourage** spin-dependent tunneling of an electron to transform two neutral donors to a D+–D– state, that is, one where two electrons orbit the same donor.
(6) The charge excess is then detected **using a** single-electron transistor.
(7) This method has two major difficulties. Firstly, the D– state has strong coupling with the environment **and (eb)** hence a short decoherence time.
(8) Secondly and perhaps more importantly, it's not clear that the D– state has a sufficiently long lifetime **to allow (eb) for** readout—the electron tunnels into the conduction band.
(9) Unlike many quantum computation schemes, the Kane quantum computer is in principle scalable to an arbitrary number of qubits. This is possible **because (e) qubits** may be individually addressed by electrical means.

Since Kane's proposal, under the guidance of Robert Clark and now Michelle Simmons, pursuing realisation of the Kane quantum computer has become the primary quantum computing effort in Australia. Currently, about 100 researchers work on the problem. Theorists have put forward a number of proposals for improved readout. Experimentally, atomic-precision deposition of phosphorus atoms has been demonstrated, using an STM technique. Detection of the movement of single electrons between small, dense clusters of phosphorus donors has also been achieved. The group remains optimistic that a practical large-scale quantum computer can be built.

NOTATION

Module One

Phosphorus donors are isotropically pure 31P, which have a nuclear spin of ½; the silicon substrate is isotropically pure 28Si which has nuclear spin 0. Using the nuclear spin of the P donors as a method to encode qubits has two major advantages secondly the qubits **may be manipulated** by applying an oscillating magnetic field, as in typical NMR proposals. By altering the voltage on the A gates, it should be possible to alter the Larmor frequency of individual donors. **This allows them** to be addressed individually, by bringing specific donors into resonance with the applied oscillating magnetic field.

G_{13} : Category one of phosphorus donors is isotropically pure 31P, which have a nuclear spin of ½; the silicon substrate is isotropically pure 28Si which has nuclear spin 0. Using the nuclear spin of the P donors as a method to encode qubits has two major advantages secondly the qubits

G_{14} : Category two of phosphorus donors is isotropically pure 31P, which have a nuclear spin of ½; the silicon substrate is isotropically pure 28Si which has nuclear spin 0. Using the nuclear spin of the P donors as a method to encode qubits has two major advantages secondly the qubits

G_{15} : Category three of phosphorus donors is isotropically pure 31P, which have a nuclear spin of ½; the silicon substrate is isotropically pure 28Si which has nuclear spin 0. Using the nuclear spin of the P donors as a method to encode qubits has two major advantages secondly the qubits

T_{13} : Category one of applying an oscillating magnetic field, as in typical NMR proposals. By altering the voltage on the A gates, it should be possible to alter the Larmor frequency of individual donors. **This allows them** to be addressed individually, by bringing specific donors into resonance with the applied oscillating magnetic field.

T_{14} : Category two of applying an oscillating magnetic field, as in typical NMR proposals. By altering the voltage on the A gates, it should be possible to alter the Larmor frequency of individual donors. **This allows them** to be addressed individually, by bringing specific donors into resonance with the applied oscillating magnetic field.

T_{15} : Category three of applying an oscillating magnetic field, as in typical NMR proposals. By altering the voltage on the A gates, it should be possible to alter the Larmor frequency of individual donors. **This allows them** to be addressed individually, by bringing specific donors into resonance with the applied oscillating magnetic field.

Module Two

Nuclear spins alone will not interact significantly with other nuclear spins 20 nm away. Nuclear spin is **useful (eb) to** perform single-qubit operations, but to make a quantum computer; two-qubit operations are also required.

G_{16} : Category one of Nuclear spins alone will not interact significantly with other nuclear spins 20 nm away. Nuclear spin

G_{17} : Category two of Nuclear spins alone will not interact significantly with other nuclear spins 20 nm away. Nuclear spin

G_{18} : Category three of Nuclear spins alone will not interact significantly with other nuclear spins 20 nm away. Nuclear spin

T_{16} : Category one of performance of single-qubit operations, but to make a quantum computer; two-qubit operations are also required.

T_{17} : Category two of performance of single-qubit operations, but to make a quantum computer; two-qubit operations are also required.

T_{18} : Category three of performance of single-qubit operations, but to make a quantum computer; two-qubit operations are also required.

Module three

This is the role of electron spin in this design. Under A-gate control, the spin is transferred from the nucleus **to (e&eb) the** donor electron

G_{20} : Category one of Under A-gate control, the spin is transferred from the nucleus

G_{21} : Category two of Under A-gate control, the spin is transferred from the nucleus

G_{22} : Category three of Under A-gate control, the spin is transferred from the nucleus

T_{20} : Category one of donor electron

T_{21} : Category two of donor electron

T_{22} : Category three of donor electron

Module four

Then, a potential is applied to the J gate, drawing adjacent donor electrons into a common region, greatly **enhancing the** interaction between the neighbouring spins. By controlling the J gate voltage, two-qubit operations are possible.

G_{24} : Category one of potential is applied to the J gate, drawing adjacent donor electrons into a common region

G_{25} : Category two of potential is applied to the J gate, drawing adjacent donor electrons into a common region

G_{26} : Category three of potential is applied to the J gate, drawing adjacent donor electrons into a common region

T_{24} : Category one of interaction between the neighbouring spins. By controlling the J gate voltage, two-qubit operations are possible.

T_{25} : Category two of interaction between the neighbouring spins. By controlling the J gate voltage, two-qubit operations are possible.

T_{26} : Category three of interaction between the neighbouring spins. By controlling the J gate voltage, two-qubit operations are possible.

Module five

Kane's proposal for readout was to apply an electric field **to encourage** spin-dependent tunneling of an electron to transform two neutral donors to a D+–D– state, that is, one where two electrons orbit the same donor

G_{28} : Category one of Kane's proposal for readout was to apply an electric field

G_{29} : Category two of Kane's proposal for readout was to apply an electric field

G_{30} : Category three of Kane's proposal for readout was to apply an electric field

T_{28} : Category one of spin-dependent tunneling of an electron to transform two neutral donors to a D+–D– state, that is, one where two electrons orbit the same donor

T_{29} : Category two of spin-dependent tunneling of an electron to transform two neutral donors to a D+–D– state, that is, one where two electrons orbit the same donor

T_{30} : Category three of spin-dependent tunneling of an electron to transform two neutral donors to a D+–D– state, that is, one where two electrons orbit the same donor

Module six

The charge excess is then detected **using a** single-electron transistor

G_{32} : Category one of single-electron transistor

G_{33} : Category two of single-electron transistor

G_{34} : Category three of single-electron transistor

T_{32} : Category one of charge excess is then detected

T_{33} : Category two of charge excess is then detected

T_{34} : Category three of charge excess is then detected

Module seven

This method has two major difficulties. Firstly, the D– state has strong coupling with the environment **and (eb)** hence a short decoherence time

G_{36} : Category one of D– state has strong coupling with the environment

G_{37} : Category two of D– state has strong coupling with the environment

G_{38} : Category three of D– state has strong coupling with the environment

T_{36} : Category one of short decoherence time

T_{37} : Category two of short decoherence time

T_{38} : Category three of short decoherence time

Module eight

Secondly and perhaps more importantly, it's not clear that the D– state has a sufficiently long lifetime **to allow (eb) for** readout—the electron tunnels into the conduction band

G_{40} : Category one of D– state has a sufficiently long lifetime

G_{41} : Category two of D– state has a sufficiently long lifetime

G_{42} : Category three of D– state has a sufficiently long lifetime

T_{40} : Category one of readout—the electron tunnels into the conduction band

T_{41} : Category two of readout—the electron tunnels into the conduction band

T_{42} : Category three of readout—the electron tunnels into the conduction band

Module Nine

Unlike many quantum computation schemes, the Kane quantum computer is in principle scalable to an arbitrary number of qubits. This is possible **because (e) qubits** may be individually addressed by electrical means

G_{44} : Category one of **qubits** may be individually addressed by electrical means

G_{45} : Category two of **qubits** may be individually addressed by electrical means

G_{46} : Category three of **qubits** may be individually addressed by electrical means

T_{44} : Category one of Kane quantum computer is in principle scalable to an arbitrary number of qubits.

T_{45} : Category two of Kane quantum computer is in principle scalable to an arbitrary number of qubits.

T_{46} : Category three of Kane quantum computer is in principle scalable to an arbitrary number of qubits.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14},t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14},t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G,t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G,t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}),t\big)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}),t\big)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}),t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}),t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27}),t\big)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27}),t\big)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)}\ \boxed{-(b''_{13})^{(1,1,)}(G,t)}\ \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)}\ \boxed{-(b''_{14})^{(1,1,)}(G,t)}\ \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)}\ \boxed{-(b''_{15})^{(1,1,)}(G,t)}\ \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category

1,2 and 3

$-(b_{20}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b_{36}'')^{(7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b_{40}'')^{(8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b_{44}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c}(a_{20}')^{(3)}+(a_{20}'')^{(3)}(T_{21},t)+(a_{16}'')^{(2,2,2)}(T_{17},t)+(a_{13}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{36}'')^{(7,7,7,7)}(T_{37},t)+(a_{40}'')^{(8,8,8,8)}(T_{41},t)+(a_{44}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c}(a_{21}')^{(3)}+(a_{21}'')^{(3)}(T_{21},t)+(a_{17}'')^{(2,2,2)}(T_{17},t)+(a_{14}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{37}'')^{(7,7,7,7)}(T_{37},t)+(a_{41}'')^{(8,8,8,8)}(T_{41},t)+(a_{45}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c}(a_{22}')^{(3)}+(a_{22}'')^{(3)}(T_{21},t)+(a_{18}'')^{(2,2,2)}(T_{17},t)+(a_{15}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{38}'')^{(7,7,7,7)}(T_{37},t)+(a_{42}'')^{(8,8,8,8)}(T_{41},t)+(a_{46}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{22} \quad (69)$$

$+(a_{20}'')^{(3)}(T_{21},t)$, $+(a_{21}'')^{(3)}(T_{21},t)$, $+(a_{22}'')^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation

coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$ *are fourth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$ *are fifth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$ *are sixth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$ *are seventh augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$ *are eighth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\,-(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\,-(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\,-(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)}\,+(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad (87)$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad (90)$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b_{28}'')^{(5,5,5)}(G_{31},t)}, \boxed{-(b_{29}'')^{(5,5,5)}(G_{31},t)}, \boxed{-(b_{30}'')^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b_{24}'')^{(4,4,4,)}(G_{27},t)}, \boxed{-(b_{25}'')^{(4,4,4,)}(G_{27},t)}, \boxed{-(b_{26}'')^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b_{45}'')^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b_{44}'')^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)}\boxed{+(a_{36}'')^{(7)}(T_{37},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)}\boxed{+(a_{37}'')^{(7)}(T_{37},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)}\boxed{+(a_{38}'')^{(7)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a_{36}'')^{(7)}(T_{37},t)}, \boxed{(a_{37}'')^{(7)}(T_{37},t)}, \boxed{(a_{38}'')^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\left[-(b''_{36})^{(7)}(G_{39},t)\right] & \left[-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\left[-(b''_{37})^{(7)}(G_{39},t)\right] & \left[-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\left[-(b''_{38})^{(7)}(G_{39},t)\right] & \left[-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)}$ 103

$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)}$ 104

$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)}$ 105

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t}$ 107

$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t}$ 108

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1$ 110

$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1$ 111

Where we suppose

$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22$ 112

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32{,}33{,}34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32{,}33{,}34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(AAAA) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36{,}37{,}38$ 131

(BBBB) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$ 132

(CCCC) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(DDDD)

$\lim_{G\to\infty}(b_i'')^{(7)}((G_{39}),t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}((G_{39}),t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(EEEE) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(FFFF) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ *and* $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\tilde{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\tilde{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad (169)$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that (170)

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that (172)

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that (174)

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(n) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $\left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right| \; e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$$
$$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$$

$$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$$

In the same way , one can obtain

$$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$

$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$

$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$

$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq$ 248

$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 250

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b'_i)^{(7)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}),(\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}),(\widetilde{T_{43}})\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)},\ \boldsymbol{if}\ (\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \ \boldsymbol{if}\ (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \boldsymbol{if}\ (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

and $\boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq v^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

406

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case,

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

407

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(n) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)} G_{14}^{*}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)} G_{14}^{*}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)} T_{14}^{*}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)} T_{14}^{*}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)} G_{17}^{*}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)} G_{17}^{*}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)} T_{17}^{*}}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)} T_{17}^{*}}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)} G_{21}^{*}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)} G_{21}^{*}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)} T_{21}^{*}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)} T_{21}^{*}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)} T_{45}^{*}}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)} T_{45}^{*}}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FORTY THREE

Quantum Control: Saltus In Demonstrando Model

INTRODUCTION—VARIABLES USED

Program: Measurement and Control for Quantum Computing Griffith University Researchers in the Centre for Quantum Computer Technology (and related researchers) at Griffith University in 2003. David Pegg (Professor), Jay Gambetta (Ph.D.), Howard Wiseman (Assoc. Professor, Program Manager)

The Postulates of Quantum Mechanics.

The basic quantum mechanical model is specified in terms of the following:

Observables.

(1) Physical quantities like position, momentum, spin, etc., **are represented** by self-adjoint operators on the Hilbert space H and are called observables.
(2) These are the noncommutative counterparts of random variables. States. A state is meant **to provide** a summary of the status of a physical system that enables the calculation of statistical quantities associated with observables.
(3) A generic state is specified by a density matrix ρ, which is a self-adjoint operator on H that is positive $\rho \geq 0$ and normalized $Tr[\rho] = 1$. This is the noncommutative counterpart of a probability density. The expectation of an observable A is given by

$(A) = (\rho,A) = Tr[\rho A]$

Pure states: $\rho = |\psi)(\psi|$, $\psi \in H$ so that

$(A) = Tr\,[|\psi)(\psi|A] = (\psi,A\psi) = (\psi|A|\psi)$

Measurement.

(4) A measurement is a physical procedure or experiment **that produces** numerical results related to observables. In any given measurement, the allowable results take values in the spectrum spec (A) of a chosen observable A. Given the state ρ, the value $a \in$ spec (A) is observed with probability Tr $[\rho Pa]$.

Conditioning.

(5) Suppose that a measurement of A **gives rise to** the observation a ∈ spec (A). Then we must **condition** the state in order to predict the outcomes of subsequent measurements, by updating the density matrix ρ using ρ ↦⟶ ρ [a] = PaρPa Tr [ρPa] .This is known as the "projection postulate".

(6) A closed (i.e. isolated) quantum system evolves in a unitary fashion: a physical quantity that is described at time t = 0 by an observable A is described at time t > 0 by [Heisenburg picture] A (t) = U (t) ∗AU (t), where U (t) is a unitary operator for each time t.

(7) The unitary is generated by the Schrödinger equation IhddtU (t) = H (t)U(t),where the (time dependent) Hamiltonian H(t) is a self-adjoint operator for each t.

(8) States evolve according to [Schrodinger picture] ρ(t) = U(t)ρU∗(t)

The two pictures are equivalent (dual):

(ρ(t),A) = (ρ,A(t))

⟨A, B⟩ = Tr [A*B], A, B ∈ B(H).

(9) In a famous experiment, Stern and Gerlach fired some silver atoms **through(e&eb) an** inhomogeneous magnetic field. They discovered two beams emerging from the magnetic field, **confirming the** discrete nature of the magnetic moment of atoms. Atoms in the upper beam are said to have "spin up", while those in the lower beam have "spin down". Hence they had measured a physical quantity called spin.

NOTATION

Module One

Physical quantities like position, momentum, spin, etc., **are represented** by self-adjoint operators on the Hilbert space H and are called observables

G_{13} : Category one of Physical quantities like position, momentum, spin, etc

G_{14} : Category two of Physical quantities like position, momentum, spin, etc

G_{15} : Category three of Physical quantities like position, momentum, spin, etc

T_{13} : Category one of self-adjoint operators on the Hilbert space H and are called observables

T_{14} : Category two of self-adjoint operators on the Hilbert space H and are called observables

T_{15} : Category three of self-adjoint operators on the Hilbert space H and are called observables

Module Two

These are the noncommutative counterparts of random variables. States. A state is meant **to provide** a summary of the status of a physical system that enables the calculation of statistical quantities associated with observables.

G_{16} : Category one of noncommutative counterparts of random variables. States. A state is meant

G_{17} : Category two of noncommutative counterparts of random variables. States. A state is meant

G_{18} : Category three of noncommutative counterparts of random variables. States. A state is meant

T_{16} : Category one of summary of the status of a physical system that enables the calculation of statistical

quantities associated with observables.

T_{17} : Category two of summary of the status of a physical system that enables the calculation of statistical quantities associated with observables.

T_{18} : Category three of summary of the status of a physical system that enables the calculation of statistical quantities associated with observables.

Module three

(1) A generic state is specified by a density matrix ρ, which is a self-adjoint operator on H that is positive $\rho \geq 0$ and normalized Tr[ρ] = 1. This is the noncommutative counterpart of a probability density. The expectation of an observable A is given by

(A) = (ρ,A) = Tr[ρA]

Pure states: $\rho = |\psi)(\psi|$, $\psi \in H$ so that

(A) = Tr $[|\psi)(\psi|A]$ = $(\psi,A\psi)$ = $(\psi|A|\psi)$

G_{20} : Category one of LHS of the equation: (A) = (ρ,A) = Tr[ρA];LHS of the equation (A) = Tr $[|\psi)(\psi|A]$ = $(\psi,A\psi)$ = $(\psi|A|\psi)$

G_{21} : Category two of LHS of the equation: (A) = (ρ,A) = Tr[ρA];LHS of the equation (A) = Tr $[|\psi)(\psi|A]$ = $(\psi,A\psi)$ = $(\psi|A|\psi)$

G_{22} : Category three of LHS of the equation: (A) = (ρ,A) = Tr[ρA];LHS of the equation (A) = Tr $[|\psi)(\psi|A]$ = $(\psi,A\psi)$ = $(\psi|A|\psi)$

T_{20} : Category one of RHS of the equation: (A) = (ρ,A) = Tr[ρA];LHS of the equation (A) = Tr $[|\psi)(\psi|A]$ = $(\psi,A\psi)$ = $(\psi|A|\psi)$

T_{21} : Category two of RHS of the equation: (A) = (ρ,A) = Tr[ρA];LHS of the equation (A) = Tr $[|\psi)(\psi|A]$ = $(\psi,A\psi)$ = $(\psi|A|\psi)$

T_{22} : Category three of RHS of the equation: (A) = (ρ,A) = Tr[ρA];LHS of the equation (A) = Tr $[|\psi)(\psi|A]$ = $(\psi,A\psi)$ = $(\psi|A|\psi)$

Module four

A measurement is a physical procedure or experiment **that produces** numerical results related to observables. In any given measurement, the allowable results take values in the spectrum spec (A) of a chosen observable A. Given the state ρ, the value $a \in$ spec (A) **is observed with** probability Tr [ρPa].
Note that the values of Tr(rou Pa) is predicted and prognosticated

G_{24} : Category one of measurement is a physical procedure or experiment; probability Tr [ρPa].

G_{25} : Category two of measurement is a physical procedure or experiment; probability Tr [ρPa].

G_{26} : Category three of measurement is a physical procedure or experiment; probability Tr [ρPa].

T_{24} : Category one of numerical results related to observables. In any given measurement, the allowable results take values in the spectrum spec (A) of a chosen observable A; state ρ, the value $a \in$ spec (A)

T_{25} : Category two of numerical results related to observables. In any given measurement, the allowable

results take values in the spectrum spec (A) of a chosen observable A; state ρ, the value a ∈ spec (A)

T_{26} : Category three of numerical results related to observables. In any given measurement, the allowable results take values in the spectrum spec (A) of a chosen observable A; state ρ, the value a ∈ spec (A)

Module five

Suppose that a measurement of A **gives rise to** the observation a ∈ spec (A). Then we must **condition** the state in order to predict the outcomes of subsequent measurements, by updating the density matrix ρ using ρ ↦⟶ ρ [a] = PaρPa Tr [ρPa] .This is known as the "projection postulate".

G_{28} : Category one of measurement of A (There may be many investigating systems and the characterstics thereof be it A or something else form the citadel of the fulcrum classification); **condition** the state; LHS of the projection postulate

G_{29} : Category two of measurement of A ; **condition** the state; LHS of the projection postulate

G_{30} : Category three of measurement of A ; **condition** the state; LHS of the projection postulate

T_{28} : Category one of observation a ∈ spec (A); predict the outcomes of subsequent measurements, by updating the density matrix ρ using ρ ↦⟶ ρ [a] = PaρPa Tr [ρPa] .This is known as the "projection postulate"; RHS of the projection postulate

T_{29} : Category two of observation a ∈ spec (A); predict the outcomes of subsequent measurements, by updating the density matrix ρ using ρ ↦⟶ ρ [a] = PaρPa Tr [ρPa] .This is known as the "projection postulate"; RHS of the projection postulate

T_{30} : Category three of observation a ∈ spec (A); predict the outcomes of subsequent measurements, by updating the density matrix ρ using ρ ↦⟶ ρ [a] = PaρPa Tr [ρPa] .This is known as the "projection postulate";RHS of the projection postulate

Module six

A closed (i.e. isolated) quantum system evolves in a unitary fashion: a physical quantity that is described at time t = 0 by an observable A is described at time t > 0 by [Heisenburg picture] A (t) = U (t) ∗AU (t), where U (t) is a unitary operator for each time t.

G_{32} : Category one of closed (i.e. isolated) quantum system; LHS of the Heisenberg picture A (t) = U (t) ∗AU (t), where U (t) is a unitary operator for each time t.

G_{33} : Category two of closed (i.e. isolated) quantum system; LHS of the Heisenberg picture A (t) = U (t) ∗AU (t), where U (t) is a unitary operator for each time t.

G_{34} : Category three of closed (i.e. isolated) quantum system; LHS of the Heisenberg picture A (t) = U (t) ∗AU (t), where U (t) is a unitary operator for each time t.

T_{32} : Category one of unitary fashion; RHS of the Heisenberg picture A (t) = U (t) ∗AU (t), where U (t) is a unitary operator for each time t.

T_{33} : Category two of unitary fashion; RHS of the Heisenberg picture A (t) = U (t) ∗AU (t), where U (t) is a unitary operator for each time t.

T_{34} : Category three of unitary fashion; RHS of the Heisenberg picture A (t) = U (t) ∗AU (t), where U (t) is

a unitary operator for each time t.

Module seven

The unitary is generated by the Schrödinger equation IhddtU (t) = H (t)U(t),where the (time dependent) Hamiltonian H(t) is a self-adjoint operator for each t.

G_{36} : Category one of Schrödinger equation IhddtU (t) = H (t)U(t),where the (time dependent) Hamiltonian H(t) is a self-adjoint operator for each t.; LHS of Schrödinger equation IhddtU (t) = H (t)U(t),where the (time dependent) Hamiltonian H(t) is a self-adjoint operator for each t.

G_{37} : Category two of Schrödinger equation IhddtU (t) = H (t)U(t),where the (time dependent) Hamiltonian H(t) is a self-adjoint operator for each t; LHS of Schrödinger equation IhddtU (t) = H (t)U(t),where the (time dependent) Hamiltonian H(t) is a self-adjoint operator for each t.

G_{38} : Category three of Schrödinger equation IhddtU (t) = H (t)U(t),where the (time dependent) Hamiltonian H(t) is a self-adjoint operator for each t; LHS of Schrödinger equation IhddtU (t) = H (t)U(t),where the (time dependent) Hamiltonian H(t) is a self-adjoint operator for each t.

T_{36} : Category one of unitary evolution; RHS of Schrödinger equation IhddtU (t) = H (t)U(t),where the (time dependent) Hamiltonian H(t) is a self-adjoint operator for each t.

T_{37} : Category two of unitary evolution; RHS of Schrödinger equation IhddtU (t) = H (t)U(t),where the (time dependent) Hamiltonian H(t) is a self-adjoint operator for each t.

T_{38} : Category three of unitary evolution; RHS of Schrödinger equation IhddtU (t) = H (t)U(t),where the (time dependent) Hamiltonian H(t) is a self-adjoint operator for each t.

Module eight

States evolve according to [Schrodinger picture] $\rho(t) = U(t)\rho U*(t)$
The two pictures are equivalent (dual):

$(\rho(t),A) = (\rho,A(t))$

$\langle A, B\rangle = Tr [A*B]$, $A, B \in B(H)$.

Note: Each term in Schrodinger picture is predicted and analyzed for stability and Solutional behaviour.

G_{40} : Category one of States evolve; LHS of the equation[Schrodinger picture] $\rho(t) = U(t)\rho U*(t)$
G_{41} : Category two of States evolve; LHS of the equation[Schrodinger picture] $\rho(t) = U(t)\rho U*(t)$
G_{42} : Category three of States evolve; LHS of the equation[Schrodinger picture] $\rho(t) = U(t)\rho U*(t)$

T_{40} : Category one of[Schrodinger picture] $\rho(t) = U(t)\rho U*(t)$;RHS of the equation[Schrodinger picture] $\rho(t) = U(t)\rho U*(t)$
T_{41} : Category two of [Schrodinger picture] $\rho(t) = U(t)\rho U*(t)$; RHS of the equation[Schrodinger picture] $\rho(t) = U(t)\rho U*(t)$
T_{42} : Category three of [Schrodinger picture] $\rho(t) = U(t)\rho U*(t)$; RHS of the equation[Schrodinger picture] $\rho(t) = U(t)\rho U*(t)$

Module Nine

In a famous experiment, Stern and Gerlach fired some silver atoms **through (e&eb) an** inhomogeneous

magnetic field. They discovered two beams emerging from the magnetic field, **confirming the** discrete nature of the magnetic moment of atoms. Atoms in the upper beam are said to have "spin up", while those in the lower beam have "spin down". Hence they had measured a physical quantity called spin.

G_{44} : Category one of two beams emerging from the magnetic field

G_{45} : Category two of two beams emerging from the magnetic field

G_{46} : Category three of two beams emerging from the magnetic field

T_{44} : Category one of discrete nature of the magnetic moment of atoms. Atoms in the upper beam are said to have "spin up", while those in the lower beam have "spin down". Hence they had measured a physical quantity called spin.

T_{45} : Category two of discrete nature of the magnetic moment of atoms. Atoms in the upper beam are said to have "spin up", while those in the lower beam have "spin down". Hence they had measured a physical quantity called spin.

T_{46} : Category three of discrete nature of the magnetic moment of atoms. Atoms in the upper beam are said to have "spin up", while those in the lower beam have "spin down". Hence they had measured a physical quantity called spin.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}$ $(a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$ $(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$ $(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\,-(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\,-(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\,-(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)}\ \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad 63$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)}\ \boxed{-(b''_{13})^{(1,1,)}(G,t)}\ \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)}\ \boxed{-(b''_{14})^{(1,1,)}(G,t)}\ \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)}\ \boxed{-(b''_{15})^{(1,1,)}(G,t)}\ \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c}(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21},t)\;\; +(a''_{16})^{(2,2,2)}(T_{17},t)\;\; +(a''_{13})^{(1,1,1,)}(T_{14},t)\\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)\;\; +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)\;\; +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a''_{36})^{(7,7,7,7)}(T_{37},t)\;\; +(a''_{40})^{(8,8,8,8)}(T_{41},t)\;\; +(a''_{44})^{(9,9,9)}(T_{45},t)\end{array}\right]G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c}(a'_{21})^{(3)}+(a''_{21})^{(3)}(T_{21},t)\;\; +(a''_{17})^{(2,2,2)}(T_{17},t)\;\; +(a''_{14})^{(1,1,1,)}(T_{14},t)\\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)\;\; +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)\;\; +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a''_{37})^{(7,7,7,7)}(T_{37},t)\;\; +(a''_{41})^{(8,8,8,8)}(T_{41},t)\;\; +(a''_{45})^{(9,9,9)}(T_{45},t)\end{array}\right]G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c}(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21},t)\;\; +(a''_{18})^{(2,2,2)}(T_{17},t)\;\; +(a''_{15})^{(1,1,1,)}(T_{14},t)\\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)\;\; +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)\;\; +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a''_{38})^{(7,7,7,7)}(T_{37},t)\;\; +(a''_{42})^{(8,8,8,8)}(T_{41},t)\;\; +(a''_{46})^{(9,9,9)}(T_{45},t)\end{array}\right]G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\,-(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\,-(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\,-(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

$Where\ \boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}\ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$

$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$

$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$

$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$

$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{c} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)}\ \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}\ \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{c} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)}\ \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}\ \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{c} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)}\ \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}\ \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad (81)$$

$Where\ \boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$And\ \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}\ are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\ are\ third\ \ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)$, $+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b_{28}')^{(5)} & -(b_{28}'')^{(5)}(G_{31},t) & -(b_{24}'')^{(4,4,)}(G_{27},t) & -(b_{32}'')^{(6,6,6)}(G_{35},t) \\ & -(b_{13}'')^{(1,1,1,1,1)}(G,t) & -(b_{16}'')^{(2,2,2,2,2)}(G_{19},t) & -(b_{20}'')^{(3,3,3,3,3)}(G_{23},t) \\ & -(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t) & -(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b_{29}')^{(5)} & -(b_{29}'')^{(5)}(G_{31},t) & -(b_{25}'')^{(4,4,)}(G_{27},t) & -(b_{33}'')^{(6,6,6)}(G_{35},t) \\ & -(b_{14}'')^{(1,1,1,1,1)}(G,t) & -(b_{17}'')^{(2,2,2,2,2)}(G_{19},t) & -(b_{21}'')^{(3,3,3,3,3)}(G_{23},t) \\ & -(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t) & -(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t) & -(b_{45}'')^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b_{30}')^{(5)} & -(b_{30}'')^{(5)}(G_{31},t) & -(b_{26}'')^{(4,4,)}(G_{27},t) & -(b_{34}'')^{(6,6,6)}(G_{35},t) \\ & -(b_{15}'')^{(1,1,1,1,1,)}(G,t) & -(b_{18}'')^{(2,2,2,2,2)}(G_{19},t) & -(b_{22}'')^{(3,3,3,3,3)}(G_{23},t) \\ & -(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t) & -(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t) & -(b_{46}'')^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b_{28}'')^{(5)}(G_{31},t)$, $-(b_{29}'')^{(5)}(G_{31},t)$, $-(b_{30}'')^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b_{24}'')^{(4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b_{32}'')^{(6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b_{13}'')^{(1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\left[-(b''_{40})^{(8)}(G_{43},t)\right] & \left[-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\left[-(b''_{41})^{(8)}(G_{43},t)\right] & \left[-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\left[-(b''_{42})^{(8)}(G_{43},t)\right] & \left[-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\,(a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\,(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)},$ are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(GGGG) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(HHHH)The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$

132

(IIII) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(JJJJ)

$\lim_{G\to\infty}(b_i'')^{(7)}((G_{39}),t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}((G_{39}),t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(KKKK) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(LLLL) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(o) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \qquad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \qquad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \qquad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(o) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FORTY FOUR

Quantum Transport: A Gesamtbedeutung Model

INTRODUCTION—VARIABLES USED

Theoretical and Numerical aspects of quantum control problems: Alfio Borzi

Application fields of quantum control methodologies

1. Quantum control: state transitions, laser induced chemistry, magnetic and optical trapping.
2. Quantum computing: qubits, data operations.
3. Quantum transport, superfluids of atoms, vortices.
4. Construction of barriers, channels, etc. for few atoms.
5. Amplification of material waves: laser of atoms.
6. Semiconductor nanostructures.

NOTATION

Module One

Quantum control models define an important class of nonlinear control mechanisms.

G_{13} : Category one of quantum mechanical models

G_{14} : Category two of quantum mechanical models

G_{15} : Category three ofquantum mechanical models

T_{13} : Category one of nonlinear control mechanisms

T_{14} : Category two of nonlinear control mechanisms

T_{15} : Category three of nonlinear control mechanisms

Module Two

Quantum mechanical models

- One-particle Schrödinger equation, $\psi = \psi(x,t)$ or $\psi = \psi(t)$

$$i\frac{\partial}{\partial t}\psi = (H_0 + V)\psi$$

- BEC Condensate Gross-Pitaevskii equation, $\psi = \psi(x,t)$

$$i\frac{\partial}{\partial t}\psi = \left(-\frac{1}{2}\nabla^2 + V + g|\psi|^2\right)\psi$$

- Time-dependent Kohn-Sham equation, $\psi_i = \psi_i(x,t)$

$$i\frac{\partial}{\partial t}\psi_i = \left(-\frac{1}{2}\nabla^2 + V_{ext} + V_{Hartree}(\rho) + V_{exc}(\rho) + V_{laser}\right)\psi_i$$

where ψ_i, $i = 1,\ldots,N$ are the K-S orbitals; $\rho = \sum_{i=1}^{N}|\psi_i|^2$ is the one-electron density.

- Multi-particle (n) Schrödinger equation, $\psi = \psi(x_1, x_2, \ldots, x_n, t)$

$$i\frac{\partial}{\partial t}\psi = \left(-\frac{1}{2}\sum_{i=1}^{n}\nabla_i^2 + \sum_{i=1}^{n}V_i + \sum_{i,j=1}^{n}U_{ij} + V_{laser}\right)\psi$$

G_{16} : Category one of LHS of the above equations

G_{17} : Category two of LHS of the above equations

G_{18} : Category three of LHS of the above equations

T_{16} : Category one of RHS of the above equations (we have dilated upon this many times. Kindly refer to earlier papers)

T_{17} : Category two of RHS of the above equations

T_{18} : Category three of RHS of the above equations

Module three

Quantum control mechanisms

Laser pulses, electric fields, and magnetic fields represent physically meaningful control mechanisms. They are represented by potentials that sum up to the stationary ones

$$V(x,t) = V_0(x) + V_{control}(x,t)$$

The dipole approximation of the electric control field modeling a laser pulse results in the form

$$V_{control}(x,t) = u(t)\,x$$

where $u : (0,T) \to \mathbb{R}$ is the modulating control amplitude.

A magnetic potential for manipulating a BEC is given by

$$V_{control}(x,\lambda(t)) = -\frac{\lambda(t)^2\, d^2}{8c}\, x^2 + \frac{1}{c}\, x^4$$



where λ is the control function.

G_{20} : Category one of LHS of the above equations

G_{21} : Category two of LHS of the above equations

G_{22} : Category three of LHS of the above equations

T_{20} : Category one of RHS of the above equations

T_{21} : Category two of RHS of the above equations

T_{22} : Category three of RHS of the above equations

Module four

Finite-level quantum models

Governed by Schrödinger-type equations for a n-component wave function $\psi : [0, T] \to \mathbb{C}^n$ as follows

$$i\dot{\psi}(t) = H(u(t))\psi(t), \qquad \psi(0) = \psi_0,$$

for $t \in [0, T]$ and $T > 0$ is a given terminal time.

The funcion $u : [0, T] \to \mathbb{C}^m$ represents the external control field.

The linear Hamiltonian $H(u) = H_0 + V(u)$, consists of

A free Hamiltonian $H_0 \in \mathbb{C}^{n\times n}$ describing the unperturbed (uncontrol system;

A control Hamiltonian $V : \mathbb{C} \to \mathbb{C}^{n\times n}$ modeling the coupling of the quantum state to the control field u.

G_{24} : Category one of LHS of the above Schrodinger type equation

G_{25} : Category two of LHS of the above Schrodinger type equation

G_{26} : Category three of LHS of the above Schrodinger type equation

T_{24} : Category one of RHS of the above Schrodinger type equation

T_{25} : Category two of RHS of the above Schrodinger type equation

T_{26} : Category three of RHS of the above Schrodinger type equation

Module five

Monotonic schemes for quantum control problems

Monotonic schemes have been initially introduced by Krotov. They are a special case of gradient based methods.
Monotonic schemes have been further developed by Tannor, Zhu & Rabitz, Maday, Turinici & Salomon.

Consider two fields u and u', and the corresponding ψ and ψ', and p and p'. We have

$$\begin{aligned}\widetilde{J}(u') - \widetilde{J}(u) &= \int_0^T \Re e(p(s)^* \cdot (V(u'(s)) - V(u(s)))\psi'(s)) \\ &\quad -\frac{\gamma}{2}\left(|u'(s)|^2 - |u(s)|^2\right) \mathrm{d}s + \frac{1}{2}\langle \psi' - \psi, \Lambda(\psi' - \psi)\rangle_{L^2(0,T;\mathbb{C}^n)}.\end{aligned}$$

G_{28} : Category one of LHS of the above equation for monotonic schemes(Krotov)

G_{29} : Category two of LHS of the above equation for monotonic schemes(Krotov)

G_{30} : Category three of LHS of the above equation for monotonic schemes(Krotov)

T_{28} : Category one of RHS of the above equation for monotonic schemes(Krotov)

T_{29} : Category two of RHS of the above equation for monotonic schemes(Krotov)

T_{30} : Category three of RHS of the above equation for monotonic schemes(Krotov)

Module six


Bose Einstein condensates model

The mean-field dynamics of the condensate is described by the Gross-Pitaevskii equation (GPE)

$$i\frac{\partial}{\partial t}\psi(\mathbf{r},t)=\left(-\frac{1}{2}\nabla^2+V(\mathbf{r},\lambda(t))+g\,|\psi(\mathbf{r},t)|^2\right)\psi(\mathbf{r},t)$$

We consider $V(\mathbf{r},\lambda(t))$ is a three-dimensional potential produced by a magnetic microtrap. The control function $\lambda(t)$ describes the variation of the confining potential with time.

Magnetic trap with optical plug



G_{32} : Category one of LHS of the above equation for Bose Einstein Model **(Please note that these models predict, project and prognosticate each and every term in the equation and makes stability analysis, Solutional behaviour and asymptotic analysis of the quantum systems and quantum mechanical systems and quantum mechanical computational systems)**

G_{33} : Category two of LHS of the above equation for Bose Einstein Model

G_{34} : Category three of LHS of the above equation for Bose Einstein Model

T_{32} : Category one of RHS of the above equation for Bose Einstein Model

T_{33} : Category two of RHS of the above equation for Bose Einstein Model

T_{34} : Category three of RHS of the above equation for Bose Einstein Model

Module seven

Total control of matter waves

Trapping and coherent manipulation of cold neutral atoms in microtra near surfaces of atomic chips is the focus of the present research towar total control of matter waves at small scales.

This achievement has boosted developments in the atomic interferometry, the construction of quantum gates, the microscopic magnetic field imaging, quantum data encoding, etc..

At the base of all these developments is the ability to manipulate Bose-Einstein condensates (BEC) subject to a control potential $V(x,\lambda(t))$ where $\lambda(t)$ parameterizes the trapping field.

We consider the problem to split and transport a BEC being confined i a single well $V(x,0)$ at $t=0$ to a double well $V(x,1)$ at time $t=T$. We have

$$V(x,\lambda(t)) = -\frac{\lambda(t)^2 d^2}{8c}x^2 + \frac{1}{c}x^4$$

where $c=40$ and d is the width of the double well potential.

G_{36} : Category one of LHS of the above equation for Bose Einstein Model(split and transport)

G_{37} : Category two of LHS of the above equation for Bose Einstein Model(split and transport)

G_{38} : Category three of LHS of the above equation for Bose Einstein Model(split and transport)

T_{36} : Category one of RHS of the above equation for Bose Einstein Model(split and transport)

T_{37} : Category two of RHS of the above equation for Bose Einstein Model(split and transport)

T_{38} : Category three of RHS of the above equation for Bose Einstein Model(split and transport)

Module eight

Optimal control formulation and optimality system

Consider a BEC at the initial state ψ_0 and a target state ψ_d.

$$J(\psi,\lambda)=\frac{1}{2}\left(1-\left|\langle\psi_d|\psi(T)\rangle\right|^2\right)+\frac{\gamma}{2}\int_0^T\left(\dot{\lambda}(t)\right)^2\,dt$$

Optimal control problem: Minimize the cost function $J(\psi,\lambda)$ subject to the condition that ψ fulfills the Gross-Pitaevskii equation.
The optimal solution is characterized by the optimality system

$$i\frac{\partial}{\partial t}\psi=\left(-\frac{1}{2}\nabla^2+V_\lambda+g|\psi|^2\right)\psi$$

$$i\frac{\partial}{\partial t}p=\left(-\frac{1}{2}\nabla^2+V_\lambda+2g|\psi|^2\right)p+g\,\psi^2\,p^*$$

$$\gamma\ddot{\lambda}=-\Re e\langle\psi|\frac{\partial V_\lambda}{\partial\lambda}|p\rangle\,,$$

where $\langle u,v\rangle=\int_\Omega u(x)^*v(x)\,dx$. We have the initial and terminal conditions

$$\psi(0)=\psi_0 \text{ and } ip(T)=-\langle\psi_d|\psi(T)\rangle\,\psi_d$$
$$\lambda(0)=0\,,\quad \lambda(T)=1\,.$$

G_{40} : Category one of LHS of the above equation for optimal control problems(optimal solution is characterised by this system)

G_{41} : Category two of LHS of the above equation for optimal control problems (optimal solution is characterised by this system)

G_{42} : Category three of LHS of the above equation for optimal control problems (optimal solution is characterised by this system)

T_{40} : Category one of RHS of the above equation for optimal control problems

T_{41} : Category two of RHS of the above equation for optimal control problems

T_{42} : Category three of RHS of the above equation for optimal control problems

Module Nine

The choice of the control space and the gradient

For a given potential $V(x, \lambda(t))$, we have a unique $\psi(\lambda) = \psi(x, t)$. In terms of λ we have the reduced objective $\hat{J}(\lambda) = J(\psi(\lambda), \lambda)$.
The Taylor series of $\hat{J}(\lambda)$ in a Hilbert space X is

$$\hat{J}(\lambda + \epsilon\varphi) = \hat{J}(\lambda) + \epsilon\left(\nabla\hat{J}(\lambda), \varphi\right)_X + \frac{\epsilon^2}{2}\left([\nabla^2\hat{J}(\lambda)]\varphi, \varphi\right)_X + O(\epsilon^3)$$

For $X = L^2(0, T; \mathbb{R})$, the reduced gradient is given

$$\nabla\hat{J}_{L^2}(\lambda) = -\gamma\ddot{\lambda} - \Re e\langle\psi|\frac{\partial V_\lambda}{\partial\lambda}|p\rangle,$$

In the case $X = H^1(0, T; \mathbb{R})$ formulation, we have that

$$-\frac{d^2}{dt^2}[\nabla\hat{J}_{H^1}(\lambda)] = -\gamma\ddot{\lambda} - \Re e\langle\psi, \frac{\partial V_\lambda}{\partial\lambda}p\rangle,$$

with $[\nabla\hat{J}(\lambda)](0) = 0$ and $[\nabla\hat{J}(\lambda)](T) = 0$.
The H^1 gradient $\nabla\hat{J}_{H^1}(\lambda)$ has the same regularity as λ, while the L^2 gradient does not.

G_{44} : Category one of LHS of the equation in the case of X=H^1(O,T;R) formulation

G_{45} : Category two of LHS of the equation in the case of X=H^1(O,T;R) formulation

G_{46} : Category three of LHS of the equation in the case of X=H^1(O,T;R) formulation

T_{44} : Category one of RHS of the equation in the case of X=H^1(O,T;R) formulation

T_{45} : Category two of RHS of the equation in the case of X=H^1(O,T;R) formulation

T_{46} : Category three of RHS of the equation in the case of X=H^1(O,T;R) formulation

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}),t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \big[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\big]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \big[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\big]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \big[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\big]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \big[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\big]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \big[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\big]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \big[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\big]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \big[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\big]G_{24} \quad 19$$
$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \big[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\big]G_{25} \quad 20$$
$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \big[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\big]G_{26} \quad 21$$
$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \big[(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27}),t\big)\big]T_{24} \quad 22$$
$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \big[(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27}),t\big)\big]T_{25} \quad 23$$
$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \big[(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27}),t\big)\big]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**
$-(b''_{24})^{(4)}\big((G_{27}),t\big) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \big[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\big]G_{28} \quad 25$$
$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \big[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\big]G_{29} \quad 26$$
$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \big[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\big]G_{30} \quad 27$$
$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \big[(b'_{28})^{(5)} - (b''_{28})^{(5)}\big((G_{31}),t\big)\big]T_{28} \quad 28$$
$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \big[(b'_{29})^{(5)} - (b''_{29})^{(5)}\big((G_{31}),t\big)\big]T_{29} \quad 29$$
$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \big[(b'_{30})^{(5)} - (b''_{30})^{(5)}\big((G_{31}),t\big)\big]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**
$-(b''_{28})^{(5)}\big((G_{31}),t\big) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \big[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\big]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}), t\right)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}), t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}), t\right)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}), t\right)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}), t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45}, t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}), t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad (55)$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad (56)$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad (57)$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $-(b_{13}'')^{(1)}(G,t)$, $-(b_{14}'')^{(1)}(G,t)$, $-(b_{15}'')^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b_{16}'')^{(2,2,)}(G_{19},t)$, $-(b_{17}'')^{(2,2,)}(G_{19},t)$, $-(b_{18}'')^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{37}'')^{(7,7,)}(G_{39},t)$, $-(b_{36}'')^{(7,7,)}(G_{39},t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8)}(G_{43},t)$ $-(b_{41}'')^{(8,8)}(G_{43},t)$ $-(b_{42}'')^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{ccc} (a_{16}')^{(2)} & +(a_{16}'')^{(2)}(T_{17},t) & +(a_{13}'')^{(1,1,)}(T_{14},t) \quad +(a_{20}'')^{(3,3,3)}(T_{21},t) \\ +(a_{24}'')^{(4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{36}'')^{(7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{ccc} (a_{17}')^{(2)} & +(a_{17}'')^{(2)}(T_{17},t) & +(a_{14}'')^{(1,1,)}(T_{14},t) \quad +(a_{21}'')^{(3,3,3)}(T_{21},t) \\ +(a_{25}'')^{(4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{37}'')^{(7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a_{18}')^{(2)} & +(a_{18}'')^{(2)}(T_{17},t) & +(a_{15}'')^{(1,1,)}(T_{14},t) \quad +(a_{22}'')^{(3,3,3)}(T_{21},t) \\ +(a_{26}'')^{(4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{38}'')^{(7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad (63)$$

Where $+(a_{16}'')^{(2)}(T_{17},t)$, $+(a_{17}'')^{(2)}(T_{17},t)$, $+(a_{18}'')^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)}\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b_{20}'')^{(3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{16}'')^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{13}'')^{(1,1,1,)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,)}(G,t)}$, $\boxed{-(b_{15}'')^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{36}'')^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b_{38}'')^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a_{24}')^{(4)} & \boxed{+(a_{24}'')^{(4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,)}(T_{29},t)} \quad \boxed{+(a_{32}'')^{(6,6,)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a_{16}'')^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{36}'')^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9)}(T_{45},t)} \end{array}\right] G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a_{25}')^{(4)} & \boxed{+(a_{25}'')^{(4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,)}(T_{29},t)} \quad \boxed{+(a_{33}'')^{(6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a_{17}'')^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{37}'')^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9)}(T_{45},t)} \end{array}\right] G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a_{26}')^{(4)} & \boxed{+(a_{26}'')^{(4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,)}(T_{29},t)} \quad \boxed{+(a_{34}'')^{(6,6,)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a_{18}'')^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{38}'')^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9)}(T_{45},t)} \end{array}\right] G_{26} \quad (75)$$

$\boxed{(a_{24}'')^{(4)}(T_{25},t)}$, $\boxed{(a_{25}'')^{(4)}(T_{25},t)}$, $\boxed{(a_{26}'')^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a_{28}'')^{(5,5,)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a_{32}'')^{(6,6,)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a_{13}'')^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{14}'')^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1)}(T_{14},t)}$ *are fourth augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a_{16}'')^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2)}(T_{17},t)}$ *are fifth augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a_{20}'')^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3)}(T_{21},t)}$

are sixth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$

are seventh augmentation coefficients for category 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \tag{76}$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \tag{77}$$

$$\frac{dT_{26}}{dt}=(b_{26})^{(4)}T_{25}-\begin{bmatrix}(b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{26} \tag{78}$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt}=(a_{28})^{(5)}G_{29}-\begin{bmatrix}(a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{28} \tag{79}$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad 81$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8\ ,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad 84$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ *are second detrition coefficients*

$for\ category$ 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ $are\ \ third\ \ detrition\ coefficients$ $for\ category$ 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt}=(a_{32})^{(6)}G_{33} \qquad 85$$

$$-\begin{bmatrix}(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t)\\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)\\ +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)\end{bmatrix}G_{32}$$

$$\frac{dG_{33}}{dt}=(a_{33})^{(6)}G_{32}-\begin{bmatrix}(a'_{33})^{(6)}+(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t)\\ +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)\\ +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)\end{bmatrix}G_{33} \qquad 86$$

$$\frac{dG_{34}}{dt}=(a_{34})^{(6)}G_{33}-\begin{bmatrix}(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t)\\ +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)\\ +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)\end{bmatrix}G_{34} \qquad 87$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ $are\ \ first\ \ augmentation\ coefficients$ $for\ category$ 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ $are\ second\ augmentation$ $coefficients\ for\ category$ 1, 2 and 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ $are\ third\ augmentation$ $coefficients\ for\ category$ 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$
seventh augmentation coefficients
$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$
Eighth augmentation coefficients
$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt}=(b_{32})^{(6)}T_{33}-\begin{bmatrix}(b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt}=(b_{33})^{(6)}T_{32}-\begin{bmatrix}(b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt}=(b_{34})^{(6)}T_{33}-\begin{bmatrix}(b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}\\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{34} \quad (90)$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3
$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3
$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3
$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3
$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2, and 3
$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{c} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{c} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third
augmentation coefficient for category 1, 2 and 3
$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth
augmentation coefficient for category 1, 2 and 3
$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth
augmentation coefficient for category 1, 2 and 3
$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth
augmentation coefficient for category 1, 2 and 3
$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh
augmentation coefficient for category 1, 2 and 3
$\boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth
augmentation coefficient for 1,2,3
$\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth
augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\boxed{-(b_{44}'')^{(9)}(G_{47},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\boxed{-(b_{45}'')^{(9)}(G_{47},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\boxed{-(b_{46}'')^{(9)}(G_{47},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{44}'')^{(9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3
$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3
$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3
$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(MMMM) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(NNNN) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(OOOO) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(PPPP)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(QQQQ) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(RRRR) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\le\ \left(\hat{P}_{13}\right)^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\le\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\le\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^{0} + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^{0} + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^{0} + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^{0} + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(p) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)} t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18}. indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16}, G_{18} and G_{16}, G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18}. The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2:

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

208

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)}$$

209

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)}$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),\left((G_{23})^{(2)},(T_{23})^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

214

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\,((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$\left|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{36})^{(7)}\left|G_{37}^{(1)}-G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a''_{36})^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0, (\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0, (\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(\nu_1)^{(6)}, (\nu_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, \nu^{(6)}, u^{(6)}$: 350

By $(\nu_1)^{(6)} > 0$, $(\nu_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)},, (\bar{\nu}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21}, t)\right) - (a_{21}'')^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \le \nu^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(p) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523

A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FORTY FIVE

Electric Dipole Control: Tanquam Ex Ungue Leonem-Technica Impendi Nationi

INTRODUCTION—VARIABLES USED

In physics, the electric dipole moment is a measure of the separation of positive and negative electrical charges in a system of charges, that is, a measure of the charge system's overall polarity. The SI units are Coulomb-meter (C m). This article is limited to static phenomena, and does not describe time-dependent or dynamic polarization.

NOTATION

Module One

BFGS on X space

With BFGS the search direction is given by $p_k = -H_k \nabla \hat{J}(\lambda_k)$.
By the Sherman-Morrison-Woodbury formula, we establish a recurrence for H.

$$H_{k+1} = H_k + \frac{s_k^\top y_k + y_k^\top H_k y_k}{(s_k^\top y_k)^2}(s_k s_k^\top) - \frac{H_k y_k s_k^\top + s_k y_k^\top H_k}{s_k^\top y_k}$$

where $s_k = \tau_k p_k$. Supposing X is either $L^2(0, T; \mathbb{R})$ or $H^1(0, T; \mathbb{R})$, the function space analog of the outer product is a dyadic operator $x \otimes y : X \to X$. The action of this operator on a third element $v \in X$ can be expressed in terms of the inner product $(x \otimes y)\, v = (y, z)_X\, v$.
From the recursion relation for H, we obtain

$$p_k = -H_0 g_k - \sum_{j=0}^{k-1} c_j [d_j (s_j, g_k)_X\, s_j - (z_j, g_k)_X\, s_j - (s_j, g_k)_X\, z_j]$$

where $c_j = (s_j, y_j)_X^{-1}$, $d_j = 1 + c_j (y_j, z_j)$, and for $z_k = H_k y_k$, we have.

$$z_k = H_0 y_k + \sum_{j=0}^{k-1} c_j \{[d_j (s_j, y_k)_X - (z_j, y_k)_X]\, s_j - (s_j, y_k)_X\, z_j\}.$$

G_{13} : Category one of LHS of the equation representative of Sherman-Morrison-Woodbury formula which establish recurrence of H

G_{14} : Category two of LHS of the equation representative of Sherman-Morrison-Woodbury formula which establish recurrence of H

G_{15} : Category three onLHS of the equation representative of Sherman-Morrison-Woodbury formula which establish recurrence of H

T_{13} : Category one of RHS of the equation representative of Sherman-Morrison-Woodbury formula which establish recurrence of H

T_{14} : Category two of RHS of the equation representative of Sherman-Morrison-Woodbury formula which establish recurrence of H

T_{15} : Category three of RHS of the equation representative of Sherman-Morrison-Woodbury formula which establish recurrence of H

Module Two

Electronic states of a charged particle in a well potential

The control of quantum electronic states has a host of applications such as control of photochemical processes and semiconductor lasers.

Consider a confining potential $V_0(x)$ with a 'well' envelope. The eigenproblem

$$\left\{-\partial_x^2 + V_0(x) - E_j\right\}\phi_j(x) = 0, \quad j = 1, 2, \ldots,$$

defines eigenfunctions representing the eigenstates with energy E_j.

A representative potential with applications in semiconductor nanostructures is the infinite barrier well potential where $V_0(x) = 0$ for $x \in (0, \ell)$ and $V_0(0) = +\infty$ and $V_0(\ell) = +\infty$.
The infinite barrier condition is equivalent to homogeneous Dirichlet boundary conditions for the wavefunction and thus we have

$$E_j = \frac{j^2\pi^2}{\ell^2} \quad \text{and} \quad \phi_j(x) = \sin(j\pi x/\ell).$$



G_{16} : Category one of LHS of the equation constitutive of infinite barrier conditions for the wave function

G_{17} : Category two of LHS of the equation constitutive of infinite barrier conditions for the wave function

G_{18} : Category three of LHS of the equation constitutive of infinite barrier conditions for the wave function

T_{16} : Category one of RHS of the equation constitutive of infinite barrier conditions for the wave function

T_{17} : Category two of RHS of the equation constitutive of infinite barrier conditions for the wave function

T_{18} : Category three of RHS of the equation constitutive of infinite barrier conditions for the wave function

Module three

Electric dipole control

Consider a control field modeling a laser pulse. Using the dipole approximation results in the following

$$V(x,t) = V_0(x) + u(t)\,x$$

where $u : (0,T) \to \mathbb{R}$ is the modulating control amplitude.
The quantum state of a charged particle subject to this potential is governed by the time-dependent Schrödinger equation $(c(\psi,u)=0)$

$$i\frac{\partial}{\partial t}\psi(x,t) = \left\{-\frac{\partial}{\partial x^2} + V(x,t)\right\}\psi(x,t), \qquad (x,t) \in Q = \Omega \times (0,T),$$

Objective of the control

$$J(\psi,u) := \frac{1}{2}\left(1 - \|P\psi(\cdot,T)\|^2_{\mathcal{H}}\right) + \frac{\gamma}{2}\|u\|^2_{\mathcal{U}}$$

where the projector $P\psi = (\psi_d,\psi)_{\mathcal{H}}\,\psi_d$ and $\|u\|^2_{\mathcal{U}} = \|u\|^2 + \alpha\|\dot{u}\|^2$



G_{20} : Category one of LHS of the equation representative and constitutive bulwark, citadel, defense, fortification of the quantum state of the charged particle subject to the potential mentioned above which is governed by the time dependent Schrõdinger equation(The model predicts, projects, forecasts, prognosticates and studies the system)

G_{21} : Category two of LHS of the equation representative and constitutive bulwark, citadel, defense, fortification of the quantum state of the charged particle subject to the potential mentioned above which is governed by the time dependent Schrõdinger equation(The model predicts, projects, forecasts, prognosticates and studies the system)

G_{22} : Category three of LHS of the equation representative and constitutive bulwark, citadel, defense, fortification of the quantum state of the charged particle subject to the potential mentioned above which is governed by the time dependent Schrõdinger equation(The model predicts, projects, forecasts, prognosticates and studies the system)

T_{20} : Category one of RHS of the equation representative and constitutive bulwark, citadel, defense, fortification of the quantum state of the charged particle subject to the potential mentioned above which is governed by the time dependent Schrõdinger equation(The model predicts, projects, forecasts, prognosticates and studies the system)

T_{21} : Category two of RHS of the equation representative and constitutive bulwark, citadel, defense, fortification of the quantum state of the charged particle subject to the potential mentioned above which is governed by the time dependent Schrõdinger equation(The model predicts, projects, forecasts, prognosticates and studies the system)

T_{22} : Category three of RHS of the equation representative and constitutive bulwark, citadel, defense,

fortification of the quantum state of the charged particle subject to the potential mentioned above which is governed by the time dependent Schrõdinger equation(The model predicts, projects, forecasts, prognosticates and studies the system)

Module four

Discrete optimality system

Let S corresponds to multiplication by i. We have that

$$\mathsf{S} = \begin{pmatrix} 0 & -\mathsf{I} \\ \mathsf{I} & 0 \end{pmatrix}, \quad (\psi_d, \psi)_{\mathcal{H}} \text{ corresponds to } \begin{pmatrix} \mathsf{y_d}^\top \\ \mathsf{y_d}^\top \mathsf{S} \end{pmatrix} \mathsf{y}$$

In this representation, we can rewrite the objective in the form

$$J(\mathsf{y},\mathsf{u}) = \frac{1}{2}\left[1 - \mathsf{y}_{N_t}^\top \begin{pmatrix} \mathsf{y_d} & -\mathsf{S}\mathsf{y_d} \end{pmatrix} \begin{pmatrix} \mathsf{y_d}^\top \\ \mathsf{y_d}^\top \mathsf{S} \end{pmatrix} \mathsf{y}_{N_t}\right] + \frac{\gamma}{2}\mathsf{u}^\top \mathsf{K}\mathsf{u}$$

The matrix K is the discretization of $I - \alpha\partial_t^2$. We have the Lagrangian

$$\mathsf{L}(\mathsf{y},\mathsf{u},\mathsf{p}) = J(\mathsf{y},\mathsf{u}) + \sum_{k=1}^{N_t} \mathsf{p}_k^\top \mathsf{c}_k(\mathsf{y},\mathsf{u})$$

Differentiating this Lagrangian with respect to its arguments and setting the derivatives to zero gives the discrete optimality system

$$\begin{aligned} \mathsf{B}_k \mathsf{y}_k &= \mathsf{B}_k^\top \mathsf{y}_{k-1} \\ \mathsf{B}_k^\top \mathsf{p}_k &= \mathsf{B}_{k+1}\mathsf{p}_{k+1} \\ \gamma \mathsf{K}\mathsf{u} &= \mathsf{f} \end{aligned}$$



G_{24} : Category one of LHS of the matrix K which is the discretization of 1-$\alpha\partial_t^2$ with the Lagrangian

G_{25} : Category two of LHS of the matrix K which is the discretization of 1-$\alpha\partial_t^2$ with the Lagrangian

G_{26} : Category three of LHS of the matrix K which is the discretization of 1-$\alpha\partial_t^2$ with the Lagrangian

T_{24} : Category one of RHS of the matrix K which is the discretization of 1-$\alpha\partial_t^2$ with the Lagrangian

T_{25} : Category two of RHS of the matrix K which is the discretization of 1-$\alpha\partial_t^2$ with the Lagrangian

T_{26} : Category three of RHS of the matrix K which is the discretization of 1-$\alpha\partial_t^2$ with the Lagrangian

Module five

Coherent quantum LQG control Hendra I. Nurdina et al, http://dx.doi.org/10.1016/j.automatica.2009.04.018,

(1) Based on a recently developed notion of physical realisability for quantum linear stochastic systems, authors formulate a quantum LQG optimal control problem **for (e&eb)** quantum linear stochastic systems where the controller itself may also be a quantum system and the plant output signal can be fully quantum.

(2) Such a control scheme is often referred to in the quantum control literature as "coherent feedback control". It distinguishes the present work from previous works on the quantum LQG problem where measurement is performed on the plant and the measurement signals **are used** as the input to a fully classical controller with no quantum degrees of freedom.
(3) The difference in formulation is the presence of additional non-linear and linear constraints on the coefficients of the sought after controller, **rendering the(=) problem** as a type of constrained controller design problem.
(4) Due to the presence of these constraints, our problem is inherently computationally hard and this also distinguishes it in an important way from the standard LQG problem. Authors propose a numerical procedure for solving this problem **based on an** alternating projections algorithm and, as an initial demonstration of the feasibility of this approach, provide fully quantum controller design examples in which numerical solutions to the problem were successfully obtained.
(5) For comparison, they also consider the case of classical linear controllers that use direct or indirect measurements, **and show that there exists** a fully quantum linear controller which offers an improvement in performance over the classical ones.

G_{28} : Category one of recently developed notion of physical realisability for quantum linear stochastic systems, quantum LQG optimal control problem based on which the problem is formulated;

G_{29} : Category two of recently developed notion of physical realisability for quantum linear stochastic systems, quantum LQG optimal control problem based on which the problem is formulated ;

G_{30} : Category three of recently developed notion of physical realisability for quantum linear stochastic systems, quantum LQG optimal control problem based on which the problem is formulated;

T_{28} : Category one of quantum linear stochastic systems where the controller itself may also be a quantum system and the plant output signal can be fully quantum.

T_{29} : Category two of quantum linear stochastic systems where the controller itself may also be a quantum system and the plant output signal can be fully quantum.

T_{30} : Category three of quantum linear stochastic systems where the controller itself may also be a quantum system and the plant output signal can be fully quantum.

Module six

Monotonically convergent optimal control theory of quantum systems with spectral constraints on the control field M. Lapert et al

(1) Authors propose a monotonically convergent algorithm which can enforce spectral constraints **on the** control field (and extends to arbitrary filters).
(2) The procedure differs from standard algorithms in that at each iteration, the control field **is taken (=) as a** linear combination of the control field (computed by the standard algorithm) and the filtered field.
(3) The parameter of the linear combination is chosen to respect the monotonic behavior of the algorithm and to be as close to the filtered field as possible. They test the efficiency of this method on molecular alignment. Using bandpass filters, **show (eb)** how to select particular rotational transitions to reach high alignment efficiency.
(4) They also consider spectral constraints corresponding to experimental conditions **using** pulse-shaping techniques. We determine an optimal solution that could be implemented experimentally with this technique.© 2009 The American Physical SocietyURL:http://link.aps.org/doi/10.1103/PhysRevA.79.063411DOI:10.1103/PhysRevA.79.0634

G_{32} : Category one of control field (and extends to arbitrary filters).

G_{33} : Category two of control field (and extends to arbitrary filters).

G_{34} : Category three of control field (and extends to arbitrary filters).

T_{32} : Category one of monotonically convergent algorithm which can enforce spectral constraints

T_{33} : Category two of monotonically convergent algorithm which can enforce spectral constraints

T_{34} : Category three of monotonically convergent algorithm which can enforce spectral constraints

Module seven

G_{36} : Category one of linear combination of the control field (computed by the standard algorithm) and the filtered field.

G_{37} : Category two of linear combination of the control field (computed by the standard algorithm) and the filtered field.

G_{38} : Category three of linear combination of the control field (computed by the standard algorithm) and the filtered field.

T_{36} : Category one of procedure differs from standard algorithms in that at each iteration, the control field

T_{37} : Category two of procedure differs from standard algorithms in that at each iteration, the control field

T_{38} : Category three of procedure differs from standard algorithms in that at each iteration, the control field

Module eight

G_{40} : Category one of parameter of the linear combination is chosen to respect the monotonic behavior of the algorithm and to be as close to the filtered field as possible. They test the efficiency of this method on molecular alignment. Using bandpass filters

G_{41} : Category two of parameter of the linear combination is chosen to respect the monotonic behavior of the algorithm and to be as close to the filtered field as possible. They test the efficiency of this method on molecular alignment. Using bandpass filters

G_{42} : Category three of parameter of the linear combination is chosen to respect the monotonic behavior of the algorithm and to be as close to the filtered field as possible. They test the efficiency of this method on molecular alignment. Using bandpass filters

T_{40} : Category one of selection of particular rotational transitions to reach high alignment efficiency.

T_{41} : Category two of selection of particular rotational transitions to reach high alignment efficiency.

T_{42} : Category three of selection of particular rotational transitions to reach high alignment efficiency.

Module Nine

G_{44} : Category one of pulse-shaping techniques towards the end of consummation of the determination of an optimal solution that could be implemented experimentally with this technique

G_{45} : Category two of pulse-shaping techniques towards the end of consummation of the determination of an optimal solution that could be implemented experimentally with this technique

G_{46} : Category three of pulse-shaping techniques towards the end of consummation of the determination of an optimal solution that could be implemented experimentally with this technique

T_{44} : Category one of spectral constraints corresponding to experimental conditions

T_{45} : Category two of spectral constraints corresponding to experimental conditions

T_{46} : Category three of spectral constraints corresponding to experimental conditions

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad 63$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition

coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{20})^{(3)}\left[+(a''_{20})^{(3)}(T_{21},t)\right]\left[+(a''_{16})^{(2,2,2)}(T_{17},t)\right]\left[+(a''_{13})^{(1,1,1,)}(T_{14},t)\right] \\ \left[+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)\right]\left[+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)\right]\left[+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{36})^{(7,7,7,7)}(T_{37},t)\right]\left[+(a''_{40})^{(8,8,8,8)}(T_{41},t)\right]\left[+(a''_{44})^{(9,9,9)}(T_{45},t)\right] \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c} (a'_{21})^{(3)}\left[+(a''_{21})^{(3)}(T_{21},t)\right]\left[+(a''_{17})^{(2,2,2)}(T_{17},t)\right]\left[+(a''_{14})^{(1,1,1,)}(T_{14},t)\right] \\ \left[+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)\right]\left[+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)\right]\left[+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{37})^{(7,7,7,7)}(T_{37},t)\right]\left[+(a''_{41})^{(8,8,8,8)}(T_{41},t)\right]\left[+(a''_{45})^{(9,9,9)}(T_{45},t)\right] \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{22})^{(3)}\left[+(a''_{22})^{(3)}(T_{21},t)\right]\left[+(a''_{18})^{(2,2,2)}(T_{17},t)\right]\left[+(a''_{15})^{(1,1,1,)}(T_{14},t)\right] \\ \left[+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)\right]\left[+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)\right]\left[+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{38})^{(7,7,7,7)}(T_{37},t)\right]\left[+(a''_{42})^{(8,8,8,8)}(T_{41},t)\right]\left[+(a''_{46})^{(9,9,9)}(T_{45},t)\right] \end{array}\right] G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for

category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad 78$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

are fourth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3
And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3
$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3
$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3
$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3
$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3
$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)} & +(a_{36}'')^{(7)}(T_{37},t) & +(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)} & +(a_{37}'')^{(7)}(T_{37},t) & +(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)} & +(a_{38}'')^{(7)}(T_{37},t) & +(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a_{36}'')^{(7)}(T_{37},t)$, $(a_{37}'')^{(7)}(T_{37},t)$, $(a_{38}'')^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)} \boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} \boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ \ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(SSSS) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(TTTT) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(UUUU) $\lim_{T_2 \to \infty} \ (a_i'')^{(7)}\,(T_{37}, t) = (p_i)^{(7)}$
(VVVV)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(WWWW) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(XXXX) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ \ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{16})^{(2)}$, $T_i^0 \leq (\hat{Q}_{16})^{(2)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(q) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on G(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right| e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right| \; e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\} ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right| e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right) d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right) - (a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right| \; e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} \} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)} - G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)};\ (G_{23})^{(2)}, (T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)} \left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)} \left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+\ (a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right) d\left(\left((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\,((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)},, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},, (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \qquad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \qquad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \qquad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}\ (v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}\ (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \qquad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \qquad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \qquad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(q) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FORTY SIX

Quantum Dots: A Poindexter Poinsettia

INTRODUCTION—VARIABLES USED

A quantum dot **is a** semiconductor whose excitons are confined in all three spatial dimensions. The electronic properties of these materials are intermediate between those of bulk semiconductors and of discrete molecules. Quantum dots were discovered in the early 1980s by Alexei Ekimov in a glass matrix and by Louis E. Brus in colloidal solutions. The term "quantum dot" was coined by Mark Reed.

Researchers have studied applications for quantum dots in transistors, solar cells, LEDs, and diode lasers. They have also investigated quantum dots as agents for medical imaging and as possible qubits in quantum computing.

(1) Quantum dots are semiconductors whose electronic characteristics are closely **related to the** size and shape of the individual crystal. Size and band gap are **inversely related** in quantum dots.
(2) For example, in fluorescent dye applications, emission frequencies increase as the size of the quantum dot decreases, **resulting in** a color shift from red to blue in the light emitted. Excitation and emission of the quantum dot are therefore highly tunable.
(3) Size of the crystals can be controlled during synthesis **using (e) which property**, the conductive properties can be carefully controlled. Quantum dots of different sizes can be assembled into a gradient multi-layer nanofilm

Electron spins in quantum dots for spintronics and quantum computation Hans-Andreas Engel et al http://dx.doi.org/10.1016/S0038-1098 (01)00110-7

(4) Coherent manipulation, filtering, and measurement of electronic spin in quantum dots and other nanostructures have **promising (e&eb) applications** in conventional and in quantum information processing and transmission.
(5) Authors present an overview of our theoretical proposal to implement a quantum computer **using (e)** electron spins in quantum dots as qubits.
(6) They discuss all necessary requirements towards a scalable quantum computer including one- and two qubit gates and read in/out tasks. They then present some concepts for promising single quantum dot devices which eventually **could be used** as building blocks for sophisticated spintronic devices.
(7) They show how a single quantum dot **can act (e) as an** efficient spin filter.
(8) Further, in combination with an ESR source, a quantum dot **can be used as** a single spin memory or as a spin pump.
(9) In addition, the sequential tunneling current through a quantum dot in the presence of an ESR field can **exhibit a** resonance whose line width is determined by the decoherence time T2 of a single dot-spin.

NOTATION

Module One

Quantum dots are semiconductors whose electronic characteristics are closely **related to the** size and shape of the individual crystal. Size and band gap are **inversely related** in quantum dots

G_{13} : Category one of quantum dots are semiconductors with electronic characteristics

G_{14} : Category two of quantum dots are semiconductors with electronic characteristics

G_{15} : Category three of quantum dots are semiconductors with electronic characteristics

T_{13} : Category one of size and shape of the individual crystal. Size and band gap are **inversely related** in quantum dots

T_{14} : Category two of size and shape of the individual crystal. Size and band gap are **inversely related** in quantum dots

T_{15} : Category three of size and shape of the individual crystal. Size and band gap are **inversely related** in quantum dots

Module Two

For example, in fluorescent dye applications, emission frequencies increase as the size of the quantum dot decreases, **resulting in** a color shift from red to blue in the light emitted. Excitation and emission of the quantum dot are therefore highly tunable

G_{16} : Category one of fluorescent dye applications, emission frequencies increase as the size of the quantum dot decreases

G_{17} : Category two of fluorescent dye applications, emission frequencies increase as the size of the quantum dot decreases

G_{18} : Category three of fluorescent dye applications, emission frequencies increase as the size of the quantum dot decreases

T_{16} : Category one of color shift from red to blue in the light emitted. Excitation and emission of the quantum dot are therefore highly tunable

T_{17} : Category two of color shift from red to blue in the light emitted. Excitation and emission of the quantum dot are therefore highly tunable

T_{18} : Category three of color shift from red to blue in the light emitted. Excitation and emission of the quantum dot are therefore highly tunable

Module three

Size of the crystals can be controlled during synthesis **using (e) which property**, the conductive properties can be carefully controlled. Quantum dots of different sizes can be assembled into a gradient multi-layer nanofilm

G_{20} : Category one of conductive properties can be carefully controlled. Quantum dots of different sizes can be assembled into a gradient multi-layer nanofilm

G_{21} : Category two of conductive properties can be carefully controlled. Quantum dots of different sizes can be assembled into a gradient multi-layer nanofilm

G_{22} : Category three of conductive properties can be carefully controlled. Quantum dots of different sizes can be assembled into a gradient multi-layer nanofilm

T_{20} : Category one of Size of the crystals can be controlled during synthesis

T_{21} : Category two of Size of the crystals can be controlled during synthesis

T_{22} : Category three of Size of the crystals can be controlled during synthesis

Module four

Coherent manipulation, filtering, and measurement of electronic spin in quantum dots and other nanostructures have **promising (e&eb) applications** in conventional and in quantum information processing and transmission.

G_{24} : Category one of Coherent manipulation, filtering, and measurement of electronic spin in quantum dots and other nanostructures

G_{25} : Category two of Coherent manipulation, filtering, and measurement of electronic spin in quantum dots and other nanostructures

G_{26} : Category three of Coherent manipulation, filtering, and measurement of electronic spin in quantum dots and other nanostructures

T_{24} : Category one of conventional and in quantum information processing and transmission.

T_{25} : Category two of conventional and in quantum information processing and transmission.

T_{26} : Category three of conventional and in quantum information processing and transmission.

Module five

Authors present an overview of our theoretical proposal to implement a quantum computer **using (e)** electron spins in quantum dots as qubits.

G_{28} : Category one of electron spins in quantum dots as qubits

G_{29} : Category two of electron spins in quantum dots as qubits

G_{30} : Category three of electron spins in quantum dots as qubits

T_{28} : Category one of theoretical proposal to implement a quantum computer

T_{29} : Category two of theoretical proposal to implement a quantum computer

T_{30} : Category three of theoretical proposal to implement a quantum computer

Module six

They discuss all necessary requirements towards a scalable quantum computers including one- and two qubit gates and read in/out tasks with some concepts for promising single quantum dot devices which eventually **could be used** as building blocks for sophisticated spintronic devices.

G_{32} : Category one of building blocks for sophisticated spintronic devices

G_{33} : Category two of building blocks for sophisticated spintronic devices

G_{34} : Category three of building blocks for sophisticated spintronic devices

T_{32} : Category one of scalable quantum computers including one- and two qubit gates and read in/out tasks

with some concepts for promising single quantum dot devices which eventually

T_{33} : Category two of scalable quantum computers including one- and two qubit gates and read in/out tasks with some concepts for promising single quantum dot devices which eventually

T_{34} : Category three of scalable quantum computers including one- and two qubit gates and read in/out tasks with some concepts for promising single quantum dot devices which eventually

Module seven

They show how a single quantum dot **can act (e) as an** efficient spin filter

G_{36} : Category one of efficient spin filter

G_{37} : Category two of efficient spin filter

G_{38} : Category three of efficient spin filter

T_{36} : Category one of single quantum dot

T_{37} : Category two of single quantum dot

T_{38} : Category three of single quantum dot

Module eight

Further, in combination with an ESR source, a quantum dot **can be used as** a single spin memory or as a spin pump

G_{40} : Category one of single spin memory or as a spin pump

G_{41} : Category two of single spin memory or as a spin pump

G_{42} : Category three of single spin memory or as a spin pump

T_{40} : Category one of combination with an ESR source, a quantum dot

T_{41} : Category two of combination with an ESR source, a quantum dot

T_{42} : Category three of combination with an ESR source, a quantum dot

Module Nine

In addition, the sequential tunneling current through a quantum dot in the presence of an ESR field can **exhibit a** resonance whose line width is determined by decoherence time T2 of a single dot-spin.

G_{44} : Category one of sequential tunneling current through a quantum dot in the presence of an ESR field

G_{45} : Category two of sequential tunneling current through a quantum dot in the presence of an ESR field

G_{46} : Category three of sequential tunneling current through a quantum dot in the presence of an ESR field

T_{44} : Category one of resonance whose line width is determined by decoherence time T2 of a single dot-spin

T_{45} : Category two of resonance whose line width is determined by decoherence time T2 of a single dot-spin

T_{46} : Category three of resonance whose line width is determined by decoherence time T2 of a single dot-spin

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}), t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}), t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}), t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{matrix}(a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{matrix}(a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{matrix}(a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{matrix}(b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}\end{matrix}\right]T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{matrix}(b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}\end{matrix}\right]T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) \end{bmatrix} G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)}\ \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\ \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\ \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{c} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\ \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\ \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\ \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{array}\right] T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{c} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\ \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\ \boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\ \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{array}\right] T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$

$are\ fourth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
$are\ fifth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
$are\ sixth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
$are\ seventh\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
$are\ eighth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

$Where\ -(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ $are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ $are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ $are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad (84)$$

$where\ \boxed{-(b_{28}'')^{(5)}(G_{31},t)}\ ,\boxed{-(b_{29}'')^{(5)}(G_{31},t)}\ ,\boxed{-(b_{30}'')^{(5)}(G_{31},t)}\ \ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{24}'')^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b_{25}'')^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b_{26}'')^{(4,4,)}(G_{27},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{32}'')^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b_{33}'')^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b_{34}'')^{(6,6,6)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{13}'')^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b_{14}'')^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b_{15}'')^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\ \boxed{-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\boxed{-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)}\ ,\boxed{-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)}\ ,\boxed{-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt}=(a_{32})^{(6)}G_{33} \qquad 85$$

$$-\left[\begin{matrix}(a_{32}')^{(6)}\boxed{+(a_{32}'')^{(6)}(T_{33},t)}\ \boxed{+(a_{28}'')^{(5,5,5)}(T_{29},t)}\ \boxed{+(a_{24}'')^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{32}$$

$$\frac{dG_{33}}{dt}=(a_{33})^{(6)}G_{32}-\left[\begin{matrix}(a_{33}')^{(6)}\boxed{+(a_{33}'')^{(6)}(T_{33},t)}\ \boxed{+(a_{29}'')^{(5,5,5)}(T_{29},t)}\ \boxed{+(a_{25}'')^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{33} \qquad 86$$

$$\frac{dG_{34}}{dt}=(a_{34})^{(6)}G_{33}-\left[\begin{matrix}(a_{34}')^{(6)}\boxed{+(a_{34}'')^{(6)}(T_{33},t)}\ \boxed{+(a_{30}'')^{(5,5,5)}(T_{29},t)}\ \boxed{+(a_{26}'')^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{34} \qquad 87$$

$\boxed{+(a_{32}'')^{(6)}(T_{33},t)}\ ,\boxed{+(a_{33}'')^{(6)}(T_{33},t)}\ ,\boxed{+(a_{34}'')^{(6)}(T_{33},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a_{28}'')^{(5,5,5)}(T_{29},t)}\ ,\boxed{+(a_{29}'')^{(5,5,5)}(T_{29},t)}\ ,\boxed{+(a_{30}'')^{(5,5,5)}(T_{29},t)}\ are\ second\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a_{24}'')^{(4,4,4,)}(T_{25},t)}\ ,\boxed{+(a_{25}'')^{(4,4,4,)}(T_{25},t)}\ ,\boxed{+(a_{26}'')^{(4,4,4,)}(T_{25},t)}\ \ are\ third\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation

coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34} \quad (90)$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$ *95*

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{c} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{c} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\ ,\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)},\ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ \ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)},\ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \le\ \ (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G\to\infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)},\ \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \le (\hat{k}_{32})^{(6)}|T_{33} - \ T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(YYYY) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(ZZZZ) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(AAAAA) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(BBBBB)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(CCCCC) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(DDDDD) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\le\ \left(\hat{P}_{13}\right)^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\le\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\le\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^{0} + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^{0} + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^{0} + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^{0} + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(r) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \tag{183}$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \tag{184}$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric (185)

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \tag{186}$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)} t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

208

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)}$$

209

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)}$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),\left((G_{23})^{(2)},(T_{23})^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

214

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{28})^{(5)}|G_{29}^{(1)}-G_{29}^{(2)}|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}|G_{28}^{(1)}-G_{28}^{(2)}|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)|G_{28}^{(1)}-G_{28}^{(2)}|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)|\;e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{31})^{(1)}-(G_{31})^{(2)}|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\,(G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right| \; e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)}G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)}G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)}T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}$, ***if*** $(v_0)^{(6)} < (v_1)^{(6)}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}$, ***if*** $(v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)}$,
and $\boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}$, ***if*** $(\bar{v}_1)^{(6)} < (v_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^{0}e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^{0}e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^{0}}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t}-e^{-(b'_{34})^{(6)}t}\right]+T_{34}^{0}e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^{0}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T_{34}^{0}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^{0}e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^{0}e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^{0}e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}}T_{44}^{0}e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^{0}}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t}-e^{-(b'_{46})^{(9)}t}\right]+T_{46}^{0}e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^{0}}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t}-e^{-(R_2)^{(9)}t}\right]+T_{46}^{0}e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^{0}}{G_{14}^{0}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)}+(C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)}=\frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad (472)$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad (473)$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad (474)$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad (475)$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad (476)$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad (477)$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad (478)$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad (479)$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad (480)$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad (481)$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad (482)$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad (483)$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad (484)$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad (484\text{A})$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: (485)

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(r) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^{*} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^{*} for which $f(T_{33}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

Definition and uniqueness of T_{37}^{*} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^{*} for which $f(T_{37}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523
A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FORTY SEVEN

Spintronics: Gotterdammerung Model

INTRODUCTION—VARIABLES USED

spintronics (a portmanteau meaning "spin transport electronics", also known as magneto electronics, is an

emerging technology exploiting both the intrinsic spin of the electron and its associated magnetic moment, in addition to its fundamental electronic charge, in solid-state devices. Ferromagnetic semiconductor sources (like manganese-doped gallium arsenide GaMnAs), increase the interface resistance with a tunnel barrier, or using hot-electron injection.

Spin detection in semiconductors is another challenge, met with the following techniques:

Faraday/Kerr rotation of transmitted/reflected photons

Circular polarization analysis of electroluminescence

Nonlocal spin valve (adapted from Johnson and Silsbee's work with metals)

Ballistic spin filtering

The latter technique was used to overcome the lack of spin-orbit interaction and materials issues to achieve spin transport in silicon, the most important semiconductor for electronics.

Because external magnetic fields (and stray fields from magnetic contacts) can cause large Hall effects and magnetoresistance in semiconductors (which mimic spin-valve effects), the only conclusive evidence of spin transport in semiconductors is demonstration of spin precession and dephasing in a magnetic field non-collinear to the injected spin orientation. This is called the Hanle effect.

Applications

Applications using spin-polarized electrical injection have shown threshold current reduction and controllable circularly polarized coherent light output. Examples include semiconductor lasers. Future applications may include a spin-based transistor having advantages over MOSFET devices such as steeper sub-threshold slope.

Magnetic-tunnel transistor: The magnetic-tunnel transistor with a single base layer, by van Dijken et al. and Jiang et al.,[has the following terminals:

Emitter (FM1): It injects spin-polarized hot electrons into the base.


Base (FM2): Spin-dependent scattering takes place in the base. It also serves as a spin filter.

Collector (GaAs): A Schottky barrier is formed at the interface. This collector regions only collects electrons when they have enough energy to overcome the Schottky barrier, and when there are states available in the semiconductor.

Electron spins in quantum dots for spintronics and quantum computation Hans-Andreas Engel et al http://dx.doi.org/10.1016/S0038-1098 (01)00110-7


(1) Finally, authors consider mobile non-local spin entangled electrons as needed for quantum communication. They propose how to create such EPR pairs **by (e&eb) means of** Andreev tunneling at a superconductor-normal junction and discuss experimental setups in which spin entanglement may be detected via transport measurements.

Figures and tables from this article:



(2) Fig. 1. Quantum dot array, is controlled **(e&eb) by** electrical gating.
(3) The electrodes (dark gray) **define** quantum dots (circles) by confining electrons.
(4) These electrons can be moved by electrical gating into the magnetized or high-g layer, **producing** locally different Zeeman splittings.
(5) Alternatively, magnetic field gradients can be applied, as e.g. **produced (e) by a** current wire (indicated on the left of the dot-array).
(6) Then, since every dot-spin is subject to a different Zeeman splitting, the spins can be addressed individually, e.g. through ESR pulses of an additional in-plane magnetic ac field with the corresponding Larmor frequency $\omega L=g\mu BB\perp/\hbar$. Such mechanisms can be **used (e) for** single-spin rotations and the initialization step, see text.
(7) The exchange coupling between the quantum dots **can be controlled** by lowering the tunnel barrier between the dots. In this figure, the two rightmost dots are drawn schematically as tunnel-coupled.

Figure options



(8) Fig. 2. (a) Implementation of a spin filter with a quantum dot coupled to unpolarized leads with chemical potentials μ1, 2. The dot is in the sequential tunneling regime, $\mu1>ES>\mu2>ES-\Delta z$, with the singlet/triplet levels ES/ET+ (counting from E↑=0) and Zeeman splitting $\Delta z=g\mu BBz$.
(9) (b) By shifting the energy levels such that $ES>\mu1>ES-\Delta z>\mu2$, the setup **can be used to** measure the single spin decoherence time T2. If initially the spin-state on the dot is $|\uparrow\rangle$, sequential tunneling is blocked by energy conservation.

NOTATION

Module One

Authors consider mobile non-local spin entangled electrons as needed for quantum communication. They propose how to create such EPR pairs **by (e&eb) means of** Andreev tunneling at a superconductor-normal junction and discuss experimental setups in which spin entanglement may be detected via transport measurements.

G_{13} : Category one of mobile non-local spins entangled electrons as needed for quantum communication. They propose how to create such EPR pairs

G_{14} : Category two of mobile non-local spins entangled electrons as needed for quantum communication. They propose how to create such EPR pairs

G_{15} : Category three of mobile non-local spins entangled electrons as needed for quantum communication. They propose how to create such EPR pairs

T_{13} : Category one of Andreev tunneling at a superconductor-normal junction and discuss experimental setups in which spin entanglement may be detected via transport measurements.

T_{14} : Category two of Andreev tunneling at a superconductor-normal junction and discuss experimental

setups in which spin entanglement may be detected via transport measurements.

T_{15} : Category three of Andreev tunneling at a superconductor-normal junction and discuss experimental setups in which spin entanglement may be detected via transport measurements.

Module Two

Quantum dot array, is controlled **(e&eb) by** electrical gating

G_{16} : Category one of Quantum dot array; electrical gating

G_{17} : Category two of Quantum dot array; electrical gating

G_{18} : Category three of Quantum dot array; electrical gating

T_{16} : Category one of electrical gating ;Quantum dot array

T_{17} : Category two of electrical gating ;Quantum dot array

T_{18} : Category three of electrical gating ;Quantum dot array

Module three

The electrodes (dark gray) **define** quantum dots (circles) by confining electrons

G_{20} : Category one of electrodes (dark gray)

G_{21} : Category two of electrodes (dark gray)

G_{22} : Category three of electrodes (dark gray)

T_{20} : Category one of quantum dots (circles) by confining electrons

T_{21} : Category two of quantum dots (circles) by confining electrons

T_{22} : Category three of quantum dots (circles) by confining electrons

Module four

These electrons can be moved by electrical gating into the magnetized or high-g layer, **producing** locally different Zeeman splittings.

G_{24} : Category one of electrons can be moved by electrical gating into the magnetized or high-g layer

G_{25} : Category two of electrons can be moved by electrical gating into the magnetized or high-g layer

G_{26} : Category three of electrons can be moved by electrical gating into the magnetized or high-g layer

T_{24} : Category one of locally different Zeeman splittings.

T_{25} : Category two of locally different Zeeman splittings.

T_{26} : Category three of locally different Zeeman splittings.

Module five

Alternatively, magnetic field gradients can be applied, as e.g. **produced (e) by a** current wire (indicated on

the left of the dot-array).

G_{28} : Category one of current wire (indicated on the left of the dot-array).

G_{29} : Category two of current wire (indicated on the left of the dot-array).

G_{30} : Category three of current wire (indicated on the left of the dot-array).

T_{28} : Category one of magnetic field gradients can be applied

T_{29} : Category two of magnetic field gradients can be applied

T_{30} : Category three of magnetic field gradients can be applied

Module six

Then, since every dot-spin is subject to a different Zeeman splitting, the spins can be addressed individually, e.g. through ESR pulses of an additional in-plane magnetic ac field with the corresponding Larmor frequency ωL=gμBB⊥/ħ. Such mechanisms can be **used (e) for** single-spin rotations and the initialization step, see text.

G_{32} : Category one of single-spin rotations and the initialization step, see text; LHS of the logarithm of the equations corresponding Larmor frequency ωL=gμBB⊥/ħ.

G_{33} : Category two of single-spin rotations and the initialization step, see text; LHS of the logarithm of the equations corresponding Larmor frequency ωL=gμBB⊥/ħ.

G_{34} : Category three of single-spin rotations and the initialization step, see text; LHS of the logarithm of the equations corresponding Larmor frequency ωL=gμBB⊥/ħ.

T_{32} : Category one of every dot-spin is subject to a different Zeeman splitting, the spins can be addressed individually, e.g. through ESR pulses of an additional in-plane magnetic ac field with the corresponding Larmor frequency ωL=gμBB⊥/ħ. Such mechanisms; RHS of LHS of the logarithm of the equations corresponding Larmor frequency ωL=gμBB⊥/ħ.

T_{33} : Category two of every dot-spin is subject to a different Zeeman splitting, the spins can be addressed individually, e.g. through ESR pulses of an additional in-plane magnetic ac field with the corresponding Larmor frequency ωL=gμBB⊥/ħ. Such mechanisms; RHS of the logarithm of the equations corresponding Larmor frequency ωL=gμBB⊥/ħ.

T_{34} : Category three of every dot-spin is subject to a different Zeeman splitting, the spins can be addressed individually, e.g. through ESR pulses of an additional in-plane magnetic ac field with the corresponding Larmor frequency ωL=gμBB⊥/ħ. Such mechanisms; RHS of the logarithm of the equations corresponding Larmor frequency ωL=gμBB⊥/ħ.

Module seven

The exchange coupling between the quantum dots **can be controlled** by lowering the tunnel barrier between the dots. In this figure, the two rightmost dots are drawn schematically as tunnel-coupled

G_{36} : Category one of exchange coupling between the quantum dots; lowering the tunnel barrier between the dots. In this figure, the two rightmost dots are drawn schematically as tunnel-coupled

G_{37} : Category two of exchange coupling between the quantum dots; lowering the tunnel barrier between the dots. In this figure, the two rightmost dots are drawn schematically as tunnel-coupled

G_{38} : Category three of exchange coupling between the quantum dots; lowering the tunnel barrier between the dots. In this figure, the two rightmost dots are drawn schematically as tunnel-coupled

T_{36} : Category one of lowering the tunnel barrier between the dots. In this figure, the two rightmost dots are drawn schematically as tunnel-coupled ;exchange coupling between the quantum dots

T_{37} : Category two of lowering the tunnel barrier between the dots. In this figure, the two rightmost dots are drawn schematically as tunnel-coupled; exchange coupling between the quantum dots

T_{38} : Category three of lowering the tunnel barrier between the dots. In this figure, the two rightmost dots are drawn schematically as tunnel-coupled ;exchange coupling between the quantum dots

Module eight

Implementation of a spin filter with a quantum dot coupled to unpolarized leads **with** chemical potentials μ1, 2. The dot is in the sequential tunneling regime, μ1>ES>μ2>ES−Δz, with the singlet/triplet levels ES/ET+ (counting from E↑=0) and Zeeman splitting Δz=gμBBz. Note the model could be applied to both tunneling regime and Zeeman splitting

G_{40} : Category one of Implementation of a spin filter with a quantum dot coupled to unpolarized leads

G_{41} : Category two of Implementation of a spin filter with a quantum dot coupled to unpolarized leads

G_{42} : Category three of Implementation of a spin filter with a quantum dot coupled to unpolarized leads

T_{40} : Category one of chemical potentials μ1, 2. The dot is in the sequential tunneling regime, μ1>ES>μ2>ES−Δz, with the singlet/triplet levels ES/ET+ (counting from E↑=0) and Zeeman splitting Δz=gμBBz.

T_{41} : Category two of chemical potentials μ1, 2. The dot is in the sequential tunneling regime, μ1>ES>μ2>ES−Δz, with the singlet/triplet levels ES/ET+ (counting from E↑=0) and Zeeman splitting Δz=gμBBz.

T_{42} : Category three of chemical potentials μ1, 2. The dot is in the sequential tunneling regime, μ1>ES>μ2>ES−Δz, with the singlet/triplet levels ES/ET+ (counting from E↑=0) and Zeeman splitting Δz=gμBBz.

Module Nine

By shifting the energy levels such that ES>μ1>ES−Δz>μ2, the setup **can be used to** measure the single spin decoherence time T2. If initially the spin-state on the dot is $|\uparrow\rangle$, sequential tunneling is blocked by energy conservation.

G_{44} : Category one of measure the single spin decoherence time T2. If initially the spin-state on the dot is $|\uparrow\rangle$, sequential tunneling is blocked by energy conservation.

G_{45} : Category two of measure the single spin decoherence time T2. If initially the spin-state on the dot is $|\uparrow\rangle$, sequential tunneling is blocked by energy conservation.

G_{46} : Category three of measure the single spin decoherence time T2. If initially the spin-state on the dot is $|\uparrow\rangle$, sequential tunneling is blocked by energy conservation.

T_{44} : Category one of shifting the energy levels such that ES>μ1>ES−Δz>μ2, the setup

T_{45} : Category two of shifting the energy levels such that ES>μ1>ES−Δz>μ2, the setup

T_{46} : Category three of shifting the energy levels such that ES>μ1>ES−Δz>μ2, the setup

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}),t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

59

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

60

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

61

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16}$$

62

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) \end{bmatrix} G_{17}$$

63

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) \end{bmatrix} G_{18}$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a_{44}'')^{(9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b_{16}')^{(2)} & -(b_{16}'')^{(2)}(G_{19},t) & -(b_{13}'')^{(1,1,)}(G,t) & -(b_{20}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{24}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{28}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{32}'')^{(6,6,6,6,6)}(G_{35},t) \\ -(b_{36}'')^{(7,7,7)}(G_{39},t) & -(b_{40}'')^{(8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b_{17}')^{(2)} & -(b_{17}'')^{(2)}(G_{19},t) & -(b_{14}'')^{(1,1,)}(G,t) & -(b_{21}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{25}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{29}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{33}'')^{(6,6,6,6,6)}(G_{35},t) \\ -(b_{37}'')^{(7,7,7)}(G_{39},t) & -(b_{41}'')^{(8,8,8)}(G_{43},t) & -(b_{45}'')^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b_{18}')^{(2)} & -(b_{18}'')^{(2)}(G_{19},t) & -(b_{15}'')^{(1,1,)}(G,t) & -(b_{22}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{26}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{30}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{34}'')^{(6,6,6,6,6)}(G_{35},t) \\ -(b_{38}'')^{(7,7,7)}(G_{39},t) & -(b_{42}'')^{(8,8,8)}(G_{43},t) & -(b_{46}'')^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad 66$$

where $-(b_{16}'')^{(2)}(G_{19},t)$, $-(b_{17}'')^{(2)}(G_{19},t)$, $-(b_{18}'')^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b_{13}'')^{(1,1,)}(G,t)$, $-(b_{14}'')^{(1,1,)}(G,t)$, $-(b_{15}'')^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b_{20}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b_{24}'')^{(4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b_{28}'')^{(5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b_{32}'')^{(6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{array}\right] T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad (75)$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$ *are fourth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$ *are fifth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$ *are sixth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$ *are seventh augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$ *are eighth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$ *are fourth detrition coefficients for category* 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$ *are fifth detrition coefficients for category* 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$

are seventh detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$

are eighth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt}=(a_{28})^{(5)}G_{29}-\left[\begin{array}{ccc}(a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt}=(a_{29})^{(5)}G_{28}-\left[\begin{array}{ccc}(a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt}=(a_{30})^{(5)}G_{29}-\left[\begin{array}{ccc}(a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt}=(b_{28})^{(5)}T_{29}-\left[\begin{array}{ccc}(b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}\end{array}\right]T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)} & +(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34} \quad 90$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition

coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37},t) & +(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37},t) & +(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37},t) & +(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a''_{36})^{(7)}(T_{37},t)$, $(a''_{37})^{(7)}(T_{37},t)$, $(a''_{38})^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation

coefficient for 1,2,3

$\frac{dT_{36}}{dt} =$ 94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$\frac{dT_{37}}{dt} =$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$\frac{dT_{38}}{dt} =$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)} G_{44}$$
$$- \left[\begin{array}{c} (a'_{45})^{(9)} \boxed{+(a''_{45})^{(9)}(T_{45},t)} \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)} G_{45}$$
$$- \left[\begin{array}{c} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)} T_{45} -$$
$$\left[\begin{array}{c} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{13}$$
$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}\left((G_{19}), t\right) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ \ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ \ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(EEEEE) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(FFFFF) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(GGGGG) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(HHHHH)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(IIIII) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(JJJJJ) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ \ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{16})^{(2)}$, $T_i^0 \leq (\hat{Q}_{16})^{(2)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)} t}$

By

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)} t}$

By

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

159

160

161

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{36})^{(7)}$, $T_i^0 \leq (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(s) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where G_i (t) $= 0$ and T_i (t) $= 0$ 200

it results

G_i (t) $\geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

T_i (t) $\geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0$ for t > 0

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(7)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}),(\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}),(\widetilde{T_{43}})\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad (333)$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad (334)$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad (335)$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad (336)$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \ \boldsymbol{if}\ (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \boldsymbol{if}\ (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

and $\boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)} (m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)} (\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

406

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case,

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

407

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t)\right) - (a_{41}'')^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(s) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FORTY EIGHT

Griffith Program: Sapientia Et Doctrina Model

INTRODUCTION—VARIABLES USED

The Griffith Program: Measurement and Control for Quantum Computing. Quantum Measurement Theory for Read-Out Devices

(1) A crucial part of quantum computing is (=) read-out.
(2) Schemes for reading out the state of a solid-state QC **include (e)** a single-electron transistor, a quantum point contact, and a micromechanical cantilever.
(3) The theory of quantum trajectories, as developed in quantum optics, has proven **very (e) useful** for understanding continuous measurement in the solid state context.
(4) However, as traditionally formulated it is a highly idealized theory, **assuming (e) that** individual quantum events (such as electrons tunnelling through a barrier) can be observed.
(5) In practice, the electronic noise and impedance of the external circuit will **mask (e)** this observation of individual quantum events (such as electrons tunnelling through a barrier).
(6) A generalization of the quantum trajectory approach **to describe** the effect of such read-out imperfections on the conditioned system state has recently been developed for photon detectors (Warszawski, Wiseman and Mabuchi, 2002). This technique is now being applied in the solid-state context.
(7) Quantum Feedback Control: Applications for quantum feedback in quantum computers **include** quantum error correction for detected errors, testing read-out devices, state engineering, and measurement engineering.
(8) State engineering **requires a** general understanding of quantum control, which will be sought.
(9) Measurement engineering by feedback **means** adaptive measurements, which could be useful in linear optics quantum computing.

NOTATION

Module One

A crucial part of quantum computing is read-out.

Electronic Read-Out of Quantum Bits Aug. 16, 2012: Science news

Quantum computers promise to reach computation speeds far beyond that of today's computers. As they would use quantum effects, however, they would also be susceptible to external interferences. Information flow into and out of the system is a critical point.

Researchers from KIT with partners from Grenoble and Strasbourg have now read out the quantum state of an atom directly by using electrodes. In the journal Nature, they report on the stable interface between classical and quantum world.

"Normally, every contact with the outer world changes information in a quantum mechanical system in a

completely uncontrolled manner," explains Professor Mario Ruben from Karlsruhe Institute of Technology. "We therefore have to keep the quantum state stable and shielded. On the other hand, information has to be read out in a controlled manner for further use."

Magnetic molecule complexes may be a solution of this dilemma. In their center, a metal atom with a pronounced magnetic moment, a spin, is located. It is surrounded by organic molecules that shield the atom. "When synthesizing this protective enclosure, we can exactly define how much the metal atom sees of the outer world," explains Ruben the trick of his research project.

The study presented is based on the metal atom terbium that was provided with an enclosure of about 100 carbon, nitrogen, and water atoms and then placed in the center of nanometer-sized, electric gold contacts. Due to the properties of the molecule, the electrodes had an effect similar to the three channels of a transistor. Electric voltage of the middle gate electrode influenced the current through the other two electrodes. In this way, the working point was set. Then, the molecule was exposed to various changing magnetic fields and the jump of the spin was reflected by the amplitude of the current curve. "By measuring current flow, we found that the nuclear spin of the metal atom is stable for up to 20 seconds," says Ruben. "For quantum mechanical processes, this is a very long time."

Ruben is sure that "the results will be of particular importance to spintronics and quantum computing." Spintronics uses the magnetic spin of single particles for information processing. The word describes the symbiosis of spin and electronics. Quantum computers use quantum mechanical effects, such as the entanglement and super-position of spins, for the parallel execution of algorithms at high speed.

G_{13} : Category one of read-out

G_{14} : Category two of read-out

G_{15} : Category three of read-out

T_{13} : Category one of crucial part of quantum computing

T_{14} : Category two of crucial part of quantum computing

T_{15} : Category three of crucial part of quantum computing

Module Two

Schemes for reading out the state of a solid-state QC **include (e)** a single-electron transistor, a quantum point contact, and a micromechanical cantilever

G_{16} : Category one of single-electron transistor, a quantum point contact, and a micromechanical cantilever

G_{17} : Category two of single-electron transistor, a quantum point contact, and a micromechanical cantilever

G_{18} : Category three of single-electron transistor, a quantum point contact, and a micromechanical cantilever

T_{16} : Category one of Schemes for reading out the state of a solid-state QC

T_{17} : Category two of Schemes for reading out the state of a solid-state QC

T_{18} : Category three of Schemes for reading out the state of a solid-state QC

Module three

The theory of quantum trajectories, as developed in quantum optics, has proven **very (e) useful** for

understanding continuous measurement in the solid state context

Some people like JunYan Luo, Lun-Wu Zhu, Xiao-Ling He, Bensheng Yun, and Shiping Ruan have studied the Continuous measurement of a quantum two-level system (qubit) by mesoscopic charge detectors **by using** Non-Markovian dynamics in continuous measurement of a solid-state charge qubit. Continuous measurement of a quantum two-level system (qubit) by mesoscopic charge detectors has attracted wide attention in the community of solid-state quantum computation. In this work, they revisit the measurement of a charge qubit by a quantum point contact, with particular attention **paid (eb) to the** unique non-Markovian measurement characteristics. The finite-frequency-support feature due to non-Markovian correlation **leads to a** remarkable suppression of the pedestal in noise spectrum, which results in a drastic violation of the Korotkov-Averin bound, the upper limit of the singleto-noise ratio imposed quantum mechanically on linear detectors. For a large measurement voltage and bandwidth, the non-Markovian effect **gives rise to** a complete inhibition of coherent jumps (quantum Zeno effect), as identified by the telegraph noise peaks in the spectrum. Mesoscale and Nanoscale Physics (cond-mat.mes-hall) arXiv: 1203.2233 [cond-mat.mes-hall]

G_{20} : Category one of continuous measurement in the solid state context; Non-Markovian dynamics in continuous measurement of a solid-state charge qubit. Continuous measurement of a quantum two-level system (qubit) by mesoscopic charge detectors; measurement of a charge qubit by a quantum point contact; finite-frequency-support feature due to non-Markovian correlation; large measurement voltage and bandwidth, the non-Markovian effect;

G_{21} : Category two of continuous measurement in the solid state context; Non-Markovian dynamics in continuous measurement of a solid-state charge qubit. Continuous measurement of a quantum two-level system (qubit) by mesoscopic charge detectors; measurement of a charge qubit by a quantum point contact; finite-frequency-support feature due to non-Markovian correlation; large measurement voltage and bandwidth, the non-Markovian effect;

G_{22} : Category three of continuous measurement in the solid state context; Non-Markovian dynamics in continuous measurement of a solid-state charge qubit. Continuous measurement of a quantum two-level system (qubit) by mesoscopic charge detectors; measurement of a charge qubit by a quantum point contact; finite-frequency-support feature due to non-Markovian correlation; large measurement voltage and bandwidth, the non-Markovian effect;

T_{20} : Category one of theory of quantum trajectories, as developed in quantum optics; Continuous measurement of a quantum two-level system (qubit) by mesoscopic charge detectors; unique non-Markovian measurement characteristics; remarkable suppression of the pedestal in noise spectrum, which results in a drastic violation of the Korotkov-Averin bound, the upper limit of the singleto-noise ratio imposed quantum mechanically on linear detectors; complete inhibition of coherent jumps (quantum Zeno effect), as identified by the telegraph noise peaks in the spectrum

T_{21} : Category two of theory of quantum trajectories, as developed in quantum optics; Continuous measurement of a quantum two-level system (qubit) by mesoscopic charge detectors; unique non-Markovian measurement characteristics; remarkable suppression of the pedestal in noise spectrum, which results in a drastic violation of the Korotkov-Averin bound, the upper limit of the singleto-noise ratio imposed quantum mechanically on linear detectors; complete inhibition of coherent jumps (quantum Zeno effect), as identified by the telegraph noise peaks in the spectrum

T_{22} : Category three of theory of quantum trajectories, as developed in quantum optics; Continuous measurement of a quantum two-level system (qubit) by mesoscopic charge detectors; unique non-Markovian measurement characteristics; remarkable suppression of the pedestal in noise spectrum, which results in a drastic violation of the Korotkov-Averin bound, the upper limit of the singleto-noise ratio imposed quantum mechanically on linear detectors; complete inhibition of coherent jumps (quantum Zeno effect), as identified

by the telegraph noise peaks in the spectrum

Quantum Trajectory Theory

Quantum trajectory theory treats the open systems encountered in quantum optics according to a scattering scenario, in which the inputs are classical fields (external fields) and the outputs are classical stochastic processes that model the scattered fields after detection (after their measurement).

The mapping from inputs to outputs is provided by a quantum stochastic process that is set up to account for a particular measurement strategy (eg., photon counting, homodyne/heterodyne detection, etc). Both the formal theory and its applications are under study. Recent work includes the development of a novel measurement scheme to correlate the quadrature amplitudes of an electromagnetic field, applications of this scheme in cavity quantum electrodynamics, and the modeling of multimode aspects of collective radiative phenomena (super-radiance).

Entanglement and Correlation in Composite Systems

Quantum optics has traditionally concerned itself with uniquely quantum mechanical aspects of optical phenomena (eg. photon anti-bunching and squeezing, violations of Bell inequalities). Attention in recent years has been focused on applications of these quantum features of light to novel schemes for processing information (so-called quantum information science). Entangled states are central to the proposed information processing protocols. Current work is directed towards understanding entangled states, and the contextual correlations they describe, in the broadest sense, ie, we are concerned with the physics of composite quantum systems in general. Specific interests include measures of entanglement for open systems, and schemes for the generation and manipulation of this entanglement. Continuous variable entanglement is of particular interest. It has been suggested by others that this form of entanglement can be accounted for within stochastic electrodynamics. The suggestion is being assessed. A quantum trajectory theory of continuous variable teleportation is being developed for comparison with the stochastic electrodynamics proposal. **From the page of Professor Howard Carmichael MSc, D. Phil (Waikato), FRSNZ The Dan Walls Chair in Theoretical Physics**

Module four

However, as traditionally formulated it is a highly idealized theory, **assuming (e) that** individual quantum events (such as electrons tunnelling through a barrier) can be observed.
In physics, a Coulomb blockade (abbreviated CB), named after Charles-Augustin de Coulomb's electrical force, is the **increased resistance** at small bias voltages of an electronic device comprising at least one low-capacitance tunnel junction. Because of the CB, the resistances of devices are not constant at low bias voltages, but increase to infinity for zero bias (i.e. no current flows).

Coulomb blockade in a tunnel junction

A tunnel junction is, in its simplest form, a thin insulating barrier between two conducting electrodes. If the electrodes are superconducting, Cooper pairs (with a charge of two elementary charges) carry the current. In the case that the electrodes are normal conducting, i.e. neither superconducting nor semiconducting, electrons (with a charge of one elementary charge) carry the current. The following reasoning is for the case of tunnel junctions with an insulating barrier between two normal conducting electrodes (NIN junctions).

According to the laws of classical electrodynamics, no current can flow through an insulating barrier. According to the laws of quantum mechanics, however, there is a nonvanishing (larger than zero) probability for an electron on one side of the barrier to reach the other side (see quantum tunnelling). When a bias

voltage is applied, this means that there will be a current, and, neglecting additional effects, the tunnelling current will be proportional to the bias voltage. In electrical terms, the tunnel junction behaves as a resistor with a constant resistance, also known as an Ohmic resistor. The resistance depends exponentially on the barrier thickness. Typical barrier thicknesses are on the order of one to several nanometers.

An arrangement of two conductors with an insulating layer in between not only has a resistance, but also a finite capacitance. The insulator is also called dielectric in this context; the tunnel junction behaves as a capacitor.

Due to the discreteness of electrical charge, current through a tunnel junction is a series of events in which exactly one electron passes (tunnels) through the tunnel barrier (we neglect cotunneling, in which two electrons tunnel simultaneously). The tunnel junction capacitor is charged with one elementary charge by the tunnelling electron, causing a voltage buildup , where is the elementary charge of 1.6×10−19 coulomb and the capacitance of the junction. If the capacitance is very small, the voltage buildup can be large enough to prevent another electron from tunnelling. The electrical current is then suppressed at low bias voltages and the resistance of the device is no longer constant. The increase of the differential resistance around zero bias is called the Coulomb blockade.

G_{24} : Category one of individual quantum events (such as electrons tunnelling through a barrier) can be observed.

G_{25} : Category two of individual quantum events (such as electrons tunnelling through a barrier) can be observed.

G_{26} : Category three of individual quantum events (such as electrons tunnelling through a barrier) can be observed.

T_{24} : Category one of Quantum Trajectory Theory is a highly idealized theory

T_{25} : Category two of Quantum Trajectory Theory is a highly idealized theory

T_{26} : Category three of Quantum Trajectory Theory is a highly idealized theory

Module five

In practice, the electronic noise and impedance of the external circuit will **mask (e)** this observation of individual quantum events (such as electrons tunnelling through a barrier).

G_{28} : Category one of observation of individual quantum events (such as electrons tunnelling through a barrier).

G_{29} : Category two of observation of individual quantum events (such as electrons tunnelling through a barrier).

G_{30} : Category three of observation of individual quantum events (such as electrons tunnelling through a barrier).

T_{28} : Category one of electronic noise and impedance of the external circuit

T_{29} : Category two of electronic noise and impedance of the external circuit

T_{30} : Category three of electronic noise and impedance of the external circuit

Module six

A generalization of the quantum trajectory approach **to describe** the effect of such read-out imperfections on the conditioned system state has recently been developed for photon detectors (Warszawski, Wiseman and Mabuchi, 2002). This technique is now being applied in the solid-state context.

The quantum trajectory approach is generalized to arbitrary coordinate systems, including curvilinear coordinates. This allows one to perform an approximate quantum trajectory propagation, which scales favorably with system size, in the same framework as standard quantum wave packet dynamics. The trajectory formulation is implemented in Jacobi coordinates for a nonrotating triatomic molecule. Wave packet reaction probabilities are computed for the O (3P) + H2 --> OH + H reaction using the approximate quantum potential. The latter is defined by the nonclassical component of the momentum operator expanded in terms of linear and exponential functions. Unlike earlier implementations with linear functions, the introduction of the exponential function provides an accurate description of asymptotic dynamics for this system and gives good agreement of approximate reaction probabilities with accurate quantum calculations. **J Phys Chem A. 2006 Apr 27; 110(16):5530-6. Quantum trajectory dynamics in arbitrary coordinates.Rassolov VA, Garashchuk S, Schatz GC.**

G_{32} : Category one of generalization of the quantum trajectory approach

G_{33} : Category two of generalization of the quantum trajectory approach

G_{34} : Category three of generalization of the quantum trajectory approach

T_{32} : Category one of effect of such read-out imperfections on the conditioned system state has recently been developed for photon detectors (Warszawski, Wiseman and Mabuchi, 2002). This technique is now being applied in the solid-state context.

T_{33} : Category two of effect of such read-out imperfections on the conditioned system state has recently been developed for photon detectors (Warszawski, Wiseman and Mabuchi, 2002). This technique is now being applied in the solid-state context.

T_{34} : Category three of effect of such read-out imperfections on the conditioned system state has recently been developed for photon detectors (Warszawski, Wiseman and Mabuchi, 2002). This technique is now being applied in the solid-state context.

Module seven

Quantum Feedback Control: Applications for quantum feedback in quantum computers **include** quantum error correction for detected errors, testing read-out devices, state engineering, and measurement engineering

Substantive limits, predicational anteority, divergential affirmation, ontological consonance, primordial exactitude, phenomenological correlates have been studied by many authors such as L. D. Tóth (See Paradigms for Quantum Feedback Control L. D. Tóth).In this review paper, we survey the main concepts and some of the recent developments in quantum feedback control. For consistency and clarity, essential ideas and notations in the theory of open quantum systems and quantum stochastic calculus, as well as continuous measurement theory are developed. We give a general description of quantum feedback control, set up a coherent model and compare it to open-loop designs. Objectives which can be achieved by feedback, such as rapid state preparation and purification or entanglement generation are formulated and

analyzed, based on the relevant literature. The connection between quantum feedback and quantum chaos is also described and unraveled which, apart from its theoretical curiosity, can shed more light on some of the intrinsic properties of this control paradigm.arXiv:1210.0360 [quant-ph]

G_{36} : Category one of quantum error correction for detected errors, testing read-out devices, state engineering, and measurement engineering

G_{37} : Category two of quantum error correction for detected errors, testing read-out devices, state engineering, and measurement engineering

G_{38} : Category three of quantum error correction for detected errors, testing read-out devices, state engineering, and measurement engineering

T_{36} : Category one of quantum feedback in quantum computers

T_{37} : Category two of quantum feedback in quantum computers

T_{38} : Category three of quantum feedback in quantum computers

Module eight

State engineering **requires a** general understanding of quantum control,

G_{40} : Category one of general understanding of quantum control

G_{41} : Category two of general understanding of quantum control

G_{42} : Category three of general understanding of quantum control

T_{40} : Category one of State engineering

T_{41} : Category two of State engineering

T_{42} : Category three of State engineering

Module Nine

Measurement engineering by feedback **means** adaptive measurements, which could be useful in linear optics quantum computing.

G_{44} : Category one of Measurement engineering by feedback

G_{45} : Category two of Measurement engineering by feedback

G_{46} : Category three of Measurement engineering by feedback

T_{44} : Category one of adaptive measurements, which could be useful in linear optics quantum computing.

T_{45} : Category two of adaptive measurements, which could be useful in linear optics quantum computing.

T_{46} : Category three of adaptive measurements, which could be useful in linear optics quantum computing.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$
$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$
$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**
$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{16})^{(2)} \; +(a''_{16})^{(2)}(T_{17},t) \; +(a''_{13})^{(1,1,)}(T_{14},t) \; +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) \; +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) \; +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) \; +(a''_{40})^{(8,8,8)}(T_{41},t) \; +(a''_{44})^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)} \; +(a''_{17})^{(2)}(T_{17},t) \; +(a''_{14})^{(1,1,)}(T_{14},t) \; +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) \; +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) \; +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) \; +(a''_{41})^{(8,8,8)}(T_{41},t) \; +(a''_{45})^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{18})^{(2)} \; +(a''_{18})^{(2)}(T_{17},t) \; +(a''_{15})^{(1,1,)}(T_{14},t) \; +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) \; +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) \; +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7)}(T_{37},t) \; +(a''_{42})^{(8,8,8)}(T_{41},t) \; +(a''_{46})^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for*

category 1, 2 and 3

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* $1,2\ 3$

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* $1,2$ *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* $1,2$ *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category $1,2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) & \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) & \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) & \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) & \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a_{32}')^{(6)} & +(a_{32}'')^{(6)}(T_{33},t) & +(a_{28}'')^{(5,5,5)}(T_{29},t) & +(a_{24}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) & \\ +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a_{33}')^{(6)} & +(a_{33}'')^{(6)}(T_{33},t) & +(a_{29}'')^{(5,5,5)}(T_{29},t) & +(a_{25}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) & \\ +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a_{34}')^{(6)} & +(a_{34}'')^{(6)}(T_{33},t) & +(a_{30}'')^{(5,5,5)}(T_{29},t) & +(a_{26}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) & \\ +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{34} \quad (87)$$

$+(a_{32}'')^{(6)}(T_{33},t)$, $+(a_{33}'')^{(6)}(T_{33},t)$, $+(a_{34}'')^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a_{28}'')^{(5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a_{24}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{ccc} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} +\ (\hat{A}_{20})^{(3)} +\ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} +\ (\hat{B}_{20})^{(3)} +\ (\hat{Q}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le\ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) =\ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(KKKKK) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(LLLLL) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(MMMMM) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(NNNNN)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(OOOOO) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(PPPPP) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\le\ \left(\hat{P}_{13}\right)^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\le\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\le\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{13})^{(1)}$, $T_i^0 \le (\hat{Q}_{13})^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 156

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{32})^{(6)}, T_i^0 \le (\hat{Q}_{32})^{(6)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(t) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13,14,15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}\leq$$
$$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{\mathrm{A}}_{16})^{(2)}+(\widehat{\mathrm{P}}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)\mathrm{d}\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)} t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

208

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)}$$

209

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)}$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),\left((G_{23})^{(2)},(T_{23})^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

214

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})},\widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})},\widetilde{(T_{31})}\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}),(\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(8)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[(\hat{P}_{44})^{(9)} + \left((\hat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{44})^{(9)}\right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}$, $if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)}$,

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}$, $if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}$, $if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}$,

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}\left((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)}\right)}\left[e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}\left((S_1)^{(3)} - (a'_{22})^{(3)}\right)}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}\left((R_1)^{(3)} - (b'_{22})^{(3)}\right)}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}\left((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)}\right)}\left[e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right) - (b''_{25})^{(4)}\left((G_{27}), t\right) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)}G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)}G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)}T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)},, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^{0}e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^{0}e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^{0}e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}}T_{44}^{0}e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^{0}}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t}-e^{-(b'_{46})^{(9)}t}\right]+T_{46}^{0}e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^{0}}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t}-e^{-(R_2)^{(9)}t}\right]+T_{46}^{0}e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^{0}}{G_{14}^{0}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)}+(C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)}=\frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)},(\nu_0)^{(7)}$:-

$$\text{For } 0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case, 423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)}+(C)^{(8)}(v_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)}-(v_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)}-(v_2)^{(8)}\right)t\right]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)}+(\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)}-(\bar{v}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)}-(\bar{v}_2)^{(8)}\right)t\right]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)}+(\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)}-(\bar{v}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)}-(\bar{v}_2)^{(8)}\right)t\right]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines** $(v_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)}\right) \leq \frac{dv^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(t) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523

A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^{*}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^{*}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}((G_{31})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FORTY NINE

Spintronics And Quantum Computation: Natura Non Contristatur-Natura Non Facit Saltum Ita Nec Lex

INTRODUCTION—VARIABLES USED

Evolutionary dynamics, in biology, is the study of the mathematical principles according to which life has evolved and continues to evolve. This is mostly achieved through the mathematical discipline of population genetics. Most population genetics considers changes in the frequencies of alleles at a small number of gene loci. When infinitesimal effects at a large number of gene loci are considered, one derives quantitative genetics. Traditional population genetic models deal with alleles and genotypes, and are frequently stochastic. In evolutionary game theory, developed first by John Maynard Smith, evolutionary biology concepts may take a deterministic mathematical form, with selection acting directly on inherited phenotypes. These models have relevance also to the generation and maintenance of tissues in mammals, since an

understanding of tissue cell kinetics, architecture, and development from adult stem cells has important implications for aging and cancer.

Spintronics Approach Enables New Quantum Technologies June 4, 2013

A team of researchers including members of the University of Chicago's Institute for Molecular Engineering highlight the power of emerging quantum technologies in two recent papers published in the Proceedings of the National Academy of Sciences (PNAS).

(1) Molecular Engineering highlight the power of emerging quantum technologies exploit quantum mechanics, the physics that dominates the atomic world, **to perform** disparate tasks such as nanoscale temperature measurement and processing quantum information with lasers.
(2) The two papers are both based on the **manipulation** of the same material, an atomic-scale defect in diamond known as the nitrogen vacancy center. Both works also leverage the intrinsic "spin" of this defect **for the** applications in temperature measurement and information processing.
(3) This spintronics approach involves **understanding and manipulating** the spin of electronics for technological advancement.

"These studies build on research efforts undertaken over the last 20 years to isolate and control single electronic spins in the solid state," said David Awschalom, a principle investigator on both papers and a Liew Family Professor in Molecular Engineering at UChicago. "Much of the initial motivation for working in this field was driven by the desire to make new computing technologies based on the principles of quantum physics. In recent years the research focus has broadened as we've come to appreciate that these same principles could enable a new generation of nanoscale sensors."

Controlling qubits with light

In one PNAS paper posted April 22 and published in the May 7 print edition, Awschalom and six co-authors at the University of California, Santa Barbara and the University of Konstanz describe a technique that offers new routes toward the eventual creation of quantum computers, which would possess far more capability than modern classical computers.

In this application, Awschalom's team has developed protocols to fully **control the** quantum state of the defect with light instead of electronics. The quantum state of interest in this defect is its electronic spin, which acts as quantum bit, or qubit, the basic unit of a quantum computer.

In classical computers, bits of information exist in one of only two states: zero or one. In the quantum mechanical realm, objects can exist in multiple states at once, enabling more complex processing.

This all-optical scheme for controlling qubits in semiconductors "**obviates the** need to have microwave circuits or electronic networks," Awschalom said. "Instead, everything can be done solely with photons, with light."

As a fully optical method, it shows promise as a more scalable approach to qubit control. In addition, this scheme is more versatile than conventional methods **and could be used to** explore quantum systems in a broad range of materials that might otherwise be difficult to develop as quantum devices.

Single spin thermometers

The quantum thermometer application, reported in a PNAS contribution posted online May 6 and published in the May 21 print edition, represents a new direction for the **manipulati**on of quantum states, which is

more commonly linked to computing, communications, and encryption.

In recent years, defect spins had also emerged **as promising candidates** for nanoscale sensing of magnetic and electric fields at room temperature. With thermometry now added to the list, Awschalom foresees the possibility of developing a multifunctional probe based on quantum physics.

"With the same sensor you could measure magnetic fields, electric fields and now temperature, all with the same probe in the same place at approximately the same time," he said. "Perhaps most importantly, since the sensor is an atomic-scale defect that **could be contained** within nanometer-scale particles of diamond, you can imagine using this system as a thermometer in challenging environments such as living cells or microfluidic circuits."

The key aspect of this innovation is the development of control techniques **for manipulating** the spin that make it a much more sensitive probe of temperature shifts. "We've been exploring the potential of defect spins for thermometry for the past few years," said David Toyli, a graduate student in physics at UCSB and lead author of the temperature sensing work.

"This latest work is exciting because we've succeeded in adapting techniques **used for** stabilizing quantum information to measuring temperature-dependent changes in the quantum states.

These techniques **minimize** the effects of environmental noise and allow us to make much more sensitive temperature measurements."

The team of researchers, also including Slava Dobrovitski of the Department of Energy's Ames Laboratory in Iowa, conducted experiments to determine the temperature range over which the spins could operate as a useful thermometer. It turns out that the particle spins can operate quite well at a wide temperature range, from room temperature to 500 degrees Kelvin (approximately 70 to 400 degrees Fahrenheit).

The chemical properties of a diamond-based thermometer also support the idea that this system could be useful for measuring temperature gradients in biological systems, such as the interior of living cells, Awschalom said. But the initial studies suggest the method is so flexible that it probably lends itself to uses yet to be imagined. "Like any new technology development, the exciting thing is what people will do with this now."

Understanding the Evolutionary Dynamics and Adaptation of Complex Networks Alireza Abbasi et al

(1) This study aims to explore the co-evolutionary processes of network structure **and s**tate for examining the evolutionary dynamic and adaptation of networks. The study addresses several major areas of network research: network structure dynamics (i.e., the longitudinal study of network structure); network evolution mechanisms (i.e., variation, selection, retention); actors' attachment logics (i.e., cumulative advantage, embedding, multi-connectivity, homophily); and self-organized network behavior.

(2) Theories and hypotheses from the intersection of network theories, evolutionary theories and complex adaptive systems are drawn and synthesized. Finally the co-evolution of local dynamics **versus** global dynamics of networks over time has been explored through proposing a research framework.


(3) Authors consider a quantum dot attached to leads in the Coulomb blockade regime which has a spin 1/2 ground state. They show that by applying an ESR field to the dot-spin the stationary current in the sequential tunneling regime **exhibits** a resonance whose line width is determined by the single-spin decoherence time T_2.

(4) The Rabi oscillations of the dot-spin are **shown to induce** coherent current oscillations from which T_2 can be deduced in the time domain.

(5) They describe a spin-inverter which **can be(e) used** to pump current through a double-dot via spin flips generated by ESR.DOI: 10.1103/PhysRevLett.86.4648 arXiv:cond-mat/0011193 [cond-mat.mes-hall](**Detection of Single Spin Decoherence in a Quantum Dot via Charge Currents Hans-Andreas Engel)**

Electron spins in quantum dots for spintronics and quantum computation Hans-Andreas Engel et al http://dx.doi.org/10.1016/S0038-1098 (01)00110-7

(6) Exciting the dot-spin via ESR (Rabi flip, draws as wavy line) the dot becomes unblocked for spin up electrons from lead 1. Finally, from the singlet, spin up or down can tunnel into lead 2. (c) Combined setup, 'spin inverter', where the spin state of dot 1 switches the current as in (b) and where dot 2 acts as a spin filter, in the regime E1S≈E2S; E1S>μ1>E1S−Δ1z; E2S>μ2>E1S−Δ2z; |tDD|<|tDL2|; and $\Delta_z^1 \not\approx \Delta_z^2$. This **allows the** transition sequence $\uparrow(\uparrow)_1(\uparrow)_2 \xrightarrow{ESR} \uparrow(\downarrow)_1(\uparrow)_2 \rightarrow (\uparrow\downarrow)_1(\uparrow)_2 \rightarrow (\uparrow)_1(\uparrow\downarrow)_2 \rightarrow (\uparrow)_1(\uparrow)_2\downarrow$.



(7) Fig. 3. In the box on the left the Andreev Entangler is sketched. The two electrons of a Cooper pair hop with amplitudeTDS from two points r1,r2 of the superconductor, SC, (distances δ r=|r1−r2|) to two dots D1,2 **by(e) means** of Andreev tunneling.

(8) The dots are coupled to normal leads L1, 2 with tunneling amplitude TDL. The chemical potentials of the SC and leads are μ S and μ 1= μ 2, resp. For efficiency, |TDL|>|TDS| and large Coulomb repulsion U is required. On the right the experimental setup for measuring noise of entangled electrons (see Section 4.2) is drawn. The entangler takes uncorrelated electrons from the Fermi leads 1′ and 2′. Pairs of entangled electrons (singlet or triplet) are produced in the entangler (e.g. singlets in the Andreev Entangler) and then injected into the leads 1 and 2, one electron per lead. The current of these two leads are mixed with a beam splitter (to induce scattering interference) **and the resulting** noise is then measured in lead 3 and 4: no noise (antibunching) for triplets, whereas we get enhanced noise (bunching) for singlets.

NOTATION

Module One

Molecular Engineering highlight the power of emerging quantum technologies exploit quantum mechanics, the physics that dominates the atomic world, **to perform** disparate tasks such as nanoscale temperature measurement and processing quantum information with lasers.

G_{13} : Category one of Molecular Engineering highlight the power of emerging quantum technologies exploit quantum mechanics, the physics that dominates the atomic world

G_{14} : Category two of Molecular Engineering highlight the power of emerging quantum technologies exploit quantum mechanics, the physics that dominates the atomic world

G_{15} : Category three of Molecular Engineering highlight the power of emerging quantum technologies

exploit quantum mechanics, the physics that dominates the atomic world

T_{13} : Category one of disparate tasks such as nanoscale temperature measurement and processing quantum information with lasers.

T_{14} : Category two of disparate tasks such as nanoscale temperature measurement and processing quantum information with lasers.

T_{15} : Category three of disparate tasks such as nanoscale temperature measurement and processing quantum information with lasers.

Module Two

The two papers are both based on the **manipulation** of the same material, an atomic-scale defect in diamond known as the nitrogen vacancy center leveraging the intrinsic "spin" of this defect **for the** applications in temperature measurement and information processing.

G_{16} : Category one of the **manipulation** of the same material, an atomic-scale defect in diamond known as the nitrogen vacancy center leveraging the intrinsic "spin" of this defect

G_{17} : Category two of the **manipulation** of the same material, an atomic-scale defect in diamond known as the nitrogen vacancy center leveraging the intrinsic "spin" of this defect

G_{18} : Category three of the **manipulation** of the same material, an atomic-scale defect in diamond known as the nitrogen vacancy center leveraging the intrinsic "spin" of this defect

T_{16} : Category one of temperature measurement and information processing

T_{17} : Category two of temperature measurement and information processing

T_{18} : Category three of temperature measurement and information processing

Module three

This all-optical scheme for controlling qubits in semiconductors "**obviates the** need to have microwave circuits or electronic networks," Awschalom said. "Instead, everything can be done solely with photons, with light."

G_{20} : Category one of need to have microwave circuits or electronic networks,". "Instead, everything can be done solely with photons, with light."

G_{21} : Category two of need to have microwave circuits or electronic networks,". "Instead, everything can be done solely with photons, with light."

G_{22} : Category three of need to have microwave circuits or electronic networks,". "Instead, everything can be done solely with photons, with light."

T_{20} : Category one of all-optical scheme for controlling qubits in semiconductors

T_{21} : Category two of all-optical scheme for controlling qubits in semiconductors

T_{22} : Category three of all-optical scheme for controlling qubits in semiconductors

Module four

With the same sensor you could measure magnetic fields, electric fields and now temperature, all with the same probe in the same place at approximately the same time," he said. "Perhaps most importantly, since the

sensor is an atomic-scale defect that **could be contained** within nanometer-scale particles of diamond, you can imagine using this system as a thermometer in challenging environments such as living cells or microfluidic circuits."

G_{24} : Category one of nanometer-scale particles of diamond, you can imagine using this system as a thermometer in challenging environments such as living cells or microfluidic circuits."

G_{25} : Category two of nanometer-scale particles of diamond, you can imagine using this system as a thermometer in challenging environments such as living cells or microfluidic circuits."

G_{26} : Category three of nanometer-scale particles of diamond, you can imagine using this system as a thermometer in challenging environments such as living cells or microfluidic circuits."

T_{24} : Category one of same sensor one could measure magnetic fields, electric fields and now temperature, all with the same probe in the same place at approximately the same time," . "Perhaps most importantly, since the sensor is an atomic-scale defect

T_{25} : Category two of same sensor one could measure magnetic fields, electric fields and now temperature, all with the same probe in the same place at approximately the same time," . "Perhaps most importantly, since the sensor is an atomic-scale defect

T_{26} : Category three of same sensor one could measure magnetic fields, electric fields and now temperature, all with the same probe in the same place at approximately the same time," . "Perhaps most importantly, since the sensor is an atomic-scale defect

Module five

These techniques **minimize** the effects of environmental noise and allow us to make much more sensitive temperature measurements

G_{28} : Category one of effects of environmental noise and allow us to make much more sensitive temperature measurements

G_{29} : Category two of effects of environmental noise and allow us to make much more sensitive temperature measurements

G_{30} : Category three of effects of environmental noise and allow us to make much more sensitive temperature measurements

T_{28} : Category one of spintronics

T_{29} : Category two of spintronics

T_{30} : Category three of spintronics

Module six

Co-evolutionary processes of network structure **and s**tate for examining the evolutionary dynamic and adaptation of networks. The study addresses several major areas of network research: network structure dynamics (i.e., the longitudinal study of network structure); network evolution mechanisms (i.e., variation, selection, retention); actors' attachment logics (i.e., cumulative advantage, embedding, multi-connectivity, homophily); and self-organized network behavior.

G_{32} : Category one of several major areas of network research: network structure dynamics (i.e., the longitudinal study of network structure); network evolution mechanisms (i.e., variation, selection, retention); actors' attachment logics (i.e., cumulative advantage, embedding, multi-connectivity, homophily); and self-organized network behavior.

G_{33} : Category two of several major areas of network research: network structure dynamics (i.e., the longitudinal study of network structure); network evolution mechanisms (i.e., variation, selection, retention); actors' attachment logics (i.e., cumulative advantage, embedding, multi-connectivity, homophily); and self-organized network behavior.

G_{34} : Category three of several major areas of network research: network structure dynamics (i.e., the longitudinal study of network structure); network evolution mechanisms (i.e., variation, selection, retention); actors' attachment logics (i.e., cumulative advantage, embedding, multi-connectivity, homophily); and self-organized network behavior.

T_{32} : Category one of co-evolutionary processes of network structure **and s**tate for examining the evolutionary dynamic and adaptation of networks

T_{33} : Category two of co-evolutionary processes of network structure **and s**tate for examining the evolutionary dynamic and adaptation of networks

T_{34} : Category three of co-evolutionary processes of network structure **and s**tate for examining the evolutionary dynamic and adaptation of networks

Module seven

Authors consider a quantum dot attached to leads in the Coulomb blockade regime which has a spin 1/2 ground state. They show that by applying an ESR field to the dot-spin the stationary current in the sequential tunneling regime **exhibits** a resonance whose line width is determined by the single-spin decoherence time T_2.

G_{36} : Category one of quantum dot attached to leads in the Coulomb blockade regime which has a spin 1/2 ground state which on application of ESR field to the dot-spin the stationary current in the sequential tunneling regime

G_{37} : Category two of quantum dot attached to leads in the Coulomb blockade regime which has a spin 1/2 ground state which on application of ESR field to the dot-spin the stationary current in the sequential tunneling regime

G_{38} : Category three of quantum dot attached to leads in the Coulomb blockade regime which has a spin 1/2 ground state which on application of ESR field to the dot-spin the stationary current in the sequential tunneling regime

T_{36} : Category one of resonance whose line width is determined by the single-spin decoherence time T_2.

T_{37} : Category two of resonance whose line width is determined by the single-spin decoherence time T_2.

T_{38} : Category three of resonance whose line width is determined by the single-spin decoherence time T_2.

Module eight

The Rabi oscillations of the dot-spin are **shown to induce** coherent current oscillations from which T_2 can be deduced in the time domain.

G_{40} : Category one of Rabi oscillations of the dot-spin

G_{41} : Category two of Rabi oscillations of the dot-spin

G_{42} : Category three of Rabi oscillations of the dot-spin

T_{40} : Category one of coherent current oscillations from which T_2 can be deduced in the time domain.

T_{41} : Category two of coherent current oscillations from which T_2 can be deduced in the time domain.

T_{42} : Category three of coherent current oscillations from which T_2 can be deduced in the time domain.

Module Nine

They describe a spin-inverter which **can be(e) used** to pump current through a double-dot via spin flips generated by ESR.DOI: 10.1103/PhysRevLett.86.4648 arXiv:cond-mat/0011193 [cond-mat.mes-hall](**Detection of Single Spin Decoherence in a Quantum Dot via Charge Currents Hans-Andreas Engel)**

G_{44} : Category one of spin-inverter

G_{45} : Category two of spin-inverter

G_{46} : Category three of spin-inverter

T_{44} : Category one of pump current through a double-dot via spin flips generated by ESR

T_{45} : Category two of pump current through a double-dot via spin flips generated by ESR

T_{46} : Category three of pump current through a double-dot via spin flips generated by ESR

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)} \ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for

category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt}=(a_{16})^{(2)}G_{17}-\left[\begin{array}{ccc}(a'_{16})^{(2)} & \boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} \quad \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)}\end{array}\right]G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt}=(a_{17})^{(2)}G_{16}-\left[\begin{array}{ccc}(a'_{17})^{(2)} & \boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} \quad \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)}\end{array}\right]G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt}=(a_{18})^{(2)}G_{17}-\left[\begin{array}{ccc}(a'_{18})^{(2)} & \boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} \quad \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)}\end{array}\right]G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt}=(b_{16})^{(2)}T_{17}-\left[\begin{array}{ccc}(b'_{16})^{(2)} & \boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} \quad \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)}\end{array}\right]T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients

for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)}&\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}&\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}&\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}&\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}&\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}&\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)}&\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}&\boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}&\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}&\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}&\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}&\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)}&\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}&\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}&\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}&\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}&\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}&\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)}&\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}&\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}&\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}&\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}&\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}&\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)}&\boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}&\boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}&\boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}&\boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}&\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}&\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category $1, 2$ *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b_{36}')^{(7)}\boxed{-(b_{36}'')^{(7)}(G_{39},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b_{37}')^{(7)}\boxed{-(b_{37}'')^{(7)}(G_{39},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b_{38}')^{(7)}\boxed{-(b_{38}'')^{(7)}(G_{39},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\,-(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\,-(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\,-(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(QQQQQ) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(RRRRR) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$

132

(SSSSS) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(TTTTT)

$\lim_{G\to\infty}(b_i'')^{(7)}((G_{39}),t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}((G_{39}),t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(UUUUU) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(VVVVV) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(u) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right) - (a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right| \; e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} \} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)} - G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)};\ (G_{23})^{(2)}, (T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\tilde{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$

$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$

$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$

$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq$ 248

$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 250

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$
$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{36})^{(7)}t}\}$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\,((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(9)}t)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)},, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad (378)$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad (379)$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad (380)$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- (381)

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 (382)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(\nu_1)^{(9)},(\nu_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},\nu^{(9)},u^{(9)}$:

By $(\nu_1)^{(9)}>0$, $(\nu_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2+(\sigma_1)^{(9)}\nu^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{\nu}_1)^{(9)},,(\bar{\nu}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)}>0$, $(\bar{\nu}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2+(\sigma_2)^{(9)}\nu^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \leq G_{46}(t) \leq \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}v^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}v^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq v^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \le \nu^{(2)}(t) \le (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \le u^{(2)}(t) \le (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(u) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FIFTY

Holographic Quantum Computation: Naturam Expellas Furca, Tamen Usque Recurret

INTRODUCTION—VARIABLES USED

Karl Tordrup et al Holographic Quantum Computing with a formidable proposition and suggestion to use a single mesoscopic ensemble of trapped polar molecules for quantum computing. A "**holographic quantum register" with hundreds of qubits is encoded in collective excitations with definite spatial phase variations**. Each phase pattern is uniquely addressed by optical Raman processes with classical optical fields, while one- and two-qubit gates and qubit readout are accomplished by transferring the qubit states to a stripline microwave cavity field and a Cooper pair box where controllable two-level unitary dynamics and detection is governed by classical microwave fields. **© 2008 The American Physical SocietyURL: http://link.aps.org/doi/10.1103/PhysRevLett.101.040501DOI: 10.1103/PhysRevLett.101.040501**

Holonomic quantum computation Paolo Zanardia http://dx.doi.org/10.1016/S0375-9601(99)00803-8

(1) Authors show that the notion of generalized Berry phase i.e., non-abelian holonomy, **can be used** for enabling quantum computation.
(2) The computational space **is realized** by an n-fold degenerate eigenspace of a family of Hamiltonians parametrized by a manifold.
(3) The point of M **represents (eb)** classical configuration of control fields and, for multi-partite systems, couplings between subsystems.
(4) Adiabatic loops in the control **induce** non trivial unitary transformations on the computational space.
(5) For a generic system it is shown that this mechanism **allows for** universal quantum computation by composing a generic pair of loops in M.

Vacuum Periodicity in Yang-Mills Quantum Theory R. Jackiw et al

(6) Authors propose a description of the vacuum in Yang-Mills theory **and (eb) arrive at** a physical interpretation of the pseudoparticle solution and the attendant violation of symmetries.
(7) The existence of topologically inequivalent classical gauge fields **gives rise to** a family of quantum mechanical vacua, parametrized by a CP-nonconserving angle.
(8) The requirement of vacuum stability against gauge transformations **renders(eb)** the vacua chirally noninvariant.URL: http://link.aps.org/doi/10.1103/PhysRevLett.37.172DOI: 10.1103/PhysRevLett.37.172

Scanning gate microscopy of localized states in wide graphene constrictions Andrei G. F. Garcia

(9) In graphene nanoconstrictions, charge puddles **due to** substrate-induced potential inhomogeneities **can form** quantum dots.

NOTATION

Module One

Authors show that the notion of generalized Berry phase i.e., non-abelian holonomy, **can be used** for enabling quantum computation.

G_{13} : Category one of quantum computation

G_{14} : Category two of quantum computation

G_{15} : Category three ofquantum computation

T_{13} : Category one of notion of generalized Berry phase i.e., non-abelian holonomy

T_{14} : Category two of notion of generalized Berry phase i.e., non-abelian holonomy

T_{15} : Category three of notion of generalized Berry phase i.e., non-abelian holonomy

Module Two

The computational space **is realized** by an n-fold degenerate eigenspace of a family of Hamiltonians parametrized by a manifold

G_{16} : Category one of n-fold degenerate eigenspace of a family of Hamiltonians parametrized by a manifold

G_{17} : Category two of n-fold degenerate eigenspace of a family of Hamiltonians parametrized by a manifold

G_{18} : Category three of n-fold degenerate eigenspace of a family of Hamiltonians parametrized by a manifold

T_{16} : Category one of computational space

T_{17} : Category two of computational space

T_{18} : Category three of computational space

Module three

The point of M **represents (eb)** classical configuration of control fields and, for multi-partite systems, couplings between subsystems

G_{20} : Category one of point of M. There are many such M points and investigating systems based on the characterstics of which the doctrine and dictum of classification is formulated.

G_{21} : Category two of point of M. There are many such M points and investigating systems based on the characterstics of which the doctrine and dictum of classification is formulated.

G_{22} : Category three of point of M. There are many such M points and investigating systems based on the characterstics of which the doctrine and dictum of classification is formulated.

T_{20} : Category one of classical configuration of control fields and, for multi-partite systems, couplings between subsystems

T_{21} : Category two of classical configuration of control fields and, for multi-partite systems, couplings between subsystems

T_{22} : Category three of classical configuration of control fields and, for multi-partite systems, couplings between subsystems

Module four

Adiabatic loops in the control **induce** non trivial unitary transformations on the computational space.

G_{24} : Category one of Adiabatic loops in the control

G_{25} : Category two of Adiabatic loops in the control

G_{26} : Category three of Adiabatic loops in the control

T_{24} : Category one of non trivial unitary transformations on the computational space

T_{25} : Category two of non trivial unitary transformations on the computational space

T_{26} : Category three of non trivial unitary transformations on the computational space

Module five

For a generic system it is shown that this mechanism **allows for** universal quantum computation by composing a generic pair of loops in M.

G_{28} : Category one of generic system it is shown that this mechanism

G_{29} : Category two of generic system it is shown that this mechanism

G_{30} : Category three of generic system it is shown that this mechanism

T_{28} : Category one of universal quantum computation by composing a generic pair of loops in M.

T_{29} : Category two of universal quantum computation by composing a generic pair of loops in M.

T_{30} : Category three of universal quantum computation by composing a generic pair of loops in M.

Module six

Authors propose a description of the vacuum in Yang-Mills theory **and (eb) arrive at** a physical interpretation of the pseudoparticle solution and the attendant violation of symmetries.

G_{32} : Category one of vacuum in Yang-Mills theory

G_{33} : Category two of vacuum in Yang-Mills theory

G_{34} : Category three of vacuum in Yang-Mills theory

T_{32} : Category one of physical interpretation of the pseudoparticle solution and the attendant violation of symmetries.

T_{33} : Category two of physical interpretation of the pseudoparticle solution and the attendant violation of symmetries.

T_{34} : Category three of physical interpretation of the pseudoparticle solution and the attendant violation of symmetries.

Module seven

The existence of topologically inequivalent classical gauge fields **gives rise to** a family of quantum mechanical vacua, parametrized by a CP-nonconserving angle

G_{36} : Category one of existence of topologically inequivalent classical gauge fields

G_{37} : Category two of existence of topologically inequivalent classical gauge fields

G_{38} : Category three of existence of topologically inequivalent classical gauge fields

T_{36} : Category one of family of quantum mechanical vacua, parametrized by a CP-nonconserving angle

T_{37} : Category two of family of quantum mechanical vacua, parametrized by a CP-nonconserving angle

T_{38} : Category three of family of quantum mechanical vacua, parametrized by a CP-nonconserving angle

Module eight

The requirement of vacuum stability against gauge transformations **renders (eb)** the vacua chirally noninvariant.

G_{40} : Category one of requirement of vacuum stability against gauge transformations

G_{41} : Category two of requirement of vacuum stability against gauge transformations

G_{42} : Category three of requirement of vacuum stability against gauge transformations

T_{40} : Category one of vacua chirally noninvariant.

T_{41} : Category two of vacua chirally noninvariant.

T_{42} : Category three of vacua chirally noninvariant.

Module Nine

In graphene nanoconstrictions, charge puddles **due to** substrate-induced potential inhomogeneities **can form** quantum dots.

G_{44} : Category one of substrate-induced potential inhomogeneities **can form** quantum dots.

G_{45} : Category two of substrate-induced potential inhomogeneities **can form** quantum dots.

G_{46} : Category three of substrate-induced potential inhomogeneities **can form** quantum dots.

T_{44} : Category one of graphene nanoconstrictions, charge puddles

T_{45} : Category two of graphene nanoconstrictions, charge puddles

T_{46} : Category three of graphene nanoconstrictions, charge puddles

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**
$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad (55)$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad (56)$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad (57)$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{20})^{(3)}\,+(a''_{20})^{(3)}(T_{21},t)\,+(a''_{16})^{(2,2,2)}(T_{17},t)\,+(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)\,+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)\,+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t)\,+(a''_{40})^{(8,8,8,8)}(T_{41},t)\,+(a''_{44})^{(9,9,9)}(T_{45},t) \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c} (a'_{21})^{(3)}\,+(a''_{21})^{(3)}(T_{21},t)\,+(a''_{17})^{(2,2,2)}(T_{17},t)\,+(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)\,+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)\,+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t)\,+(a''_{41})^{(8,8,8,8)}(T_{41},t)\,+(a''_{45})^{(9,9,9)}(T_{45},t) \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{22})^{(3)}\,+(a''_{22})^{(3)}(T_{21},t)\,+(a''_{18})^{(2,2,2)}(T_{17},t)\,+(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)\,+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)\,+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t)\,+(a''_{42})^{(8,8,8,8)}(T_{41},t)\,+(a''_{46})^{(9,9,9)}(T_{45},t) \end{array}\right] G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{c} (b'_{20})^{(3)}\,-(b''_{20})^{(3)}(G_{23},t)\,-(b''_{16})^{(2,2,2)}(G_{19},t)\,-(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)\,-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)\,-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t)\,-(b''_{40})^{(8,8,8,8)}(G_{43},t)\,-(b''_{44})^{(9,9,9)}(G_{47},t) \end{array}\right] T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{c} (b'_{21})^{(3)}\,-(b''_{21})^{(3)}(G_{23},t)\,-(b''_{17})^{(2,2,2)}(G_{19},t)\,-(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)\,-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)\,-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t)\,-(b''_{41})^{(8,8,8,8)}(G_{43},t)\,-(b''_{45})^{(9,9,9)}(G_{47},t) \end{array}\right] T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$

$are\ fourth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
$are\ fifth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
$are\ sixth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
$are\ seventh\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
$are\ eighth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

$Where\ -(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ $are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ $are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ $are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad (84)$$

$where\ \boxed{-(b''_{28})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{29})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{30})^{(5)}(G_{31},t)}\ \ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}\ ,\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}\ ,\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt}=(a_{32})^{(6)}G_{33} \qquad 85$$

$$-\begin{bmatrix}(a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{32}$$

$$\frac{dG_{33}}{dt}=(a_{33})^{(6)}G_{32}-\begin{bmatrix}(a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{33} \qquad 86$$

$$\frac{dG_{34}}{dt}=(a_{34})^{(6)}G_{33}-\begin{bmatrix}(a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{34} \qquad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}\ ,\boxed{+(a''_{33})^{(6)}(T_{33},t)}\ ,\boxed{+(a''_{34})^{(6)}(T_{33},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}\ ,\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\ ,\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\ are\ second\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}\ \ are\ third\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation

coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34} \quad (90)$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$ *95*

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{c} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{c} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \le (r_i)^{(1)} \le (b_i')^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \le (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(WWWWW) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(XXXXX) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(YYYYY) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(ZZZZZ)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(AAAAAA) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(BBBBBB) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\le\ \left(\hat{P}_{13}\right)^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\le\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\le\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(v) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)} t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

208

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)}$$

209

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)}$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),\left((G_{23})^{(2)},(T_{23})^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

214

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad (222)$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad (223)$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself (224)

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric (225)

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40{,}41{,}42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 275

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(8)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\,((G_{43})(t),t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}\,(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}$, $if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)}$,

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}$, $if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}$, $if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}$,

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\leq G_{15}(t)\leq\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\leq T_{13}(t)\leq T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\leq T_{13}(t)\leq\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\leq T_{15}(t)\leq \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\leq-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(\mathrm{T}_{17},t)+(a''_{17})^{(2)}(\mathrm{T}_{17},t)\leq-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\leq-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\leq-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)}G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)}G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)}T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(\nu_1)^{(6)}, (\nu_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, \nu^{(6)}, u^{(6)}$: 350

By $(\nu_1)^{(6)} > 0$, $(\nu_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)},, (\bar{\nu}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, $\boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad (354)$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad (355)$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad (356)$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^{0}e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^{0}e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^{0}}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t}-e^{-(b'_{34})^{(6)}t}\right]+T_{34}^{0}e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^{0}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T_{34}^{0}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $v^{(4)}$:- $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

For $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(v) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523

A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FIFTY ONE

Quantum Information Theory In The Field Of Quantum Gravity: A Hauptvermutung

INTRODUCTION—VARIABLES USED

August 2001, Volume 33, Issue 8, pp 1305-1318 Quantum Computation Toward Quantum Gravity. A. Zizzi General Theory and Gravitation

(1) The aim of this paper is to enlighten the emerging relevance of Quantum Information Theory in the field of Quantum Gravity. As it was suggested by J. A. Wheeler, information theory must play a **relevant role** in understanding the foundations of Quantum Mechanics (the "It from bit" proposal). Here we suggest that quantum information **must play a relevant role** in Quantum Gravity (the "It from qubit" proposal).

(2) The conjecture is that Quantum Gravity, the theory which will reconcile Quantum Mechanics with General Relativity, **can be formulated in terms of** quantum bits of information (qubits) stored in space at the Planck scale.

(3) This conjecture **is based on the** following arguments: a) The holographic principle, b) The loop quantum gravity approach and spin networks, c) Quantum geometry and black hole entropy.

(4) From the above arguments, as they stand in the literature, **it follows that** the edges of spin networks pierce the black hole horizon and excite curvature degrees of freedom on the surface.

(5) These excitations **are** micro-states of Chern-Simons theory and account of the black hole entropy which turns out to be a quarter of the area of the horizon, (in units of Planck area), in accordance with the holographic principle.

(6) Moreover, the states which dominate the counting **correspond to** punctures of spin j = 1/2 and one can in fact visualize each micro-state as a bit of information.

(7) The obvious generalization of this result **is to consider** open spin networks with edges labeled by the spin −1/ 2 representation of SU (2) in a superposed state of spin "on" and spin "down."

(8) The micro-state corresponding to such a puncture **will be a** pixel of area which is "on" and "off" at the same time, and it will encode a qubit of information.

(9) This picture, when applied to quantum cosmology, **describes an** early inflationary universe which is a discrete version of the de Sitter universe.

NOTATION

Module One

The aim of this paper is to enlighten the emerging relevance of Quantum Information Theory in the field of Quantum Gravity. As it was suggested by J. A. Wheeler, information theory must play a **relevant role** in understanding the foundations of Quantum Mechanics (the "It from bit" proposal). Here we suggest that quantum information **must play a relevant role** in Quantum Gravity (the "It from qubit" proposal).

G_{13} : Category one of quantum information; Quantum Gravity

G_{14} : Category two of quantum information; Quantum Gravity

G_{15} : Category three ofquantum information ;Quantum Gravity

T_{13} : Category one of Quantum Gravity ;quantum information

T_{14} : Category two of Quantum Gravity; quantum information

T_{15} : Category three of Quantum Gravity; quantum information

Module Two

The conjecture is that Quantum Gravity, the theory which will reconcile Quantum Mechanics with General Relativity, **can be formulated in terms of** quantum bits of information (qubits) stored in space at the Planck scale.

G_{16} : Category one of quantum bits of information (qubits) stored in space at the Planck scale.

G_{17} : Category two of quantum bits of information (qubits) stored in space at the Planck scale.

G_{18} : Category three of quantum bits of information (qubits) stored in space at the Planck scale.

T_{16} : Category one of conjecture is that Quantum Gravity, the theory which will reconcile Quantum Mechanics with General Relativity

T_{17} : Category two of conjecture is that Quantum Gravity, the theory which will reconcile Quantum Mechanics with General Relativity

T_{18} : Category three of conjecture is that Quantum Gravity, the theory which will reconcile Quantum Mechanics with General Relativity

Module three

This conjecture **is based on the** following arguments: a) The holographic principle, b) The loop quantum gravity approach and spin networks, c) Quantum geometry and black hole entropy

G_{20} : Category one of a) The holographic principle, b) The loop quantum gravity approach and spin networks, c) Quantum geometry and black hole entropy

G_{21} : Category two of a) The holographic principle, b) The loop quantum gravity approach and spin networks, c) Quantum geometry and black hole entropy

G_{22} : Category three of a) The holographic principle, b) The loop quantum gravity approach and spin networks, c) Quantum geometry and black hole entropy

T_{20} : Category one of Quantum Gravity, the theory which will reconcile Quantum Mechanics with General Relativity, **can be formulated in terms of** quantum bits of information (qubits) stored in space at the Planck scale.

T_{21} : Category two of Quantum Gravity, the theory which will reconcile Quantum Mechanics with General Relativity, **can be formulated in terms of** quantum bits of information (qubits) stored in space at the Planck scale.

T_{22} : Category three of Quantum Gravity, the theory which will reconcile Quantum Mechanics with General Relativity, **can be formulated in terms of** quantum bits of information (qubits) stored in space at the Planck scale.

Module four

From the above arguments, as they stand in the literature, **it follows that** the edges of spin networks pierce the black hole horizon and excite curvature degrees of freedom on the surface

G_{24} : Category one of above arguments, as they stand in the literature

G_{25} : Category two of above arguments, as they stand in the literature

G_{26} : Category three of above arguments, as they stand in the literature

T_{24} : Category one of edges of spin networks pierce the black hole horizon and excite curvature degrees of freedom on the surface

T_{25} : Category two of edges of spin networks pierce the black hole horizon and excite curvature degrees of freedom on the surface

T_{26} : Category three of edges of spin networks pierce the black hole horizon and excite curvature degrees of freedom on the surface

Module five

These excitations **are** micro-states of Chern-Simons theory and account of the black hole entropy which turns out to be a quarter of the area of the horizon, (in units of Planck area), in accordance with the holographic principle

G_{28} : Category one of micro-states of Chern-Simons theory and account of the black hole entropy which turns out to be a quarter of the area of the horizon, (in units of Planck area), in accordance with the holographic principle

G_{29} : Category two of micro-states of Chern-Simons theory and account of the black hole entropy which turns out to be a quarter of the area of the horizon, (in units of Planck area), in accordance with the holographic principle

G_{30} : Category three of micro-states of Chern-Simons theory and account of the black hole entropy which turns out to be a quarter of the area of the horizon, (in units of Planck area), in accordance with the holographic principle

T_{28} : Category one of These excitations

T_{29} : Category two of These excitations

T_{30} : Category three of These excitations

Module six

Moreover, the states which dominate the counting **correspond to** punctures of spin j = 1/2 and one can in fact visualize each micro-state as a bit of information.

G_{32} : Category one of states which dominate the counting; punctures of spin j = 1/2 and one can in fact visualize each micro-state as a bit of information.

G_{33} : Category two of states which dominate the counting; punctures of spin j = 1/2 and one can in fact visualize each micro-state as a bit of information.

G_{34} : Category three of states which dominate the counting; punctures of spin j = 1/2 and one can in fact visualize each micro-state as a bit of information.

T_{32} : Category one of punctures of spin j = 1/2 and one can in fact visualize each micro-state as a bit of

information. ; states which dominate the counting

T_{33} : Category two of punctures of spin j = 1/2 and one can in fact visualize each micro-state as a bit of information. ;states which dominate the counting

T_{34} : Category three of punctures of spin j = 1/2 and one can in fact visualize each micro-state as a bit of information. ;states which dominate the counting

Module seven

The obvious generalization of this result **is to consider** open spin networks with edges labeled by the spin −1/ 2 representation of SU (2) **in a** superposed state of spin "on" and spin "down."

G_{36} : Category one of superposed state of spins "on" and spins "down."

G_{37} : Category two of superposed state of spins "on" and spins" down."

G_{38} : Category three of superposed state of spins "on" and spins "down."

T_{36} : Category one of open spin networks with edges labeled by the spin −1/ 2 representation of SU (2)

T_{37} : Category two of open spin networks with edges labeled by the spin −1/ 2 representation of SU (2)

T_{38} : Category three of open spin networks with edges labeled by the spin −1/ 2 representation of SU (2)

Module eight

The micro-state corresponding to such a puncture **will be a** pixel of area which is "on" and "off" at the same time, and it will encode a qubit of information

G_{40} : Category one of pixel of area which is "on" and "off" at the same time, and it will encode a qubit of information

G_{41} : Category two of pixel of area which is "on" and "off" at the same time, and it will encode a qubit of information

G_{42} : Category three of pixel of area which is "on" and "off" at the same time, and it will encode a qubit of information

T_{40} : Category one of micro-state corresponding to such a puncture

T_{41} : Category two of micro-state corresponding to such a puncture

T_{42} : Category three of micro-state corresponding to such a puncture

Module Nine

This picture, when applied to quantum cosmology, **describes an** early inflationary universe which is a discrete version of the de Sitter universe.

G_{44} : Category one of picture, when applied to quantum cosmology

G_{45} : Category two of picture, when applied to quantum cosmology

G_{46} : Category three of picture, when applied to quantum cosmology. In theoretical physics, quantum cosmology **is a field** attempting to study the effect of quantum mechanics **on (e&eb) the** formation of the universe, or its early evolution, especially just after the Big Bang. Despite many attempts, such as the Wheeler-DeWitt equation and more recently loop quantum cosmology, the field remains a rather speculative branch of quantum gravity. An important problem in this field **is the** origin of physical information in the universe

T_{44} : Category one of early inflationary universe which is a discrete version of the de Sitter universe.

T_{45} : Category two of early inflationary universe which is a discrete version of the de Sitter universe.

T_{46} : Category three of early inflationary universe which is a discrete version of the de Sitter universe.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients
Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G,t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G,t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}),t\big)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}),t\big)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}),t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}),t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27}),t\big)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27}),t\big)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27}),t\big)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\big((G_{27}),t\big) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}),t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}),t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}),t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}),t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}),t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}),t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad 63$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition

coefficients for category 1,2 and 3

$-(b_{28}'')^{(5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b_{32}'')^{(6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b_{36}'')^{(7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b_{40}'')^{(8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b_{44}'')^{(9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c}(a_{20}')^{(3)}+(a_{20}'')^{(3)}(T_{21},t)\ +(a_{16}'')^{(2,2,2)}(T_{17},t)\ +(a_{13}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)\ +(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)\ +(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{36}'')^{(7,7,7,7)}(T_{37},t)\ +(a_{40}'')^{(8,8,8,8)}(T_{41},t)\ +(a_{44}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c}(a_{21}')^{(3)}+(a_{21}'')^{(3)}(T_{21},t)\ +(a_{17}'')^{(2,2,2)}(T_{17},t)\ +(a_{14}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)\ +(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)\ +(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{37}'')^{(7,7,7,7)}(T_{37},t)\ +(a_{41}'')^{(8,8,8,8)}(T_{41},t)\ +(a_{45}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c}(a_{22}')^{(3)}+(a_{22}'')^{(3)}(T_{21},t)\ +(a_{18}'')^{(2,2,2)}(T_{17},t)\ +(a_{15}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)\ +(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)\ +(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{38}'')^{(7,7,7,7)}(T_{37},t)\ +(a_{42}'')^{(8,8,8,8)}(T_{41},t)\ +(a_{46}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{22} \quad 69$$

$+(a_{20}'')^{(3)}(T_{21},t)$, $+(a_{21}'')^{(3)}(T_{21},t)$, $+(a_{22}'')^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a_{44}'')^{(9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for

category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad 78$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

are fourth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$

are fifth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$

are sixth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$

are seventh detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$

are eighth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)} & +(a_{36}'')^{(7)}(T_{37},t) & +(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)} & +(a_{37}'')^{(7)}(T_{37},t) & +(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)} & +(a_{38}'')^{(7)}(T_{37},t) & +(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a_{36}'')^{(7)}(T_{37},t)$, $(a_{37}'')^{(7)}(T_{37},t)$, $(a_{38}'')^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)} G_{44}$$
$$- \left[\begin{array}{ccc} (a'_{45})^{(9)} & \boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \quad \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)} G_{45}$$
$$- \left[\begin{array}{ccc} (a'_{46})^{(9)} & \boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \quad \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)} T_{45} -$$
$$\left[\begin{array}{ccc} (b'_{44})^{(9)} & \boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \quad \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{13}$$
$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, \ (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ \ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(CCCCCC) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(DDDDDD) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(EEEEEE) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(FFFFFF)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(GGGGGG) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(HHHHHH) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ \ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{13})^{(1)}$, $T_i^0 \leq (\hat{Q}_{13})^{(1)}$, 155

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{16})^{(2)}$, $T_i^0 \leq (\hat{Q}_{16})^{(2)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^{0} + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^{0} + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^{0} + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(w) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} +$$

$$G_{13}^{(2)} \left|(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)} \left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} t} \leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}} \left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)} (T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)} \left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)} \left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)};\ (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\tilde{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(9)}t)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},, (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad (333)$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad (334)$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad (335)$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad (336)$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\leq T_{28}(t)\leq T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \qquad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\leq T_{28}(t)\leq \frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \qquad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\leq T_{30}(t)\leq \qquad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\leq -(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\leq -(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\leq -(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\leq -(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)} (m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)} (\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

406

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case,

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

407

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t)\right) - (a_{41}'')^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(w) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FIFTY TWO

Interference Fringes Of Electrons In Backscattering: A Posse Ad Esse

INTRODUCTION—VARIABLES USED

Backscatter of waves in physical space

Backscattering occurs in quite different physical situations. The incoming waves or particles can be deflected from their original direction by quite different mechanisms:

(a) Diffuse reflection from large particles and Mie scattering, causing alpenglow and gegenschein, and showing up in weather radar;
(b) Inelastic collisions between electromagnetic waves and the transmitting medium (Brillouin scattering and Raman scattering), important in fiber optics, see below;
(c) Elastic collisions between accelerated ions and a sample (Rutherford backscattering)
(d) Bragg diffraction from crystals, used in inelastic scattering experiments (neutron backscattering, X-ray backscattering spectroscopy);

Compton scattering, used in Backscatter X-ray imaging.

Sometimes, the scattering is more or less isotropic, i. e. the incoming particles are scattered randomly in various directions, with no particular preference for backward scattering. In these cases, the term "backscattering" just designates the detector location chosen for some practical reasons:

(1) in X-ray imaging, backscattering means just the opposite of transmission imaging;
(2) in inelastic neutron or X-ray spectroscopy, backscattering geometry is chosen because it optimizes the energy resolution;
(3) In astronomy, backscattered light is that which is reflected with a phase angle of less than 90°.
(4) In other cases, the scattering intensity is enhanced in backward direction.

This can have different reasons:

(i) In alpenglow, red light prevails because the blue part of the spectrum is depleted by Rayleigh scattering.
(ii) In gegenschein, constructive interference might play a role (this needs verification).
(iii) Coherent backscattering is observed in random media; for visible light most typically in suspensions like milk. Due to weak localization, enhanced multiple scattering is observed in back direction.
(iv) The Back Scattering Alignment (BSA) coordinate system is often used in radar applications
(v) The Forward Scattering Alignment (FSA) coordinate system is primarily used in optical applications
(vi) Radar, especially weather radar
(vii) Backscattering is the principle behind radar systems.

In weather radar, backscattering is proportional to the 6th power of the diameter of the target multiplied by its inherent reflective properties. Water is almost 4 times more reflective than ice but droplets are much

smaller than snowflakes or hail stones. So the backscattering is dependent on a mix of these two factors. The strongest backscatter comes from hail and large graupel (solid ice) due to their sizes. Another strong return is from melting snow or wet sleet, as they combine size and water reflectivity. They often show up as much higher rates of precipitation than actually occurring in what is called a bright band. Rain is a moderate backscatter, being stronger with large drops (such as from a thunderstorm) and much weaker with small droplets (such as mist or drizzle). Snow has rather weak backscatter.

Backscatter in waveguides

The backscattering method is also employed in fiber optics applications to detect optical faults. Light propagating through a fiber optic cable gradually attenuates due to Rayleigh scattering. Faults are thus detected by monitoring the variation of part of the Rayleigh backscattered light. Since the backscattered light attenuates exponentially as it travels along the optical fiber cable, the attenuation characteristic is represented in a logarithmic scale graph. If the slope of the graph is steep, then power loss is high. If the slope is gentle, then optical fiber has a satisfactory loss characteristic.

The loss measurement by the backscattering method allows measurement of a fiber optic cable at one end without cutting the optical fiber hence it can be conveniently used for the construction and maintenance of optical fibers.

Backscatter in photography

Main article: Orb The term backscatter in photography refers to light from a flash or strobe reflecting back from particles in the lens's field of view causing specks of light to appear in the photo. This gives rise to what are sometimes referred to as orb artifacts. Photographic backscatter can result from snowflakes, rain or mist, or airborne dust. Due to the size limitations of the modern compact and ultra-compact cameras, especially digital cameras, the distance between the lens and the built-in flash has decreased, thereby decreasing the angle of light reflection to the lens and increasing the likelihood of light reflection off normally sub-visible particles. Hence, the orb artifact is commonplace with small digital or film camera photographs

Scanning gate microscopy of localized states in wide graphene constrictions Andrei G. F. Garcia

(1) Authors use a scanning gate microscope to probe a short, wide graphene constriction **and observe** a set of ring-shaped resonances in maps of the graphene conductance as a function of the local potential perturbation, a signature of quantum dot-mediated transport.
(2) By varying the scanning tip height and applied tip voltage, **they observe (e) the** screening effect of the nearby graphene. Introducing a simple capacitive model to account for screening, they model the conduction of the graphene constriction and how the tip gates the quantum dots. We identify the locations of some of the quantum dots and extract their parameters.URL:http://link.aps.org/doi/10.1103/PhysRevB.87.085446DOI: 10.1103/PhysRevB.87.085446

A A Kozikov et al 2013 New J. Phys. 15 013056 doi:10.1088/1367-2630/15/1/013056 Interference of electrons in backscattering through a quantum point contact

(3) Scanning gate microscopy is used to locally investigate electron transport in a high-mobility two-dimensional electron gas **formed (eb) in a** GaAs/AlGaAs heterostructure.
(4) Using quantum point contacts, they observe branches **caused (e) by** electron backscattering decorated with interference fringes similar to previous observations by Topinka et al (2000 Science 289 2323).

(5) Authors investigate the branches at different points of a conductance plateau as well as between plateaus, and **demonstrate that** the most dramatic changes in branch pattern occur at the low-energy side of the conductance plateaus.

(6) The branches disappear at magnetic fields as low as 50 mT, **demonstrating** the importance of backscattering for the observation of the branching effect. The spacing between the interference fringes varies by more than 50% for different branches across scales of microns. Several scenarios are discussed to explain this observation.

Shota Kuwahara et al 2009 Nanotechnology 20 225702 doi:10.1088/0957-4484/20/22/225702 A new AFM–HRTEM combined technique for probing isolated carbon nanotubes

(7) A new technique has been developed for characterizing individual carbon nanotubes (CNTs). This technique combines atomic force microscopy (AFM) measurements with an independent determination of the CNT structure obtained **by (e) using** thin carbon grid membranes for high-resolution transmission electron microscopy (HRTEM) measurements.

(8) The membrane grids are easily transferred onto silicon oxide substrates, where they are observed by AFM and an optical charged-coupled device. Electrostatic force microscopy measurements are also performed on single- and double-walled CNTs whose structures had previously been determined by HRTEM. **The results reveal** that the diameter and outer tube length have a greater effect **on (eb) determining the** capacitance of the nanotubes than the presence of the inner tube of double-walled CNTs.

Non-Abelian anyons and topological quantum computation Chetan Nayak et al

(9) Topological quantum computation has emerged as one of the most exciting approaches **to constructing** a fault-tolerant quantum computer.

NOTATION

Module One

Authors use a scanning gate microscope probe a short, wide graphene constriction **and observe** a set of ring-shaped resonances in maps of the graphene conductance as a function of the local potential perturbation, a signature of quantum dot-mediated transport.

G_{13} : Category one of scanning gate microscope probe a short, wide graphene constriction

G_{14} : Category two of scanning gate microscope probe a short, wide graphene constriction

G_{15} : Category three of scanning gate microscope probe a short, wide graphene constriction

T_{13} : Category one of set of ring-shaped resonances in maps of the graphene conductance as a function of the local potential perturbation, a signature of quantum dot-mediated transport.

T_{14} : Category two of set of ring-shaped resonances in maps of the graphene conductance as a function of the local potential perturbation, a signature of quantum dot-mediated transport.

T_{15} : Category three of set of ring-shaped resonances in maps of the graphene conductance as a function of the local potential perturbation, a signature of quantum dot-mediated transport.

Module Two

By varying the scanning tip height and applied tip voltage, **they observe (eb) the** screening effect of the nearby graphene. Introducing a simple capacitive model to account for screening, they model the conduction of the graphene constriction and how the tip gates the quantum dots. They identify the locations of some of the quantum dots and extract their parameters.

G_{16} : Category one of variation of scanning tip height and applied tip voltage

G_{17} : Category two of variation of scanning tip height and applied tip voltage

G_{18} : Category three of variation of scanning tip height and applied tip voltage

T_{16} : Category one of screening effect of the nearby graphene.

T_{17} : Category two of screening effect of the nearby graphene.

T_{18} : Category three of screening effect of the nearby graphene.

Module three

Scanning gate microscopy **is (e) used** to locally investigate electron transport in a high-mobility two-dimensional electron gas **formed (eb) in a** GaAs/AlGaAs heterostructure

G_{20} : Category one of locally investigate electron transport in a high-mobility two-dimensional electron gas **formed (eb) in a** GaAs/AlGaAs heterostructure

G_{21} : Category two of locally investigate electron transport in a high-mobility two-dimensional electron gas **formed (eb) in a** GaAs/AlGaAs heterostructure

G_{22} : Category three of locally investigate electron transport in a high-mobility two-dimensional electron gas **formed (eb) in a** GaAs/AlGaAs heterostructure

T_{20} : Category one of Scanning gate microscopy Various investigating systems prevail. Characterstics of those systems are taken in to consideration in the consummation of the stratification consolidation profile.

T_{21} : Category two of Scanning gate microscopy

T_{22} : Category three of Scanning gate microscopy

Module four

Using quantum point contacts, they observe branches **caused (e) by** electron backscattering decorated with interference fringes similar to previous observations by Topinka et al (2000 Science 289 2323).

G_{24} : Category one of electron backscattering decorated with interference fringes similar to previous observations by Topinka et al (2000 Science 289 2323).

G_{25} : Category two of electron backscattering decorated with interference fringes similar to previous observations by Topinka et al (2000 Science 289 2323).

G_{26} : Category three of electron backscattering decorated with interference fringes similar to previous observations by Topinka et al (2000 Science 289 2323).

T_{24} : Category one of (using) quantum point contacts ,branches could be observed

T_{25} : Category two of (using) quantum point contacts ,branches could be observed

T_{26} : Category three of(using) quantum point contacts ,branches could be observed

Module five

Authors investigate the branches at different points of a conductance plateau as well as between plateaus, and **demonstrate that** the most dramatic changes in branch pattern occur at the low-energy side of the

conductance plateaus

Coherent Branched Flow in a Two-Dimensional Electron Gas M.A.Topinka et al

Locus of essence, sense and perception and presuppositioal resemblances, and primordial exactitude of Semiconductor nanostructures based on two dimensional electron gases (2DEGs) have been studied by **M.A.Topinka et al**(Coherent Branched Flow in Two-Dimensional Electron Gas Semiconductor nanostructures based on two dimensional electron gases (2DEGs))potential to provide new approaches to sensing, information processing, and quantum computation. Much is known about electron transport in 2DEG nanostructures and many remarkable phenomena have been discovered (e.g. weak localization, quantum chaos, universal conductance fluctuations) 1, 2 - yet a fundamental aspect of these devices, namely how electrons move through them, has never been clarified. Important details about the actual pattern of electron flow are not specified by statistical measures such as the mean free path. Scanned probe microscope (SPM) measurements **allow** spatial investigations of nanostructures, and it has **recently become possible** to directly image electron flow through 2DEG devices using newly developed SPM techniques3-13. Here we present SPM images of electron flow from a quantum point contact (QPC) which shows unexpected dynamical channeling - the electron flow forms persistent, narrow, branching channels rather than smoothly spreading fans. Theoretical study of this flow, including electron scattering by impurities and donor atoms, **shows that** the channels are not due to deep valleys in the potential, but rather are **caused by the** indirect cumulative effect of small angle scattering. Surprisingly, the channels are decorated by interference fringes well beyond where the simplest thermal averaging arguments **suggest they** should be found. These findings may have important implications for 2DEG physics and for the design of future nanostructure devices. Images of electron flow from the quantum point contact (QPC) are obtained by raster scanning a negatively charged SPM tip above the surface of the device and simultaneously measuring the position-dependent conductance of the device. The negatively charged tip capacitively couples to the 2DEG, creating a depletion region that backscatters electron waves. When the tip is positioned over areas with high electron flow from the QPC the conductance is decreased, whereas when the tip is over areas of relatively low electron flow the conductance is unmodified. By raster scanning the tip over the sample and simultaneously recording the effect the tip has on device conductance, a two dimensional image of electron flow can be obtained. (**Coherent Branched Flow in a Two-Dimensional Electron Gas M.A.Topinka et al)**

G_{28} : Category one of branches at different points of a conductance plateau as well as between plateaus

G_{29} : Category two of branches at different points of a conductance plateau as well as between plateaus

G_{30} : Category three of branches at different points of a conductance plateau as well as between plateaus

T_{28} : Category one of most dramatic changes in branch pattern occur at the low-energy side of the conductance plateaus

T_{29} : Category two of most dramatic changes in branch pattern occur at the low-energy side of the conductance plateaus

T_{30} : Category three of most dramatic changes in branch pattern occur at the low-energy side of the conductance plateaus

Module six

The branches disappear at magnetic fields as low as 50 mT, **demonstrating** the importance of backscattering for the observation of the branching effect. The spacing between the interference fringes varies by more than

50% for different branches across scales of microns. Several scenarios are discussed to explain this observation

G_{32} : Category one of branches disappear at magnetic fields as low as 50 mT

G_{33} : Category two of branches disappear at magnetic fields as low as 50 mT

G_{34} : Category three of branches disappear at magnetic fields as low as 50 mT

T_{32} : Category one of importance of backscattering for the observation of the branching effect. The spacing between the interference fringes varies by more than 50% for different branches across scales of microns.

T_{33} : Category two of importance of backscattering for the observation of the branching effect. The spacing between the interference fringes varies by more than 50% for different branches across scales of microns.

T_{34} : Category three of importance of backscattering for the observation of the branching effect. The spacing between the interference fringes varies by more than 50% for different branches across scales of microns.

Module seven

A new technique has been developed for characterizing individual carbon nanotubes (CNTs). This technique combines atomic force microscopy (AFM) measurements with an independent determination of the CNT structure obtained **by (e) using** thin carbon grid membranes for high-resolution transmission electron microscopy (HRTEM) measurements

G_{36} : Category one of thin carbon grid membranes for high-resolution transmission electron microscopy (HRTEM) measurements

G_{37} : Category two of thin carbon grid membranes for high-resolution transmission electron microscopy (HRTEM) measurements

G_{38} : Category three of thin carbon grid membranes for high-resolution transmission electron microscopy (HRTEM) measurements

T_{36} : Category one of new technique has been developed for characterizing individual carbon nanotubes (CNTs). This technique combines atomic force microscopy (AFM) measurements with an independent determination of the CNT structure obtained

T_{37} : Category two of new technique has been developed for characterizing individual carbon nanotubes (CNTs). This technique combines atomic force microscopy (AFM) measurements with an independent determination of the CNT structure obtained

T_{38} : Category three of new technique has been developed for characterizing individual carbon nanotubes (CNTs). This technique combines atomic force microscopy (AFM) measurements with an independent determination of the CNT structure obtained

Module eight

The membrane grids are easily transferred onto silicon oxide substrates, where they are observed by AFM and an optical charged-coupled device. Electrostatic force microscopy measurements are also performed on single- and double-walled CNTs whose structures had previously been determined by HRTEM. **The results**

reveal that the diameter and outer tube length have a greater effect **on (eb) determining the** capacitance of the nanotubes than the presence of the inner tube of double-walled CNTs.

G_{40} : Category one of membrane grids are easily transferred onto silicon oxide substrates, where they are observed by AFM and an optical charged-coupled device. Electrostatic force microscopy measurements are also performed on single- and double-walled CNTs whose structures had previously been determined by HRTEM

G_{41} : Category two of membrane grids are easily transferred onto silicon oxide substrates, where they are observed by AFM and an optical charged-coupled device. Electrostatic force microscopy measurements are also performed on single- and double-walled CNTs whose structures had previously been determined by HRTEM

G_{42} : Category three of membrane grids are easily transferred onto silicon oxide substrates, where they are observed by AFM and an optical charged-coupled device. Electrostatic force microscopy measurements are also performed on single- and double-walled CNTs whose structures had previously been determined by HRTEM

T_{40} : Category one of diameter and outer tube length have a greater effect **on (eb) determining the** capacitance of the nanotubes than the presence of the inner tube of double-walled CNTs.

T_{41} : Category two of diameter and outer tube length have a greater effect **on (eb) determining the** capacitance of the nanotubes than the presence of the inner tube of double-walled CNTs.

T_{42} : Category three of diameter and outer tube length have a greater effect **on (eb) determining the** capacitance of the nanotubes than the presence of the inner tube of double-walled CNTs.

Module Nine

(1) Topological quantum computation has emerged as one of the most exciting approaches **to constructing** a fault-tolerant quantum computer
(2) A topological quantum computer is a theoretical quantum computer that **employs t**wo-dimensional quasiparticles called anyons, whose world lines cross over one another to form braids in a three-dimensional spacetime (i.e., one temporal plus two spatial dimensions).
(3) These braids **form the** logic gates that make up the computer.
(4) The advantage of a quantum computer **based on** quantum braids over using trapped quantum particles is that the former is much more stable.
(5) The smallest perturbations **can cause** a quantum particle to decohere and introduce errors in the computation, but such small perturbations do not change the topological properties of the braids.
(6) This is like the effort required to cut a string and reattach the ends to form a different braid, as opposed to a ball (representing an ordinary quantum particle in four-dimensional spacetime) simply bumping into a wall. The original proposal for topological quantum computation is due **to Alexei Kitaev in 1997**.
(7) Elements of a topological quantum computer originate in a purely mathematical realm, experiments in Fractional quantum Hall systems **indicate these** elements may be created in the real world using semiconductors made of gallium arsenide near absolute zero and subjected to strong magnetic fields.

G_{44} : Category one of Topological quantum computation has emerged as one of the most exciting approaches; braids; smallest perturbations; Elements of a topological quantum computer originate in a purely mathematical realm, experiments in Fractional quantum Hall systems

G_{45} : Category two of Topological quantum computation has emerged as one of the most exciting approaches; braids; smallest perturbations; Elements of a topological quantum computer originate in a purely mathematical realm, experiments in Fractional quantum Hall systems

G_{46} : Category three of Topological quantum computation has emerged as one of the most exciting approaches; braids; smallest perturbations; Elements of a topological quantum computer originate in a purely mathematical realm, experiments in Fractional quantum Hall systems

T_{44} : Category one of fault-tolerant quantum computer; logic gates that make up the computer; quantum particle to decohere and introduce errors in the computation, but such small perturbations do not change the topological properties of the braids; elements may be created in the real world using semiconductors made of gallium arsenide near absolute zero and subjected to strong magnetic fields.

T_{45} : Category two of fault-tolerant quantum computer logic gates that make up the computer; quantum particle to decohere and introduce errors in the computation, but such small perturbations do not change the topological properties of the braids; elements may be created in the real world using semiconductors made of gallium arsenide near absolute zero and subjected to strong magnetic fields.

T_{46} : Category three of fault-tolerant quantum computer; logic gates that make up the computer; quantum particle to decohere and introduce errors in the computation, but such small perturbations do not change the topological properties of the braids; elements may be created in the real world using semiconductors made of gallium arsenide near absolute zero and subjected to strong magnetic fields.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)} \ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) \quad +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{ccc} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) \quad +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right]G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) \quad +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right]G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for

category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt}=(a_{16})^{(2)}G_{17}-\begin{bmatrix}(a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t)\\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t)\end{bmatrix}G_{16} \quad 61$$

$$\frac{dG_{17}}{dt}=(a_{17})^{(2)}G_{16}-\begin{bmatrix}(a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t)\\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t)\end{bmatrix}G_{17} \quad 62$$

$$\frac{dG_{18}}{dt}=(a_{18})^{(2)}G_{17}-\begin{bmatrix}(a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t)\\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t)\end{bmatrix}G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt}=(b_{16})^{(2)}T_{17}-\begin{bmatrix}(b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t)\\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t)\\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t)\end{bmatrix}T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients

for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)}&\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}&\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}&\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}&\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}&\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}&\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)}&\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}&\boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}&\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}&\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}&\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}&\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)}&\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}&\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}&\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}&\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}&\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}&\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)}&\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}&\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}&\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}&\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}&\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}&\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)}&\boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}&\boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}&\boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}&\boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}&\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}&\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad 78$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad 81$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\,-(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\,-(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\,-(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)}$ 103

$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)}$ 104

$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)}$ 105

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t}$ 107

$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t}$ 108

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1$ 110

$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1$ 111

Where we suppose

$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22$ 112

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with (116)
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \quad (117)$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32{,}33{,}34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32{,}33{,}34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(IIIIII) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36{,}37{,}38$ 131

(JJJJJJ) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$

132

(KKKKKK) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(LLLLLL)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(MMMMMM) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(NNNNNN) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad 146\ A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{20})^{(3)}, T_i^0 \le (\hat{Q}_{20})^{(3)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad (169)$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that (170)

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that (172)

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that (174)

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(x) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}ds_{(16)} +$

$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\tilde{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right) \quad (248)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also G_{33} *and* G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t} \; and \; (\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2 \; and \; \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37} \; and \; G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},, (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)},\ \boldsymbol{if}\ (\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)},\ \boldsymbol{if}\ (v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \boldsymbol{if}\ (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$
and $\boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)},\ \boldsymbol{if}\ (\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}$, $\boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}$, $\boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(x) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FIFTY THREE

Non-Abelian Anyons: Ad Augusta Per Angusta

INTRODUCTION—VARIABLES USED

Monolayer doping (MLD) is a well controlled, wafer-scale surface doping technique first developed at the University of California, Berkeley, in 2007. This work is aimed for attaining controlled doping of semiconductor materials with atomic accuracy, especially at nanoscale, which is not easily obtained by other existing technologies. This technique is currently **used for** fabricating ultrashallow junctions (USJs) as the heavily doped source/drain (S/D) contacts of metal-oxide-semiconductor field effect transistors (MOSFETs) as well as enabling dopant profiling of nanostructures.

This MLD technique **utilizes the** crystalline nature of semiconductors and its self-limiting surface reaction properties to form highly uniform, self-assembled, covalently bonded dopant-containing monolayers followed by a subsequent annealing step for the incorporation and diffusion of dopants.

The monolayer formation reaction is self-limiting, thereby, **resulting i**n the deterministic coverage of dopant atoms on the surface.

MLD differs from other conventional doping techniques such as spin-on-dopants (SODs) and gas phase doping techniques **in the way of** dopant dose control.

Such control in MLD is much more precise **due to the** self-limiting formation of covalently attached dopants on the surface while the SODs just rely on the thickness control of the spin-on oxide and the gas phase technique depends on the control of dopant gas flow rate; therefore, the excellent dose control in MLD **can yield** the exact tuning of the resulting dopant profile.

Compared to ion-implantation, MLD **does not involve** the energetic introduction of dopant species into the semiconductor lattice where crystal damages are induced.

In the case of implantation, defects such as interstitials and vacancies are inevitably **generated,** which interact with the dopants to further broaden the junction profile. This is known as the transient-enhanced diffusion (TED), which **limits the** formation of good quality of USJs.

Also, stochastic variation in the dopant positioning and sever stoichiometric imbalance are thus induced for binary and tertiary compound semiconductors by the implantation techniques. In contrast, all MLD dopant atoms are thermally diffused from the crystal surface to the bulk and the dopant profile can be easily controlled by the thermal budget. Since the MLD system can be classified as a limited source model, this is desirable for controlled USJ fabrication with high uniformity and low stochastic variation. Combined with the excellent dopant dose uniformity and coverage in MLD, **it is especially attractive for** doping nonplanar

devices such as fin-FETs and nanowires. As a result, high quality sub-5 nm ultra-shallow junction has been demonstrated in silicon via the **use of** this MLD technique**.(Wikipedia, J. C. Ho, R. Yerushalmi, Z. A. Jacobson, Z. Fan, R. L. Alley, A. Javey, "Controlled nanoscale doping of semiconductors via molecular monolayers", Nature Materials, 7 (1), 62-67, 2008.)**

An anyon is a type of particle that occurs only in two-dimensional systems, with properties much **less restricted than** fermions and bosons; the operation of exchanging two identical particles **may cause a** global phase shift but cannot affect observables. Anyons are generally classified as abelian or non-abelian, as explained below. Abelian anyons have been detected and **play a major role** in the fractional quantum Hall effect. Non-abelian anyons have not been definitively detected although this is an active area of research.

Non-abelian anyons

Is topological order stable at non-zero temperature? We assume the statement and give a model (Unsolved model: Both the possibilities are discussed). **Unsolved problem**

(1) Topological order is **stable (e)** at non zero temperature. At non zero temperature **there exists (eb)** stable topological order**.**
(2) Topological order is **not (e) s**table at non zero temperature. At non zero temperature there does not(e) exist stable topological order

In 1988, Jürg Fröhlich showed that it was possible in a valid particle theory for the particle exchange operation **to be** non-commutative (non-Abelian statistics). Gregory Moore, Nicholas Read, and Xiao-Gang Wen pointed out that non-Abelian statistics **can be realized** in fractional quantum Hall effect. While at first non-abelian anyons were generally considered a mathematical curiosity, physicists began pushing toward their discovery when Alexei Kitaev showed that non-abelian anyons **could be used to** construct a topological quantum computer. As of 2012, no experiment has conclusively **demonstrated the** existence of non-abelian anyons although promising hints are emerging in the study of the ν = 5/2 FQHE state Moore, Gregory; Read, Nicholas (1991). **("Nonabelions in the fractional quantum hall effect". Nuclear Physics B 360 (2–3): 362. Doi: 10.1016/0550-3213(91)90407-O.; Xiao-Gang Wen Non-Abelian Statistics in the FQH states Phys. Rev. Lett. 66, 802 (1991); ^ Stern, Ady (2010). "Non-Abelian states of matter". Nature 464 (7286): 187–93. Doi: 10.1038/nature08915. PMID 20220836. ^ Sanghun an; Jiang; Choi; Kang; Simon; Pfeiffer; West; Baldwin (2011). "Braiding of Abelian and Non-Abelian Anyons in the Fractional Quantum Hall Effect". ArXiv: 1112.3400 [cond-mat.mes-hall].)**

Non-Abelian anyons and topological quantum computation Chetan Nayak et al

(1) The proposal relies on the existence of topological states of matter whose quasiparticle excitations are neither bosons nor fermions, **but (=) are** particles known as non-Abelian anyons, meaning that they obey non-Abelian braiding statistics.
(2) Quantum information is stored in states with multiple quasiparticles, **which (=) have** a topological degeneracy.
(3) The unitary gate operations **that are necessary** for quantum computation are carried out by **braiding (e&eb)** quasiparticles and then measuring the multiquasiparticle states.
(4) The fault tolerance of a topological quantum computer **arises (e) from** the nonlocal encoding of the quasiparticle states, which makes them immune to errors caused by local perturbations.
(5) To date, the only such topological states thought **to have been (eb) found in** nature are fractional quantum Hall states, most prominently the ν=5⁄2 state, although several other prospective candidates have been proposed in systems as disparate as ultracold atoms in optical lattices and thin-film superconductors.

(6) In this review article, current research in this field is described, focusing on the general theoretical concepts of non-Abelian statistics **as it relates to** topological quantum computation, on understanding non-Abelian quantum Hall states, on proposed experiments to detect non-Abelian anyons, and on proposed architectures for a topological quantum computer.

(7) Both the mathematical underpinnings of topological quantum computation **and (e&eb) the** physics of the subject are addressed, using the ν=5⁄2 fractional quantum Hall state as the archetype of a non-Abelian topological state enabling fault-tolerant quantum computation.URL: http://link.aps.org/doi/10.1103/RevModPhys.80.1083DOI: 10.1103/RevModPhys.80.1083PACS: 05.30.Pr, 03.67.Lx, 03.67.Pp, 73.43.−f

Atomic-Scale, All Epitaxial In-Plane Gated Donor Quantum Dot in Silicon Fuhrer et al Nano Lett., 2009, 9 (2), pp 707–710 DOI: 10.1021/nl803196f January 2, 2009 Copyright © 2009 American Chemical Society

(8) Doping profiles **can be (e) specified by the** product of n functions. n is the dimension of the simulation, i.e. n = 1, 2 or 3. Each function depends only on one coordinate.

(9) The doping profile is independent of the regions specified before. The function **is (e&eb) applied to** the region given by the specifier only-region. The doping concentration is at the position specified by the specifier position. The function is normalized such that the result is doping concentration at position. The impurity number specifies the kind of impurities used in the profile.

NOTATION

Module One

Is topological order stable at non-zero temperature? We assume the statement and give a model (Unsolved model: Both the possibilities are discussed). **Unsolved problem**

Topological order is **stable (e)** at non zero temperature. At non zero temperature **there exists (eb)** stable topological order**.**

G_{13} : Category one of At non zero temperature

G_{14} : Category two of At non zero temperature

G_{15} : Category three of At non zero temperature

T_{13} : Category one of stable topological order**.**

T_{14} : Category two of stable topological order**.**

T_{15} : Category three of stable topological order**.**

Module Two (Case two)

Topological order is **not (e) s**table at non zero temperature. At non zero temperature there **does not(e)** exist stable topological order

G_{16} : Category one of existence of stable topological order

G_{17} : Category two of existence of stable topological order

G_{18} : Category three of existence of stable topological order

T_{16} : Category one of non zero temperature

T_{17} : Category two of non zero temperature

T_{18} : Category three of non zero temperature

Module three

The proposal relies on the existence of topological states of matter whose quasiparticle excitations are neither bosons nor fermions, **but are** particles known as non-Abelian anyons, meaning that they obey non-Abelian braiding statistics.

G_{20} : Category one of particles known as non-Abelian anyons, meaning that they obey non-Abelian braiding statistics.

G_{21} : Category two of particles known as non-Abelian anyons, meaning that they obey non-Abelian braiding statistics.

G_{22} : Category three of particles known as non-Abelian anyons, meaning that they obey non-Abelian braiding statistics.

T_{20} : Category one of existence of topological states of matter whose quasiparticle excitations are neither bosons nor fermions,

T_{21} : Category two of existence of topological states of matter whose quasiparticle excitations are neither bosons nor fermions

T_{22} : Category three of existence of topological states of matter whose quasiparticle excitations are neither bosons nor fermions

Module four

Quantum information is stored in states with multiple quasiparticles, **which have** a topological degeneracy

G_{24} : Category one of topological degeneracy

G_{25} : Category two of topological degeneracy

G_{26} : Category three of topological degeneracy

T_{24} : Category one of Quantum information is stored in states with multiple quasiparticles

T_{25} : Category two of Quantum information is stored in states with multiple quasiparticles

T_{26} : Category three of Quantum information is stored in states with multiple quasiparticles

Module five

The unitary gate operations **that are necessary** for quantum computation are carried out by **braiding (e&eb)** quasiparticles and then measuring the multiquasiparticle states

G_{28} : Category one of unitary gate operations **that are necessary** for quantum computation are carried out; quasiparticles and then measuring the multiquasiparticle states

G_{29} : Category two of unitary gate operations **that are necessary** for quantum computation are carried out; quasiparticles and then measuring the multiquasiparticle states

G_{30} : Category three of unitary gate operations **that are necessary** for quantum computation are carried out; quasiparticles and then measuring the multiquasiparticle states

T_{28} : Category one of quasiparticles and then measuring the multiquasiparticle states ;unitary gate operations **that are necessary** for quantum computation are carried out

T_{29} : Category two of quasiparticles and then measuring the multiquasiparticle states; unitary gate operations **that are necessary** for quantum computation are carried out

T_{30} : Category three of quasiparticles and then measuring the multiquasiparticle states; unitary gate operations **that are necessary** for quantum computation are carried out

Module six

The fault tolerance of a topological quantum computer **arises (e) from** the nonlocal encoding of the quasiparticle states, which makes them immune to errors caused by local perturbations. In this review article, current research in this field is described, focusing on the general theoretical concepts of non-Abelian statistics **as it relates to** topological quantum computation, on understanding non-Abelian quantum Hall states, on proposed experiments to detect non-Abelian anyons, and on proposed architectures for a topological quantum computer.

G_{32} : Category one of nonlocal encoding of the quasiparticle states, which makes them immune to errors caused by local perturbations.

G_{33} : Category two of nonlocal encoding of the quasiparticle states, which makes them immune to errors caused by local perturbations.

G_{34} : Category three of nonlocal encoding of the quasiparticle states, which makes them immune to errors caused by local perturbations.

T_{32} : Category one of fault tolerance of a topological quantum computer;

T_{33} : Category two of fault tolerance of a topological quantum computer;

T_{34} : Category three of fault tolerance of a topological quantum computer;

Module seven

G_{36} : Category one of fractional quantum Hall states, most prominently the ν=5⁄2 state, although several other prospective candidates have been proposed in systems as disparate as ultracold atoms in optical lattices and thin-film superconductors

G_{37} : Category two of fractional quantum Hall states, most prominently the ν=5⁄2 state, although several other prospective candidates have been proposed in systems as disparate as ultracold atoms in optical lattices and thin-film superconductors

G_{38} : Category three of fractional quantum Hall states, most prominently the ν=5⁄2 state, although several other prospective candidates have been proposed in systems as disparate as ultracold atoms in optical lattices and thin-film superconductors

T_{36} : Category one of topological states thought **to have been (eb) found in** nature

T_{37} : Category two of topological states thought **to have been (eb) found in** nature

T_{38} : Category three of topological states thought **to have been (eb) found in** nature

Module eight

Both the mathematical underpinnings of topological quantum computation **and (e&eb) the** physics of the subject are addressed, **using(E)** the ν=5⁄2 fractional quantum Hall state as the archetype of a non-Abelian topological state enabling fault-tolerant quantum computation.URL: http://link.aps.org/doi/10.1103/RevModPhys.80.1083DOI:

G_{40} : Category one of ν=5⁄2 fractional quantum Hall state as the archetype of a non-Abelian topological state enabling fault-tolerant quantum computation

G_{41} : Category two of ν=5⁄2 fractional quantum Hall state as the archetype of a non-Abelian topological state enabling fault-tolerant quantum computation

G_{42} : Category three of ν=5⁄2 fractional quantum Hall state as the archetype of a non-Abelian topological state enabling fault-tolerant quantum computation

T_{40} : Category one of mathematical underpinnings of topological quantum computation **and (e&eb) the** physics of the subject are addressed

T_{41} : Category two of mathematical underpinnings of topological quantum computation **and (e&eb) the** physics of the subject are addressed

T_{42} : Category three of mathematical underpinnings of topological quantum computation **and (e&eb) the** physics of the subject are addressed

Module Nine

The doping profile is independent of the regions specified before. The function **is (e&eb) applied to** the region given by the specifier only-region. The doping concentration is at the position specified by the specifier position. The function is normalized such that the result is doping concentration at position. The impurity number specifies the kind of impurities used in the profile

G_{44} : Category one of doping profile is independent of the regions specified before. The function (Doping profiles **can be (e) specified by the** product of n functions. n is the dimension of the simulation, i.e. n = 1, 2 or 3. Each function depends only on one coordinate)

G_{45} : Category two of doping profile is independent of the regions specified before. The function (Doping profiles **can be (e) specified by the** product of n functions. n is the dimension of the simulation, i.e. n = 1, 2 or 3. Each function depends only on one coordinate)

G_{46} : Category three of doping profile is independent of the regions specified before. The function (Doping profiles **can be (e) specified by the** product of n functions. n is the dimension of the simulation, i.e. n = 1, 2 or 3. Each function depends only on one coordinate)

T_{44} : Category one of region given by the specifier only-region

T_{45} : Category two of region given by the specifier only-region

T_{46} : Category three of region given by the specifier only-region

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}),t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$
$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$
$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$
$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**
$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$
$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

59

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

60

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

61

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16}$$

62

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17}$$

63

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18}$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a_{44}'')^{(9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b_{16}')^{(2)} & -(b_{16}'')^{(2)}(G_{19},t) & -(b_{13}'')^{(1,1,)}(G,t) & -(b_{20}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{24}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{28}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{32}'')^{(6,6,6,6,6)}(G_{35},t) \\ -(b_{36}'')^{(7,7,7)}(G_{39},t) & -(b_{40}'')^{(8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b_{17}')^{(2)} & -(b_{17}'')^{(2)}(G_{19},t) & -(b_{14}'')^{(1,1,)}(G,t) & -(b_{21}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{25}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{29}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{33}'')^{(6,6,6,6,6)}(G_{35},t) \\ -(b_{37}'')^{(7,7,7)}(G_{39},t) & -(b_{41}'')^{(8,8,8)}(G_{43},t) & -(b_{45}'')^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b_{18}')^{(2)} & -(b_{18}'')^{(2)}(G_{19},t) & -(b_{15}'')^{(1,1,)}(G,t) & -(b_{22}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{26}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{30}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{34}'')^{(6,6,6,6,6)}(G_{35},t) \\ -(b_{38}'')^{(7,7,7)}(G_{39},t) & -(b_{42}'')^{(8,8,8)}(G_{43},t) & -(b_{46}'')^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad 66$$

where $-(b_{16}'')^{(2)}(G_{19},t)$, $-(b_{17}'')^{(2)}(G_{19},t)$, $-(b_{18}'')^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b_{13}'')^{(1,1,)}(G,t)$, $-(b_{14}'')^{(1,1,)}(G,t)$, $-(b_{15}'')^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b_{20}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b_{24}'')^{(4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b_{28}'')^{(5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b_{32}'')^{(6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt}=(a_{20})^{(3)}G_{21}-\left[\begin{array}{ccc}(a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}\\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}\end{array}\right]G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt}=(a_{21})^{(3)}G_{20}-\left[\begin{array}{ccc}(a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}\\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}\end{array}\right]G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt}=(a_{22})^{(3)}G_{21}-\left[\begin{array}{ccc}(a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}\\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}\end{array}\right]G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt}=(b_{20})^{(3)}T_{21}-\left[\begin{array}{ccc}(b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,1,)}(G,t)}\\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}\end{array}\right]T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1,2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1,2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1,2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category $1,2$ *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category $1,2$ *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$

are sixth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$

are seventh detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$

are eighth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt}=(a_{28})^{(5)}G_{29}-\left[\begin{array}{ccc}(a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt}=(a_{29})^{(5)}G_{28}-\left[\begin{array}{ccc}(a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt}=(a_{30})^{(5)}G_{29}-\left[\begin{array}{ccc}(a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt}=(b_{28})^{(5)}T_{29}-\left[\begin{array}{ccc}(b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}\end{array}\right]T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)} & +(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34} \quad (90)$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition

coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37},t) & +(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37},t) & +(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37},t) & +(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a''_{36})^{(7)}(T_{37},t)$, $(a''_{37})^{(7)}(T_{37},t)$, $(a''_{38})^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation

coefficient for 1,2,3

$\frac{dT_{36}}{dt} =$ 94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$\frac{dT_{37}}{dt} =$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$\frac{dT_{38}}{dt} =$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)} G_{44}$$
$$- \left[\begin{array}{c} (a'_{45})^{(9)} \boxed{+(a''_{45})^{(9)}(T_{45},t)} \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)} G_{45}$$
$$- \left[\begin{array}{c} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)} T_{45} -$$
$$\left[\begin{array}{c} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{13}$$
$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ \ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(OOOOOO) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(PPPPPP) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(QQQQQQ) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(RRRRRR)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(SSSSSS) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(TTTTTT) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and \ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ \ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{16})^{(2)}$, $T_i^0 \leq (\hat{Q}_{16})^{(2)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

By

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\tilde{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\tilde{M}_{20})^{(3)} t}$

By

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

159

160

161

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^{0} + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^{0} + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^{0} + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(y) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : \left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}ds_{(16)} +$

$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})($and\ not\ on\ \ t$) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+\ (a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right) d\left(\left((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(6)}$ and $(b''_i)^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(6)} - (a''_i)^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b'_i)^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t} \; and \; (\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2 \; and \; \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37} \; and \; G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad (333)$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad (334)$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad (335)$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad (336)$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}$, $\boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}$, $\boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t)\right) - (a_{41}'')^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(y) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FIFTY FOUR

Monolayer Doping (MLD): Victoria Concordia Crescit

INTRODUCTION—VARIABLES USED

Source: Wikipedia

(1) Monolayer doping (MLD) is a well controlled, wafer-scale surface doping technique first developed at the University of California, Berkeley, in 2007. This work is aimed **for (e) attaining** controlled doping of semiconductor materials with atomic accuracy, especially at nanoscale, which is not easily obtained by other existing technologies.

(2) This technique is currently **used for** fabricating ultrashallow junctions (USJs) as the heavily doped source/drain (S/D) contacts of metal-oxide-semiconductor field effect transistors (MOSFETs) as well as enabling dopant profiling of nanostructures.

(3) This MLD technique utilizes the crystalline nature of semiconductors and its self-limiting surface reaction properties **to form** highly uniform, self-assembled, covalently bonded dopant-containing monolayers followed by a subsequent annealing step for the incorporation and diffusion of dopants.

(4) The monolayer formation reaction is self-limiting, thereby, **resulting (eb) in the** deterministic coverage of dopant atoms on the surface. MLD differs from other conventional doping techniques such as spin-on-dopants (SODs) and gas phase doping techniques in the way of dopant dose control.

(5) Such control in MLD is much more precise **due to the** self-limiting formation of covalently attached dopants on the surface

(6) SODs just rely on the thickness control of the spin-on oxide and the gas phase technique **depends on** the control of dopant gas flow rate; therefore, the excellent dose control in MLD can yield the exact tuning of the resulting dopant profile.

(7) Compared to ion-implantation, MLD **does (e) not** involve the energetic introduction of dopant species into the semiconductor lattice where crystal damages are induced.

(8) In the case of implantation, defects such as interstitials and vacancies are inevitably generated, which interact with the dopants **to further (eb) broaden** the junction profile. This is known as the transient-enhanced diffusion (TED), which limits the formation of good quality of USJs.

(9) Also, stochastic variation in the dopant positioning and sever stoichiometric imbalance are **thus induced** for binary and tertiary compound semiconductors by the implantation techniques.

(10) In contrast, all MLD dopant atoms are thermally diffused from the crystal surface to the bulk and the dopant profile **can be easily controlled** by the thermal budget. Since the MLD system

can be classified as a limited source model, this is desirable for controlled USJ fabrication with high uniformity and low stochastic variation. Combined with the excellent dopant dose uniformity and coverage in MLD, it is especially attractive for doping nonplanar devices such as fin-FETs and nanowires. As a result, high quality sub-5 nm ultra-shallow junction has been demonstrated in silicon via the use of this MLD technique(Wikipedia)

NOTATION

Module One

Monolayer doping (MLD) is a well controlled, wafer-scale surface doping technique first developed at the University of California, Berkeley, in 2007. This work is aimed **for (eb) attaining** controlled doping of semiconductor materials with atomic accuracy, especially at nanoscale, which is not easily obtained by other existing technologies

G_{13} : Category one of Monolayer doping (MLD) is a well controlled, wafer-scale surface doping technique

G_{14} : Category two of Monolayer doping (MLD) is a well controlled, wafer-scale surface doping technique

G_{15} : Category three of Monolayer doping (MLD) is a well controlled, wafer-scale surface doping technique

T_{13} : Category one of controlled doping of semiconductor materials with atomic accuracy, especially at nanoscale, which is not easily obtained by other existing technologies

T_{14} : Category two of controlled doping of semiconductor materials with atomic accuracy, especially at nanoscale, which is not easily obtained by other existing technologies

T_{15} : Category three of controlled doping of semiconductor materials with atomic accuracy, especially at nanoscale, which is not easily obtained by other existing technologies

Module Two

This technique is currently **used for** fabricating ultrashallow junctions (USJs) as the heavily doped source/drain (S/D) contacts of metal-oxide-semiconductor field effect transistors (MOSFETs) as well as enabling dopant profiling of nanostructures

G_{16} : Category one of fabricating ultrashallow junctions (USJs) as the heavily doped source/drain (S/D) contacts of metal-oxide-semiconductor field effect transistors (MOSFETs) as well as enabling dopant profiling of nanostructures

G_{17} : Category two of fabricating ultrashallow junctions (USJs) as the heavily doped source/drain (S/D) contacts of metal-oxide-semiconductor field effect transistors (MOSFETs) as well as enabling dopant profiling of nanostructures

G_{18} : Category three of fabricating ultrashallow junctions (USJs) as the heavily doped source/drain (S/D) contacts of metal-oxide-semiconductor field effect transistors (MOSFETs) as well as enabling dopant profiling of nanostructures

T_{16} : Category one of Monolayer doping (MLD) is a well controlled, wafer-scale surface doping technique

T_{17} : Category two of Monolayer doping (MLD) is a well controlled, wafer-scale surface doping technique

T_{18} : Category three of Monolayer doping (MLD) is a well controlled, wafer-scale surface doping technique

Module three

This MLD technique utilizes the crystalline nature of semiconductors and its self-limiting surface reaction properties **to form** highly uniform, self-assembled, covalently bonded dopant-containing monolayers followed by a subsequent annealing step for the incorporation and diffusion of dopants

G_{20} : Category one of MLD technique utilizes the crystalline nature of semiconductors and its self-limiting surface reaction properties

G_{21} : Category two of MLD technique utilizes the crystalline nature of semiconductors and its self-limiting surface reaction properties

G_{22} : Category three of MLD technique utilizes the crystalline nature of semiconductors and its self-limiting surface reaction properties

T_{20} : Category one of highly uniform, self-assembled, covalently bonded dopant-containing monolayers followed by a subsequent annealing step for the incorporation and diffusion of dopants

T_{21} : Category two of highly uniform, self-assembled, covalently bonded dopant-containing monolayers followed by a subsequent annealing step for the incorporation and diffusion of dopants

T_{22} : Category three of highly uniform, self-assembled, covalently bonded dopant-containing monolayers followed by a subsequent annealing step for the incorporation and diffusion of dopants

Module four

The monolayer formation reaction is self-limiting, thereby, **resulting (eb) in the** deterministic coverage of dopant atoms on the surface. MLD differs from other conventional doping techniques such as spin-on-dopants (SODs) and gas phase doping techniques in the way of dopant dose control.

G_{24} : Category one of monolayer formation reaction is self-limiting

G_{25} : Category two of monolayer formation reaction is self-limiting

G_{26} : Category three of monolayer formation reaction is self-limiting

T_{24} : Category one of deterministic coverage of dopant atoms on the surface. MLD differs from other conventional doping techniques such as spin-on-dopants (SODs) and gas phase doping techniques in the way of dopant dose control.

T_{25} : Category two of deterministic coverage of dopant atoms on the surface. MLD differs from other conventional doping techniques such as spin-on-dopants (SODs) and gas phase doping techniques in the way of dopant dose control.

T_{26} : Category three of deterministic coverage of dopant atoms on the surface. MLD differs from other conventional doping techniques such as spin-on-dopants (SODs) and gas phase doping techniques in the way of dopant dose control.

Module five

Such control in MLD is much more precise **due to the** self-limiting formation of covalently attached dopants on the surface

G_{28} : Category one of self-limiting formation of covalently attached dopants on the surface

G_{29} : Category two of self-limiting formation of covalently attached dopants on the surface

G_{30} : Category three of self-limiting formation of covalently attached dopants on the surface

T_{28} : Category one of Such control in MLD is much more precise

T_{29} : Category two of Such control in MLD is much more precise

T_{30} : Category three of Such control in MLD is much more precise

Module six

SODs just rely on the thickness control of the spin-on oxide and the gas phase technique **depends on** the control of dopant gas flow rate; therefore, the excellent dose control in MLD can yield the exact tuning of the resulting dopant profile.

G_{32} : Category one of constant(control of dopant gas flow rate; therefore, the excellent dose control in MLD can yield the exact tuning of the resulting dopant profile)

G_{33} : Category two of constant(control of dopant gas flow rate; therefore, the excellent dose control in MLD can yield the exact tuning of the resulting dopant profile)

G_{34} : Category three of constant(control of dopant gas flow rate; therefore, the excellent dose control in MLD can yield the exact tuning of the resulting dopant profile)

T_{32} : Category one of SODs just rely on the thickness control of the spin-on oxide and the gas phase technique

T_{33} : Category two of SODs just rely on the thickness control of the spin-on oxide and the gas phase technique

T_{34} : Category three of SODs just rely on the thickness control of the spin-on oxide and the gas phase technique

Module seven

Compared to ion-implantation, MLD **does (e) not** involve the energetic introduction of dopant species into the semiconductor lattice where crystal damages are induced.

G_{36} : Category one of energetic introduction of dopant species into the semiconductor lattice where crystal damages are induced.

G_{37} : Category two of energetic introduction of dopant species into the semiconductor lattice where crystal damages are induced.

G_{38} : Category three of energetic introduction of dopant species into the semiconductor lattice where crystal damages are induced.

T_{36} : Category one of Compared to ion-implantation, MLD

T_{37} : Category two of Compared to ion-implantation, MLD

T_{38} : Category three of Compared to ion-implantation, MLD

Module eight

In the case of implantation, defects such as interstitials and vacancies are inevitably generated, which interact with the dopants **to further (eb) broaden** the junction profile. This is known as the transient-enhanced diffusion (TED), which limits the formation of good quality of USJs.

G_{40} : Category one of case of implantation, defects such as interstitials and vacancies are inevitably generated, which interact with the dopants

G_{41} : Category two of case of implantation, defects such as interstitials and vacancies are inevitably generated, which interact with the dopants

G_{42} : Category three of case of implantation, defects such as interstitials and vacancies are inevitably generated, which interact with the dopants

T_{40} : Category one of junction profile. This is known as the transient-enhanced diffusion (TED), which limits the formation of good quality of USJs.

T_{41} : Category two of junction profile. This is known as the transient-enhanced diffusion (TED), which limits the formation of good quality of USJs.

T_{42} : Category three of junction profile. This is known as the transient-enhanced diffusion (TED), which limits the formation of good quality of USJs.

Module Nine

stochastic variation in the dopant positioning and sever stoichiometric imbalance are **thus induced** for binary and tertiary compound semiconductors by the implantation techniques

G_{44} : Category one of stochastic variation in the dopant positioning and sever stoichiometric imbalance

G_{45} : Category two of stochastic variation in the dopant positioning and sever stoichiometric imbalance

G_{46} : Category three of stochastic variation in the dopant positioning and sever stoichiometric imbalance

T_{44} : Category one of binary and tertiary compound semiconductors by the implantation techniques

T_{45} : Category two of binary and tertiary compound semiconductors by the implantation techniques

T_{46} : Category three of binary and tertiary compound semiconductors by the implantation techniques

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$

$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)} \ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) \quad +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{ccc} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) \quad +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right]G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) \quad +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right]G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\left[-(b''_{13})^{(1)}(G,t)\right] & \left[-(b''_{16})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\left[-(b''_{14})^{(1)}(G,t)\right] & \left[-(b''_{17})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\left[-(b''_{15})^{(1)}(G,t)\right] & \left[-(b''_{18})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & \boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} \quad \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{array}\right] G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & \boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} \quad \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{array}\right] G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & \boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} \quad \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{array}\right] G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition*

coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\left[\begin{array}{ccc}(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t)\\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t)\\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t)\end{array}\right]G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\left[\begin{array}{ccc}(a'_{25})^{(4)}+(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t)\\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t)\\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t)\end{array}\right]G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\left[\begin{array}{ccc}(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t)\\ +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t)\\ +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t)\end{array}\right]G_{26} \quad (75)$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* $1,2\ 3$

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* $1,2$ *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* $1,2$ *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category $1,2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
are eighth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad (81)$$

$Where\ \boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$And\ \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}\ are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\ are\ third\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} & \boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} \\ \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} & \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} \\ \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} \\ \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} & & \end{array}\right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} & \boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} \\ \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} & \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} \\ \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} \\ \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} & & \end{array}\right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} & \boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} \\ \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} & \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} \\ \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} \\ \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} & & \end{array}\right] T_{30} \quad 84$$

$where\ \boxed{-(b''_{28})^{(5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5)}(G_{31},t)}\ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients

for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[\begin{matrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)}\ \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}\ \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[\begin{matrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)}\ \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\ \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[\begin{matrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\ \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\ \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{34} \quad (87)$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad (90)$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad (91)$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)}\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad (92)$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)}\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$\frac{dG_{38}}{dt}$ *93*

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{36}}{dt} =$ 94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$\frac{dT_{37}}{dt} =$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$\frac{dT_{38}}{dt} =$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third

augmentation coefficient for category 1, 2 and 3
$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3
$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$,$+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$,$+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$,$+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$,$+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\left[-(b''_{40})^{(8)}(G_{43},t)\right] & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\left[-(b''_{41})^{(8)}(G_{43},t)\right] & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\left[-(b''_{42})^{(8)}(G_{43},t)\right] & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3
$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3
$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients

for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{44})^{(9)} & -(b''_{44})^{(9)}(G_{47},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t) \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)} & -(b''_{45})^{(9)}(G_{47},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t) \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)} & -(b''_{46})^{(9)}(G_{47},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t) \end{bmatrix} T_{15}$$

Where $-(b''_{44})^{(9)}(G_{47},t)$, $-(b''_{45})^{(9)}(G_{47},t)$, $-(b''_{46})^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15$ *97*

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$(a_i'')^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$

$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$

$lim_{T_2 \to \infty}(a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)}$ 98

$\lim_{G \to \infty}(b_i'')^{(1)}(G, t) = (r_i)^{(1)}$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$

$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$

$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}\ (T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}\ (G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G \to \infty}(b_i'')^{(4)}\big((G_{27}), t\big) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$

Where we suppose

(UUUUUU) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(VVVVVV) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$

132

(WWWWWW) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(XXXXXX)

$\lim_{G \to \infty} (b_i'')^{(7)}\big((G_{39}), t\big) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\big((G_{39}), t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(YYYYYY) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(ZZZZZZ) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants

$(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \tag{136}$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: (137)

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \tag{138}$$

$$(b_i'')^{(8)}((G_{43}), t) \leq \ (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \tag{139}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)} (T_{41}, t) = (p_i)^{(8)} \tag{140}$$

$$\lim_{G \to \infty} (b_i'')^{(8)} \left((G_{43}), t\right) = \ (r_i)^{(8)} \tag{141}$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \tag{142}$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}\left((G_{43}), t\right)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \tag{143}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}} \ , \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \tag{144}$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}, i = 40,41,42,$

Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \qquad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, \ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq \ (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = \ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , $T_i : \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{13})^{(1)}$, $T_i^0 \leq (\hat{Q}_{13})^{(1)}$, 155

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 156

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)},\ T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{24})^{(4)}$, $T_i^0 \leq (\hat{Q}_{24})^{(4)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166 A

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)}\left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)}(\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)}\left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)}\left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)}\left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(z) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \tag{180}$$

$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right] \tag{181}$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose $\tag{182}$

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \tag{183}$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \le (\hat{Q}_{13})^{(1)} \tag{184}$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric $\tag{185}$

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\le \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\,G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)},i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 19 to 24 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\le\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t),t)) = (b'_{14})^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b''_{15})^{(1)}\,(G(t),t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\le$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t)=0$ and $T_i(t)=0$ 200

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(2)}-(a_i'')^{(2)}(T_{17}(s_{(16)}),s_{(16)})\}ds_{(16)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(2)}t)}>0\quad\text{for } t>0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1,\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16}<(\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt}\le\left((\widehat{M}_{16})^{(2)}\right)_1-(a'_{17})^{(2)}G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a'_{17})^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a'_{18})^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t),t)) = (b'_{17})^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t),t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t),t\right) = (b'_{18})^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)} + \left((\widehat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|{G_{23}}^{(1)}-{G_{23}}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\le \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ and\ (\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)},i=20{,}21{,}22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 216

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(3)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1,\left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If $G_{20}\ is$ bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t),t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t = log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b''_{22})^{(3)}\left((G_{23})(t),t\right) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)} + \left((\hat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{24})^{(4)}\right] \leq (\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\le \qquad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}\ and\ (\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 228

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}),s_{(24)})\}ds_{(24)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(4)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1,\left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b'_{25})^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)} + \left((\widehat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\widehat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{28})^{(5)}\right] \leq (\widehat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})},\widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})},\widetilde{(T_{31})}\right)=\mathcal{A}^{(5)}((G_{31}),(T_{31}))$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\le \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}\ and\ (\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ 239

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(5)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1,\left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)}G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a'_{29})^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\ ((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right),\quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\le \quad 248$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}\ and\ (\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i=32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ 250

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(6)}-(a_i'')^{(6)}(T_{33}(s_{(32)}),s_{(32)})\}ds_{(32)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(6)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1,\left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)}G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1/(a'_{33})^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right),\ \ t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i , T_i satisfying

Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:(\widetilde{(G_{39})},\widetilde{(T_{39})})=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\le\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\le \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)})d\left(((G_{39})^{(1)},(T_{39})^{(1)};(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i=36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ 261

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(7)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

268

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{40})^{(8)}\right]\leq(\hat{Q}_{40})^{(8)}$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval [0, t] 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)} t)} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_1 / (a_{41}')^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_2 / (a_{42}')^{(8)}$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[(\hat{P}_{44})^{(9)} + \left((\hat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{44})^{(9)}\right]\leq(\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 99 to 44 it results

$$G_i\,(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\geq 0$$

$$T_i\,(t)\geq T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2$ *and* $\left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} *and* G_{46} . indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a'_{45})^{(9)} G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)},,(\bar{v}_2)^{(1)},(\bar{u}_1)^{(1)},(\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}$, $(m_2)^{(1)}$, $(\mu_1)^{(1)},(\mu_2)^{(1)},(\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}$, $(m_2)^{(1)}$, $(\mu_1)^{(1)},(\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}\ , if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(\ m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}\ , if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\ \mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t}\right] + G_{15}^0 e^{-(S_2)^{(1)}t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- 290

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying
$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(\mathrm{T}_{17},t)+(a''_{17})^{(2)}(\mathrm{T}_{17},t)\le-(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: 295

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2+(\tau_2)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ 302

Definition of $(m_1)^{(2)},(m_2)^{(2)},(\mu_1)^{(2)},(\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)},(m_2)^{(2)},(\mu_1)^{(2)},(\mu_2)^{(2)}$ by 304

$(m_2)^{(2)}=(\nu_0)^{(2)},(m_1)^{(2)}=(\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)}<(\nu_1)^{(2)}$ 305

$(m_2)^{(2)}=(\nu_1)^{(2)},(m_1)^{(2)}=(\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)}<(\nu_0)^{(2)}<(\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)}=\frac{\mathrm{G}_{16}^{0}}{\mathrm{G}_{17}^{0}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)},\ \boldsymbol{if}\ (u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \boldsymbol{if}\ (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G^0_{20}e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{20}(t) \le G^0_{20}e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}}G^0_{20}e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}}G^0_{20}e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)}G_{20}^{0}}{(m_1)^{(3)}((S_1)^{(3)}-(p_{20})^{(3)}-(S_2)^{(3)})}\left[e^{((S_1)^{(3)}-(p_{20})^{(3)})t}-e^{-(S_2)^{(3)}t}\right]+G_{22}^{0}e^{-(S_2)^{(3)}t}\le G_{22}(t)\le\right. \quad 324$$

$$\left.\frac{(a_{22})^{(3)}G_{20}^{0}}{(m_2)^{(3)}((S_1)^{(3)}-(a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t}-e^{-(a'_{22})^{(3)}t}\right]+G_{22}^{0}e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^{0}e^{(R_1)^{(3)}t}\le T_{20}(t)\le T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^{0}e^{(R_1)^{(3)}t}\le T_{20}(t)\le\frac{1}{(\mu_2)^{(3)}}T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^{0}}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t}-e^{-(b'_{22})^{(3)}t}\right]+T_{22}^{0}e^{-(b'_{22})^{(3)}t}\le T_{22}(t)\le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^{0}}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t}-e^{-(R_2)^{(3)}t}\right]+T_{22}^{0}e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)},(S_2)^{(3)},(R_1)^{(3)},(R_2)^{(3)}$:- 328

Where $(S_1)^{(3)}=(a_{20})^{(3)}(m_2)^{(3)}-(a'_{20})^{(3)}$

$(S_2)^{(3)}=(a_{22})^{(3)}-(p_{22})^{(3)}$

$(R_1)^{(3)}=(b_{20})^{(3)}(\mu_2)^{(3)}-(b'_{20})^{(3)}$

$(R_2)^{(3)}=(b'_{22})^{(3)}-(r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)},(\sigma_2)^{(4)},(\tau_1)^{(4)},(\tau_2)^{(4)}$:

$(\sigma_1)^{(4)},(\sigma_2)^{(4)},(\tau_1)^{(4)},(\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)}\le-(a'_{24})^{(4)}+(a'_{25})^{(4)}-(a''_{24})^{(4)}(T_{25},t)+(a''_{25})^{(4)}(T_{25},t)\le-(\sigma_1)^{(4)}$

$-(\tau_2)^{(4)}\le-(b'_{24})^{(4)}+(b'_{25})^{(4)}-(b''_{24})^{(4)}\big((G_{27}),t\big)-(b''_{25})^{(4)}\big((G_{27}),t\big)\le-(\tau_1)^{(4)}$

Definition of $(\nu_1)^{(4)},(\nu_2)^{(4)},(u_1)^{(4)},(u_2)^{(4)},\nu^{(4)},u^{(4)}$: 329

By $(\nu_1)^{(4)}>0$, $(\nu_2)^{(4)}<0$ and respectively $(u_1)^{(4)}>0$, $(u_2)^{(4)}<0$ the roots of the equations $(a_{25})^{(4)}\left(\nu^{(4)}\right)^2+(\sigma_1)^{(4)}\nu^{(4)}-(a_{24})^{(4)}=0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2+(\tau_1)^{(4)}u^{(4)}-(b_{24})^{(4)}=0$ and

Definition of $(\bar{\nu}_1)^{(4)},,(\bar{\nu}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)}>0$, $(\bar{\nu}_2)^{(4)}<0$ and respectively $(\bar{u}_1)^{(4)}>0$, $(\bar{u}_2)^{(4)}<0$ the roots of the equations $(a_{25})^{(4)}\left(\nu^{(4)}\right)^2+(\sigma_2)^{(4)}\nu^{(4)}-(a_{24})^{(4)}=0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2+(\tau_2)^{(4)}u^{(4)}-(b_{24})^{(4)}=0$

Definition of $(m_1)^{(4)},(m_2)^{(4)},(\mu_1)^{(4)},(\mu_2)^{(4)},(\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)},(m_2)^{(4)},(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)},\ \boldsymbol{if}\ (\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}\ , \boldsymbol{if}\ (\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)},$

and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(\ m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}\ , \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\ \mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)}, , (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{\left((S_1)^{(5)} - (p_{28})^{(5)}\right)t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)}-(p_{28})^{(5)}-(S_2)^{(5)})}\left[e^{((S_1)^{(5)}-(p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, , (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}$, ***if*** $(v_0)^{(6)} < (v_1)^{(6)}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}$, ***if*** $(v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)}$,
and $\boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}$, ***if*** $(\bar{v}_1)^{(6)} < (v_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$
$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0$, $(\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0$, $(u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0$, $(\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0$, $(\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0$, $(\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0$, $(\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}$, ***if*** $(\nu_0)^{(8)} < (\nu_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, ***if*** $(\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,
and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, ***if*** $(\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, ***if*** $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, ***if*** $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T^0_{40}}{T^0_{41}}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}$, ***if*** $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G^0_{40} e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} \le G_{40}(t) \le G^0_{40} e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)}-(p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)}-(p_{40})^{(8)}-(S_2)^{(8)})}\left[e^{((S_1)^{(8)}-(p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}\left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t}\right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$ and $(b_{45})^{(9)}\left(u^{(9)}\right)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}$, $(\mu_2)^{(9)}$, $(v_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}$, $(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}$, $\boldsymbol{if}\ (v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}$, $\boldsymbol{if}\ (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)}$, and $\boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, $\boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, $\boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$, and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{\left((S_1)^{(9)} - (p_{44})^{(9)}\right)t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{\left((S_1)^{(9)} - (p_{44})^{(9)}\right)t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}\left((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)}\right)}\left[e^{\left((S_1)^{(9)} - (p_{44})^{(9)}\right)t} - e^{-(S_2)^{(9)}t}\right] + G_{46}^0 e^{-(S_2)^{(9)}t} \le G_{46}(t) \le \right.$$

$$\left.\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}\left((S_1)^{(9)} - (a'_{46})^{(9)}\right)}\left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}\right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}\right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{\left((R_1)^{(9)} + (r_{44})^{(9)}\right)t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{\left((R_1)^{(9)} + (r_{44})^{(9)}\right)t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}\left((R_1)^{(9)} - (b'_{46})^{(9)}\right)}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}\left((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)}\right)}\left[e^{\left((R_1)^{(9)} + (r_{44})^{(9)}\right)t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \leq \frac{dv^{(1)}}{dt} \leq -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \leq (\bar{v}_1)^{(1)}$$

386

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{d\nu^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)}+(\mathrm{C})^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}{1+(\mathrm{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{\mathrm{C}})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{\mathrm{C}})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{\mathrm{C}})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{\mathrm{G}_{16}^0}{\mathrm{G}_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(\mathrm{C})^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(\mathrm{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{\mathrm{C}})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{\mathrm{C}})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{\mathrm{G}_{16}^0}{\mathrm{G}_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{\mathrm{C}})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{\mathrm{C}})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right) - (a''_{25})^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\quad \boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_0)^{(4)}\right)t\right]}}{4+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)}-(\nu_0)^{(4)}}{(\nu_0)^{(4)}-(\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{4+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)}-(\nu_0)^{(4)}}{(\nu_0)^{(4)}-(\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \le \nu^{(4)}(t) \le (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \le u^{(4)}(t) \le (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{28})^{(5)} = (a''_{29})^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

415

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{32}'')^{(6)} = (a_{33}'')^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b_{32}'')^{(6)} = (b_{33}'')^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a_{36}')^{(7)} - (a_{37}')^{(7)} + (a_{36}'')^{(7)}(T_{37}, t)\right) - (a_{37}'')^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\quad \boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition

$(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $v^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\quad \boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)}\right) \leq \frac{dv^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$v^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq v^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \le \nu^{(9)}(t) \le (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0$ 475

$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0$ 476

$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0$ 477

$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0$ 478

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(z) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T_{14}^{*} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^{*} for which $f(T_{14}^{*}) = 0$. With this value , we obtain from the three first

equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that

there exists a unique T_{37}^{*} for which $f(T_{37}^{*})=0$. With this value , we obtain from the three first equations

$$G_{36}=\frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}=\frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0)<0, f(\infty)>0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*})=0$. With this value , we obtain from the three first equations

$$G_{40}=\frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}=\frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0)<0, f(\infty)>0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*})=0$. With this value , we obtain from the three first equations

$$G_{44}=\frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46}=\frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G)=(b'_{13})^{(1)}(b'_{14})^{(1)}-(b_{13})^{(1)}(b_{14})^{(1)}-$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G)+(b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right]+(b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G)=0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0)>0$, $\varphi(\infty)<0$ it follows that there exists a unique G_{14}^{*} such that $\varphi(G^{*})=0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19})=(b'_{16})^{(2)}(b'_{17})^{(2)}-(b_{16})^{(2)}(b_{17})^{(2)}-$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19})+(b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19})=0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0)>0$, $\varphi(\infty)<0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*})=0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23})=(b'_{20})^{(3)}(b'_{21})^{(3)}-(b_{20})^{(3)}(b_{21})^{(3)}-$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{21} such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$$

$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^*)]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G^*_{17} given by $\varphi((G_{19})^*) = 0$, T^*_{17} given by $f(T^*_{17}) = 0$ and 512

$$G^*_{16} = \frac{(a_{16})^{(2)}G^*_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G^*_{18} = \frac{(a_{18})^{(2)}G^*_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \quad 513$$

$$T^*_{16} = \frac{(b_{16})^{(2)}T^*_{17}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T^*_{18} = \frac{(b_{18})^{(2)}T^*_{17}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G^*_{21} given by $\varphi((G_{23})^*) = 0$, T^*_{21} given by $f(T^*_{21}) = 0$ and

$$G^*_{20} = \frac{(a_{20})^{(3)}G^*_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G^*_{22} = \frac{(a_{22})^{(3)}G^*_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

$$T^*_{20} = \frac{(b_{20})^{(3)}T^*_{21}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T^*_{22} = \frac{(b_{22})^{(3)}T^*_{21}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G^*_{25} given by $\varphi(G_{27}) = 0$, T^*_{25} given by $f(T^*_{25}) = 0$ and

$$G^*_{24} = \frac{(a_{24})^{(4)}G^*_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G^*_{26} = \frac{(a_{26})^{(4)}G^*_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

$T_{24}^{*} = \frac{(b_{24})^{(4)}T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]}$, $T_{26}^{*} = \frac{(b_{26})^{(4)}T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]}$ 517

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$G_{28}^{*} = \frac{(a_{28})^{(5)}G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]}$, $G_{30}^{*} = \frac{(a_{30})^{(5)}G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and A

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)} T_{13}^* \mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)} T_{14}^* \mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)} T_{15}^* \mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$\frac{\partial(a_{17}'')^{(2)}}{\partial \mathrm{T}_{17}}(\mathrm{T}_{17}^{*}) = (q_{17})^{(2)}$, $\frac{\partial(b_i'')^{(2)}}{\partial \mathrm{G}_j}(\,(G_{19})^{*}\,) = s_{ij}$ 533

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{\mathrm{d}\mathbb{G}_{16}}{\mathrm{dt}} = -\big((a_{16}')^{(2)} + (p_{16})^{(2)}\big)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}\mathrm{G}_{16}^{*}\mathbb{T}_{17} \quad 534$$

$$\frac{\mathrm{d}\mathbb{G}_{17}}{\mathrm{dt}} = -\big((a_{17}')^{(2)} + (p_{17})^{(2)}\big)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}\mathrm{G}_{17}^{*}\mathbb{T}_{17} \quad 535$$

$$\frac{\mathrm{d}\mathbb{G}_{18}}{\mathrm{dt}} = -\big((a_{18}')^{(2)} + (p_{18})^{(2)}\big)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}\mathrm{G}_{18}^{*}\mathbb{T}_{17} \quad 536$$

$$\frac{\mathrm{d}\mathbb{T}_{16}}{\mathrm{dt}} = -\big((b_{16}')^{(2)} - (r_{16})^{(2)}\big)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \textstyle\sum_{j=16}^{18}\big(s_{(16)(j)}\mathrm{T}_{16}^{*}\mathbb{G}_j\big) \quad 537$$

$$\frac{\mathrm{d}\mathbb{T}_{17}}{\mathrm{dt}} = -\big((b_{17}')^{(2)} - (r_{17})^{(2)}\big)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \textstyle\sum_{j=16}^{18}\big(s_{(17)(j)}\mathrm{T}_{17}^{*}\mathbb{G}_j\big) \quad 538$$

$$\frac{\mathrm{d}\mathbb{T}_{18}}{\mathrm{dt}} = -\big((b_{18}')^{(2)} - (r_{18})^{(2)}\big)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \textstyle\sum_{j=16}^{18}\big(s_{(18)(j)}\mathrm{T}_{18}^{*}\mathbb{G}_j\big) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}\ and\ (b_i'')^{(3)}$ Belong to $C^{(3)}(\,\mathbb{R}_{+})$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^{*}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}(\,(G_{23})^{*}\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\big((a_{20}')^{(3)} + (p_{20})^{(3)}\big)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^{*}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\big((a_{21}')^{(3)} + (p_{21})^{(3)}\big)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^{*}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\big((a_{22}')^{(3)} + (p_{22})^{(3)}\big)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^{*}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\big((b_{20}')^{(3)} - (r_{20})^{(3)}\big)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \textstyle\sum_{j=20}^{22}\big(s_{(20)(j)}T_{20}^{*}\mathbb{G}_j\big) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\big((b_{21}')^{(3)} - (r_{21})^{(3)}\big)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \textstyle\sum_{j=20}^{22}\big(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\big) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\big((b_{22}')^{(3)} - (r_{22})^{(3)}\big)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \textstyle\sum_{j=20}^{22}\big(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\big) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \ , \ \frac{\partial (b_i'')^{(4)}}{\partial G_j}((G_{27})^* \ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\big((a_{24}')^{(4)} + (p_{24})^{(4)}\big)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^*\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\big((a_{25}')^{(4)} + (p_{25})^{(4)}\big)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\big((a_{26}')^{(4)} + (p_{26})^{(4)}\big)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\big((b_{24}')^{(4)} - (r_{24})^{(4)}\big)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\big(s_{(24)(j)}T_{24}^*\mathbb{G}_j\big) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\big((b_{25}')^{(4)} - (r_{25})^{(4)}\big)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\big(s_{(25)(j)}T_{25}^*\mathbb{G}_j\big) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\big((b_{26}')^{(4)} - (r_{26})^{(4)}\big)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\big(s_{(26)(j)}T_{26}^*\mathbb{G}_j\big) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \ , \ \frac{\partial (b_i'')^{(5)}}{\partial G_j}(\ (G_{31})^* \ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\big((a_{28}')^{(5)} + (p_{28})^{(5)}\big)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\big((a_{29}')^{(5)} + (p_{29})^{(5)}\big)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\big((a_{30}')^{(5)} + (p_{30})^{(5)}\big)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\big((b_{28}')^{(5)} - (r_{28})^{(5)}\big)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\big(s_{(28)(j)}T_{28}^*\mathbb{G}_j\big) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{37})^{(7)}}{\partial T_{37}}(T^*_{37}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a'_{36})^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G^*_{36}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a'_{37})^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G^*_{37}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a'_{38})^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G^*_{38}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b'_{36})^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T^*_{36}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b'_{37})^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T^*_{37}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b'_{38})^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T^*_{38}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{41})^{(8)}}{\partial T_{41}}(T^*_{41}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a'_{40})^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G^*_{40}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a'_{41})^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G^*_{41}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a'_{42})^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G^*_{42}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b'_{40})^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T^*_{40}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586

ASYMPTOTIC STABILITY ANALYSIS A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}(\,(G_{47})^*\,) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\big((a_{44}')^{(9)} + (p_{44})^{(9)}\big)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \qquad 586\,B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\big((a_{45}')^{(9)} + (p_{45})^{(9)}\big)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \qquad 586\,C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\big((a_{46}')^{(9)} + (p_{46})^{(9)}\big)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \qquad 586\,D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\big((b_{44}')^{(9)} - (r_{44})^{(9)}\big)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\big(s_{(44)(j)}T_{44}^*\mathbb{G}_j\big) \qquad 586\,E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\big((b_{45}')^{(9)} - (r_{45})^{(9)}\big)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\big(s_{(45)(j)}T_{45}^*\mathbb{G}_j\big) \qquad 586\,F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\big((b_{46}')^{(9)} - (r_{46})^{(9)}\big)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\big(s_{(46)(j)}T_{46}^*\mathbb{G}_j\big) \qquad 586\,G$$

The characteristic equation of this system is 587

$$\big((\lambda)^{(1)} + (b_{15}')^{(1)} - (r_{15})^{(1)}\big)\{\big((\lambda)^{(1)} + (a_{15}')^{(1)} + (p_{15})^{(1)}\big)$$

$$\Big[\big((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}\big)(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^*\big)\Big]$$

$$\big((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\big)s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^*\big)$$

$$+\big((\lambda)^{(1)} + (a_{14}')^{(1)} + (p_{14})^{(1)}\big)(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^*\big)$$

$$\big((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\big)s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^*\big)$$

$$\big((\lambda)^{(1)}\big)^2 + \big(\,(a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\big)\,(\lambda)^{(1)}\big)$$

$$\big((\lambda)^{(1)}\big)^2 + \big(\,(b_{13}')^{(1)} + (b_{14}')^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\big)\,(\lambda)^{(1)}\big)$$

$$+\Big(\big((\lambda)^{(1)}\big)^2 + \big(\,(a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\big)\,(\lambda)^{(1)}\Big)(q_{15})^{(1)}G_{15}$$

$$+\big((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}\big)\,\big((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*\big)$$

$$\big((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\big)s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^*\big)\} = 0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)})\{((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)})$$

$$\left[\left(((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)})(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\right]$$

$$\left(((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)})s_{(17),(17)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\right)$$

$$+\left(((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)})(q_{16})^{(2)}\mathrm{G}^{*}_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\right)$$

$$\left(((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)})s_{(17),(16)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\right)$$

$$\left(((\lambda)^{(2)})^{2}+(\,(a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)})\,(\lambda)^{(2)}\right)$$

$$\left(((\lambda)^{(2)})^{2}+(\,(b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)})\,(\lambda)^{(2)}\right)$$

$$+\left(((\lambda)^{(2)})^{2}+(\,(a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)})\,(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left(((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)})s_{(17),(18)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)})\{((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)})$$

$$\left[\left(((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)})(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\right]$$

$$\left(((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)})s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\right)$$

$$+\left(((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)})(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left(((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)})s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\right)$$

$$\left(((\lambda)^{(3)})^{2}+(\,(a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)})\,(\lambda)^{(3)}\right)$$

$$\left(((\lambda)^{(3)})^{2}+(\,(b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)})\,(\lambda)^{(3)}\right)$$

$$+\left(((\lambda)^{(3)})^{2}+(\,(a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)})\,(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)})s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\right)\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\right)\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\right)\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\right)\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^{2}+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(38)}T^{*}_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^{2}+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(46)}T^{*}_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FIFTY FIVE

Quantum Geometry And Black Hole Entropy: Aeì Ho Theòs Ho Mégas Geōmetreî Tò Sýmpan

INTRODUCTION—VARIABLES USED

Quantum Geometry of Isolated Horizons and Black Hole Entropy A. Ashtekar, J. Baez, K. Krasnov

(1) Using the earlier developed classical Hamiltonian framework as the point of departure, authors carry out a non-perturbative quantization of the sector of general relativity, coupled to matter, **admitting (e)** non-rotating isolated horizons as inner boundaries.
(2) The emphasis is on the quantum geometry of the horizon. Polymer excitations of the bulk quantum geometry pierce the horizon **endowing (eb)** it with area.
(3) The intrinsic geometry of the horizon **is then (e) described by** the quantum Chern-Simons theory of a U (1) connection on a punctured 2-sphere, the horizon.
(4) Subtle mathematical features of the quantum Chern-Simons theory **turn out to (e) are important** for the existence of a coherent quantum theory of the horizon geometry.
(5) Heuristically, the intrinsic geometry is flat everywhere **except (e: flatness eaten at) a**t the punctures.
(6) The distributional curvature of the U (1) connection at the punctures **gives rise to** quantized deficit angles which account for the overall curvature.
(7) For macroscopic black holes, the logarithm of the number of these horizon microstates **is proportional** to the area, irrespective of the values of (non-gravitational) charges.
(8) Thus, the black hole entropy **can be (e) accounted** for entirely by the quantum states of the horizon geometry.
(9) Our analysis is applicable to all non-rotating black holes**, including the** astrophysically interesting ones which are very far from extremality.

Furthermore, cosmological horizons (to which statistical mechanical considerations are known to apply) are naturally incorporated. An effort has been made to make the paper self-contained by including short reviews

of the background material. Adv.Theor.Math.Phys. 4 (2000) 1-94Report number: NSF-ITP-99-153: arXiv: gr-qc/0005126

NOTATION

Module One

Using the earlier developed classical Hamiltonian framework as the point of departure, authors carry out a non-perturbative quantization of the sector of general relativity, coupled to matter, **admitting (e)** non-rotating isolated horizons as inner boundaries

G_{13} : Category one of non-rotating isolated horizons as inner boundaries

G_{14} : Category two of non-rotating isolated horizons as inner boundaries

G_{15} : Category three ofnon-rotating isolated horizons as inner boundaries

T_{13} : Category one of classical Hamiltonian framework as the point of departure, authors carry out a non-perturbative quantization of the sector of general relativity, coupled to matter

T_{14} : Category two of classical Hamiltonian framework as the point of departure, authors carry out a non-perturbative quantization of the sector of general relativity, coupled to matter

T_{15} : Category three of classical Hamiltonian framework as the point of departure, authors carry out a non-perturbative quantization of the sector of general relativity, coupled to matter

Module Two

The emphasis is on the quantum geometry of the horizon. Polymer excitations of the bulk quantum geometry pierce the horizon **endowing (eb)** it with area.

G_{16} : Category one of emphasis is on the quantum geometry of the horizon. Polymer excitations of the bulk quantum geometry pierce the horizon

G_{17} : Category two of emphasis is on the quantum geometry of the horizon. Polymer excitations of the bulk quantum geometry pierce the horizon

G_{18} : Category three of emphasis is on the quantum geometry of the horizon. Polymer excitations of the bulk quantum geometry pierce the horizon

T_{16} : Category one of area.

T_{17} : Category two of area.

T_{18} : Category three of area.

Module three

Neuron DNA gives Mind like Higgs Boson gives mass

G_{20} : Category one of Neuron DNA(individual, collective, nature's general ledger)

G_{21} : Category two of Neuron DNA

G_{22} : Category three of Neuron DNA

T_{20} : Category one of Mind (anagrammatic field acting upon the storage of information and receptacle of disinformation or misinformation or indoctrinated one at all three levels and the concomitant fantasies, dreams et,., even pathological gratifications and deprivations. Please se earlier papers)

T_{21} : Category two of Mind

T_{22} : Category three of Mind

Module four

The intrinsic geometry of the horizon **is then (e) described by** the quantum Chern-Simons theory of a U (1) connection on a punctured 2-sphere, the horizon.

G_{24} : Category one of quantum Chern-Simons theory of a U (1) connection on a punctured 2-sphere, the horizon.

G_{25} : Category two of quantum Chern-Simons theory of a U (1) connection on a punctured 2-sphere, the horizon.

G_{26} : Category three of quantum Chern-Simons theory of a U (1) connection on a punctured 2-sphere, the horizon.

T_{24} : Category one of intrinsic geometry of the horizon

T_{25} : Category two of intrinsic geometry of the horizon

T_{26} : Category three of intrinsic geometry of the horizon

Module five

Subtle mathematical features of the quantum Chern-Simons theory **turn out to (e) be important** for the existence of a coherent quantum theory of the horizon geometry

G_{28} : Category one of existence of a coherent quantum theory of the horizon geometry

G_{29} : Category two of existence of a coherent quantum theory of the horizon geometry

G_{30} : Category three of existence of a coherent quantum theory of the horizon geometry

T_{28} : Category one of Subtle mathematical features of the quantum Chern-Simons theory

T_{29} : Category two of Subtle mathematical features of the quantum Chern-Simons theory

T_{30} : Category three of Subtle mathematical features of the quantum Chern-Simons theory

Module six

Heuristically, the intrinsic geometry is flat everywhere **except (e: flatness eaten at) a**t the punctures

G_{32} : Category one of punctures

G_{33} : Category two of punctures

G_{34} : Category three of punctures

T_{32} : Category one of intrinsic geometry is flat everywhere

T_{33} : Category two of intrinsic geometry is flat everywhere

T_{34} : Category three of intrinsic geometry is flat everywhere

Module seven

The distributional curvature of the U (1) connection at the punctures **gives rise to** quantized deficit angles which account for the overall curvature

G_{36} : Category one of distributional curvature of the U (1) connection at the punctures

G_{37} : Category two of distributional curvature of the U (1) connection at the punctures

G_{38} : Category three of distributional curvature of the U (1) connection at the punctures

T_{36} : Category one of quantized deficit angles which account for the overall curvature

T_{37} : Category two of quantized deficit angles which account for the overall curvature

T_{38} : Category three of quantized deficit angles which account for the overall curvature

Module eight

Thus, the black hole entropy **can be (e) accounted** for entirely by the quantum states of the horizon geometry

G_{40} : Category one of entirely by the quantum states of the horizon geometry

G_{41} : Category two of entirely by the quantum states of the horizon geometry

G_{42} : Category three of entirely by the quantum states of the horizon geometry

T_{40} : Category one of black hole entropy

T_{41} : Category two of black hole entropy

T_{42} : Category three of black hole entropy

Module Nine

For macroscopic black holes, the logarithm of the number of these horizon microstates **is proportional** to the area, irrespective of the values of (non-gravitational) charges

G_{44} : Category one of constant multiplied by the area, irrespective of the values of (non-gravitational) charges

G_{45} : Category two of constant multiplied by the area, irrespective of the values of (non-gravitational) charges

G_{46} : Category three of constant multiplied by the area, irrespective of the values of (non-gravitational) charges

T_{44} : Category one of macroscopic black holes, the logarithm of the number of these horizon microstates

T_{45} : Category two of macroscopic black holes, the logarithm of the number of these horizon microstates

T_{46} : Category three of macroscopic black holes, the logarithm of the number of these horizon microstates

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**
$-(b''_{28})^{(5)}\big((G_{31}),t\big) =$ **First detritions factor**
Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \big[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\big]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \big[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\big]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \big[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\big]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \big[(b'_{32})^{(6)} - (b''_{32})^{(6)}\big((G_{35}),t\big)\big]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \big[(b'_{33})^{(6)} - (b''_{33})^{(6)}\big((G_{35}),t\big)\big]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \big[(b'_{34})^{(6)} - (b''_{34})^{(6)}\big((G_{35}),t\big)\big]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**
Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \big[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\big]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \big[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\big]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \big[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\big]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \big[(b'_{36})^{(7)} - (b''_{36})^{(7)}\big((G_{39}),t\big)\big]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \big[(b'_{37})^{(7)} - (b''_{37})^{(7)}\big((G_{39}),t\big)\big]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \big[(b'_{38})^{(7)} - (b''_{38})^{(7)}\big((G_{39}),t\big)\big]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**
Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \big[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\big]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \big[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\big]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \big[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\big]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \big[(b'_{40})^{(8)} - (b''_{40})^{(8)}\big((G_{43}),t\big)\big]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \big[(b'_{41})^{(8)} - (b''_{41})^{(8)}\big((G_{43}),t\big)\big]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \big[(b'_{42})^{(8)} - (b''_{42})^{(8)}\big((G_{43}),t\big)\big]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \big[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\big]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \big[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\big]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \big[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\big]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \big[(b'_{44})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big)\big]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \big[(b'_{45})^{(9)} - (b''_{45})^{(9)}\big((G_{47}),t\big)\big]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b_{44}'')^{(9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt}=(a_{20})^{(3)}G_{21}-\left[\begin{array}{c}(a_{20}')^{(3)}\,+(a_{20}'')^{(3)}(T_{21},t)\,+(a_{16}'')^{(2,2,2)}(T_{17},t)\,+(a_{13}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)\,+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)\,+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{36}'')^{(7,7,7,7)}(T_{37},t)\,+(a_{40}'')^{(8,8,8,8)}(T_{41},t)\,+(a_{44}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt}=(a_{21})^{(3)}G_{20}-\left[\begin{array}{c}(a_{21}')^{(3)}\,+(a_{21}'')^{(3)}(T_{21},t)\,+(a_{17}'')^{(2,2,2)}(T_{17},t)\,+(a_{14}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)\,+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)\,+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{37}'')^{(7,7,7,7)}(T_{37},t)\,+(a_{41}'')^{(8,8,8,8)}(T_{41},t)\,+(a_{45}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt}=(a_{22})^{(3)}G_{21}-\left[\begin{array}{c}(a_{22}')^{(3)}\,+(a_{22}'')^{(3)}(T_{21},t)\,+(a_{18}'')^{(2,2,2)}(T_{17},t)\,+(a_{15}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)\,+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)\,+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{38}'')^{(7,7,7,7)}(T_{37},t)\,+(a_{42}'')^{(8,8,8,8)}(T_{41},t)\,+(a_{46}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{22} \quad (69)$$

$+(a_{20}'')^{(3)}(T_{21},t)$, $+(a_{21}'')^{(3)}(T_{21},t)$, $+(a_{22}'')^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a_{44}'')^{(9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt}=(b_{20})^{(3)}T_{21}-\left[\begin{array}{c}(b_{20}')^{(3)}\,-(b_{20}'')^{(3)}(G_{23},t)\,-(b_{16}'')^{(2,2,2)}(G_{19},t)\,-(b_{13}'')^{(1,1,1,)}(G,t)\\ -(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)\,-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)\,-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)\\ -(b_{36}'')^{(7,7,7,7)}(G_{39},t)\,-(b_{40}'')^{(8,8,8,8)}(G_{43},t)\,-(b_{44}'')^{(9,9,9)}(G_{47},t)\end{array}\right]T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt}=(b_{21})^{(3)}T_{20}-\left[\begin{array}{c}(b_{21}')^{(3)}\,-(b_{21}'')^{(3)}(G_{23},t)\,-(b_{17}'')^{(2,2,2)}(G_{19},t)\,-(b_{14}'')^{(1,1,1,)}(G,t)\\ -(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t)\,-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)\,-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t)\\ -(b_{37}'')^{(7,7,7,7)}(G_{39},t)\,-(b_{41}'')^{(8,8,8,8)}(G_{43},t)\,-(b_{45}'')^{(9,9,9)}(G_{47},t)\end{array}\right]T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$

$are\ fourth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
$are\ fifth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
$are\ sixth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
$are\ seventh\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
$are\ eighth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

$Where\ -(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ $are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ $are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ $are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad (84)$$

$where\ \boxed{-(b''_{28})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{29})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{30})^{(5)}(G_{31},t)}\ \ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}\ ,\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}\ ,\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt}=(a_{32})^{(6)}G_{33} \qquad 85$$

$$-\begin{bmatrix}(a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{32}$$

$$\frac{dG_{33}}{dt}=(a_{33})^{(6)}G_{32}-\begin{bmatrix}(a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{33} \qquad 86$$

$$\frac{dG_{34}}{dt}=(a_{34})^{(6)}G_{33}-\begin{bmatrix}(a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{34} \qquad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}\ ,\boxed{+(a''_{33})^{(6)}(T_{33},t)}\ ,\boxed{+(a''_{34})^{(6)}(T_{33},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}\ ,\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\ ,\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\ are\ second\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}\ \ are\ third\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation

coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34} \quad (90)$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$ *95*

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{c} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{c} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \le (r_i)^{(1)} \le (b_i')^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \le (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} +\ (\hat{A}_{20})^{(3)} +\ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} +\ (\hat{B}_{20})^{(3)} +\ (\hat{Q}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le\ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) =\ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(AAAAAAA) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(BBBBBBB) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(CCCCCCC) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(DDDDDDD)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(EEEEEEE) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(FFFFFFF) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\le\ (\hat{P}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\le\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\le\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{13})^{(1)}$, $T_i^0 \le (\hat{Q}_{13})^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 156

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(aa) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$

In the same way , one can obtain

$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18}. indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16}, G_{18} and G_{16}, G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18}. The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2:

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

208

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)}$$

209

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)}$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

214

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$\left|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{36})^{(7)}\left|G_{37}^{(1)}-G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a''_{36})^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\leq \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0, (\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0, (\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}\left((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)}\right)}\left[e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}\left((S_1)^{(3)} - (a'_{22})^{(3)}\right)}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}\left((R_1)^{(3)} - (b'_{22})^{(3)}\right)}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}\left((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)}\right)}\left[e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right) - (b''_{25})^{(4)}\left((G_{27}), t\right) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, $\boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^{0}e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^{0}e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^{0}}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t}-e^{-(b'_{34})^{(6)}t}\right]+T_{34}^{0}e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^{0}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T_{34}^{0}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, $\boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, $\boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, $\boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, $\boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)}+(C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21}, t)\right) - (a_{21}'')^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)}=\frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(aa) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523

A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FIFTY SIX

Laws Of Black Hole Mechanics : Deus Ex Machina

INTRODUCTION—VARIABLES USED

The laws of black hole mechanics

The four laws of black hole mechanics are physical properties that black holes are believed to satisfy. The laws, analogous to the laws of thermodynamics, were discovered by Brandon, Stephen Hawking and James Bardeen.

Statement of the laws

The laws of black hole mechanics are expressed in geometrized units.

The Zeroth Law

(1) The horizon **has** constant surface gravity for a stationary black hole.

The First Law

(2) Change of mass is related to change of area, angular momentum, and electric charge by:

$$dM = \frac{\kappa}{8\pi}\, dA + \Omega\, dJ + \Phi\, dQ,$$

where M is the mass, κ is the surface gravity, A is the horizon area, Ω is the angular velocity, J is the angular momentum, Φ is the electrostatic potential and Q is the electric charge.

The Second Law

(3) The horizon area is, assuming the weak energy condition, a non-decreasing function of time,

$$\frac{dA}{dt} \geq 0.$$

This "law" was superseded by Hawking's discovery that black holes radiate, which causes both the black hole's mass and the area of its horizon to decrease over time.

The Third Law

(4) It is not possible **to (eb) form** a black hole with vanishing surface gravity. $\kappa = 0$ is not possible to achieve.

Discussion of the laws

The Zeroth Law

(5) The zeroth law is analogous to the zeroth law of thermodynamics which states that the temperature is constant at **all points** and throughout a body in thermal equilibrium. It suggests that the surface gravity **is analogous to** temperature. T constant for thermal equilibrium for a normal system is analogous to κ constant over the horizon of a stationary black hole.

The First Law

(7)The left hand side, dM, is the change in mass/energy. Although the first term does not have an immediately obvious physical interpretation, the second and third terms on the right hand side **represent** changes in energy due to rotation and electromagnetism. Analogously, the first law of thermodynamics is a statement of energy conservation, which **contains on its** right hand side the term T dS.

The Second Law

(6) The second law is the statement of Hawking's area theorem. Analogously, the second law of thermodynamics states that the change in entropy in an isolated system **will be greater than or equal** to 0 for a spontaneous process, suggesting a link between entropy and the area of a black hole horizon. However, this version violates the second law of thermodynamics by matter losing (its) entropy as it falls in, giving a decrease in entropy. Generalized second law introduced as total

entropy = black hole entropy + outside entropy.

The Third Law

(7) Extremal black holes have vanishing surface gravity. Stating that κ cannot go to zero **is(=)analogous** to the third law of thermodynamics which states, the entropy of a system at absolute zero is a well-defined constant **because(e)** a system at zero temperature exists in its ground state. Furthermore, ΔS will reach zero at 0 kelvins, but S itself will also reach zero, at least for perfect crystalline substances. No experimentally verified violations of the laws of thermodynamics are known.

Interpretation of the laws

(8) The four laws of black hole mechanics **suggest that** one should identify the surface gravity of a black hole with temperature and the area of the event horizon with entropy, at least up to some multiplicative constants.

If one only considers black holes classically, then they have zero temperature and, by the no hair theorem, zero entropy, and the laws of black hole mechanics remain an analogy. However, when quantum mechanical effects are taken into account, one finds that black holes emit thermal radiation (Hawking radiation) at temperature

$$T_{\mathrm{H}} = \frac{\kappa}{2\pi}.$$

From the first law of black hole mechanics, this determines the multiplicative constant of the Bekenstein-Hawking entropy which is

$$S_{\mathrm{BH}} = \frac{A}{4}.$$

Beyond black holes

(9) Hawking and Page have shown that black hole thermodynamics is more general than black holes, that cosmological event horizons **also(e) have** an entropy and temperature.

(10) More fundamentally, 't Hooft and Susskind used the laws of black hole thermodynamics **to(eb) argue** for a general Holographic Principle of nature, which asserts that consistent theories of gravity and quantum mechanics must be lower dimensional.

(11) Though not yet fully understood in general, the holographic principle **is(e) central to** theories like the AdS/CFT correspondence.

NOTATION

Module One

The horizon **has** constant surface gravity for a stationary black hole

$$dM = \frac{\kappa}{8\pi}\, dA + \Omega\, dJ + \Phi\, dQ,$$

G_{13} : Category one of constant surface gravity for a stationary black hole **;**LHS of the first law**;**

G_{14} : Category two of constant surface gravity for a stationary black hole; LHS of the first law;

G_{15} : Category three of constant surface gravity for a stationary black hole ;LHS of the first law;

T_{13} : Category one of horizon; RHS of the first law

T_{14} : Category two of horizon ; RHS of the first law;

T_{15} : Category three of horizon; RHS of the first law

Module Two

$$\frac{dA}{dt} \geq 0.$$

G_{16} : Category one of LHS of the above equation

G_{17} : Category two of LHS of the above equation

G_{18} : Category three of LHS of the above equation

T_{16} : Category one of RHS of the above equation

T_{17} : Category two of RHS of the above equation

T_{18} : Category three of RHS of the above equation

Module three

It is not possible **to (eb) form** a black hole with vanishing surface gravity. $\kappa = 0$ is not possible to achieve.

G_{20} : Category one of vanishing surface gravity. $\kappa = 0$ is not possible to achieve

G_{21} : Category two of vanishing surface gravity. $\kappa = 0$ is not possible to achieve

G_{22} : Category three of vanishing surface gravity. $\kappa = 0$ is not possible to achieve

T_{20} : Category one of impossibility of the formation of the black hole

T_{21} : Category two of impossibility of the formation of the black hole

T_{22} : Category three of impossibility of the formation of the black hole

Module four

The zeroth law is analogous to the zeroth law of thermodynamics which states that the temperature is constant at **all points** and throughout a body in thermal equilibrium. It suggests that the surface gravity **is analogous to** temperature. T constant for thermal equilibrium for a normal system is analogous to κ constant over the horizon of a stationary black hole

G_{24} : Category one of temperature. T constant for thermal equilibrium for a normal system is analogous to κ constant over the horizon of a stationary black hole

G_{25} : Category two of temperature. T constant for thermal equilibrium for a normal system is analogous to κ constant over the horizon of a stationary black hole

G_{26} : Category three of temperature. T constant for thermal equilibrium for a normal system is analogous

to κ constant over the horizon of a stationary black hole

T_{24} : Category one of zeroth law is analogous to the zeroth law of thermodynamics which states that the temperature is constant at **all points** and throughout a body in thermal equilibrium. It suggests that the surface gravity

T_{25} : Category two of zeroth law is analogous to the zeroth law of thermodynamics which states that the temperature is constant at **all points** and throughout a body in thermal equilibrium. It suggests that the surface gravity

T_{26} : Category three of zeroth law is analogous to the zeroth law of thermodynamics which states that the temperature is constant at **all points** and throughout a body in thermal equilibrium. It suggests that the surface gravity

Module five

The second law is the statement of Hawking's area theorem. Analogously, the second law of thermodynamics states that the change in entropy in an isolated system **will be greater than or equal** to 0 for a spontaneous process, suggesting a link between entropy and the area of a black hole horizon. However, this version violates the second law of thermodynamics by matter losing (its) entropy as it falls in, giving a decrease in entropy. **Generalized second law introduced as total entropy = black hole entropy + outside entropy**

G_{28} : Category one of LHS of the equation Generalized second law introduced as total entropy = black hole entropy + outside entropy

G_{29} : Category two of LHS of the equation Generalized second law introduced as total entropy = black hole entropy + outside entropy

G_{30} : Category three of LHS of the equation Generalized second law introduced as total entropy = black hole entropy + outside entropy

T_{28} : Category one of RHS of the equation Generalized second law introduced as total entropy = black hole entropy + outside entropy

T_{29} : Category two of RHS of the equation Generalized second law introduced as total entropy = black hole entropy + outside entropy

T_{30} : Category three of RHS of the equation Generalized second law introduced as total entropy = black hole entropy + outside entropy

Module six

$$T_{\mathrm{H}} = \frac{\kappa}{2\pi}. \qquad S_{\mathrm{BH}} = \frac{A}{4}.$$

G_{32} : Category one of LHS of the above equations

G_{33} : Category two of LHS of the above equations

G_{34} : Category three of LHS of the above equations

T_{32} : Category one of RHS of the above equations

T_{33} : Category two of RHS of the above equations

T_{34} : Category three of RHS of the above equations

Module seven

Hawking and Page have shown that black hole thermodynamics is more general than black holes, that cosmological event horizons **also(e) have** an entropy and temperature

G_{36} : Category one of entropy and temperature

G_{37} : Category two of entropy and temperature

G_{38} : Category three of entropy and temperature

T_{36} : Category one of black hole thermodynamics is more general than black holes, that cosmological event horizons

T_{37} : Category two of black hole thermodynamics is more general than black holes, that cosmological event horizons

T_{38} : Category three of black hole thermodynamics is more general than black holes, that cosmological event horizons

Module eight

More fundamentally, 't Hooft and Susskind used the laws of black hole thermodynamics **to(eb) argue** for a general Holographic Principle of nature, which asserts that consistent theories of gravity and quantum mechanics must be lower dimensional.

G_{40} : Category one of, 't Hooft and Susskind used the laws of black hole thermodynamics

G_{41} : Category two of, 't Hooft and Susskind used the laws of black hole thermodynamics

G_{42} : Category three of, 't Hooft and Susskind used the laws of black hole thermodynamics

T_{40} : Category one of general Holographic Principle of nature, which asserts that consistent theories of gravity and quantum mechanics must be lower dimensional.

T_{41} : Category two of general Holographic Principle of nature, which asserts that consistent theories of gravity and quantum mechanics must be lower dimensional.

T_{42} : Category three of general Holographic Principle of nature, which asserts that consistent theories of gravity and quantum mechanics must be lower dimensional.

Module Nine

Though not yet fully understood in general, the holographic principle **is (e) central to** theories like the AdS/CFT correspondence.

G_{44} : Category one of. AdS/CFT correspondence

G_{45} : Category two of AdS/CFT correspondence

G_{46} : Category three of AdS/CFT correspondence

T_{44} : Category one of holographic principle is used in the construction of theories

T_{45} : Category two of holographic principle is used in the construction of theories

T_{46} : Category three of holographic principle is used in the construction of theories

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G,t)\right]T_{14}$ 5

$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15}$ 6

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16}$ 7

$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17}$ 8

$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18}$ 9

$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}),t\right)\right]T_{16}$ 10

$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17}$ 11

$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18}$ 12

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20}$ 13

$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21}$ 14

$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22}$ 15

$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20}$ 16

$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21}$ 17

$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22}$ 18

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24}$ 19

$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25}$ 20

$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26}$ 21

$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24}$ 22

$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25}$ 23

$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26}$ 24

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}), t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}), t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition

coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)}\ \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\ \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\ \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for

category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

are fourth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$

are fifth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$

are sixth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$

are seventh detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$

are eighth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad (84)$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ *are second detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b_{46}'')^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)} & +(a_{36}'')^{(7)}(T_{37},t) & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{40}'')^{(8,8,8,8,8,8,8,)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)} & +(a_{37}'')^{(7)}(T_{37},t) & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)} & +(a_{38}'')^{(7)}(T_{37},t) & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a_{36}'')^{(7)}(T_{37},t)$, $(a_{37}'')^{(7)}(T_{37},t)$, $(a_{38}'')^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)} G_{44}$$
$$- \left[\begin{array}{c} (a'_{45})^{(9)} \boxed{+(a''_{45})^{(9)}(T_{45},t)} \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)} G_{45}$$
$$- \left[\begin{array}{c} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)} T_{45} -$$
$$\left[\begin{array}{c} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{13}$$
$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}\left((G_{19}), t\right) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ \ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(GGGGGGG) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(HHHHHHH) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(IIIIIII) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(JJJJJJJ)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(KKKKKKK) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(LLLLLLL) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and \ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ \ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)}$, 155

$0 \leq G_i(t) - G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \leq T_i(t) - T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{16})^{(2)}$, $T_i^0 \leq (\hat{Q}_{16})^{(2)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^{0} + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^{0} + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^{0} + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(bb) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+\ (a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ (228)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}),s_{(24)})\}ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(4)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(7)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\,((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \le \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \le \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \ge (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \ge \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \ge \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$
Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \qquad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \qquad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \qquad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

406

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case,

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

407

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(bb) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)} G_{14}^{*}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)} G_{14}^{*}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)} T_{14}^{*}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)} T_{14}^{*}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)} G_{17}^{*}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)} G_{17}^{*}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)} T_{17}^{*}}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)} T_{17}^{*}}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)} G_{21}^{*}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)} G_{21}^{*}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)} T_{21}^{*}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)} T_{21}^{*}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FIFTY SEVEN

Correlation Space And Extremal Entanglement: Labor Omnia Vincit-Leges Humanae Nascuntur, Vivunt, Et Moriuntur

INTRODUCTION—VARIABLES USED

Preserving Coherence in Quantum Computation by Pairing Quantum Bits Lu-Ming Duan et al

(1) A scheme for protecting quantum states **from (eb) both** independent and cooperative decoherence is proposed.
(2) The scheme operates by pairing each qubit (two-state quantum system) **with an** ancilla qubit and by encoding the states of the qubits into corresponding coherence-preserving states of qubit pairs.
(3) In this scheme, amplitude damping (loss of energy) as well as phase damping (dephasing) **is(e) prevented** by a strategy called "free-Hamiltonian elimination."
(4) Authors further extend the scheme to include quantum gate operations **and show that** loss and decoherence during such operations can also be prevented.© 1997 The American Physical Society URL:http://link.aps.org/doi/10.1103/PhysRevLett.79.1953 DOI:10.1103/PhysRevLett.79.1953

Quantum Computation in Correlation Space and Extremal Entanglement J.-M. Cai et al

(5) Recently, a framework was established to systematically **construct (eb)** novel universal resource states for measurement-based quantum computation using techniques involving finitely correlated states.
(6) With these methods, universal states were found which are in certain ways much less entangled **than the** original cluster-state model, and **it was hence (eb) believed that** with this approach, many of the extremal entanglement features of the cluster states could be relaxed.
(7) The new resources were constructed as "computationally universal" states—i.e., they allow one to efficiently reproduce the classical output of each quantum computation—whereas the cluster states **are** universal in a stronger sense since they are "universal state preparators."
(8) Here, authors show that the new resources are universal state preparators after all, and must therefore **exhibit a** whole class of extremal entanglement features, similar to the cluster states.© 2009 The American Physical SocietyURL:http://link.aps.org/doi/10.1103/PhysRevLett.103.050503DOI:10.1103/PhysRevLett.103.050503

Nature Physics, 19 - 26 2 January 2009 | doi: 10.1038/nphys1157 H. J. Briegel et al Measurement-based quantum computation

(9) Quantum computation offers a promising new kind of information processing, **where (EB)** the non-classical features of quantum mechanics are harnessed and exploited.

NOTATION

Module One

A scheme for protecting quantum states **from (eb) both** independent and cooperative decoherence is proposed

G_{13} : Category one of protection of quantum states

G_{14} : Category two of protection of quantum states

G_{15} : Category three of protection of quantum states

T_{13} : Category one of independent and cooperative decoherence

T_{14} : Category two of independent and cooperative decoherence

T_{15} : Category three of independent and cooperative decoherence

Module Two

The scheme operates by pairing each qubit (two-state quantum system) **with an** ancilla qubit and by encoding the states of the qubits into corresponding coherence-preserving states of qubit pairs

G_{16} : Category one of scheme operates by pairing each qubit (two-state quantum system)

G_{17} : Category two of scheme operates by pairing each qubit (two-state quantum system)

G_{18} : Category three of scheme operates by pairing each qubit (two-state quantum system)

T_{16} : Category one of ancilla qubit and by encoding the states of the qubits into corresponding coherence-preserving states of qubit pairs

T_{17} : Category two of ancilla qubit and by encoding the states of the qubits into corresponding coherence-preserving states of qubit pairs

T_{18} : Category three of ancilla qubit and by encoding the states of the qubits into corresponding coherence-preserving states of qubit pairs

Module three

In this scheme, amplitude damping (loss of energy) as well as phase damping (dephasing) **is(e) prevented** by a strategy called "free-Hamiltonian elimination"

G_{20} : Category one of "free-Hamiltonian elimination"

G_{21} : Category two of "free-Hamiltonian elimination"

G_{22} : Category three of "free-Hamiltonian elimination"

T_{20} : Category one of amplitude damping (loss of energy) as well as phase damping (dephasing)

T_{21} : Category two of amplitude damping (loss of energy) as well as phase damping (dephasing)

T_{22} : Category three of amplitude damping (loss of energy) as well as phase damping (dephasing)

Module four

Authors further extend the scheme to include quantum gate operations **and show that** loss and decoherence during such operations can also be prevented

G_{24} : Category one of extension of the scheme to include quantum gate operations

G_{25} : Category two of extension of the scheme to include quantum gate operations

G_{26} : Category three of extension of the scheme to include quantum gate operations

T_{24} : Category one of loss and decoherence during such operations can also be prevented

T_{25} : Category two of loss and decoherence during such operations can also be prevented

T_{26} : Category three of loss and decoherence during such operations can also be prevented

Module five

Recently, a framework was established to systematically **construct (eb)** novel universal resource states for measurement-based quantum computation using techniques involving finitely correlated states.

G_{28} : Category one of measurement-based quantum computation using techniques involving finitely correlated states

G_{29} : Category two of measurement-based quantum computation using techniques involving finitely correlated states

G_{30} : Category three of measurement-based quantum computation using techniques involving finitely correlated states

T_{28} : Category one of framework was established to systematically **construct (eb)** novel universal resource states

T_{29} : Category two of framework was established to systematically **construct (eb)** novel universal resource states

T_{30} : Category three of framework was established to systematically **construct (eb)** novel universal resource states

Module six

With these methods, universal states were found which are in certain ways much less entangled **than the** original cluster-state model, and **it was hence (eb) believed that** with this approach, many of the extremal entanglement features of the cluster states could be relaxed

G_{32} : Category one of universal states were found which are in certain ways much less entangled **than the** original cluster-state model,

G_{33} : Category two of universal states were found which are in certain ways much less entangled **than the** original cluster-state model,

G_{34} : Category three of universal states were found which are in certain ways much less entangled **than the** original cluster-state model,

T_{32} : Category one of many of the extremal entanglement features of the cluster states could be relaxed

T_{33} : Category two of many of the extremal entanglement features of the cluster states could be relaxed

T_{34} : Category three of many of the extremal entanglement features of the cluster states could be relaxed

Module seven

The new resources were constructed as "computationally universal" states—i.e., they allow one to efficiently reproduce the classical output of each quantum computation—whereas the cluster states **are** universal in a stronger sense since they are "universal state preparators."

G_{36} : Category one of universal in a stronger sense since they are "universal state preparators."

G_{37} : Category two of universal in a stronger sense since they are "universal state preparators."

G_{38} : Category three of universal in a stronger sense since they are "universal state preparators."

T_{36} : Category one of new resources were constructed as "computationally universal" states—i.e., they allow one to efficiently reproduce the classical output of each quantum computation—whereas the cluster states

T_{37} : Category two of new resources were constructed as "computationally universal" states—i.e., they allow one to efficiently reproduce the classical output of each quantum computation—whereas the cluster states

T_{38} : Category three of new resources were constructed as "computationally universal" states—i.e., they allow one to efficiently reproduce the classical output of each quantum computation—whereas the cluster states

Module eight

G_{40} : Category one of "universal state preparators."

G_{41} : Category two of "universal state preparators."

G_{42} : Category three of "universal state preparators."

T_{40} : Category one of cluster states

T_{41} : Category two of cluster states

T_{42} : Category three of cluster states

Module Nine

Here, authors show that the new resources are universal state preparators after all, and must therefore **exhibit a** whole class of extremal entanglement features, similar to the cluster states

G_{44} : Category one of new resources are universal state preparators after all, and must therefore

G_{45} : Category two of new resources are universal state preparators after all, and must therefore

G_{46} : Category three of new resources are universal state preparators after all, and must therefore

T_{44} : Category one of whole class of extremal entanglement features, similar to the cluster states

T_{45} : Category two of whole class of extremal entanglement features, similar to the cluster states

T_{46} : Category three of whole class of extremal entanglement features, similar to the cluster states

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}), t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}), t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}), t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) \end{bmatrix} G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{20})^{(3)}\,+(a''_{20})^{(3)}(T_{21},t)\,+(a''_{16})^{(2,2,2)}(T_{17},t)\,+(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)\,+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)\,+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t)\,+(a''_{40})^{(8,8,8,8)}(T_{41},t)\,+(a''_{44})^{(9,9,9)}(T_{45},t) \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c} (a'_{21})^{(3)}\,+(a''_{21})^{(3)}(T_{21},t)\,+(a''_{17})^{(2,2,2)}(T_{17},t)\,+(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)\,+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)\,+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t)\,+(a''_{41})^{(8,8,8,8)}(T_{41},t)\,+(a''_{45})^{(9,9,9)}(T_{45},t) \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{22})^{(3)}\,+(a''_{22})^{(3)}(T_{21},t)\,+(a''_{18})^{(2,2,2)}(T_{17},t)\,+(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)\,+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)\,+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t)\,+(a''_{42})^{(8,8,8,8)}(T_{41},t)\,+(a''_{46})^{(9,9,9)}(T_{45},t) \end{array}\right] G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{c} (b'_{20})^{(3)}\,-(b''_{20})^{(3)}(G_{23},t)\,-(b''_{16})^{(2,2,2)}(G_{19},t)\,-(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)\,-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)\,-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t)\,-(b''_{40})^{(8,8,8,8)}(G_{43},t)\,-(b''_{44})^{(9,9,9)}(G_{47},t) \end{array}\right] T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{c} (b'_{21})^{(3)}\,-(b''_{21})^{(3)}(G_{23},t)\,-(b''_{17})^{(2,2,2)}(G_{19},t)\,-(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)\,-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)\,-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t)\,-(b''_{41})^{(8,8,8,8)}(G_{43},t)\,-(b''_{45})^{(9,9,9)}(G_{47},t) \end{array}\right] T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$

$are\ fourth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
$are\ fifth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
$are\ sixth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
$are\ seventh\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
$are\ eighth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

$Where\ -(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ $are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ $are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ $are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad (84)$$

$where\ \boxed{-(b''_{28})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{29})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{30})^{(5)}(G_{31},t)}\ \ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}\ ,\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}\ ,\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt}=(a_{32})^{(6)}G_{33} \qquad 85$$

$$-\left[\begin{matrix}(a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)}\ \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}\ \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{32}$$

$$\frac{dG_{33}}{dt}=(a_{33})^{(6)}G_{32}-\left[\begin{matrix}(a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)}\ \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\ \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{33} \qquad 86$$

$$\frac{dG_{34}}{dt}=(a_{34})^{(6)}G_{33}-\left[\begin{matrix}(a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\ \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\ \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{34} \qquad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}\ ,\boxed{+(a''_{33})^{(6)}(T_{33},t)}\ ,\boxed{+(a''_{34})^{(6)}(T_{33},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}\ ,\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\ ,\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\ are\ second\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}\ \ are\ third\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation

coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34} \quad 90$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$ *95*

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{ccc} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\ ,\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)},\ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ \ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)},\ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \le\ \ (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G\to\infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)},\ \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \le (\hat{k}_{32})^{(6)}|T_{33} - \ T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(MMMMMMM) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(NNNNNNN) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(OOOOOOO) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(PPPPPPP)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(QQQQQQQ) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(RRRRRRR) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\leq\ \left(\hat{P}_{13}\right)^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\leq\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\leq\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\leq\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(cc) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$

In the same way , one can obtain

$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)} t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18}. indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16}, G_{18} and G_{16}, G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18}. The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2:

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

208

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)}$$

209

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)}$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

214

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\widehat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{36})^{(7)}\right]\leq(\widehat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$\left|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{36})^{(7)}\left|G_{37}^{(1)}-G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a''_{36})^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0, (\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0, (\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)}$ (293)

$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)}$ (294)

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$$ 318

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$ and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$ and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,

and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(\nu_1)^{(6)}, (\nu_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, \nu^{(6)}, u^{(6)}$: 350

By $(\nu_1)^{(6)} > 0$, $(\nu_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)},, (\bar{\nu}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, $\boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^{0}e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^{0}e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^{0}}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t}-e^{-(b'_{34})^{(6)}t}\right]+T_{34}^{0}e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^{0}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T_{34}^{0}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)}+(C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21}, t)\right) - (a_{21}'')^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)},(v_0)^{(7)}$:-

$$\text{For } 0<\boxed{(v_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(v_1)^{(7)}<(\bar{v}_1)^{(7)}$$

$$v^{(7)}(t)\geq\frac{(v_1)^{(7)}+(C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(v_1)^{(7)}-(v_0)^{(7)}}{(v_0)^{(7)}-(v_2)^{(7)}}}$$

it follows $(v_0)^{(7)}\leq v^{(7)}(t)\leq(v_1)^{(7)}$

In the same manner , we get 417

$$v^{(7)}(t)\leq\frac{(\bar{v}_1)^{(7)}+(\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{v}_1)^{(7)}-(v_0)^{(7)}}{(v_0)^{(7)}-(\bar{v}_2)^{(7)}}}$$

From which we deduce $(v_0)^{(7)}\leq v^{(7)}(t)\leq(\bar{v}_1)^{(7)}$

If $0<(v_1)^{(7)}<(v_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)}\leq\frac{(v_1)^{(7)}+(C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_2)^{(7)}\right)t\right]}}\leq v^{(7)}(t)\leq$$

$$\frac{(\bar{v}_1)^{(7)}+(\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}\leq(\bar{v}_1)^{(7)}$$

If $0<(v_1)^{(7)}\leq(\bar{v}_1)^{(7)}\leq\boxed{(v_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)}\leq v^{(7)}(t)\leq\frac{(\bar{v}_1)^{(7)}+(\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}\leq(v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)}\leq v^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{v^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

423

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case,

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

424

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(cc) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523

A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FIFTY EIGHT

Extreme Spin Squeezing: Crede Quod Habes, Et Habes

INTRODUCTION—VARIABLES USED

Nature Physics, 19 - 26 2 January 2009 | doi: 10.1038/nphys1157 H. J. Briegel et al Measurement-based quantum computation

(1) A number of models of quantum computation exist. These models have been **shown to be** formally equivalent, but their underlying elementary concepts and the requirements for their practical realization can differ significantly.
(2) A particularly exciting paradigm is that of measurement-based quantum computation, where the processing of quantum information **takes (EB) place** by rounds of simple measurements on qubits prepared in a highly entangled state. Authors review recent developments in measurement-based quantum computation **with a view to both** fundamental and practical issues, in particular the power of quantum computation, the protection against noise (fault tolerance) and steps towards experimental realization. Finally, they highlight a number of connections between this field and other branches of physics and mathematics.

Entanglement and Extreme Spin Squeezing Anders S. Sørensen and Klaus Mølmer

(3) For any mean value of a Cartesian component of a spin vector authors **identify (eb)** the smallest possible uncertainty in any of the orthogonal components.

(4) The corresponding states are optimal for spectroscopy and atomic clocks. Authors show that the results for different spin J **can be used** to identify entanglement and to quantify the depth of entanglement in systems with many particles.

(5) With the procedure developed in this Letter, collective spin measurements on an ensemble of particles **can be (e) used** as an experimental proof of multiparticle entanglement. © 2001 The American Physical SocietyURL:http://link.aps.org/doi/10.1103/PhysRevLett.86.4431DOI:10.1103/PhysRevLett.86.4431

The Smithsonian/NASA Astrophysics Data System Quantum Field Theory and Decoherence in the Early Universe Koksma, J. F. Ph.D. Thesis, University Utrecht, The Netherlands, 2011

(6) Quantum field theory(In theoretical physics, quantum field theory (QFT) **is(=) a** theoretical framework for constructing quantum mechanical models of subatomic particles in particle physics and quasiparticles in condensed matter physics, by treating a particle as an excited state of an underlying physical field. These excited states are called **field quanta**. For example, quantum electrodynamics (QED) **has** one electron field and one photon field, quantum chromodynamics (QCD) has one field for each type of quark, and in condensed matter there is an atomic displacement field **that gives rise to** phonon particles.

(7) In QFT, interactions between particles in quantum mechanics **are (e)represented by** interaction terms between the underlying fields.

(8) QFT interaction terms **are similar in spirit to those** between electric and magnetic fields in Maxwell's equations. However unlike the classical fields of Maxwell's theory, fields in QFT generally **exist in** quantum superpositions of states and obey the laws of quantum mechanics) **is (E) indispensable for** understanding many aspects of cosmology, both in the early Universe and today.

(9) For example, quantum processes (Quantum Processes: Quantum properties **dominate the** fields of atomic and molecular physics. Radiation is quantized such that for a given frequency of radiation, there can be only one value of quantum energy for the photons of that radiation. The energy levels of atoms and molecules can have only certain quantized values. Transitions between these quantized states occur by the photon processes **absorption, emission, and stimulated** emission. All of these processes require that the photon energy given by the Planck relationship is equal to the energy separation of the participating pair of quantum energy states) c**ould be (e) paramount to understand** the nature of the mysterious dark energy resulting in the Universe's recently observed accelerated expansion

NOTATION

Module One

A number of models of quantum computation exist. These models have been **shown to be** formally equivalent, but their underlying elementary concepts and the requirements for their practical realization can differ significantly

G_{13} : Category one of models of quantum computation exist. These models have been **shown to be** formally equivalent,

G_{14} : Category two of models of quantum computation exist. These models have been **shown to be** formally equivalent,

G_{15} : Category three of models of quantum computation exist. These models have been **shown to be**

formally equivalent,

T_{13} : Category one of underlying elementary concepts and the requirements for their practical realization can differ significantly

T_{14} : Category two of underlying elementary concepts and the requirements for their practical realization can differ significantly

T_{15} : Category three of underlying elementary concepts and the requirements for their practical realization can differ significantly

Module Two

A particularly exciting paradigm is that of measurement-based quantum computation, where the processing of quantum information **takes (EB) place** by rounds of simple measurements on qubits prepared in a highly entangled state. Authors review recent developments in measurement-based quantum computation **with a view to both** fundamental and practical issues, in particular the power of quantum computation, the protection against noise (fault tolerance) and steps towards experimental realization. Finally, they highlight a number of connections between this field and other branches of physics and mathematics.

G_{16} : Category one of particularly exciting paradigm is that of measurement-based quantum computation, where the processing of quantum information

G_{17} : Category two of particularly exciting paradigm is that of measurement-based quantum computation, where the processing of quantum information

G_{18} : Category three of particularly exciting paradigm is that of measurement-based quantum computation, where the processing of quantum information

T_{16} : Category one of rounds of simple measurements on qubits prepared in a highly entangled state. Authors review recent developments in measurement-based quantum computation **with a view to both** fundamental and practical issues, in particular the power of quantum computation, the protection against noise (fault tolerance) and steps towards experimental realization. Finally, they highlight a number of connections between this field and other branches of physics and mathematics

T_{17} : Category two of rounds of simple measurements on qubits prepared in a highly entangled state. Authors review recent developments in measurement-based quantum computation **with a view to both** fundamental and practical issues, in particular the power of quantum computation, the protection against noise (fault tolerance) and steps towards experimental realization

T_{18} : Category three of rounds of simple measurements on qubits prepared in a highly entangled state. Authors review recent developments in measurement-based quantum computation **with a view to both** fundamental and practical issues, in particular the power of quantum computation, the protection against noise (fault tolerance) and steps towards experimental realization

Module three

For any mean value of a Cartesian component of a spin vector authors **identify (eb)** the smallest possible uncertainty in any of the orthogonal components

G_{20} : Category one of mean value of a Cartesian component of a spin vector

G_{21} : Category two of mean value of a Cartesian component of a spin vector

G_{22} : Category three of mean value of a Cartesian component of a spin vector

T_{20} : Category one of smallest possible uncertainty in any of the orthogonal components

T_{21} : Category two of smallest possible uncertainty in any of the orthogonal components

T_{22} : Category three of smallest possible uncertainty in any of the orthogonal components

Module four

Corresponding states are optimal for spectroscopy and atomic clocks. Authors show that the results for different spin J **can be used** to identify entanglement and to quantify the depth of entanglement in systems with many particles.

G_{24} : Category one of identify entanglement and to quantify the depth of entanglement in systems with many particles.

G_{25} : Category two of identify entanglement and to quantify the depth of entanglement in systems with many particles.

G_{26} : Category three of identify entanglement and to quantify the depth of entanglement in systems with many particles.

T_{24} : Category one of Corresponding states is optimal for spectroscopy and atomic clocks. Authors show that the results for different spin J

T_{25} : Category two of Corresponding states are optimal for spectroscopy and atomic clocks. Authors show that the results for different spin J

T_{26} : Category three of Corresponding states are optimal for spectroscopy and atomic clocks. Authors show that the results for different spin J

Module five

With the procedure developed in this Letter, collective spin measurements on an ensemble of particles **can be (e) used** as an experimental proof of multiparticle entanglement

G_{28} : Category one of experimental proof of multiparticle entanglement

G_{29} : Category two of experimental proof of multiparticle entanglement

G_{30} : Category three of experimental proof of multiparticle entanglement

T_{28} : Category one of procedure developed in this Letter, collective spin measurements on an ensemble of particles

T_{29} : Category two of procedure developed in this Letter, collective spin measurements on an ensemble of particles

T_{30} : Category three of procedure developed in this Letter, collective spin measurements on an ensemble of particles

Module six

Quantum field theory(In theoretical physics, quantum field theory (QFT) **is(=) a** theoretical framework for constructing quantum mechanical models of subatomic particles in particle physics and quasiparticles in

condensed matter physics, by treating a particle as an excited state of an underlying physical field. These excited states are called **field quanta**. For example, quantum electrodynamics (QED) **has** one electron field and one photon field, quantum chromodynamics (QCD) has one field for each type of quark, and in condensed matter there is an atomic displacement field **that gives rise to** phonon particles

G_{32} : Category one of quantum electrodynamics (QED) **has** one electron field and one photon field, quantum chromodynamics (QCD) has one field for each type of quark, and in condensed matter there is an atomic displacement field

G_{33} : Category two of quantum electrodynamics (QED) **has** one electron field and one photon field, quantum chromodynamics (QCD) has one field for each type of quark, and in condensed matter there is an atomic displacement field

G_{34} : Category three of quantum electrodynamics (QED) **has** one electron field and one photon field, quantum chromodynamics (QCD) has one field for each type of quark, and in condensed matter there is an atomic displacement field

T_{32} : Category one of phonon particles

T_{33} : Category two of phonon particles

T_{34} : Category three of phonon particles

Module seven

In QFT, interactions between particles in quantum mechanics **are (e) represented by** interaction terms between the underlying fields.

G_{36} : Category one of interaction terms between the underlying fields.

G_{37} : Category two of interaction terms between the underlying fields.

G_{38} : Category three of interaction terms between the underlying fields.

T_{36} : Category one of QFT, interactions between particles in quantum mechanics (systems which satisfy the QFT axiomatics and whose characterstics form the pillar and post of classification. This does not mean there are three QFT. It is like Newtonian gravity. Numerous systems could be studied and they could be classified based on some parameters).

T_{37} : Category two of QFT, interactions between particles in quantum mechanics

T_{38} : Category three of QFT, interactions between particles in quantum mechanics

Module eight

QFT interaction terms **are similar in spirit to those** between electric and magnetic fields in Maxwell's equations. However unlike the classical fields of Maxwell's theory, fields in QFT generally **exist in** quantum superpositions of states and obey the laws of quantum mechanics) **is (E) indispensable for** understanding many aspects of cosmology, both in the early Universe and today.

G_{40} : Category one of understanding many aspects of cosmology, both in the early Universe and today.

G_{41} : Category two of understanding many aspects of cosmology, both in the early Universe and today.

G_{42} : Category three of understanding many aspects of cosmology, both in the early Universe and today.

T_{40} : Category one of QFT interaction terms **are similar in spirit to those** between electric and magnetic fields in Maxwell's equations. However unlike the classical fields of Maxwell's theory, fields in QFT generally **exist in** quantum superpositions of states and obey the laws of quantum mechanics)

T_{41} : Category two of QFT interaction terms **are similar in spirit to those** between electric and magnetic fields in Maxwell's equations. However unlike the classical fields of Maxwell's theory, fields in QFT generally **exist in** quantum superpositions of states and obey the laws of quantum mechanics)

T_{42} : Category three of QFT interaction terms **are similar in spirit to those** between electric and magnetic fields in Maxwell's equations. However unlike the classical fields of Maxwell's theory, fields in QFT generally **exist in** quantum superpositions of states and obey the laws of quantum mechanics)

Module Nine

For example, quantum processes (Quantum Processes: Quantum properties **dominate the** fields of atomic and molecular physics. Radiation is quantized such that for a given frequency of radiation, there can be only one value of quantum energy for the photons of that radiation. The energy levels of atoms and molecules can have only certain quantized values. Transitions between these quantized states occur by the photon processes **absorption, emission, and stimulated** emission. All of these processes require that the photon energy given by the Planck relationship is equal to the energy separation of the participating pair of quantum energy states) c**ould be (e) paramount to understand** the nature of the mysterious dark energy resulting in the Universe's recently observed accelerated expansion

G_{44} : Category one of nature of the mysterious dark energy resulting in the Universe's recently observed accelerated expansion

G_{45} : Category two of nature of the mysterious dark energy resulting in the Universe's recently observed accelerated expansion

G_{46} : Category three of nature of the mysterious dark energy resulting in the Universe's recently observed accelerated expansion

T_{44} : Category one of Transitions between these quantized states occurs by the photon processes **absorption, emission, and stimulated** emission. All of these processes require that the photon energy given by the Planck relationship is equal to the energy separation of the participating pair of quantum energy states)

T_{45} : Category two of Transitions between these quantized states occur by the photon processes **absorption, emission, and stimulated** emission. All of these processes require that the photon energy given by the Planck relationship is equal to the energy separation of the participating pair of quantum energy states)

T_{46} : Category three of Transitions between these quantized states occur by the photon processes **absorption, emission, and stimulated** emission. All of these processes require that the photon energy given by the Planck relationship is equal to the energy separation of the participating pair of quantum energy states)

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$
$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$
$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**
$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)}\ \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{array}\right] G_{18} \quad 63$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for*

category 1, 2 and 3

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* $1,2\ 3$

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* $1,2$ *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* $1,2$ *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category $1,2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a_{32}')^{(6)} & +(a_{32}'')^{(6)}(T_{33},t) & +(a_{28}'')^{(5,5,5)}(T_{29},t) & +(a_{24}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a_{33}')^{(6)} & +(a_{33}'')^{(6)}(T_{33},t) & +(a_{29}'')^{(5,5,5)}(T_{29},t) & +(a_{25}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a_{34}')^{(6)} & +(a_{34}'')^{(6)}(T_{33},t) & +(a_{30}'')^{(5,5,5)}(T_{29},t) & +(a_{26}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a_{32}'')^{(6)}(T_{33},t)$, $+(a_{33}'')^{(6)}(T_{33},t)$, $+(a_{34}'')^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a_{28}'')^{(5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a_{24}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{ccc} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $-(b_{36}'')^{(7)}(G_{39},t)$, $-(b_{37}'')^{(7)}(G_{39},t)$, $-(b_{38}'')^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)} & +(a_{44}'')^{(9)}(T_{45},t) & +(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & & +(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & & +(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)} & +(a_{45}'')^{(9)}(T_{45},t) & +(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & & +(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & & +(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)} & +(a_{46}'')^{(9)}(T_{37},t) & +(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & & +(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & & +(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{15}$$

Where $+(a_{44}'')^{(9)}(T_{45},t)$, $+(a_{45}'')^{(9)}(T_{45},t)$, $+(a_{46}'')^{(9)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \le \ (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \le (\hat{k}_{32})^{(6)}|T_{33} - \ T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(SSSSSSS) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(TTTTTTT) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(UUUUUUU) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(VVVVVVV)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(WWWWWWW) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(XXXXXXX) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\leq\ \left(\hat{P}_{13}\right)^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\leq\ (\hat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\leq\ (\hat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\leq\ (\hat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{13})^{(1)}$, $T_i^0 \le (\hat{Q}_{13})^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 156

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^{0} + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^{0} + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^{0} + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^{0} + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(dd) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

208

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)}$$

209

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)}$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

214

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$

In the same way , one can obtain

$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\,((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\hat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\hat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\hat{A}_{36})^{(7)}+(\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\hat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0, (\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0, (\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)}G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)}G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)}T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(\nu_1)^{(6)}, (\nu_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, \nu^{(6)}, u^{(6)}$: 350

By $(\nu_1)^{(6)} > 0$, $(\nu_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)},, (\bar{\nu}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad (354)$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad (355)$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad (356)$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)},, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

$$\Big(\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t}\Big)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)}+(C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21}, t)\right) - (a_{21}'')^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain

412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)}=\frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

423

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case,

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

424

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(dd) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523

A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FIFTY NINE

Quantum Feedback Control: Multum In Parvo

INTRODUCTION—VARIABLES USED

2003 Annual Report on the Research Program Measurement and Control in Quantum Computing.

Program Manager
A/Prof Howard Wiseman, GU

Measurement and Control Researchers
Students: Mr Neil Oxtoby (PhD), Mr Kenneth Pregnell (PhD), Ms Laura Thomsen (PhD), Mr Daniel Atkins (PhD), Mr Joshua Combes (Honours)
Staff: Prof David Pegg, Dr Kurt Jacobs, Mr Jay Gambetta, Dr He-Bi Sun (UQ)

Collaborating Centre Researchers
University of Queensland: Prof Gerard Milburn, A/Prof Tim Ralph

Other Collaborators
Macquarie University / University of Queensland: Dr Stephen Bartlett
University of Hertfordshire: Dr John Vaccaro
California Institute of Technology: A/Prof Hideo Mabuchi, Ms Charlene Ahn, Mr Asa Hopkins
Los Alamos National Laboratory: Dr Salman Habib, Dr Tanmoy Bhattacharya, Dr Daniel Steck
University of Maryland: Dr Keith Schwab

Program Description
The aim of this program is to understand the measurement and control of devices operating on a quantum scale and to apply this understanding to quantum information processing. The program has four sub-programs: Quantum Measurement Theory for Read-Out Devices; Quantum Feedback Control; Quantum Information -- Measurement Interface; Quantum Computing -- Measurement Interface.

Quantum Measurement Theory for Read-Out Devices

(1) The aim of this sub-program is to develop methods for determining the state of a qubit (or register) conditioned on the results of monitoring the qubit using a realistic read-out device. That is, for example, the measurement record is filtered and has noise **added by the** external circuit containing the detector that couples to the qubit [Oxtoby et al, J. Phys. C **15**, 8055-64 (2003)].

(2) The main outcome for this year [Oxtoby et al, submitted] is that we have formulated a rigorous method **for deriving** a quantum trajectory (that is, the stochastic evolution for the conditioned system state) for a large class of solid-state detectors in a simplified external circuit, as shown in Figure 1.

(3) We have also investigated realistic detection in a cavity QED context**, to see its effect on** estimating an unknown parameter (the position of the atom) from the detection record [Warszawski et al, submitted]. This is relevant for quantum computing in that domain.

(4) Figure 1. Circuit diagram for a "realistic" read-out device. The read-out on which the state of the CQD (coupled quantum dots) qubit is conditioned **is the** current through the Ammeter A. This represents the output of a Current Amplifier which adds noise as it amplifies. The bias voltage also contains noise. The Detector could be a DC SET (Single Electron Transistor) or a QPC (Quantum Point Contact).

Quantum Feedback Control

(5) The aim of this sub-program is to investigate feedback control of quantum systems **useful for** quantum information processing, such as error correction of quantum registers, or cooling a system down to its lowest two quantum states.

(6) This year we have published a paper establishing a protocol **to protect(e)** a quantum register **against** detected errors using only a single additional qubit [Ahn et al, Physical Review A **67**, 052310 (2003)].

(7) Work is underway to generalize this for more than one type of detected error per qubit. In addition,

we have considered motional cooling **to (e)close** to the ground state in both a cavity QED system [Steck et al, submitted] and a nanomechanical system (see Figure 2) [Hopkins et al, SPIE **5276** (2003)].

Figure 2. Diagram for the cooling of a nanomechanical resonator close to its ground state using feedback based on continuous monitoring by an SET. This device is currently being built at the Laboratory for Physical Sciences, University of Maryland.

Quantum Information -- Measurement Interface

(8) The aim of this sub-program is to study questions at the interface of quantum information and quantum measurement. The main outcome for this year has been (i) to generalize the concept of super-selection rules (SSRs) **to describe** a large class of restrictions on measurements and other quantum operations

(9) Physical Review Letters **91**, 097903 (2003)] is significant for quantum information processing (QIP) in systems subject to SSRs, such as Bose-Einstein condensation, or NMR ensemble quantum computing. Another outcome has been **to derive** fundamental inequalities **relating to** information extraction and the purification of conditioned states in measurements on a quantum system drawn randomly from an ensemble [Jacobs, Physical Review A **68**, 054302 (2003)].

NOTATION

Module One

The aim of this sub-program is to develop methods for determining the state of a qubit (or register) conditioned on the results of monitoring the qubit using a realistic read-out device. That is, for example, the measurement record is filtered and has noise **added by the** external circuit containing the detector that couples to the qubit [Oxtoby et al, J. Phys. C **15**, 8055-64 (2003)].

G_{13} : Category one of determining the state of a qubit (or register) conditioned on the results of monitoring the qubit using a realistic read-out device. That is, for example, the measurement record is filtered and has noise

G_{14} : Category two of determining the state of a qubit (or register) conditioned on the results of monitoring the qubit using a realistic read-out device. That is, for example, the measurement record is filtered and has noise

G_{15} : Category three of determining the state of a qubit (or register) conditioned on the results of monitoring the qubit using a realistic read-out device. That is, for example, the measurement record is filtered and has noise

T_{13} : Category one of external circuit containing the detector that couples to the qubit [Oxtoby et al, J. Phys. C **15**, 8055-64 (2003)].

T_{14} : Category two of external circuit containing the detector that couples to the qubit [Oxtoby et al, J. Phys. C **15**, 8055-64 (2003)].

T_{15} : Category three of external circuit containing the detector that couples to the qubit [Oxtoby et al, J. Phys. C **15**, 8055-64 (2003)].

Module Two

The main outcome for this year [Oxtoby et al, submitted] is that we have formulation of a rigorous method

for deriving a quantum trajectory (that is, the stochastic evolution for the conditioned system state) for a large class of solid-state detectors in a simplified external circuit, as shown in Figure 1. Figure deleted for spatial constraints

G_{16} : Category one of formulation of a rigorous method

G_{17} : Category two of formulation of a rigorous method

G_{18} : Category three of formulation of a rigorous method

T_{16} : Category one of quantum trajectory (that is, the stochastic evolution for the conditioned system state) for a large class of solid-state detectors in a simplified external circuit

T_{17} : Category two of quantum trajectory (that is, the stochastic evolution for the conditioned system state) for a large class of solid-state detectors in a simplified external circuit

T_{18} : Category three of quantum trajectory (that is, the stochastic evolution for the conditioned system state) for a large class of solid-state detectors in a simplified external circuit

Module three

We have also investigated realistic detection in a cavity QED context**, to see its effect on** estimating an unknown parameter (the position of the atom) from the detection record [Warszawski et al, submitted]. This is relevant for quantum computing in that domain.

G_{20} : Category one of realistic detection in a cavity QED context; estimating an unknown parameter (the position of the atom) from the detection record [Warszawski et al, submitted]. This is relevant for quantum computing in that domain.

G_{21} : Category two of realistic detection in a cavity QED context; estimating an unknown parameter (the position of the atom) from the detection record [Warszawski et al, submitted]. This is relevant for quantum computing in that domain.

G_{22} : Category three of realistic detection in a cavity QED context; estimating an unknown parameter (the position of the atom) from the detection record [Warszawski et al, submitted]. This is relevant for quantum computing in that domain.

T_{20} : Category one of estimating an unknown parameter (the position of the atom) from the detection record [Warszawski et al, submitted]. This is relevant for quantum computing in that domain.; realistic detection in a cavity QED context

T_{21} : Category two of estimating an unknown parameter (the position of the atom) from the detection record [Warszawski et al, submitted]. This is relevant for quantum computing in that domain.; realistic detection in a cavity QED context

T_{22} : Category three of estimating an unknown parameter (the position of the atom) from the detection record [Warszawski et al, submitted]. This is relevant for quantum computing in that domain.; realistic detection in a cavity QED context

Module four

The read-out on which the state of the CQD (coupled quantum dots) qubit is conditioned **is the** current through the Ammeter A. This represents the output of a Current Amplifier which adds noise as it amplifies. The bias voltage also contains noise. The Detector could be a DC SET (Single Electron Transistor) or a QPC (Quantum Point Contact).

G_{24} : Category one of current through the Ammeter A. This represents the output of a Current Amplifier which adds noise as it amplifies. The bias voltage also contains noise. The Detector could be a DC SET

(Single Electron Transistor) or a QPC (Quantum Point Contact).

G_{25} : Category two of current through the Ammeter A. This represents the output of a Current Amplifier which adds noise as it amplifies. The bias voltage also contains noise. The Detector could be a DC SET (Single Electron Transistor) or a QPC (Quantum Point Contact).

G_{26} : Category three of current through the Ammeter A. This represents the output of a Current Amplifier which adds noise as it amplifies. The bias voltage also contains noise. The Detector could be a DC SET (Single Electron Transistor) or a QPC (Quantum Point Contact).

T_{24} : Category one of read-out on which the state of the CQD (coupled quantum dots) qubit is conditioned

T_{25} : Category two of read-out on which the state of the CQD (coupled quantum dots) qubit is conditioned

T_{26} : Category three of read-out on which the state of the CQD (coupled quantum dots) qubit is conditioned

Module five

The aim of this sub-program is to investigate feedback control of quantum systems **useful for** quantum information processing, such as error correction of quantum registers, or cooling a system down to its lowest two quantum states.

G_{28} : Category one of quantum information processing, such as error correction of quantum registers, or cooling a system down to its lowest two quantum states.

G_{29} : Category two of quantum information processing, such as error correction of quantum registers, or cooling a system down to its lowest two quantum states.

G_{30} : Category three of quantum information processing, such as error correction of quantum registers, or cooling a system down to its lowest two quantum states.

T_{28} : Category one of feedback control of quantum systems

T_{29} : Category two of feedback control of quantum systems

T_{30} : Category three of feedback control of quantum systems

Module six

This year we have published a paper establishing a protocol **to protect(e)** a quantum register **against** detected errors using only a single additional qubit [Ahn et al, Physical Review A **67**, 052310 (2003)].

G_{32} : Category one of detected errors using only a single additional qubit

G_{33} : Category two of detected errors using only a single additional qubit

G_{34} : Category three of detected errors using only a single additional qubit

T_{32} : Category one of protocol **to protect(e)** a quantum register

T_{33} : Category two of protocol **to protect(e)** a quantum register

T_{34} : Category three of protocol **to protect(e)** a quantum register

Module seven

Work is underway to generalize this for more than one type of detected error per qubit. In addition, we have considered motional cooling **to (e)close** to the ground state in both a cavity QED system [Steck et al, submitted] and a nanomechanical system (see Figure 2) [Hopkins et al, SPIE **5276** (2003)].

G_{36} : Category one of ground state in both a cavity QED system and a nanomechanical system

G_{37} : Category two of ground state in both a cavity QED system and a nanomechanical system

G_{38} : Category three of ground state in both a cavity QED system and a nanomechanical system

T_{36} : Category one of motional cooling

T_{37} : Category two of motional cooling

T_{38} : Category three of motional cooling

Module eight

The aim of this sub-program is to study questions at the interface of quantum information and quantum measurement. The main outcome for this year has been (i) to generalization and the conceptualization of super-selection rules (SSRs) **to describe** a large class of restrictions on measurements and other quantum operations

G_{40} : Category one of generalization and the conceptualization of super-selection rules (SSRs)

G_{41} : Category two of generalization and the conceptualization of super-selection rules (SSRs)

G_{42} : Category three of generalization and the conceptualization of super-selection rules (SSRs)

T_{40} : Category one of large class of restrictions on measurements and other quantum operations

T_{41} : Category two of large class of restrictions on measurements and other quantum operations

T_{42} : Category three of large class of restrictions on measurements and other quantum operations

Module Nine

Physical Review Letters **91**, 097903 (2003)] is significant for quantum information processing (QIP) in systems subject to SSRs, such as Bose-Einstein condensation, or NMR ensemble quantum computing. Another outcome has been **to derive** fundamental inequalities **relating to** information extraction and the purification of conditioned states in measurements on a quantum system drawn randomly from an ensemble [Jacobs, Physical Review A **68**, 054302 (2003)].

G_{44} : Category one of Physical Review Letters **91**, 097903 (2003)] is significant for quantum information processing (QIP) in systems subject to SSRs, such as Bose-Einstein condensation, or NMR ensemble quantum computing. Another outcome has been **to derive** fundamental inequalities

G_{45} : Category two of Physical Review Letters **91**, 097903 (2003)] is significant for quantum information processing (QIP) in systems subject to SSRs, such as Bose-Einstein condensation, or NMR ensemble quantum computing. Another outcome has been **to derive** fundamental inequalities

G_{46} : Category three of Physical Review Letters **91**, 097903 (2003)] is significant for quantum information

processing (QIP) in systems subject to SSRs, such as Bose-Einstein condensation, or NMR ensemble quantum computing. Another outcome has been **to derive** fundamental inequalities

T_{44} : Category one of information extraction and the purification of conditioned states in measurements on a quantum system drawn randomly from an ensemble [Jacobs, Physical Review A **68**, 054302 (2003)].

T_{45} : Category two of information extraction and the purification of conditioned states in measurements on a quantum system drawn randomly from an ensemble [Jacobs, Physical Review A **68**, 054302 (2003)].

T_{46} : Category three of information extraction and the purification of conditioned states in measurements on a quantum system drawn randomly from an ensemble [Jacobs, Physical Review A **68**, 054302 (2003)].

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients
Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27}), t\big)\right]T_{25} \quad 23$$

$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26}$ 24

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28}$ 25

$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29}$ 26

$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30}$ 27

$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28}$ 28

$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29}$ 29

$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30}$ 30

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32}$ 31

$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33}$ 32

$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34}$ 33

$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32}$ 34

$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33}$ 35

$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34}$ 36

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36}$ 37

$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37}$ *38*

$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38}$ *39*

$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36}$ *40*

$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37}$ *41*

$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38}$ *42*

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40}$ *43*

$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41}$ *44*

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category

1,2 and 3

$-(b_{20}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b_{36}'')^{(7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b_{40}'')^{(8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b_{44}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c}(a_{20}')^{(3)}+(a_{20}'')^{(3)}(T_{21},t)+(a_{16}'')^{(2,2,2)}(T_{17},t)+(a_{13}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{36}'')^{(7,7,7,7)}(T_{37},t)+(a_{40}'')^{(8,8,8,8)}(T_{41},t)+(a_{44}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c}(a_{21}')^{(3)}+(a_{21}'')^{(3)}(T_{21},t)+(a_{17}'')^{(2,2,2)}(T_{17},t)+(a_{14}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{37}'')^{(7,7,7,7)}(T_{37},t)+(a_{41}'')^{(8,8,8,8)}(T_{41},t)+(a_{45}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c}(a_{22}')^{(3)}+(a_{22}'')^{(3)}(T_{21},t)+(a_{18}'')^{(2,2,2)}(T_{17},t)+(a_{15}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{38}'')^{(7,7,7,7)}(T_{37},t)+(a_{42}'')^{(8,8,8,8)}(T_{41},t)+(a_{46}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{22} \quad (69)$$

$+(a_{20}'')^{(3)}(T_{21},t)$, $+(a_{21}'')^{(3)}(T_{21},t)$, $+(a_{22}'')^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation

coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$ *are fourth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$ *are fifth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$ *are sixth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$ *are seventh augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$ *are eighth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b_{32}'')^{(6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b_{13}'')^{(1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b_{16}'')^{(2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b_{20}'')^{(3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b_{46}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a_{28}')^{(5)} & +(a_{28}'')^{(5)}(T_{29},t) & +(a_{24}'')^{(4,4,)}(T_{25},t) & +(a_{32}'')^{(6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1)}(T_{14},t) & +(a_{16}'')^{(2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3)}(T_{21},t) \\ & +(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a_{29}')^{(5)} & +(a_{29}'')^{(5)}(T_{29},t) & +(a_{25}'')^{(4,4,)}(T_{25},t) & +(a_{33}'')^{(6,6,6)}(T_{33},t) \\ & +(a_{14}'')^{(1,1,1,1,1)}(T_{14},t) & +(a_{17}'')^{(2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3)}(T_{21},t) \\ & +(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a_{30}')^{(5)} & +(a_{30}'')^{(5)}(T_{29},t) & +(a_{26}'')^{(4,4,)}(T_{25},t) & +(a_{34}'')^{(6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1)}(T_{14},t) & +(a_{18}'')^{(2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3)}(T_{21},t) \\ & +(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a_{28}'')^{(5)}(T_{29},t)$, $+(a_{29}'')^{(5)}(T_{29},t)$, $+(a_{30}'')^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a_{24}'')^{(4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a_{32}'')^{(6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad (85)$$

$$- \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)}\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad (87)$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad (90)$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b_{28}'')^{(5,5,5)}(G_{31},t)}, \boxed{-(b_{29}'')^{(5,5,5)}(G_{31},t)}, \boxed{-(b_{30}'')^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b_{24}'')^{(4,4,4,)}(G_{27},t)}, \boxed{-(b_{25}'')^{(4,4,4,)}(G_{27},t)}, \boxed{-(b_{26}'')^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b_{45}'')^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b_{44}'')^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)}\boxed{+(a_{36}'')^{(7)}(T_{37},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)}\boxed{+(a_{37}'')^{(7)}(T_{37},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)}\boxed{+(a_{38}'')^{(7)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a_{36}'')^{(7)}(T_{37},t)}, \boxed{(a_{37}'')^{(7)}(T_{37},t)}, \boxed{(a_{38}'')^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\,-(b''_{36})^{(7)}(G_{39},t) & -(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\,-(b''_{37})^{(7)}(G_{39},t) & -(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\,-(b''_{38})^{(7)}(G_{39},t) & -(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)}$ 103

$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)}$ 104

$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)}$ 105

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t}$ 107

$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t}$ 108

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1$ 110

$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1$ 111

Where we suppose

$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22$ 112

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(YYYYYYY) $\quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(ZZZZZZZ) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$ 132

(AAAAAAAA) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(BBBBBBBB)

$\lim_{G\to\infty}(b_i'')^{(7)}((G_{39}),t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}((G_{39}),t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(CCCCCCCC) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(DDDDDDDD) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad (169)$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that (170)

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that (172)

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that (174)

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(ee) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \le (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : \left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \le \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on G(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(3)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \le \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \le \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \ge (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \ge \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \ge \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$
Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)}, , (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad (333)$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad (334)$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad (335)$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad (336)$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(ee) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)} G_{14}^{*}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)} G_{14}^{*}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)} T_{14}^{*}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)} T_{14}^{*}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)} G_{17}^{*}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)} G_{17}^{*}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)} T_{17}^{*}}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)} T_{17}^{*}}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)} G_{21}^{*}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)} G_{21}^{*}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)} T_{21}^{*}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)} T_{21}^{*}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SIXTY

Measurement And Control For Quantum Computing: Affirmanti Non Neganti Incumbit Probatio

INTRODUCTION—VARIABLES USED

New qubit control bodes well for future of quantum computing By Eric Gershon January 11, 2013

Yale University scientists have found a way to observe quantum information while preserving its integrity, an achievement that offers researchers greater control in the volatile realm of quantum mechanics and greatly improves the prospects of quantum computing.

Quantum computers would be exponentially faster than the most powerful computers of today.

“Our experiment is a dress rehearsal for a type of process essential for quantum computing,” said Michel Devoret, the Frederick William Beinecke Professor of Applied Physics & Physics at Yale and principal investigator of research published Jan. 11 in the journal Science. “What this experiment really allows is an active understanding of quantum mechanics. It’s one thing to stare at a theoretical formula and it’s another thing to be able to control a real quantum object.”

In quantum systems, microscopic units called qubits represent information. Qubits can assume either of two states — “0” or “1” — or both simultaneously. Correctly recognizing, interpreting, and tracking their state is necessary for quantum computing. However, the act of monitoring them usually damages their information content.

The Yale physicists successfully devised a new, non-destructive measurement system for observing, tracking and documenting all changes in a qubit’s state, thus preserving the qubit’s informational value. In principle, the scientists said, this should allow them to monitor the qubit’s state in order to correct for random errors.

“As long as you know what error process has occurred, you can correct,” Devoret said. “And then everything’s fine. You can basically undo the errors.”

An innovation by Yale University physicists offers scientists greater control in the volatile realm of quantum mechanics and greatly improves the prospects of quantum computing. Quantum computers would be exponentially faster than the most powerful computers of today.

“That’s the key,” said Michael Hatridge, a postdoctoral associate in physics at Yale and lead author of the Science paper, “the ability to talk to the qubit and hear what it’s telling you.”

He continued: "A major problem with quantum computing is the finite lifetime of information stored in the qubits, which steadily decays and which must be corrected. We now know that it is possible to do this correction by feedback involving a continuous measurement. Our work advances the prospects of large-scale quantum computers by opening the door to continuous measurement-based quantum feedback."

The Yale physicists successfully measured one qubit. The challenge ahead is to measure and control many at once, and the team is developing ultra-fast digital electronics for this purpose.

"We are on the threshold between the ability to measure and control one or two qubits, and many," Hatridge said.

Other authors of the paper are S. Shankar, M. Mirrahimi, F. Schackert, K. Geerlings, T. Brecht, K.M. Sliwa, B. Abdo, L. Frunzio, S.M. Girvin, and R.J. Schoelkopf.

Support for the research was provided by the National Science Foundation, the United States Army Research Office, the Intelligence Advanced Research Projects Activity, the Agency National de Recherche, and the College de France.

The Griffith Program: Measurement and Control for Quantum Computing. Quantum Measurement Theory for Read-Out Devices

(1) Quantum Information - Measurement Interface: Dissipation is the transfer of energy from the system to the environment. It is now well understood that what is more relevant for quantum computers **is** decoherence, the transfer of information from the system to the environment.
(2) This information is quantum information, **but it can be turned** into classical information by measuring the environment. This new perspective on quantum measurement theory is a fruitful field for investigation.
(3) Quantum Computing - Measurement Interface: This part of the program will seek new schemes for quantum measurement as they may be useful for quantum computing. On the flip-side, quantum algorithms **should also suggest** new measurement schemes.
(4) Thirdly, we have examined the question of the validity of optical continuous variable quantum teleportation, **and concluded that** a laser should be regarded as a clock, not a quantum channel [Wiseman, J. Opt. B (to be published)]. Finally, we have investigated the preparation of spin-squeezed states for the purposes of interferometric measurement of phase shifts [Combes et al, unpublished]. We find that the maximally spin squeezed state (shown in Figure 3: deleted due to spatial constraints) **is not the** optimal state for this purpose, and we identify a simple scheme to produce the optimal state.
(5) Figure 3. Wigner function for the maximally spin squeezed state of an ensemble of 20 atoms, plotted as a function of polar co-ordinates for the spin direction. Note the "ripples" (positive and negative fluctuations) in the tails of the distribution. These make this state sub-optimal for single-shot measurement of the shift in the phase f (which could be produced by a magnetic field, for example)

Quantum Information -- Computing Interface

(6) The aim of this sub-program is to study questions at the interface of quantum information and quantum computing. The main outcome for this year has been the discovery of a general scheme to retrodict quantum states in optics **using only** coherent states and beam splitters [Pregnell and Pegg, J. Mod. Opt. (to be published)].
(7) That is, it is possible to engineer an optical device such that one particular measurement result corresponds to a given (arbitrary) element in a positive operator measure (POM).

(8) Authors have also discovered a simpler scheme for the specific problem where the POM is that (e) arising from measurement of the Pegg-Barnett phase operator [Pregnell and Pegg, Physical Review A 67, 063814 (2003)].

(9) This scheme (shown in Figure 4: deleted due to spatial constraints) **requires a** binomial input state as a reference field, but we show that this can be approximated by a squeezed state for low photon numbers.

NOTATION

Module One

Quantum Information - Measurement Interface: Dissipation is the transfer of energy from the system to the environment. It is now well understood that what is more relevant for quantum computers **is** decoherence, the transfer of information from the system to the environment

G_{13} : Category one of decoherence, the transfer of information from the system to the environment

G_{14} : Category two of decoherence, the transfer of information from the system to the environment

G_{15} : Category three of decoherence, the transfer of information from the system to the environment

T_{13} : Category one of Dissipation is the transfer of energy from the system to the environment. It is now well understood that what is more relevant for quantum computers

T_{14} : Category two of Dissipation is the transfer of energy from the system to the environment. It is now well understood that what is more relevant for quantum computers

T_{15} : Category three of Dissipation is the transfer of energy from the system to the environment. It is now well understood that what is more relevant for quantum computers

Module Two

This information is quantum information, **but it can be turned** into classical information by measuring the environment. This new perspective on quantum measurement theory is a fruitful field for investigation.

G_{16} : Category one of information which is quantum information

G_{17} : Category two of information which is quantum information

G_{18} : Category three of information which is quantum information

T_{16} : Category one of classical information by measuring the environment.

T_{17} : Category two of classical information by measuring the environment.

T_{18} : Category three of classical information by measuring the environment.

Module three

Quantum Computing - Measurement Interface: This part of the program will seek new schemes for quantum measurement as they may be useful for quantum computing. On the flip-side, quantum algorithms **should also suggest** new measurement schemes

G_{20} : Category one of quantum algorithms

G_{21} : Category two of quantum algorithms

G_{22} : Category three of quantum algorithms

T_{20} : Category one of new measurement schemes

T_{21} : Category two of new measurement schemes

T_{22} : Category three of new measurement schemes

Module four

Thirdly, we have examined the question of the validity of optical continuous variable quantum teleportation, **and concluded that** a laser should be regarded as a clock, not a quantum channel [Wiseman, J. Opt. B (to be published)]. Finally, we have investigated the preparation of spin-squeezed states for the purposes of interferometric measurement of phase shifts [Combes et al, unpublished]. We find that the maximally spin squeezed state (shown in Figure 3: deleted due to spatial constraints) **is not the** optimal state for this purpose, and we identify a simple scheme to produce the optimal state.

G_{24} : Category one of validity of optical continuous variable quantum teleportation

G_{25} : Category two of validity of optical continuous variable quantum teleportation

G_{26} : Category three of validity of optical continuous variable quantum teleportation

T_{24} : Category one of laser should be regarded as a clock, not a quantum channel

T_{25} : Category two of laser should be regarded as a clock, not a quantum channel

T_{26} : Category three of laser should be regarded as a clock, not a quantum channel

Module five

G_{28} : Category one of clock optimal state for this purpose, and author identifies a simple scheme to produce the optimal state.

G_{29} : Category two of clock optimal state for this purpose, and authors identify a simple scheme to produce the optimal state.

G_{30} : Category three of clock optimal state for this purpose, and authors identify a simple scheme to produce the optimal state.

T_{28} : Category one of laser maximally spin squeezed state (shown in Figure 3: deleted due to spatial constraints)

T_{29} : Category two of laser maximally spin squeezed state (shown in Figure 3: deleted due to spatial constraints)

T_{30} : Category three of laser maximally spin squeezed state (shown in Figure 3: deleted due to spatial constraints)

Module six

Wigner function for the maximally spin squeezed state of an ensemble of 20 atoms, plotted as a function of polar co-ordinates for the spin direction; "ripples" (positive and negative fluctuations) in the tails of the distribution **which(eb) make** this state sub-optimal for single-shot measurement of the shift in the phase f (which could be produced by a magnetic field, for example)

G_{32} : Category one of Wigner function for the maximally spin squeezed state of an ensemble of 20 atoms, plotted as a function of polar co-ordinates for the spin direction; "ripples" (positive and negative

fluctuations) in the tails of the distribution

G_{33} : Category two of Wigner function for the maximally spin squeezed state of an ensemble of 20 atoms, plotted as a function of polar co-ordinates for the spin direction; "ripples" (positive and negative fluctuations) in the tails of the distribution

G_{34} : Category three of Wigner function for the maximally spin squeezed state of an ensemble of 20 atoms, plotted as a function of polar co-ordinates for the spin direction; "ripples" (positive and negative fluctuations) in the tails of the distribution

T_{32} : Category one of state sub-optimal for single-shot measurement of the shift in the phase f (which could be produced by a magnetic field, for example)

T_{33} : Category two of state sub-optimal for single-shot measurement of the shift in the phase f (which could be produced by a magnetic field, for example)

T_{34} : Category three of state sub-optimal for single-shot measurement of the shift in the phase f (which could be produced by a magnetic field, for example)

Module seven

The aim of this sub-program is to study questions at the interface of quantum information and quantum computing. The main outcome for this year has been the discovery of a general scheme to retrodict quantum states in optics **using only** coherent states and beam splitters [Pregnell and Pegg, J. Mod. Opt. (to be published)].

G_{36} : Category one of coherent states and beam splitters [Pregnell and Pegg, J. Mod. Opt. (to be published)].

G_{37} : Category two of coherent states and beam splitters [Pregnell and Pegg, J. Mod. Opt. (to be published)].

G_{38} : Category three of coherent states and beam splitters [Pregnell and Pegg, J. Mod. Opt. (to be published)].

T_{36} : Category one of general scheme to retrodict quantum states in optics

T_{37} : Category two of general scheme to retrodict quantum states in optics

T_{38} : Category three of general scheme to retrodict quantum states in optics

Module eight

Authors have also discovered a simpler scheme for the specific problem where the POM is that (e) arising from measurement of the Pegg-Barnett phase operator [Pregnell and Pegg, Physical Review A 67, 063814 (2003)].

G_{40} : Category one of arising from measurement of the Pegg-Barnett phase operator [Pregnell and Pegg, Physical Review A 67, 063814 (2003)].

G_{41} : Category two of arising from measurement of the Pegg-Barnett phase operator [Pregnell and Pegg, Physical Review A 67, 063814 (2003)].

G_{42} : Category three of arising from measurement of the Pegg-Barnett phase operator [Pregnell and Pegg, Physical Review A 67, 063814 (2003)].

T_{40} : Category one of specific problem where the POM

T_{41} : Category two of specific problem where the POM

T_{42} : Category three of specific problem where the POM

Module Nine

This scheme (shown in Figure 4: deleted due to spatial constraints) **requires a** binomial input state as a reference field, but we show that this can be approximated by a squeezed state for low photon numbers.

G_{44} : Category one of binomial input state as a reference field, but we show that this can be approximated by a squeezed state for low photon numbers

G_{45} : Category two of binomial input state as a reference field, but we show that this can be approximated by a squeezed state for low photon numbers

G_{46} : Category three of binomial input state as a reference field, but we show that this can be approximated by a squeezed state for low photon numbers

T_{44} : Category one of scheme for the specific problem where the POM is that (e) arising from measurement of the Pegg-Barnett phase operator [Pregnell and Pegg, Physical Review A 67, 063814 (2003)].

T_{45} : Category two of scheme for the specific problem where the POM is that (e) arising from measurement of the Pegg-Barnett phase operator [Pregnell and Pegg, Physical Review A 67, 063814 (2003)].

T_{46} : Category three of scheme for the specific problem where the POM is that (e) arising from measurement of the Pegg-Barnett phase operator [Pregnell and Pegg, Physical Review A 67, 063814 (2003)].

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$

$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation

coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\left[-(b''_{13})^{(1)}(G,t)\right] & \left[-(b''_{16})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\left[-(b''_{14})^{(1)}(G,t)\right] & \left[-(b''_{17})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\left[-(b''_{15})^{(1)}(G,t)\right] & \left[-(b''_{18})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad 73$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad 74$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad 76$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)} & \boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category $1, 2$ *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b_{36}')^{(7)}\boxed{-(b_{36}'')^{(7)}(G_{39},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b_{37}')^{(7)}\boxed{-(b_{37}'')^{(7)}(G_{39},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b_{38}')^{(7)}\boxed{-(b_{38}'')^{(7)}(G_{39},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$

$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)}$ 98

$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$

$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$\frac{1}{(\hat{M}_{13})^{(1)}}[\,(a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\,(\hat{k}_{13})^{(1)}] < 1$

$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)}$ 102

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(EEEEEEEE) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(FFFFFFFF) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(GGGGGGGG) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(HHHHHHHH)

$\lim_{G\to\infty}(b_i'')^{(7)}\big((G_{39}),t\big) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\big((G_{39}),t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(IIIIIIII) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(JJJJJJJJ) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad 146\ A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(ff) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $\left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on G(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right| \; e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$$
$$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$$

$$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$$

In the same way , one can obtain

$$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right) - (a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right| \; e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} \} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)} - G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)};\ (G_{23})^{(2)}, (T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(3)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)} \left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)} \left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(6)}$ and $(b''_i)^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(6)} - (a''_i)^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$
$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{36})^{(7)}t}\}$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t} \; and \; (\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2 \; and \; \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37} \; and \; G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(9)}t)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \le \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \le \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \ge (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \ge \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \ge \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \qquad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \qquad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \qquad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(ff) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SIXTY ONE

Quantum Control Engineering: Omnia Causa Fiunt

INTRODUCTION—VARIABLES USED

(1) In quantum mechanics, a complete set of commuting observables (CSCO) **is a** set of commuting operators whose eigenvalues completely specify the state of a system (Gasiorowicz 1974, p. 119).

(2) For example, in the case of the hydrogen atom, the Hamiltonian H, the angular momentum L and its projection Lz along any arbitrary z axis form a CSCO (if one ignores the spin of the proton and of the electron as well as the movement of the center of mass). Since each pair of observables in the set commutes, the observables are all compatible so that the measurement of one observable **has no** effect on the result of measuring another observable in the set.

(3) It is therefore not necessary **to (EB) specify** the order in which the different observables are measured.

(4) Measurement of the complete set of observables **constitutes (EB) a** complete measurement, in the sense that it projects the quantum state of the system onto a unique and known vector in the basis defined by the set of operators.

(5) This specifies the conditions for the preparation of a well-defined quantum state. Also, the states which form the basis of the CSCO are **completely (E) defined by** specifying the eigenvalues λ_1, λ_2, ..., associated with each observable O_1, O_2, ... of the CSCO. In the above example of the hydrogen atom, the basis states are defined by the quantum numbers $|n, l, m\rangle$, and it is not necessary to include the wave function or the atomic orbital.

(6) In quantum physics, the relation between system state and the value of an observable **requires** some basic linear algebra for its description.

(7) In the mathematical formulation of quantum mechanics, states are given by non-zero vectors in a Hilbert space V (where two vectors are considered to specify the same state if, and only if, they are scalar multiples of each other) and observables are given by self-adjoint operators on V. However, as indicated below, not every self-adjoint operator **corresponds to** a physically meaningful observable.

(8) For the case of a system of particles, the space V **consists of** functions called wave functions or state vectors. In the case of transformation laws in quantum mechanics, the requisite automorphisms are unitary (or antiunitary) linear transformations of the Hilbert space V.

(9) Under Galilean relativity or special relativity, the mathematics of frames of reference is particularly simple, and in fact restricts considerably the set of physically meaningful observables.

(10) In quantum mechanics, measurement of observables **exhibits** some seemingly unintuitive properties.

(11) Specifically, if a system is in a state **described by a** vector in a Hilbert space, the measurement process affects the state in a non-deterministic, but statistically predictable way.

(12) In particular, after a measurement is applied, the state description by a single vector may be destroyed, **being replaced by** a statistical ensemble.

(13) The irreversible nature of measurement operations in quantum physics is sometimes **referred to as the** measurement problem and is described mathematically by quantum operations. By the structure of quantum operations, this description is mathematically **equivalent(=) to that** offered by relative state interpretation where the original system is regarded as a subsystem of a larger system and the

state of the original system is given by the partial trace of the state of the larger system.

(14) In quantum mechanics each dynamical variable (e.g. position, translational momentum, orbital angular momentum, spin, total angular momentum, energy, etc.) **is associated with** a Hermitian operator that acts on the state of the quantum system and whose eigenvalues correspond to the possible values of the dynamical variable.

(15) For example, suppose $|a\rangle$ is an eigenket (eigenvector) of the observable $\mathbf{A}$, with eigenvalue a, and exists in a d-dimensional Hilbert space. Then

$$\mathbf{A}|a\rangle = a|a\rangle.$$

This eigenket equation says that if a measurement of the observable $\mathbf{A}$ is made while the system of interest is in the state $|a\rangle$, then the observed value of that particular measurement **must return the** eigenvalue a with certainty.

(16) However, if the system of interest is in the general state $|\phi\rangle \in \mathcal{H}$, then the eigenvalue a is returned with probability $|\langle a|\phi\rangle|^2$ **(Born rule**). One must note that the above definition is somewhat dependent upon our convention of choosing real numbers to represent real physical quantities. Indeed, just because dynamical variables are "real" and not "unreal" in the metaphysical sense does not mean that they must correspond to real numbers in the mathematical sense.

To be more precise, the dynamical variable/observable is a (not necessarily bounded) Hermitian operator in a Hilbert Space and thus is represented by a Hermitian matrix if the space is finite-dimensional. In an infinite-dimensional Hilbert space, the observable is represented by a symmetric operator, which may not be defined everywhere (i.e. its domain is not the whole space - there exist some states that are not in the domain of the operator). The reason for such a change is that in an infinite-dimensional Hilbert space, the operator **becomes** unbounded, which means that it no longer has a largest eigenvalue. This is not the case in a finite-dimensional Hilbert space, where every operator is bounded - it has a largest eigenvalue. For example, if we consider the position of a point particle moving along a line, this particle's position variable can take on any number on the real-line, which is uncountably infinite. Since the eigenvalue of an observable represents a real physical quantity for that particular dynamical variable, then we must conclude that there is no largest eigenvalue for the position observable in this uncountably infinite-dimensional Hilbert space, since the field we're working over consists of the real-line. Nonetheless, whether we are working in an infinite-dimensional or finite-dimensional Hilbert space, the role of an observable in quantum mechanics is to assign real numbers to outcomes of *particular measurements*; this means that only certain measurements can determine the value of an observable for some state of a quantum system. In classical mechanics, *any* measurement can be made to determine the value of an observable.

Incompatibility of observables in quantum mechanics

A crucial difference between classical quantities and quantum mechanical observables is that the latter may not be simultaneously measurable. This is mathematically expressed by non-commutativity of the corresponding operators, to the effect that

$$\mathbf{AB} - \mathbf{BA} \neq \mathbf{0}.$$


This inequality expresses a dependence of measurement results **on (α) the** order in which measurements of observables $\mathbf{A}$ and $\mathbf{B}$ are performed. Observables corresponding to non-commutative operators are called *incompatible*.

QUANTUM CONTROL:: Matt James ANU

This is the html version of the file http://users.cecs.anu.edu.au/~trumpf/LSslides/Matt_James.pdf. Google automatically generates html versions of documents as we crawl the web.

- Robust control
  - H-infinity synthesis with classical, quantum or mixed classical/quantum controllers

[James, Nurdin, Petersen 2006]



- LQG [James, Nurdin, Petersen; current]

Quantum Mechanics non-commuting observables

[Q, P] = QP − PQ = ihI

(2) expectation

(Q) = ∫ q|ψ(q, t)|2dq

(3) Heisenburg uncertainty

ΔQΔP ≥12|(i[Q, P])| =h2

(4) Schrodinger equation

ih∂ψ(q, t)∂t= −h2/2m∂2ψ(q, t)∂q2+ V (q)ψ(q, t)

quantum probability

(J ,P)

Some mathematical preliminaries.

Hilbert space with inner product (·,·)

(5) We take H = Cn, n-dimensional complex vectors, (ψ,φ) = ∑nk=1ψ∗kφk.Vectors are written (Dirac's kets)

φ = |φ) ∈ H

(6) Dual vectors are called bras, ψ = (ψ| ∈ H∗ ≡ H, so that

(ψ,φ) = (ψ||φ) = (ψ|φ)

Let S (H) be the Banach space of bounded operators A : H → H.

For any A ∈ S(H) its adjoint A∗ ∈ S(H) is an operator defined by

(7) (A∗ψ, φ) = (ψ, Aφ) for all ψ, φ ∈ H.

Also define

(8) (A, B) = Tr[A∗B], A, B ∈ S(H)

(9) An operator A ∈ S (H) is called normal if AA∗ = A∗A. Two important types of normal operators are self-adjoint (A = A∗), and unitary (A∗ = A−1). The spectral theorem for a self-adjoint operator A says that (finite dimensional case) it has a finite number of real eigenvalues and that A can be written as A = ∑a∈spec (A) Pa where Pa is the projection onto the eigenspace corresponding to the eigenvalue a (diagonal representation).

NOTATION

Module One

QUANTUM CONTROL:: **Matt James ANU**

Quantum Mechanics non-commuting observables
[Q, P] = QP − PQ = ihI

Probe Model: system X interacts with probe M and gives **result or measurement** Y

G_{13} : Category one of LHS of the equation of Quantum Mechanics non-commuting observables ;interaction of system X with probe M

G_{14} : Category two of LHS of the equation of Quantum Mechanics non-commuting observables; interaction of system X with probe M

G_{15} : Category three of LHS of the equation of Quantum Mechanics non-commuting observables; interaction of system X with probe M

T_{13} : Category one of RHS of the equation of Quantum Mechanics non-commuting observables; **result or measurement** Y

T_{14} : Category two of RHS of the equation of Quantum Mechanics non-commuting observables; **result or measurement** Y

T_{15} : Category three of RHS of the equation of Quantum Mechanics non-commuting observables; **result or measurement** Y

Module Two

QUANTUM CONTROL:: **Matt James ANU**

Expectation:

$(Q) = \int q|\psi(q, t)|2dq$

Quantum physics shows an inherent statistical behaviour: The measured outcome of an experiment will generally not be the same if the experiment is repeated several times. Only the statistical mean of the measured values, averaged over a large number of runs of the experiment, is a repeatable quantity. Quantum theory does not, in fact, predict the result of individual measurements, but only their statistical mean. This predicted mean value is called the expectation value. While the computation of the mean value of experimental results is very much the same as in classical statistics, its mathematical representation in the formalism of quantum theory differs significantly from classical measure theory.

Filtering



Classical: Wiener, Kalman, Kushner, Stratonovich, Duncan, Mortensen, Zakai

$$\begin{aligned} d\hat{\xi} &= A\hat{\xi}dt + PC^T(dy - C\hat{\xi}dt) \\ \dot{P} &= AP + PA^T - PC^TCP + BB^T \end{aligned}$$

Kalman filter

Quantum: - general quantum filtering - Belavkin
- generalised measurement theory - Davis,...
- stochastic master equations - Carmichael, ...

$$d\rho_t = -\frac{i}{\hbar}[H,\rho_t]dt + \mathcal{D}[L]\rho_t dt + \mathcal{H}[L]\rho_t(dy_t - \mathrm{tr}[(L+L^\dagger)\rho_t]dt)$$

quantum filter
(stochastic master equation)

QUANTUM CONTROL:: **Matt James ANU**

G_{16} : Category one of LHS of the equation for expectation, Kallman filter and quantum filter

G_{17} : Category two of LHS of the equation for expectation Kallman filter and quantum filter

G_{18} : Category three of LHS of the equation for expectation Kallman filter and quantum filter

T_{16} : Category one of RHS of the equation for expectation Kallman filter and quantum filter

T_{17} : Category two of RHS of the equation for expectation Kallman filter and quantum filter

T_{18} : Category three of RHS of the equation for expectation Kallman filter and quantum filter

Module three

QUANTUM CONTROL:: **Matt James ANU**

Heisenburg uncertainty

$\Delta Q \Delta P \geq 12|(i[Q, P])| = h2$

Quantum control

- The quantum probability framework is well-suited to quantum control engineering
- There is a theory of quantum stochastic differential equations

$$dU_t = \left\{ LdA_t^* - L^* dA_t - \frac{1}{2} L^* L dt - iH(u(t))dt \right\} U_t, \quad U_0 = I$$

- The quantum filtering theory facilitates the design of classical controllers for quantum systems
- Not much is known about the design of quantum controllers for quantum systems

G_{20} : Category one of LHS of the Heisenburg uncertainty and quantum stochastic differential equation

G_{21} : Category two of LHS of the Heisenburg uncertainty and quantum stochastic differential equation

G_{22} : Category three of LHS of the Heisenburg uncertainty and quantum stochastic differential equation

T_{20} : Category one of RHS of the Heisenburg uncertainty and quantum stochastic differential equation

T_{21} : Category two of RHS of the Heisenburg uncertainty and quantum stochastic differential equation

T_{22} : Category three of RHS of the Heisenburg uncertainty and quantum stochastic differential equation

Module four

QUANTUM CONTROL:: **Matt James ANU**

Schrodinger equation

$ih\partial\psi(q, t)\partial t = -h2/2m\partial 2\psi(q, t)\partial q2 + V(q)\psi(q, t)$

Probe measurement generates a commutative algebra

$$\mathscr{Y} = \text{alg}\{U^*(I \otimes M)U\}$$

System observables belong to commutant:

$$U^*(X \otimes I)U \in \mathscr{Y}'$$

Therefore the conditional expectations

$$\mathbb{P}[U^*(X \otimes I)U|\mathscr{Y}]$$

are well defined.

This allows statistical estimation for system observables given measurement dat

The von Neumann "projection postulate" is a special case.

In continuous time, this leads to *quantum filtering*.

G_{24} : Category one of LHS of the Schrodinger equation; probe measurement generation of commutative algebra

G_{25} : Category two of LHS of the Schrodinger equation; probe measurement generation of commutative algebra

G_{26} : Category three of LHS of the Schrodinger equation; probe measurement generation of commutative algebra

T_{24} : Category one of RHS of the Schrodinger equation; quantum filtering

T_{25} : Category two of RHS of the Schrodinger equation; quantum filtering

T_{26} : Category three of RHS of the Schrodinger equation; quantum filtering

Module five

QUANTUM CONTROL:: Matt James ANU

We take H = Cn, n-dimensional complex vectors, (ψ,φ) = ∑nk=1ψ∗kφk.Vectors are written (Dirac's kets)

φ = |φ) ∈ H

For example, suppose $|a\rangle$ is an eigenket (eigenvector) of the observable $\mathbf{A}$, with eigenvalue a, and exists in a d-dimensional Hilbert space. Then

$$\mathbf{A}|a\rangle = a|a\rangle.$$

This eigenket equation says that if a measurement of the observable $\mathbf{A}$ is made while the system of interest is in the state $|a\rangle$, then the observed value of that particular measurement **must return the** eigenvalue a with

certainty.

) Dual vectors are called bras, $\psi = (\psi| \in H* \equiv H$, so that

$(\psi,\varphi) = (\psi||\varphi) = (\psi|\varphi)$

Let S (H) be the Banach space of bounded operators A : H → H.

Some of our recent and current work in quantum control

- Optimal quantum feedback control
  - risk-sensitive [James, 2004, 2005]

$$J^{\mu}(\mathrm{K}) = \mathbb{P}[R^*(T)e^{\mu C_2(T)}R(T)]$$

$$\frac{dR(t)}{dt} = \frac{\mu}{2}C_1(t)R(t), \quad R(0) = I$$

$$\mathrm{K}^{\mu,\star} : \begin{cases} d\sigma_t^{\mu}(X) &= \sigma_t^{\mu}\big(\mathcal{L}^{\mu,u(t)}(X)\big)dt + \sigma_t^{\mu}(L^*X + XL)dY_t \\ u(t) &= \mathrm{u}^{\mu,\star}(\sigma_t^{\mu}, t) \end{cases}$$

[separation structure]

  - QLEQG [D'Helon, Doherty, James, Wilson 2006]

[QLEQG ≠ CLEQG]

QUANTUM CONTROL:: **Matt James ANU**

G_{28} : Category one of LHS of the above equations and that of optimal feedback quantum control(James: two equations)

G_{29} : Category two of LHS of the above equations; optimal feedback quantum control(James: two equations)

G_{30} : Category three of LHS of the above equations; optimal feedback quantum control(James: two equations)

T_{28} : Category one of RHS of the above equations; optimal feedback quantum control(James: two equations)

T_{29} : Category two of RHS of the above equations; optimal feedback quantum control(James: two equations)

T_{30} : Category three of RHS of the above equations; optimal feedback quantum control(James: two equations)

Module six

QUANTUM CONTROL:: **Matt James ANU**

Incompatibility of observables in quantum mechanics

A crucial difference between classical quantities and quantum mechanical observables is that the latter may not be simultaneously measurable. This is mathematically expressed by non-commutativity of the corresponding operators, to the effect that

$$\mathbf{AB} - \mathbf{BA} \neq \mathbf{0}.$$

This inequality expresses a dependence of measurement results **on (α) the** order in which measurements of observables $\mathbf{A}$ and $\mathbf{B}$ are performed. Observables corresponding to non-commutative operators are called *incompatible*. In consideration with the incompatibility and inequality of the measurement problem the only two possibilities are that AB is either added to BA or BA subtracted from AB. We deal with the inequality in the following model .This is important in the sense it tackles **"measurement problem". This means AB is eating BA orAB getting eaten up by BA**

G_{32} : Category one of AB;BA

G_{33} : Category two of AB;BA

G_{34} : Category three of AB;BA

T_{32} : Category one of BA;AB

T_{33} : Category two of BA;AB

T_{34} : Category three of BA;AB

Module seven

QUANTUM CONTROL:: **Matt James ANU**

For any A ∈ S(H) its adjoint A∗ ∈ S(H) is an operator defined by

(A∗ψ, φ) = (ψ, Aφ) for all ψ, φ ∈ H.

G_{36} : Category one of LHS of the equation for the operator

G_{37} : Category two of LHS of the equation for the operator

G_{38} : Category three of LHS of the equation for the operator

T_{36} : Category one of RHS of the equation for the operator

T_{37} : Category two of RHS of the equation for the operator

T_{38} : Category three of RHS of the equation for the operator

Module eight

QUANTUM CONTROL:: **Matt James ANU**

(A, B) = Tr[A∗B], A, B ∈ S(H)

G_{40} : Category one of LHS of the above equation for the Trace operator

G_{41} : Category two of LHS of the above equation for the Trace operator

G_{42} : Category three of LHS of the above equation for the Trace operator

T_{40} : Category one of RHS of the above equation for the Trace operator

T_{41} : Category two of RHS of the above equation for the Trace operator

T_{42} : Category three of RHS of the above equation for the Trace operator

Module Nine

QUANTUM CONTROL:: **Matt James ANU**

An operator A ∈ S (H) is called normal if AA∗ = A∗A. Two important types of normal operators are self-adjoint (A = A∗), and unitary (A∗ = A−1). The spectral theorem for a self-adjoint operator A says that (finite dimensional case) it has a finite number of real eigenvalues and that A can be written as A = ∑a∈spec (A) Pa where Pa is the projection onto the eigenspace corresponding to the eigenvalue a (diagonal representation).

G_{44} : Category one of LHS of the equation for the spectral operator with conservation and preservation of normality

G_{45} : Category two of LHS of the equation for the spectral operator with conservation and preservation of normality

G_{46} : Category three of LHS of the equation for the spectral operator with conservation and preservation of normality

T_{44} : Category one of RHS of the equation for the spectral operator with conservation and preservation of normality

T_{45} : Category two of RHS of the equation for the spectral operator with conservation and preservation of normality

T_{46} : Category three of RHS of the equation for the spectral operator with conservation and preservation of normality

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$
$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$
$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**
$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{37}'')^{(7,7,)}(G_{39},t)$, $-(b_{36}'')^{(7,7,)}(G_{39},t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8)}(G_{43},t)$ $-(b_{41}'')^{(8,8)}(G_{43},t)$ $-(b_{42}'')^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17},t) + (a_{13}'')^{(1,1,)}(T_{14},t) + (a_{20}'')^{(3,3,3)}(T_{21},t) \\ + (a_{24}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{28}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{32}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{36}'')^{(7,7,7)}(T_{37},t) + (a_{40}'')^{(8,8,8)}(T_{41},t) + (a_{44}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17},t) + (a_{14}'')^{(1,1,)}(T_{14},t) + (a_{21}'')^{(3,3,3)}(T_{21},t) \\ + (a_{25}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{29}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{33}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{37}'')^{(7,7,7)}(T_{37},t) + (a_{41}'')^{(8,8,8)}(T_{41},t) + (a_{45}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17},t) + (a_{15}'')^{(1,1,)}(T_{14},t) + (a_{22}'')^{(3,3,3)}(T_{21},t) \\ + (a_{26}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{30}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{34}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{38}'')^{(7,7,7)}(T_{37},t) + (a_{42}'')^{(8,8,8)}(T_{41},t) + (a_{46}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad 63$$

Where $+(a_{16}'')^{(2)}(T_{17},t)$, $+(a_{17}'')^{(2)}(T_{17},t)$, $+(a_{18}'')^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for*

category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category $1, 2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a_{32}')^{(6)} & +(a_{32}'')^{(6)}(T_{33},t) & +(a_{28}'')^{(5,5,5)}(T_{29},t) & +(a_{24}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a_{33}')^{(6)} & +(a_{33}'')^{(6)}(T_{33},t) & +(a_{29}'')^{(5,5,5)}(T_{29},t) & +(a_{25}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a_{34}')^{(6)} & +(a_{34}'')^{(6)}(T_{33},t) & +(a_{30}'')^{(5,5,5)}(T_{29},t) & +(a_{26}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a_{32}'')^{(6)}(T_{33},t)$, $+(a_{33}'')^{(6)}(T_{33},t)$, $+(a_{34}'')^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a_{28}'')^{(5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a_{24}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{c} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{c} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \le (r_i)^{(1)} \le (b_i')^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \le (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(KKKKKKKK) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(LLLLLLLL) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(MMMMMMMM) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(NNNNNNNN)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(OOOOOOOO) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(PPPPPPPP) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \le (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$
satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\leq\ \left(\hat{P}_{13}\right)^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\leq\ (\hat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\leq\ (\hat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\leq\ (\hat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 156

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(gg) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)} t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

208

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)}$$

209

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)}$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),\left((G_{23})^{(2)},(T_{23})^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

214

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40{,}41{,}42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 275

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(8)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\,((G_{43})(t),t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}\,(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^{0}e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^{0}e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^{0}}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t}-e^{-(b'_{34})^{(6)}t}\right]+T_{34}^{0}e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^{0}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T_{34}^{0}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)},(S_2)^{(6)},(R_1)^{(6)},(R_2)^{(6)}$:- 359

Where $(S_1)^{(6)}=(a_{32})^{(6)}(m_2)^{(6)}-(a'_{32})^{(6)}$

$(S_2)^{(6)}=(a_{34})^{(6)}-(p_{34})^{(6)}$

$(R_1)^{(6)}=(b_{32})^{(6)}(\mu_2)^{(6)}-(b'_{32})^{(6)}$

$(R_2)^{(6)}=(b'_{34})^{(6)}-(r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)},(\sigma_2)^{(7)},(\tau_1)^{(7)},(\tau_2)^{(7)}$:

$(\sigma_1)^{(7)},(\sigma_2)^{(7)},(\tau_1)^{(7)},(\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)}+(a'_{37})^{(7)}-(a''_{36})^{(7)}(T_{37},t)+(a''_{37})^{(7)}(T_{37},t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)}+(b'_{37})^{(7)}-(b''_{36})^{(7)}\big((G_{39}),t\big)-(b''_{37})^{(7)}\big((G_{39}),t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)},(v_2)^{(7)},(u_1)^{(7)},(u_2)^{(7)},v^{(7)},u^{(7)}$: 361

By $(v_1)^{(7)}>0,(v_2)^{(7)}<0$ and respectively $(u_1)^{(7)}>0,(u_2)^{(7)}<0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}=0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2+(\tau_1)^{(7)}u^{(7)}-(b_{36})^{(7)}=0$ and

Definition of $(\bar{v}_1)^{(7)},,(\bar{v}_2)^{(7)},(\bar{u}_1)^{(7)},(\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)}>0,(\bar{v}_2)^{(7)}<0$ and respectively $(\bar{u}_1)^{(7)}>0,(\bar{u}_2)^{(7)}<0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}=0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2+(\tau_2)^{(7)}u^{(7)}-(b_{36})^{(7)}=0$

Definition of $(m_1)^{(7)},(m_2)^{(7)},(\mu_1)^{(7)},(\mu_2)^{(7)},(v_0)^{(7)}$:-

If we define $(m_1)^{(7)},(m_2)^{(7)},(\mu_1)^{(7)},(\mu_2)^{(7)}$ by

$(m_2)^{(7)}=(v_0)^{(7)},(m_1)^{(7)}=(v_1)^{(7)}$, ***if*** $(v_0)^{(7)}<(v_1)^{(7)}$

$(m_2)^{(7)}=(v_1)^{(7)},(m_1)^{(7)}=(\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)}<(v_0)^{(7)}<(\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(gg) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^{*} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^{*} for which $f(T_{33}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

Definition and uniqueness of T_{37}^{*} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^{*} for which $f(T_{37}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523
A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^{*}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^{*}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}((G_{31})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SIXTY TWO

General Relativity And Quantum Cosmology Computational Complexity: Generalia Specialibus Non Derogant

@@@

Abstract: In the bewildering flux of ideas and tentative experimentation, it is but natural that we stand baffled befuddled, craved, corroded, belabored and burgeoned about the thought of the interrelationships and contiguous similarities and incompossible antigeneralities of computer generations, granularity, non-equilibrium thermodynamics, quantum computing, acausal processes. Like a man standing on the threshold of infinity trying to ponder what lies beyond the veil which separates the seen from unseen, human beings stand both curious and mouth gaped and confounded. Nowadays, we are using mainly computer of fourth generation, and we are designing fifth-generation computers. It is reasonable to ask: what is the perspective? What will the computers of generation omega look like? As the speed of data processing increases, we face a **natural (E) limitation of** causality, according to which the speed of all processes is limited by the speed of light. Lately, a new area of acausal (causality violating) processes has entered mainstream physics. This area has important astrophysical applications. In this paper, authors show how non-equilibrium thermodynamics **makes (EB) these** processes consistent, how these processes **can (E) be used in** computations, and how the very possibility of these processes leads to the granularity of the physical world. Computer generations, granularity, non-equilibrium thermodynamics, quantum computing, acausal processes. **(Towards Computers of Generation Omega - Non-Equilibrium Thermodynamics, Granularity, and Acausal Processes: A Brief Survey (1997) M. Koshelev, V. Kreinovich Emphasis and some literary additions mine)**

INTRODUCTION—VARIABLES USED

Relativistic quantum information and time machines DOI:10.1080/00107514.2011.640146 Timothy C.

Ralpha & Tony G. Downesa Contemporary Physics

(1) Relativistic quantum information combines the informational approach **to** understanding and using quantum mechanical systems – quantum information – with the relativistic view of the Universe. In this introductory review authors examine key results **to emerge from this** new field of research in physics and discuss future directions.
(2) A particularly active area recently has been the question of what happens when quantum systems **interact with** general relativistic closed timelike curves – effectively time machines. They discuss two different approaches that have been suggested for modelling such situations.
(3) It is argued that the approach **based on** matching the density operator of the quantum state between the future and past most consistently **avoids (e)the** paradoxes usually associated with time travel.

Computation and Spacetime Structure Mike Stannett

(4) Authors investigate the relationship between computation and spacetime structure, focusing on the role of closed timelike curves (CTCs) **in promoting** computational speedup.
(5) They note first that CTC traversal can **be (eb&e) interpreted** in two distinct ways, depending on one's understanding of spacetime.
(6) Focusing on one interpretation **leads us to** develop a toy universe in which no CTC can be traversed more than once, whence no computational speedup is possible.
(7) Focusing on the second (and more standard) interpretation **leads to the** surprising conclusion that CTCs act as perfect information repositories:
(8) Just as black holes have entropy, **so (eb) do** CTCs.
(9) If we also assume that P is not equal to NP, we find that all observers agree that, even if unbounded time travel existed in their youth, this capability eventually **vanishes (e) a**s they grow older.
(10) Thus the computational assumption "P is not NP" is also an assumption **concerning**(e&eb)cosmological structure arXiv: 1103.1127 [gr-qc]

Unconventional Computation Lecture Notes in Computer Science Volume 6714, 2011, pp 201-211 Computation with Narrow CTCsA. C. Cem Say, Abuzer Yakaryılmaz

(11) Authors examine some variants of computation with closed timelike curves (CTCs), where various **restrictions (e) are imposed** on the memory of the computer, and the information carrying capacity and range of the CTC.
(12) They give full characterizations of the classes of languages recognized by polynomial time probabilistic and quantum computers that can send a single classical bit **to (e&eb)** their own past. Such narrow CTCs are demonstrated to add the power of limited nondeterminism to deterministic computers, **and lead to** exponential speedup in constant-space probabilistic and quantum computation.

Wormholes and Time Travel Shortcut through Spacetime General Relativity: Wormholes and Time Travel Shortcut through Spacetime - If Traversable Wormholes Exist, They Could Allow Time Travel

(1) Einstein's Theory of Relativity **says that** time travel is perfectly possible — if you're going forward.
(2) Finding a way to travel backwards **requires** breaking the speed of light, which so far seems impossible.
(3) But now, strange-but-true phenomena such as quantum non-locality, where particles instantly teleport across vast distances **may give (eb) us a way** to make the dream of traveling back and forth through time a reality.
(4) There is no observational evidence for wormholes, but on a theoretical level **there (=) are** valid solutions to the equations of the theory of general relativity which contain wormholes.

(5) Travelling faster than the speed of light **(eb)** time travel
(6) The use of cosmic strings and black **holes (eb)** time travel
(7) Wormholes and Alcubierre drive (**eb)** time travel
(8) The first type of wormhole solution discovered was the Schwarzschild wormhole which would be present in the Schwarzschild metric describing an eternal black hole, but it was found that this type of wormhole **would collapse** (basket case bankruptcy, **cataclysmic catastrophe**, cave-in, conk out, **crackup crash**, debacle, **dis integrational destruction, disorganizational disruption**, exhaustion, failure, faint, flop, **pros rational ruination**, ruining, smash, **smashup subsidence, undoing wreck)too** quickly for anything to cross from one end to the other.
(9) Wormholes which could actually be crossed, known as traversable wormholes, would only be possible if exotic matter with negative energy density **could be used** to stabilize them (many physicists such as Stephen Hawking, Kip Thorne, and others believe that the Casimir effect is evidence that negative energy densities are possible in nature).

NOTATION

Module One

Relativistic quantum information combines the informational approach **to** understanding and using quantum mechanical systems – quantum information – with the relativistic view of the Universe. In this introductory review authors examine key results **to emerge from this** new field of research in physics and discuss future directions.

A particularly active area recently has been the question of what happens when quantum systems **interact with** general relativistic closed timelike curves – effectively time machines. They discuss two different approaches that have been suggested for modelling such situations.

It is argued that the approach **based on** matching the density operator of the quantum state between the future and past most consistently **avoids (e)the** paradoxes usually associated with time travel.

Authors investigate the relationship between computation and spacetime structure, focusing on the role of closed timelike curves (CTCs) **in promoting** computational speedup.

They note first that CTC traversal can **be (eb&e) interpretation** in two distinct ways, depending on one understands of spacetime.

Focusing on one interpretation **leads us to** develop a toy universe in which no CTC can be traversed more than once, whence no computational speedup is possible

Focusing on the second (and more standard) interpretation **leads to the** surprising conclusion that CTCs act as perfect information repositories
black holes have entropy, **so (eb) do** CTCs have entropy

If we also assume that P is not equal to NP, we find that all observers agree that, even if unbounded time travel existed in their youth, this capability eventually **vanishes (e) a**s they grow older.

Thus the computational assumption "P is not NP" is also an assumption **concerning**(e&eb)cosmological structure arXiv: 1103.1127 [gr-qc]

G_{13} : Category one of Relativistic quantum information combines the informational approach ; quantum systems; paradoxes usually associated with time travel; computation and spacetime structure, focusing on the role of closed timelike curves (CTCs); CTC traversal; one interpretation; second (and more

standard) interpretation; black holes have entropy(ac=dc-0); **a**s they grow older; computational assumption "P is not NP" is also an assumption

G_{14} : Category two of Relativistic quantum information combines the informational approach ; quantum systems; paradoxes usually associated with time travel; computation and spacetime structure, focusing on the role of closed timelike curves (CTCs); CTC traversal; one interpretation; second (and more standard) interpretation; black holes have entropy(ac=dc-0); **a**s they grow older; computational assumption "P is not NP" is also an assumption

G_{15} : Category three of Relativistic quantum information combines the informational approach; quantum systems ;paradoxes usually associated with time travel; computation and spacetime structure, focusing on the role of closed timelike curves (CTCs);CTC traversal ;one interpretation ;second (and more standard) interpretation ;black holes have entropy(ac=dc-0);**a**s they grow older; computational assumption "P is not NP" is also an assumption

T_{13} : Category one of understanding and using quantum mechanical systems – quantum information – with the relativistic view of the Universe; general relativistic closed timelike curves – effectively time machines; approach **based on** matching the density operator of the quantum state between the future and past most consistently; computational speedup; **interpretation** in two distinct ways, depending on one understands of spacetime; develop a toy universe in which no CTC can be traversed more than once, whence no computational speedup is possible; surprising conclusion that CTCs act as perfect information repositories; CTCs have entropy; assume that P is not equal to NP, we find that all observers agree that, even if unbounded time travel existed in their youth, this capability eventually; cosmological structure arXiv: 1103.1127 [gr-qc]

T_{14} : Category two of understanding and using quantum mechanical systems – quantum information – with the relativistic view of the Universe; general relativistic closed timelike curves – effectively time machines; approach **based on** matching the density operator of the quantum state between the future and past most consistently; computational speedup; **interpretation** in two distinct ways, depending on one understands of spacetime; develop a toy universe in which no CTC can be traversed more than once, whence no computational speedup is possible; surprising conclusion that CTCs act as perfect information repositories; CTCs have entropy; assume that P is not equal to NP, we find that all observers agree that, even if unbounded time travel existed in their youth, this capability eventually; cosmological structure arXiv: 1103.1127 [gr-qc]

T_{15} : Category three of understanding and using quantum mechanical systems – quantum information – with the relativistic view of the Universe; general relativistic closed timelike curves – effectively time machines; approach **based on** matching the density operator of the quantum state between the future and past most consistently; computational speedup; **interpretation** in two distinct ways, depending on one understands of spacetime; develop a toy universe in which no CTC can be traversed more than once, whence no computational speedup is possible; surprising conclusion that CTCs act as perfect information repositories; CTCs have entropy; assume that P is not equal to NP, we find that all observers agree that, even if unbounded time travel existed in their youth, this capability eventually; cosmological structure arXiv: 1103.1127 [gr-qc]

Module Two

Authors examine some variants of computation with closed timelike curves (CTCs), where various **restrictions (e) are imposed** on the memory of the computer, and the information carrying capacity and range of the CTC.

G_{16} : Category one of memory of the computer, and the information carrying capacity and range of the CTC.

G_{17} : Category two of memory of the computer, and the information carrying capacity and range of the CTC.

G_{18} : Category three of memory of the computer, and the information carrying capacity and range of the CTC.

T_{16} : Category one of variants of computation with closed timelike curves (CTCs),

T_{17} : Category two of variants of computation with closed timelike curves (CTCs),

T_{18} : Category three of variants of computation with closed timelike curves (CTCs),

Module three

They give full characterizations of the classes of languages recognized by polynomial time probabilistic and quantum computers that can send a single classical bit **to (e&eb)** their own past

G_{20} : Category one of characterizations of the classes of languages recognized by polynomial time probabilistic and quantum computers that can send a single classical bit; their own past

G_{21} : Category two of characterizations of the classes of languages recognized by polynomial time probabilistic and quantum computers that can send a single classical bit; their own past

G_{22} : Category three of characterizations of the classes of languages recognized by polynomial time probabilistic and quantum computers that can send a single classical bit; their own past

T_{20} : Category one of their own past ;characterizations of the classes of languages recognized by polynomial time probabilistic and quantum computers that can send a single classical bit

T_{21} : Category two of their own past ;characterizations of the classes of languages recognized by polynomial time probabilistic and quantum computers that can send a single classical bit

T_{22} : Category three of their own past ;characterizations of the classes of languages recognized by polynomial time probabilistic and quantum computers that can send a single classical bit

Module four

Such narrow CTCs are demonstrated to add the power of limited nondeterminism to deterministic computers, **and lead to** exponential speedup in constant-space probabilistic and quantum computation.

G_{24} : Category one of narrow CTCs are demonstrated to add the power of limited nondeterminism to deterministic computers

G_{25} : Category two of narrow CTCs are demonstrated to add the power of limited nondeterminism to deterministic computers

G_{26} : Category three of narrow CTCs are demonstrated to add the power of limited nondeterminism to deterministic computers

T_{24} : Category one of exponential speedup in constant-space probabilistic and quantum computation

T_{25} : Category two of exponential speedup in constant-space probabilistic and quantum computation

T_{26} : Category three of exponential speedup in constant-space probabilistic and quantum computation

Module five

Einstein's Theory of Relativity **says that** time travel is perfectly possible — if you're going forward

G_{28} : Category one of Einstein's Theory of Relativity. We are talking of the systems to which such a theory is applied. Characterstics from the bastion of the stratification schema and doxa in the present frame work of analytical treatment and coextensive representations.

G_{29} : Category two of Einstein's Theory of Relativity

G_{30} : Category three of Einstein's Theory of Relativity

T_{28} : Category one of time travel is perfectly possible — if you're going forward

T_{29} : Category two of time travel is perfectly possible — if you're going forward

T_{30} : Category three of time travel is perfectly possible — if you're going forward

Module six

Finding a way to travel backwards **requires** breaking the speed of light, which so far seems impossible

Wormholes and Alcubierre drive (**eb)** time travel

G_{32} : Category one of breaking the speed of light; Wormholes and Alcubierre drive

G_{33} : Category two of breaking the speed of light; Wormholes and Alcubierre drive

G_{34} : Category three of breaking the speed of light; Wormholes and Alcubierre drive

T_{32} : Category one of way to travel backwards. I have my own directions and limits as you would have a different one; time travel

T_{33} : Category two of way to travel backwards; time travel

T_{34} : Category three of way to travel backwards; time travel

Module seven

But now, strange-but-true phenomena such as quantum non-locality, where particles instantly teleport across vast distances **may give (eb) us a way** to make the dream of traveling back and forth through time a reality

G_{36} : Category one of phenomena such as quantum non-locality, where particles instantly teleport across vast distances

G_{37} : Category two of phenomena such as quantum non-locality, where particles instantly teleport across vast distances

G_{38} : Category three of phenomena such as quantum non-locality, where particles instantly teleport across vast distances

T_{36} : Category one of traveling back and forth through time a reality

T_{37} : Category two of traveling back and forth through time a reality

T_{38} : Category three of traveling back and forth through time a reality

Module eight

There is no observational evidence for wormholes, but on a theoretical level **there are** valid solutions to the equations of the theory of general relativity which contain wormholes

G_{40} : Category one of observational evidence for wormholes is lacking

G_{41} : Category two of observational evidence for wormholes is lacking

G_{42} : Category three of observational evidence for wormholes is lacking

T_{40} : Category one of valid solutions to the equations of the theory of general relativity which contain wormholes

T_{41} : Category two of valid solutions to the equations of the theory of general relativity which contain wormholes

T_{42} : Category three of valid solutions to the equations of the theory of general relativity which contain wormholes

Module Nine

The first type of wormhole solution discovered was the Schwarzschild wormhole which would be present in the Schwarzschild metric describing an eternal black hole, but it was found that this type of wormhole **would collapse** (basket case bankruptcy, **cataclysmic catastrophe**, cave-in, conk out, **crackup crash**, debacle, **dis integrational destruction, disorganizational disruption**, exhaustion, failure, faint, flop, **pros rational ruination**, ruining, smash, **smashup subsidence, undoing wreck)too** quickly for anything to cross from one end to the other.

G_{44} : Category one of quickly for anything to cross from one end to the other.

G_{45} : Category two of quickly for anything to cross from one end to the other.

G_{46} : Category three of quickly for anything to cross from one end to the other.

T_{44} : Category one of wormhole solution discovered was the Schwarzschild wormhole which would be present in the Schwarzschild metric describing an eternal black hole, but it was found that this type of wormhole

T_{45} : Category two of wormhole solution discovered was the Schwarzschild wormhole which would be present in the Schwarzschild metric describing an eternal black hole, but it was found that this type of wormhole

T_{46} : Category three of wormhole solution discovered was the Schwarzschild wormhole which would be present in the Schwarzschild metric describing an eternal black hole, but it was found that this type of wormhole

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27}),t\big)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27}),t\big)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27}),t\big)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\big((G_{27}),t\big) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\big((G_{31}),t\big)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\big((G_{31}),t\big)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\big((G_{31}),t\big)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\big((G_{31}),t\big) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$
$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$
$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**
$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{37}'')^{(7,7,)}(G_{39},t)$, $-(b_{36}'')^{(7,7,)}(G_{39},t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8)}(G_{43},t)$ $-(b_{41}'')^{(8,8)}(G_{43},t)$ $-(b_{42}'')^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17},t) + (a_{13}'')^{(1,1,)}(T_{14},t) + (a_{20}'')^{(3,3,3)}(T_{21},t) \\ + (a_{24}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{28}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{32}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{36}'')^{(7,7,7)}(T_{37},t) + (a_{40}'')^{(8,8,8)}(T_{41},t) + (a_{44}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17},t) + (a_{14}'')^{(1,1,)}(T_{14},t) + (a_{21}'')^{(3,3,3)}(T_{21},t) \\ + (a_{25}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{29}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{33}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{37}'')^{(7,7,7)}(T_{37},t) + (a_{41}'')^{(8,8,8)}(T_{41},t) + (a_{45}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17},t) + (a_{15}'')^{(1,1,)}(T_{14},t) + (a_{22}'')^{(3,3,3)}(T_{21},t) \\ + (a_{26}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{30}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{34}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{38}'')^{(7,7,7)}(T_{37},t) + (a_{42}'')^{(8,8,8)}(T_{41},t) + (a_{46}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad 63$$

Where $+(a_{16}'')^{(2)}(T_{17},t)$, $+(a_{17}'')^{(2)}(T_{17},t)$, $+(a_{18}'')^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for*

category 1, 2 and 3

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* $1,2\ 3$

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* $1,2$ *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* $1,2$ *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category $1,2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a_{32}')^{(6)} & +(a_{32}'')^{(6)}(T_{33},t) & +(a_{28}'')^{(5,5,5)}(T_{29},t) & +(a_{24}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a_{33}')^{(6)} & +(a_{33}'')^{(6)}(T_{33},t) & +(a_{29}'')^{(5,5,5)}(T_{29},t) & +(a_{25}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a_{34}')^{(6)} & +(a_{34}'')^{(6)}(T_{33},t) & +(a_{30}'')^{(5,5,5)}(T_{29},t) & +(a_{26}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a_{32}'')^{(6)}(T_{33},t)$, $+(a_{33}'')^{(6)}(T_{33},t)$, $+(a_{34}'')^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a_{28}'')^{(5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a_{24}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{ccc} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T_{21}'| e^{-(\hat{M}_{20})^{(3)} t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} ||G_{23} - G_{23}'|| e^{-(\hat{M}_{20})^{(3)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(QQQQQQQQ) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(RRRRRRRR) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(SSSSSSSS) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(TTTTTTTT) $\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(UUUUUUUU) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(VVVVVVVV) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\le\ (\hat{P}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\le\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\ \ G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\ \ T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\le\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\ \ G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\ \ T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(hh) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18}. indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16}, G_{18} and G_{16}, G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18}. The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2:

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad (208)$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad (209)$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself (210)

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric (211)

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t}\}$$

Indeed if we denote (212)

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results (213)

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad (214)$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. (215)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$

In the same way , one can obtain

$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25}\ \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25}\ \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\widehat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{36})^{(7)}\right]\leq(\widehat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(\nu_1)^{(6)}, (\nu_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, \nu^{(6)}, u^{(6)}$: 350

By $(\nu_1)^{(6)} > 0$, $(\nu_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)},, (\bar{\nu}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21}, t)\right) - (a_{21}'')^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}$, $\boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}$, $\boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)}=\frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

423

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case,

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

424

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(hh) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523
A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^{*}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^{*}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}((G_{31})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SIXTY THREE

Quantum Transport: Hannibal is at the Gate

@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@

Abstract: Disjunctive synthesis, divergential affirmation Predicational anteriorities, phenomenological correlates and naturalism of physics and interrogation problem and the nature of problem as the subject state has been widely discussed by various philosophers like Kant, Kierkegaard and Deluze. Hameroff recently has written a paper that begins with a general introduction to the nature of human consciousness and outlines several different philosophical approaches. A critique of traditional reductionist and dualist positions is offered and it is suggested that consciousness **should be viewed as** an emergent property of physical systems. However, although consciousness has its origin in distributed brain processes **it has** macroscopic properties - most notably the `unitary sense of self', non-deterministic free will, and non-algorithmic `intuitive' processing - which can best be described by quantum-mechanical principles. (**Quantum coherence in microtubules: A neural basis for emergent consciousness? Author: Hameroff, Stuart R. Source: Journal of Consciousness Studies, Volume 1, Number 1, 1994, pp. 91-118(28))** Observation of coextensive representations and axiomatic predications of coherent quantum dynamics have been studied by various authors and has become one of the prominent and predominant focus and locus of essence, sense and perception, in the subterranean realm and ceratoid dualism of both consciousness and quantum transport. Coherent time evolution between two quantum states of a superconducting flux qubit comprising three Josephson junctions in a loop with the superposition of the two states carrying opposite macroscopic persistent currents is manipulated resonant microwave pulses is done by **Chiorescu1 et al.** Readout **by**

means of switching-event measurement with an attached superconducting quantum interference device revealed quantum-state oscillations with high fidelity. Under strong microwave driving, it was possible to induce hundreds of coherent oscillations. Pulsed operations on this first sample yielded a relaxation time of 900 nanoseconds and a free-induction dephasing time of 20 nanoseconds. These results are promising for future solid-state quantum computing. (**Published Online February 13 2003 Science 21 March 2003: Vol. 299 no. 5614 pp. 1869-1871 DOI: 10.1126/science.1081045 Coherent Quantum Dynamics of a Superconducting Flux Qubit I. Chiorescu1 et al).** In this section we concentrate upon the quantum transport and its wide amplitudinal ramifications on quantum computation and analogous emergent consciousness. Emergent consciousness is also discussed somewhere in the paper, with the theories of start Hameroff and Penrose.

INTRODUCTION—VARIABLES USED

A supramolecular assembly or "super molecule" **is a well defined** complex of molecules held together by noncovalent bonds. While a supramolecular assembly can be simply **composed of two** molecules (e.g., a DNA double helix or an inclusion compound), it is more often used to denote larger complexes of molecules **that form** sphere-, rod-, or sheet-like species. The dimensions of supramolecular assemblies can range from nanometers to micrometers. Thus they allow access to nanoscale objects **using a** bottom-up approach in far fewer steps than a single molecule of similar dimensions. The process by which a supramolecular assembly forms is called molecular self-assembly. Some try to distinguish self-assembly as the process by which individual molecules form the defined aggregate. Self-organization, then, is the process **by which** those aggregates **create** higher-order structures. This can become useful when talking about liquid crystals and block copolymers. Supramolecular assemblies are being investigated as new materials in a variety of contexts. For instance, Samuel I. Stupp and coworkers at Northwestern University **showed that** a supramolecular assembly of peptide amphiphiles in the form of nanofibers could be used to promote the growth of neurons. A great advantage to this supramolecular approach is that the **nanofibers** will degrade back into the individual peptide molecules that can be broken down by the body. Another example with implications at the biology/materials science interface is of self-assembling dendritic dipeptides, which form hollow cylindrical supramolecular assemblies in solution and in bulk. The cylindrical assemblies possess internal helical order and self-organize into columnar liquid crystalline lattices. When inserted into vesicular membranes, the porous cylindrical assemblies mediate transport of protons across the membrane. Self-assembling dendrons has also been **used to** generate arrays of nanowires. Electron donor-acceptor complexes comprise the core of the cylindrical supramolecular assemblies, which further self-organize into two-dimensional columnar liquid crystalline lattices. Each cylindrical supramolecular assembly functions as an individual wire. High charge carrier Mobilities for holes and electrons were obtained.(Wikipedia)

NATURE CHEMISTRY | ARTICLE Utilizing redox-chemistry to elucidate the nature of exciton transitions in supramolecular dye nanotubes D. M. Eisele et al Nature Chemistry 4, 655–662 (2012) doi:10.1038/nchem.1380

(1) Supramolecular assemblies that interact with light have **recently garnered (eb) much interest** as well-defined nanoscale materials for electronic excitation energy collection and transport.
(2) However, to control such complex systems it is essential to understand how their various parts interact and whether these interactions **result in** coherently shared excited states (excitons) or in diffusive energy transport between them.
(3) Here, authors address this by studying a model system consisting of two concentric cylindrical dye aggregates **in (eb) a** light-harvesting nanotube.
(4) Through selective chemistry authors are able to unambiguously **determine (eb) the** supramolecular origin of the observed excitonic transitions.
(5) **Determination the** supramolecular origin of the observed excitonic transitions required **(e) the**

development of a new theoretical model of the supramolecular structure of the assembly.

(6) **determination the** supramolecular origin of the observed excitonic transitions **demonstrate** that the two cylinders of the nanotube have distinct spectral responses and are best described as two separate, weakly coupled excitonic systems.

(7) Understanding such interactions **is (e) critical to** the control of energy transfer on a molecular scale, a goal in various applications ranging from artificial photosynthesis to molecular electronics.

Nature of Exciton−Exciton Annihilation in an Aggregated Cyanine Dye Serdar Özçelik and Daniel L. Akins J. Phys. Chem. B, 1997, 101 (16), pp 3021–3024 DOI: 10.1021/jp970429l April 17, 1997

(8) For aggregated 1,1‘,3,3‘-tetraethyl-5,5‘,6,6‘-tetrachlorobenzimidazolocarbocyanine (TTBC) adsorbed onto silica, authors have **ascertained** that coherent optical dynamical processes **play(e) key roles** in defining the temporal response and spectral profile of the fluorescence.

(9) They have also **determined (eb)** the coherent processes involved **(e&eb)** in what might generally be termed exciton−exciton annihilation. Additionally, they provide in the TTBC system a concrete example of room temperature superradiance.

NOTATION

Module One

Supramolecular assemblies that interact with light have **recently garnered (eb) much interest** as well-defined nanoscale materials for electronic excitation energy collection and transport

G_{13} : Category one of Supramolecular assemblies that interact with light have

G_{14} : Category two of Supramolecular assemblies that interact with light have

G_{15} : Category three of Supramolecular assemblies that interact with light have

T_{13} : Category one of nanoscale materials for electronic excitation energy collection and transport

T_{14} : Category two of nanoscale materials for electronic excitation energy collection and transport

T_{15} : Category three of nanoscale materials for electronic excitation energy collection and transport

Module Two

However, to control such complex systems it is essential to understand how their various parts interact and whether these interactions **result in** coherently shared excited states (excitons) or in diffusive energy transport between them

G_{16} : Category one of control of such complex systems consummation calls for comprehension of the interactions of the various parts

G_{17} : Category two of control of such complex systems consummation calls for comprehension of the interactions of the various parts

G_{18} : Category three of control of such complex systems consummation calls for comprehension of the interactions of the various parts

T_{16} : Category one of coherently shared excited states (excitons) or in diffusive energy transport between them

T_{17} : Category two of coherently shared excited states (excitons) or in diffusive energy transport between them

T_{18} : Category three of coherently shared excited states (excitons) or in diffusive energy transport between

them

Module three

Here, authors address this by studying a model system consisting of two concentric cylindrical dye aggregates **in (eb) a** light-harvesting nanotube

G_{20} : Category one of two concentric cylindrical dye aggregates

G_{21} : Category two of two concentric cylindrical dye aggregates

G_{22} : Category three of two concentric cylindrical dye aggregates

T_{20} : Category one of light-harvesting nanotube

T_{21} : Category two of light-harvesting nanotube

T_{22} : Category three of light-harvesting nanotube

Module four

Through selective chemistry authors are able to unambiguously **determine (eb) the** supramolecular origin of the observed excitonic transitions.

G_{24} : Category one of selective chemistry

G_{25} : Category two of selective chemistry

G_{26} : Category three of selective chemistry

T_{24} : Category one of supramolecular origin of the observed excitonic transitions.

T_{25} : Category two of supramolecular origin of the observed excitonic transitions.

T_{26} : Category three of supramolecular origin of the observed excitonic transitions.

Module five

determination the supramolecular origin of the observed excitonic transitions required **(e) the** development of a new theoretical model of the supramolecular structure of the assembly

G_{28} : Category one of development of a new theoretical model of the supramolecular structure of the assembly

G_{29} : Category two of development of a new theoretical model of the supramolecular structure of the assembly

G_{30} : Category three of development of a new theoretical model of the supramolecular structure of the assembly

T_{28} : Category one of **determination the** supramolecular origin of the observed excitonic transitions

T_{29} : Category two of **determination the** supramolecular origin of the observed excitonic transitions

T_{30} : Category three of **determination the** supramolecular origin of the observed excitonic transitions

Module six

determination the supramolecular origin of the observed excitonic transitions **demonstrate** that the two cylinders of the nanotube have distinct spectral responses and are best described as two separate, weakly coupled excitonic systems.
Understanding such interactions **is (e) critical to** the control of energy transfer on a molecular scale, a goal in various applications ranging from artificial photosynthesis to molecular electronics
G_{32} : Category one of **determination the** supramolecular origin of the observed excitonic transitions ; control of energy transfer on a molecular scale, a goal in various applications ranging from artificial photosynthesis to molecular electronics

G_{33} : Category two of **determination the** supramolecular origin of the observed excitonic transitions ; control of energy transfer on a molecular scale, a goal in various applications ranging from artificial photosynthesis to molecular electronics

G_{34} : Category three of **determination the** supramolecular origin of the observed excitonic transitions ; control of energy transfer on a molecular scale, a goal in various applications ranging from artificial photosynthesis to molecular electronics

T_{32} : Category one of cylinders of the nanotube have distinct spectral responses and are best described as two separate, weakly coupled excitonic systems; Understanding such interactions

T_{33} : Category two of cylinders of the nanotube have distinct spectral responses and are best described as two separate, weakly coupled excitonic systems; Understanding such interactions

T_{34} : Category three of cylinders of the nanotube have distinct spectral responses and are best described as two separate, weakly coupled excitonic systems; Understanding such interactions

Module seven

For aggregated 1,1‘,3,3‘-tetraethyl-5,5‘,6,6‘-tetrachlorobenzimidazolocarbocyanine (TTBC) adsorbed onto silica, authors have **ascertained** that coherent optical dynamical processes **play(e) key roles** in defining the temporal response and spectral profile of the fluorescence

They have also **determined (eb)** the coherent processes involved **(e&eb)** in what might generally be termed exciton−exciton annihilation. Additionally, they provide in the TTBC system a concrete example of room temperature superradiance.

G_{36} : Category one of coherent optical dynamical processes **play(e) key roles** in defining the temporal response and spectral profile of the fluorescence; coherent processes; exciton−exciton annihilation

G_{37} : Category two of coherent optical dynamical processes **play(e) key roles** in defining the temporal response and spectral profile of the fluorescence; coherent processes; exciton−exciton annihilation

G_{38} : Category three of coherent optical dynamical processes **play(e) key roles** in defining the temporal response and spectral profile of the fluorescence; coherent processes; exciton−exciton annihilation

T_{36} : Category one of aggregated 1,1‘,3,3‘-tetraethyl-5,5‘,6,6‘-tetrachlorobenzimidazolocarbocyanine (TTBC) adsorbed onto silica, authors have **ascertained** that coherent optical dynamical processes exciton−exciton annihilation;; coherent processes;

T_{37} : Category two of aggregated 1,1‘,3,3‘-tetraethyl-5,5‘,6,6‘-tetrachlorobenzimidazolocarbocyanine

(TTBC) adsorbed onto silica, authors have **ascertained** that coherent optical dynamical processes; exciton−exciton annihilation; coherent processes;

T_{38} : Category three of aggregated 1,1‘,3,3‘-tetraethyl-5,5‘,6,6‘-tetrachlorobenzimidazolocarbocyanine (TTBC) adsorbed onto silica, authors have **ascertained** that coherent optical dynamical processes; exciton−exciton annihilation ;coherent processes

Module eight

Consciousness **should be viewed as** an emergent property of physical systems (**Quantum coherence in microtubules: A neural basis for emergent consciousness? Author: Hameroff, Stuart R. Source: Journal of Consciousness Studies, Volume 1, Number 1, 1994, pp. 91-118(28))**

G_{40} : Category one of emergent property of physical systems

G_{41} : Category two of emergent property of physical systems

G_{42} : Category three of emergent property of physical systems

T_{40} : Category one of Consciousness (we can call that as three different individual general ledgers or individual, collective and cosmic consciousness which are again individual, collective and nature’s general ledgers. Analogy ends there. Rest it is to be noted that Stuart’s definition could be adopted instead of our definition as the storage of information which is anagrammatically acted upon by the “mind” field resulting in phantasmagoria, dreams, and other pathological repercussions too. Here we are reviewing Stuart’s paper.)

T_{41} : Category two of Consciousness (we can call that as three different individual general ledgers or individual, collective and cosmic consciousness which are again individual, collective and nature’s general ledgers)

T_{42} : Category three of Consciousness (we can call that as three different individual general ledgers or individual, collective and cosmic consciousness which are again individual, collective and nature’s general ledgers)

Module Nine

However, although consciousness has its origin in distributed brain processes **it has** macroscopic properties - most notably the `unitary sense of self', non-deterministic free will, and non-algorithmic `intuitive' processing - which can best be described by quantum-mechanical principles. Note that the model can be applied to each sentence, each equation of any paper. In the quantum mechanical processes there could be creation and destruction of interactional elemental endowment of quantum particles, and it is necessary to maintain a progressive quantum transport register to obtain the correct picture rather than obtention of the final record. Same thing holds good for LHC. Progressive like the one maintained in the bank helps comprehension of the changes that has taken place and the transactions and interactions. This is formidably most important point to be noted in experimental observations.

G_{44} : Category one of macroscopic properties - most notably the `unitary sense of self', non-deterministic free will, and non-algorithmic `intuitive' processing - which can best be described by quantum-mechanical principles

G_{45} : Category two of macroscopic properties - most notably the `unitary sense of self', non-deterministic free will, and non-algorithmic `intuitive' processing - which can best be described by quantum-mechanical

principles

G_{46} : Category three of macroscopic properties - most notably the `unitary sense of self', non-deterministic free will, and non-algorithmic `intuitive' processing - which can best be described by quantum-mechanical principles

T_{44} : Category one of consciousness has its origin in distributed brain processes

T_{45} : Category two of consciousness has its origin in distributed brain processes

T_{46} : Category three of consciousness has its origin in distributed brain processes

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}),t\big)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}),t\big)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}),t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}),t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27}),t\big)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27}),t\big)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27}),t\big)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\big((G_{27}),t\big) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$
$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$
$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$
$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)\right]T_{28} \quad 28$$
$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)\right]T_{29} \quad 29$$
$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}), t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition

coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)}\ \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\ \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\ \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for

category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{16}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{17}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{18}'')^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{20}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{21}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{22}'')^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{46}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{45}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{44}'')^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{28}')^{(5)} & \boxed{+(a_{28}'')^{(5)}(T_{29},t)} & \boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)} \quad \boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a_{29}')^{(5)} & \boxed{+(a_{29}'')^{(5)}(T_{29},t)} & \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)} \quad \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{30}')^{(5)} & \boxed{+(a_{30}'')^{(5)}(T_{29},t)} & \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)} \quad \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad (81)$$

Where $\boxed{+(a_{28}'')^{(5)}(T_{29},t)}, \boxed{+(a_{29}'')^{(5)}(T_{29},t)}, \boxed{+(a_{30}'')^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3
And $\boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3
$\boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3
$\boxed{+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3
$\boxed{+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3
$\boxed{+(a_{40}'')^{(8,8\,,8,8,8,8)}(T_{41},t)}, \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3
$\boxed{+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) \\ -(b''_{32})^{(6,6,6)}(G_{35},t) & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) \\ -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) \\ -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) & & \end{array}\right] T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) \\ -(b''_{33})^{(6,6,6)}(G_{35},t) & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) \\ -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) \\ -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) & & \end{array}\right] T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) \\ -(b''_{34})^{(6,6,6)}(G_{35},t) & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) \\ -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) \\ -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) & & \end{array}\right] T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) \\ +(a''_{24})^{(4,4,4,)}(T_{25},t) & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) \\ +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) \\ +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) & & \end{array}\right] G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) \\ +(a''_{25})^{(4,4,4,)}(T_{25},t) & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) \\ +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) \\ +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) & & \end{array}\right] G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b_{46}'')^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)} & +(a_{36}'')^{(7)}(T_{37},t) & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{40}'')^{(8,8,8,8,8,8,8,)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)} & +(a_{37}'')^{(7)}(T_{37},t) & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)} & +(a_{38}'')^{(7)}(T_{37},t) & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a_{36}'')^{(7)}(T_{37},t)$, $(a_{37}'')^{(7)}(T_{37},t)$, $(a_{38}'')^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)} G_{44}$$
$$- \left[\begin{array}{ccc} (a'_{45})^{(9)} \boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)} G_{45}$$
$$- \left[\begin{array}{ccc} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)} T_{45} -$$
$$\left[\begin{array}{ccc} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{13}$$
$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(\mathrm{p_i})^{(2)}$, $(\mathrm{r_i})^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}\left((G_{19}), t\right) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ \ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ \ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(WWWWWWWW) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(XXXXXXXX) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(YYYYYYYY) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(ZZZZZZZZ)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(AAAAAAAAA) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(BBBBBBBBB) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{16})^{(2)}$, $T_i^0 \leq (\hat{Q}_{16})^{(2)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^{0} + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^{0} + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^{0} + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{40})^{(8)} \text{ , } T_i^0 \le (\hat{Q}_{40})^{(8)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(ii) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ and\ (\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(7)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)},, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad (333)$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad (334)$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad (335)$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad (336)$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \qquad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le \frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \qquad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \qquad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le -(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le -(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \qquad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \qquad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \qquad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(\nu_1)^{(9)},(\nu_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},\nu^{(9)},u^{(9)}$:

By $(\nu_1)^{(9)}>0$, $(\nu_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2+(\sigma_1)^{(9)}\nu^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{\nu}_1)^{(9)},,(\bar{\nu}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)}>0$, $(\bar{\nu}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2+(\sigma_2)^{(9)}\nu^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}$, $\boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}$, $\boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(ii) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)} G_{14}^{*}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)} G_{14}^{*}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)} T_{14}^{*}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)} T_{14}^{*}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)} G_{17}^{*}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)} G_{17}^{*}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)} T_{17}^{*}}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)} T_{17}^{*}}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)} G_{21}^{*}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)} G_{21}^{*}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)} T_{21}^{*}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)} T_{21}^{*}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\Big)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\Big)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SIXTY FOUR

Quantum Topological Computation: Multum In Parvo : Much In Little

@@

Abstract: Quantum computing—the manipulation of a quantum mechanical system **to do** information processing—has attracted considerable recent attention, largely triggered by Shor's proposed algorithm for finding prime factors in polynomial instead of exponential time. The importance of this problem **has also led to** numerous attempts to realization, consummation and further consolidation and consubstantiation of quantum computers, including systems such as trapped ions and quantum dots. In their Research Article, Gershenfeld and Chuang propose the use of a much less exotic system—nuclear magnetic resonance (NMR) of molecules in a room-temperature solution. They demonstrate that such a "bulk spin-resonance" system **is (eb)capable in** principle of doing quantum computation, and they discuss the generation of 6 to 10 quantum bits ("qubits"), which would be a daunting, but not impossible task with today's technology. Of course, solution NMR was used in the 1950s to study equally small molecules, yet today we study proteins with thousands of spins. If an NMR quantum computer were ultimately scalable to larger numbers of qubits (say 100), the implications for computational science would be exciting. There is doubt, however, that solution NMR quantum computing will ever be useful. Many-body entangled quantum states studied in condensed matter physics **can be primary resources for** quantum information, allowing any quantum computation to be realized using measurements alone, on the state. Such a universal state would be remarkably valuable, if only it were thermodynamically stable and experimentally accessible, **by virtue(e) of being** the unique ground state of a physically reasonable Hamiltonian made of two-body, nearest-neighbor interactions. Xie Chen et al introduce such a state, **composed of** six-state particles on a hexagonal lattice, and describe a general method for analyzing its properties based on its projected entangled pair state representation.**(© 2009 The American Physical SocietyURL: http://link.aps.org/doi/10.1103/PhysRevLett.102.220501DOI**: 10.1103/PhysRevLett.102.220501 **Gapped Two-Body Hamiltonian Whose Unique Ground State Is Universal for One-Way Quantum Computation Xie Chen et al)** Ensembles of uncoupled two-level systems (magnetic resonance or any other form) have quite classical dynamics, as shown by Feynman. Thus the clock cycles for any nonclassical dynamics, **including all of the** computing operations in the report and in any other conceivable treatment, require times on the order of the reciprocal of the spin-spin couplings (≈200 Hz for directly bonded atoms, ≈10 Hz for protons on nearest-neighbor carbons) per step. Many such steps would be needed for logic

operations between two separated spins. Dipolar couplings (for example, in solids) **can increase the** couplings by another factor of 10, but then the eigenstates are not the simple spin product states, and each logical manipulation will be much more complex. The slowest limit of speed estimated by Gershenfeld and Chuang (10 logic gates per second) is thus grossly overoptimistic for a reasonably sized molecule. **(Science 12 September 1997: Vol. 277 no. 5332 pp. 1688-1690 DOI: 10.1126/science.277.5332.1688 TECHNICAL COMMENTS: The Usefulness of NMR Quantum Computing Warren S. Warren: emphasis mine)**

INTRODUCTION—VARIABLES USED

A topological quantum computer **is a** theoretical quantum computer **that employs two**-dimensional quasiparticles called anyons, whose world lines cross over one another **to form** braids in a three-dimensional spacetime (i.e., one temporal plus two spatial dimensions). These braids **form the** logic gates that make up the computer. The advantage of a quantum computer based on quantum braids over using trapped quantum particles is that the former is much more stable. The smallest perturbations can cause a quantum particle to decohere and introduce errors in the computation, but such small perturbations do not change the topological properties of the braids. This is like the effort required to cut a string and reattach the ends to form a different braid, as opposed to a ball (representing an ordinary quantum particle in four-dimensional spacetime) simply bumping into a wall. The original proposal for topological quantum computation is due to Alexei Kitaev in 1997. While the elements of a topological quantum computer **originate in a** purely mathematical realm, experiments in Fractional quantum Hall systems indicate these elements **may be created** in the real world using semiconductors made of gallium arsenide near absolute zero and subjected to strong magnetic fields. The theory of quantum computation **can be constructed from the** abstract study of anyonic systems. In mathematical terms, **these are** unitary topological modular functors. Anyonic systems underlie the Jones polynomial and anyonic systems **arise in** Witten-Chern-Simons theory. The braiding and fusion of anyonic excitations in quantum Hall electron liquids and 2D-magnets **are modeled by** modular functors, opening a new possibility for the realization of quantum computers. The chief advantage of anyonic computation **would be** physical error correction: An error rate scaling like $e^ {-\a\l} $, where $\l$ is a length scale, and α is some positive constant. In contrast, the $\q$presumptive" qubit-model of quantum computation, **which repairs** errors combinatorically, **requires a** fantastically low initial error rate (about $10^ {-4}$) before computation can be stabilized. (**Topological Quantum Computation Michael H. Freedman et al** ArXiv: quant-ph/0101025)

Correlation phenomena observed by CMS may shed light on exotic state of matter Julia Velkovska and Wei Li 2013-05-31 (Figures and a few paragraphs are deleted attributable to spatial constraints)

(1) The Large Hadron Collider conducted a short pilot run in September 2012, in which protons and lead nuclei were collided for the first time at centre-of-mass energies of 5.02 TeV. This pilot run set the stage for a longer, sustained run of proton-lead (pPb) collisions in early 2013 as part of the LHC's heavy-ion programme. The new run **yielded** (eb) 30000 times more data compared to the pilot run and allowed CMS to explore rare events that only happen once in every 10 million proton-lead interactions.

(2) Of particular interest were collisions **that (eb) produce** a large number of particles (more than 50), since it was in these collisions that intriguing correlations were observed during the pilot run (reported in Phys. Lett. B 718 (2013) 795). CMS has submitted a new paper to Phys.Lett. B (arXiv: 1305.0609), bringing important insights essential for unraveling the origin of these mysterious particle correlations.

Multiplicity and transverse-momentum dependence of two- and four-particle correlations in pPb and PbPb collisions CMS Collaboration

(3) Measurements of two- **and** four-particle angular correlations for charged particles emitted in pPb collisions are presented over a wide range in pseudorapidity and full azimuth.

(4) The data, corresponding to an integrated luminosity of approximately 31 inverse nanobarns, **were (e) collected** during the 2013 LHC pPb run at a nucleon-nucleon center-of-mass energy of 5.02 TeV by the CMS experiment.

(5) The results are compared to 2.76 TeV semi-peripheral PbPb collision data, collected during the 2011 PbPb run, covering a similar range of particle multiplicities. The observed correlations **are (e) characterized by** the near-side (abs (Delta (phi) ~0) associated pair yields and the azimuthal anisotropy Fourier harmonics (v[n]).

(6) The second-order (v [2]) and third-order (v [3]) anisotropy harmonics are extracted **using (e) the** two-particle azimuthal correlation technique.

(7) A four-particle correlation method is also applied to obtain the value of v[2] and further explore the multi-particle nature of the correlations. Both associated pair yields and anisotropy harmonics are studied **as a (e&eb) function** of particle multiplicity and transverse momentum.

(8) The associated pair yields, the four-particle v [2], and the v [3] become apparent at about the same multiplicity. A remarkable similarity in the v [3] signal **as a (e&eb) function of** multiplicity is observed between the pPb and PbPb systems. Predictions based on the color glass condensate and hydrodynamic models are compared to the experimental results. Report number: CMS-HIN-13-002, CERN-PH-EP-2013-077 arXiv:1305.0609 [nucl-ex]

(9) This phenomenon **is a** correlation between pairs of particles that is enhanced at small relative azimuthal angle $\Delta\varphi$ over a wide range of rapidity separation (see $\Delta\eta$ in Image 3: deleted due to spatial restrictions), producing a ridge-like structure in data plots.

NOTATION

Module One

The Large Hadron Collider conducted a short pilot run in September 2012, in which protons and lead nuclei were collided for the first time at centre-of-mass energies of 5.02 TeV. This pilot run set the stage for a longer, sustained run of proton-lead (pPb) collisions in early 2013 as part of the LHC's heavy-ion programme. The new run **yielded** (eb) 30000 times more data compared to the pilot run and allowed CMS to explore rare events that only happen once in every 10 million proton-lead interactions.

Measurements of two- **and** four-particle angular correlations for charged particles emitted in pPb collisions are presented over a wide range in pseudorapidity and full azimuth.

G_{13} : Category one of pilot run set the stage for a longer, sustained run of proton-lead (pPb) collisions; wide range in pseudorapidity and full azimuth

G_{14} : Category two of pilot run set the stage for a longer, sustained run of proton-lead (pPb) collisions; wide range in pseudorapidity and full azimuth

G_{15} : Category three of pilot run set the stage for a longer, sustained run of proton-lead (pPb) collisions ;wide range in pseudorapidity and full azimuth

T_{13} : Category one of30000 times more data compared to the pilot run and allowed CMS to explore rare events that only happen once in every 10 million proton-lead interactions; Measurements of two- **and** four-particle angular correlations for charged particles emitted in pPb collisions

T_{14} : Category two of 30000 times more data compared to the pilot run and allowed CMS to explore rare events that only happen once in every 10 million proton-lead interactions; Measurements of two- **and** four-particle angular correlations for charged particles emitted in pPb collisions

T_{15} : Category three of 30000 times more data compared to the pilot run and allowed CMS to explore rare events that only happen once in every 10 million proton-lead interactions; Measurements of two- **and** four-particle angular correlations for charged particles emitted in pPb collisions

Module Two

Multiplicity and transverse-momentum dependence of two- and four-particle correlations in pPb and Pb Pb collisions

G_{16} : Category one of two- and four-particle correlations in pPb and Pb Pb collisions

G_{17} : Category two of two- and four-particle correlations in pPb and Pb Pb collisions

G_{18} : Category three of two- and four-particle correlations in pPb and Pb Pb collisions

T_{16} : Category one of Multiplicity and transverse-momentum

T_{17} : Category two of Multiplicity and transverse-momentum

T_{18} : Category three of Multiplicity and transverse-momentum

Module three

The data, corresponding to an integrated luminosity of approximately 31 inverse nanobarns, **were (e) collected** during the 2013 LHC pPb run at a nucleon-nucleon center-of-mass energy of 5.02 TeV by the CMS experiment

G_{20} : Category one of LHC pPb run at a nucleon-nucleon center-of-mass energy of 5.02 TeV by the CMS experiment

G_{21} : Category two of LHC pPb run at a nucleon-nucleon center-of-mass energy of 5.02 TeV by the CMS experiment

G_{22} : Category three of LHC pPb run at a nucleon-nucleon center-of-mass energy of 5.02 TeV by the CMS experiment

T_{20} : Category one of data, corresponding to an integrated luminosity of approximately 31 inverse nanobarns

T_{21} : Category two of data, corresponding to an integrated luminosity of approximately 31 inverse nanobarns

T_{22} : Category three of data, corresponding to an integrated luminosity of approximately 31 inverse nanobarns

Module four

The results are compared to 2.76 TeV semi-peripheral PbPb collision data, collected during the 2011 PbPb run, covering a similar range of particle multiplicities. The observed correlations **are (e) characterized by** the near-side (abs (Delta (phi)~0) associated pair yields and the azimuthal anisotropy Fourier harmonics (v[n]).

G_{24} : Category one of near-side (abs (Delta (phi)~0) associated pair yields and the azimuthal anisotropy Fourier harmonics (v[n]).

G_{25} : Category two of near-side (abs (Delta (phi)~0) associated pair yields and the azimuthal anisotropy Fourier harmonics (v[n]).

G_{26} : Category three of near-side (abs (Delta (phi)~0) associated pair yields and the azimuthal anisotropy Fourier harmonics (v[n]).

T_{24} : Category one of observed correlations

T_{25} : Category two of observed correlations

T_{26} : Category three of observed correlations

Module five

The second-order (v [2]) and third-order (v [3]) anisotropy harmonics are extracted **using (e) the** two-particle azimuthal correlation technique.

Extraction of jet properties from two particle azimuthal correlations in pp and AuAu collisions at root of S (NN)=200 Gev (Paul Constantine)

Two particle azimuthal correlation functions measured in pp and $AuAu$ collisions at $\sqrt{s_{NN}}$ =200 GeV at midrapidity with the central arms of the PHENIX detector are analyzed in order to extract the properties of hadronic jets produced in QCD vacuum and highly excited QCD media.

Already, the published results on hadron spectra at high transverse momentum (p_T) in pp [13], dAu [16] and $AuAu$ [14] collisions at RHIC energies gave definitive proof for the discovery of hadronic jet quenching.

We use the method of two particle azimuthal correlation functions to gain more detailed information about this phenomenon, by extracting quantities like the jet shape parameters j_T and k_T, the jet conditional yields (number of associated hadrons per high p_T trigger hadron from jet fragmentation) and the jet fragmentation function.

The analysis of pp data starts with the measurement of the vacuum fragmentation function from which the p_T dependence of the mean fragmentation momentum fraction $\langle z \rangle$ is extracted. We obtain a constant value of $\langle z \rangle$= 0.74±0.02 for p_T above 3 GeV/c. Soft (non-perturbative) parton fragmentation becomes significant below 3 GeV/c and the slope of the fragmentation function is p_T dependent in this region. Gaussian fits to jet induced azimuthal correlations are employed to measure the mean jet fragmentation transverse momentum $\langle |j_{Ty}| \rangle$= 359 ± 11 (stat) ± 6 (syst) MeV/c and the mean partonic transverse momentum $\langle |k_{Ty}| \rangle$= 964 ± 49 (stat) ± 16 (syst) MeV/c.

The analysis of $AuAu$ data is based on azimuthal correlation functions between charged hadrons in the 1.5-3 GeV/c and 3-5 GeV/c p_T regions in five classes of collision centrality. The mean jet fragmentation transverse momentum $\langle |j_{Ty}| \rangle$ is centrality independent and consistent with the value in pp data mentioned above. Even though the statistical and systematical errors associated with the extracted mean partonic transverse momentum $\langle |k_{Ty}| \rangle$ (multiplied by $\langle z_{trigg} \rangle$) are rather large, a strong broadening with the

centrality is observed. This is an important finding because hadronic jet quenching through energy loss (medium induced gluon radiation) is expected to be accompanied by a broadening effect.

Another important finding is that the jet conditional yields in both the near ($\Delta\phi \sim 0$) and away ($\Delta\phi \sim \pi$) regions exhibit a slightly raising trend with centrality. This was also expected to accompany the jet quenching found at higher p_T based on simple energy conservation considerations.

This detailed study of hadronic jet properties is another piece of the Quark-Gluon Plasma puzzle that RHIC is trying to solve.

G_{28} : Category one of two-particle azimuthal correlation technique.

G_{29} : Category two of two-particle azimuthal correlation technique.

G_{30} : Category three of two-particle azimuthal correlation technique.

T_{28} : Category one of second-order (v [2]) and third-order (v [3]) anisotropy harmonics

T_{29} : Category two of second-order (v [2]) and third-order (v [3]) anisotropy harmonics

T_{30} : Category three of second-order (v [2]) and third-order (v [3]) anisotropy harmonics

Module six

two particle azimuthal correlation functions measured in pp and AuAu collsions are analysed **to(eb) extract** properties of hadronic jets produced in QCD vaccum and highly excited QD media

G_{32} : Category one of two particle azimuthal correlation functions measured in pp and AuAu collsions

G_{33} : Category two of two particle azimuthal correlation functions measured in pp and AuAu collsions

G_{34} : Category three of two particle azimuthal correlation functions measured in pp and AuAu collsions

T_{32} : Category one of properties of hadronic jets produced in QCD vaccum and highly excited QD media

T_{33} : Category two of properties of hadronic jets produced in QCD vaccum and highly excited QD media

T_{34} : Category three of properties of hadronic jets produced in QCD vaccum and highly excited QD media

Module seven

A four-particle correlation method is also applied to obtain the value of v[2] and further explore the multi-particle nature of the correlations; both associated pair yields and anisotropy harmonics are studied **as a (e&eb) function** of particle multiplicity and transverse momentum.

The CMS experiment at the Large Hadron Collider has recently presented data on the average transverse momentum of protons, kaons and pions as a function of particle multiplicity. Authors relate the particle multiplicity to saturation momentum **using** recently published computations of the interaction radius determined from the theory of the Color Glass Condensate. They show that the pp and the pA experimental data scale **in terms of** these saturation momenta. Computing transverse momentum spectra for identified particles **using** Boltzmann-type distributions and relating different associated multiplicities using geometric

scaling, these simple distributions reproduce the observed dependence of the mean transverse momentum on particle multiplicities seen in both pp and pA interactions for pions to good accuracy, and to fair agreement for protons and kaons.arXiv:1306.2350 [hep-ph] **(Transverse Momentum of Protons, Pions and Kaons in High Multiplicity pp and pA Collisions: Evidence for the Color Glass Condensate? Larry McLerran et al)**

G_{36} : Category one of four-particle correlation method is also applied to obtain the value of v[2] and further explore the multi-particle nature of the correlations; both associated pair yields and anisotropy harmonics recently published computations of the interaction radius determined from the theory of the Color Glass Condensate; Boltzmann-type distributions and relating different associated multiplicities using geometric scaling, these simple distributions reproduce the observed dependence of the mean transverse momentum on particle multiplicities seen in both pp and pA interactions for pions to good accuracy, and to fair agreement for protons and kaons;

G_{37} : Category two of four-particle correlation method is also applied to obtain the value of v[2] and further explore the multi-particle nature of the correlations; both associated pair yields and anisotropy harmonics; recently published computations of the interaction radius determined from the theory of the Color Glass Condensate; Boltzmann-type distributions and relating different associated multiplicities using geometric scaling, these simple distributions reproduce the observed dependence of the mean transverse momentum on particle multiplicities seen in both pp and pA interactions for pions to good accuracy, and to fair agreement for protons and kaons;

G_{38} : Category three of four-particle correlation method is also applied to obtain the value of v[2] and further explore the multi-particle nature of the correlations; both associated pair yields and anisotropy harmonics; recently published computations of the interaction radius determined from the theory of the Color Glass Condensate; Boltzmann-type distributions and relating different associated multiplicities using geometric scaling, these simple distributions reproduce the observed dependence of the mean transverse momentum on particle multiplicities seen in both pp and pA interactions for pions to good accuracy, and to fair agreement for protons and kaons;

T_{36} : Category one of particle multiplicity and transverse momentum; CMS experiment at the Large Hadron Collider has recently presented data on the average transverse momentum of protons, kaons and pions as a function of particle multiplicity. Authors relate the particle multiplicity to saturation momentum; Computing transverse momentum spectra for identified particles;

T_{37} : Category two of particle multiplicity and transverse momentum; CMS experiment at the Large Hadron Collider has recently presented data on the average transverse momentum of protons, kaons and pions as a function of particle multiplicity. Authors relate the particle multiplicity to saturation momentum; Computing transverse momentum spectra for identified particles

T_{38} : Category three of particle multiplicity and transverse momentum; CMS experiment at the Large Hadron Collider has recently presented data on the average transverse momentum of protons, kaons and pions as a function of particle multiplicity. Authors relate the particle multiplicity to saturation momentum; Computing transverse momentum spectra for identified particles

Module eight

The associated pair yields, the four-particle v [2], and the v [3] become apparent at about the same multiplicity; remarkable similarity in the v [3] signal **as a (e&eb) function of** multiplicity is observed between the pPb and PbPb systems. Predictions based on the color glass condensate and hydrodynamic

models are compared to the experimental results

G_{40} : Category one of associated pair yields, the four-particle v [2], and the v [3] become apparent at about the same multiplicity; remarkable similarity in the v [3] signal;

G_{41} : Category two of associated pair yields, the four-particle v [2], and the v [3] become apparent at about the same multiplicity; remarkable similarity in the v [3] signal;

G_{42} : Category three of associated pair yields, the four-particle v [2], and the v [3] become apparent at about the same multiplicity; remarkable similarity in the v [3] signal;

T_{40} : Category one of multiplicity is observed between the pPb and PbPb systems; associated pair yields, the four-particle v [2], and the v [3] become apparent at about the same multiplicity; remarkable similarity in the v [3] signal;

T_{41} : Category two of associated pair yields, the four-particle v [2], and the v [3] become apparent at about the same multiplicity; multiplicity is observed between the pPb and PbPb systems; remarkable similarity in the v [3] signal;

T_{42} : Category three of associated pair yields, the four-particle v [2], and the v [3] become apparent at about the same multiplicity; multiplicity is observed between the pPb and PbPb systems; remarkable similarity in the v [3] signal;

Module Nine

This phenomenon **is a** correlation between pairs of particles that is enhanced at small relative azimuthal angle Δφ over a wide range of rapidity separation (see Δη in Image 3: deleted due to spatial restrictions), producing a ridge-like structure in data plots.

G_{44} : Category one of correlation between pairs of particles that is enhanced at small relative azimuthal angle Δφ over a wide range of rapidity separation (see Δη in Image 3: deleted due to spatial restrictions), producing a ridge-like structure in data plots

G_{45} : Category two of correlation between pairs of particles that is enhanced at small relative azimuthal angle Δφ over a wide range of rapidity separation (see Δη in Image 3: deleted due to spatial restrictions), producing a ridge-like structure in data plots

G_{46} : Category three of correlation between pairs of particles that is enhanced at small relative azimuthal angle Δφ over a wide range of rapidity separation (see Δη in Image 3: deleted due to spatial restrictions), producing a ridge-like structure in data plots

T_{44} : Category one of four-particle correlation method is also applied to obtain the value of v[2] and further explore the multi-particle nature of the correlations; both associated pair yields and anisotropy harmonics are studied **as a (e&eb) function** of particle multiplicity and transverse momentum

T_{45} : Category two of four-particle correlation method is also applied to obtain the value of v[2] and further explore the multi-particle nature of the correlations; both associated pair yields and anisotropy harmonics are studied **as a (e&eb) function** of particle multiplicity and transverse momentum

T_{46} : Category three of four-particle correlation method is also applied to obtain the value of v[2] and further explore the multi-particle nature of the correlations; both associated pair yields and anisotropy harmonics are

studied **as a (e&eb) function** of particle multiplicity and transverse momentum

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients
Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$
$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$
$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$
$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**
$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$
$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\ \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\ \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\ \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\ \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\ \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\ \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\ \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\ \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\ \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\ \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\ \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\ \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\ \boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\ \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

59

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

60

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

61

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16}$$

62

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) \end{bmatrix} G_{17}$$

63

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) \end{bmatrix} G_{18}$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a_{44}'')^{(9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b_{16}')^{(2)} & -(b_{16}'')^{(2)}(G_{19},t) & -(b_{13}'')^{(1,1,)}(G,t) & -(b_{20}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{24}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{28}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{32}'')^{(6,6,6,6,6)}(G_{35},t) \\ -(b_{36}'')^{(7,7,7)}(G_{39},t) & -(b_{40}'')^{(8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b_{17}')^{(2)} & -(b_{17}'')^{(2)}(G_{19},t) & -(b_{14}'')^{(1,1,)}(G,t) & -(b_{21}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{25}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{29}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{33}'')^{(6,6,6,6,6)}(G_{35},t) \\ -(b_{37}'')^{(7,7,7)}(G_{39},t) & -(b_{41}'')^{(8,8,8)}(G_{43},t) & -(b_{45}'')^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b_{18}')^{(2)} & -(b_{18}'')^{(2)}(G_{19},t) & -(b_{15}'')^{(1,1,)}(G,t) & -(b_{22}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{26}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{30}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{34}'')^{(6,6,6,6,6)}(G_{35},t) \\ -(b_{38}'')^{(7,7,7)}(G_{39},t) & -(b_{42}'')^{(8,8,8)}(G_{43},t) & -(b_{46}'')^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad 66$$

where $-(b_{16}'')^{(2)}(G_{19},t)$, $-(b_{17}'')^{(2)}(G_{19},t)$, $-(b_{18}'')^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b_{13}'')^{(1,1,)}(G,t)$, $-(b_{14}'')^{(1,1,)}(G,t)$, $-(b_{15}'')^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b_{20}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b_{24}'')^{(4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b_{28}'')^{(5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b_{32}'')^{(6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b_{36}'')^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b_{40}'')^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b_{44}'')^{(9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{ccc} (a_{20}')^{(3)}\boxed{+(a_{20}'')^{(3)}(T_{21},t)} & \boxed{+(a_{16}'')^{(2,2,2)}(T_{17},t)} & \boxed{+(a_{13}'')^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{36}'')^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{ccc} (a_{21}')^{(3)}\boxed{+(a_{21}'')^{(3)}(T_{21},t)} & \boxed{+(a_{17}'')^{(2,2,2)}(T_{17},t)} & \boxed{+(a_{14}'')^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{37}'')^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{ccc} (a_{22}')^{(3)}\boxed{+(a_{22}'')^{(3)}(T_{21},t)} & \boxed{+(a_{18}'')^{(2,2,2)}(T_{17},t)} & \boxed{+(a_{15}'')^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{38}'')^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{22} \quad (69)$$

$\boxed{+(a_{20}'')^{(3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a_{14}'')^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{36}'')^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a_{37}'')^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a_{38}'')^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{40}'')^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{44}'')^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b_{20}')^{(3)}\boxed{-(b_{20}'')^{(3)}(G_{23},t)} & \boxed{-(b_{16}'')^{(2,2,2)}(G_{19},t)} & \boxed{-(b_{13}'')^{(1,1,1,)}(G,t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{36}'')^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b_{40}'')^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b_{44}'')^{(9,9,9)}(G_{47},t)} \end{array}\right] T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1,2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1,2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1,2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category $1,2$ *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category $1,2$ *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$

are sixth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$

are seventh detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$

are eighth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt}=(a_{28})^{(5)}G_{29}-\left[\begin{array}{ccc}(a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt}=(a_{29})^{(5)}G_{28}-\left[\begin{array}{ccc}(a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt}=(a_{30})^{(5)}G_{29}-\left[\begin{array}{ccc}(a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt}=(b_{28})^{(5)}T_{29}-\left[\begin{array}{ccc}(b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}\end{array}\right]T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)} & +(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34} \quad (90)$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition

coefficients for category 1, 2, and 3
$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3
$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3
$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37},t) & +(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37},t) & +(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37},t) & +(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a''_{36})^{(7)}(T_{37},t)$, $(a''_{37})^{(7)}(T_{37},t)$, $(a''_{38})^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3
$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3
$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3
$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3
$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation

coefficient for 1,2,3

$\frac{dT_{36}}{dt} =$ 94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$\frac{dT_{37}}{dt} =$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$\frac{dT_{38}}{dt} =$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)} G_{44}$$
$$- \left[\begin{array}{c} (a'_{45})^{(9)} \boxed{+(a''_{45})^{(9)}(T_{45},t)} \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)} G_{45}$$
$$- \left[\begin{array}{c} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)} T_{45} -$$
$$\left[\begin{array}{c} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{13}$$
$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(\mathrm{p_i})^{(2)}$, $(\mathrm{r_i})^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}\left((G_{19}), t\right) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ \ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ \ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(CCCCCCCCC) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(DDDDDDDDD) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(EEEEEEEEE) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(FFFFFFFFF)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(GGGGGGGGG) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(HHHHHHHHH) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)}$, 155

$0 \leq G_i(t) - G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \leq T_i(t) - T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^{0} + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^{0} + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^{0} + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{36})^{(7)}$, $T_i^0 \leq (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(jj) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ and\ (\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)},\ if\ (v_0)^{(1)} < (v_1)^{(1)}$

$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, if\ (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$

and $\boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)},\ if\ (\bar{v}_1)^{(1)} < (v_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad (333)$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad (334)$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad (335)$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad (336)$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \qquad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \qquad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \qquad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

406

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case,

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

407

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t)\right) - (a_{41}'')^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(jj) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}$, $\frac{\partial (b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\big((a_{32}')^{(6)} + (p_{32})^{(6)}\big)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\big((a_{33}')^{(6)} + (p_{33})^{(6)}\big)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\big((a_{34}')^{(6)} + (p_{34})^{(6)}\big)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\big((b_{32}')^{(6)} - (r_{32})^{(6)}\big)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(32)(j)}T_{32}^*\mathbb{G}_j\big) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\big((b_{33}')^{(6)} - (r_{33})^{(6)}\big)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\big(s_{(33)(j)}T_{33}^*\mathbb{G}_j\big) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\big((b_{34}')^{(6)} - (r_{34})^{(6)}\big)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(34)(j)}T_{34}^*\mathbb{G}_j\big) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial (b_i'')^{(7)}}{\partial G_j}\big((G_{39})^{**} \big) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\big((a_{36}')^{(7)} + (p_{36})^{(7)}\big)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\big((a_{37}')^{(7)} + (p_{37})^{(7)}\big)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\big((a_{38}')^{(7)} + (p_{38})^{(7)}\big)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\big((b_{36}')^{(7)} - (r_{36})^{(7)}\big)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(36)(j)}T_{36}^*\mathbb{G}_j\big) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\big((b_{37}')^{(7)} - (r_{37})^{(7)}\big)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\big(s_{(37)(j)}T_{37}^*\mathbb{G}_j\big) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\big((b_{38}')^{(7)} - (r_{38})^{(7)}\big)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(38)(j)}T_{38}^*\mathbb{G}_j\big) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

Advances in Physics Theories and Applications
ISSN 2224-719X (Paper) ISSN 2225-0638 (Online)
Vol 14, 2013

SECTION SIXTY FIVE

LQG And Quantum Geometry: Dona Clandestina Sunt Semper Suspiciosa Clandestine Gifts And Sybaritic, Luxuriant Trade Offs

Abstract: Problem of describing the quantum behavior of gravity, and thus understanding quantum spacetime, is still open in all its wide ranging amplitudinal forms and manifestations with its specifications and speculations about internal differentiation, structural morphology, imperative incompatibilities, phenomenological correlates and predicational correlates. Loop quantum gravity is a well-developed approach to this problem. It is a mathematically well-defined background-independent quantization of general relativity, with its conventional matter couplings. Today research in loop quantum gravity forms a vast area, ranging from mathematical foundations to physical applications. Among the most significant results obtained so far are: (i) the computation of the spectra of geometrical quantities such as area and volume, which yield tentative quantitative predictions for Planck-scale physics. (ii) A physical picture of the microstructure of quantum spacetime, characterized by Planck-scale discreteness. Discreteness **emerges** as a standard quantum effect from the discrete spectra, and provides a mathematical realization of Wheeler's "spacetime foam" intuition. (iii) Control of spacetime singularities, such as those in the interior of black holes and the cosmological one. This, in particular, has opened up the possibility of a theoretical investigation into the very early universe and the spacetime regions beyond the Big Bang. (iv) A derivation of the Bekenstein–Hawking black-hole entropy. (v) Low-energy calculations, yielding n-point functions well defined in a background-independent context. The theory is at the roots of, or strictly related to, a number of formalisms that have been developed for describing background-independent quantum field theory, such as spin foams, group field theory, causal spin networks, and other .The mathematics of loop quantum gravity is solidly defined, and is understood from several alternative points of view. Long-standing problems, such as the lack of a scalar product, the difficult **of controlling** the over completeness of the loop basis, and the problem of implementing the **reality condition** in the quantum theory, have been successfully solved or sidestepped. The kinematics **is given by** a well-defined Hilbert space H that carries a representation of the basic operators. A convenient orthonormal basis in H is provided by the spin-network states, the diffeomorphism-invariant states are given by the s-knot states, and the structure and properties of the (diff-invariant) quantum states of the geometry are quite well understood. These states **give a description** of quantum spacetime in terms of discrete excitations of the geometry. More precisely, in terms of elementary excitations carrying discretized quanta of area. (**Loop Quantum Gravity Carlo Rovelli emphasis mine)** New representation for quantum general relativity is described, **which is defined in terms of** functionals of sets of loops in three-space. In these representation exact solutions of the quantum constraints may be obtained, the result is **related to the** simplification of the constraints in Ashtekar new formalism. Closed form the general solution of the diffeomorphism constraints and a large class of solutions of the full set of constraints. **Are classified** by the knot and link classes of the spatial three-manifold.© 1988 The American

Physical SocietyURL: http://link.aps.org/doi/10.1103/PhysRevLett.61.1155DOI: 10.1103/PhysRevLett.61.1155 **Knot Theory and Quantum Gravity Carlo Rovelli**

INTRODUCTION—VARIABLES USED

(1) Loop quantum gravity (LQG) is a theory that attempts **to describe** the quantum properties of gravity.
(2) LQG **is also a** theory of quantum space and quantum time, **because,** as discovered with general relativity, the geometry of spacetime is a manifestation of gravity.
(3) LQG is an attempt to merge and adapt standard quantum mechanics **and** standard general relativity.
(4) The main output of the theory **is a** physical picture of space where space is granular.
(5) The granularity is a direct **consequence of the** quantization.
(6) It has the same nature of the granularity of the photons in the quantum theory of electromagnetism or the discrete levels of the energy of the atoms. But, here, it is space itself which is discrete. More precisely, space can be viewed as an extremely fine fabric or network "woven" **of** finite loops. These networks of loops **are called** spin networks. The evolution of a spin network over time is called spin foam. The predicted size of this structure is the Planck length, which are approximately 10−35 meters. According to the theory, there is no meaning to distance at scales smaller than the Planck scale. Therefore LQG predicts that not just matter, but also space itself, **has an** atomic structure. Today LQG is a vast area of research, developed in several directions, which involves about 50 research groups worldwide. They all share the basic physical assumptions and the mathematical description of quantum space. The full development of the theory is pursued in two directions: the more traditional canonical loop quantum gravity **and the** younger covariant loop quantum gravity, more commonly called spin foam theory.

Several research directions study the physical consequences of the theory. Among these, the most developed is the application of LQG to cosmology, called Loop quantum cosmology (LQC). LQC applies LQG ideas to the study of the early universe and the physics of the Big Bang. Its most spectacular consequence **is that** the evolution of the universe can be continued beyond the Big Bang. The Big Bang appears thus to be replaced by a sort of cosmic Big Bounce.

(7) In theoretical physics, general covariance **is the** invariance of the form of physical laws under arbitrary differentiable coordinate transformations. The essential idea is that coordinates are only artifices **used in** describing nature, and hence should play no role in the formulation of fundamental physical laws. A more significant requirement is the principle of General Relativity that states that the laws of physics take the same form in all reference systems. This is a generalization of the principle of special relativity which states that the laws of physics take the same form in all inertial frames. In mathematics, a diffeomorphism **is an** isomorphism in the category of smooth manifolds. It is an invertible function that maps one differentiable manifold to another, such that both the function and its inverse are smooth. These are the defining symmetry transformations of General Relativity since the theory is formulated only in terms of a differentiable manifold.
(8) In General Relativity, General covariance is intimately related to "diffeomorphism invariance". This symmetry is one of the defining features of the theory. However, it is a common misunderstanding that "diffeomorphism invariance" refers to the invariance of physical laws under arbitrary coordinate transformations; this is untrue and in fact every physical theory is invariant under coordinate transformations. Diffeomorphisms, as mathematicians define them, correspond to something much more radical; intuitively a way they can be envisaged is as simultaneously dragging all the physical fields (including the gravitational field) over the bare differentiable manifold while staying in the same coordinate system; diffeomorphisms are the true symmetry transformations of General Relativity, and come about from the assertion that the formulation of the

theory is based on a bare differentiable manifold, but not on any prior geometry - the theory is background-independent (this is a profound shift, as all physical theories before general relativity had as part of their formulation a prior geometry). What is preserved under such transformations are the coincidences between the values the gravitational field take at such and such a `place' and the values the matter fields take there, from these relationships one can form a notion of matter being located with respect to the gravitational field, or vice versa. This is what Einstein discovered, physical entities are located with respect to one another only and not with respect to the spacetime manifold - as Carlo Rovelli puts it: "**No more fields on spacetime: just fields on fields.".** This is the true meaning of the saying "The stage disappears **and becomes(=)** one of the actors"; space-time as a `container' over which physics takes place has no objective physical meaning and instead the gravitational interaction is represented as just one of the fields forming the world. This is known as the relationalism interpretation of space-time. The realization by Einstein that General Relativity should be interpreted this way is the origin of his remark "Beyond my wildest expectations".

In LQG this aspect of General Relativity is taken seriously and this symmetry is preserved by requiring that the physical states remain invariant under the generators of diffeomorphisms. The interpretation of this condition is well understood for purely spatial diffeomorphisms. However, the understanding of diffeomorphisms involving time (the Hamiltonian constraint) is more subtle because it is related to dynamics and the so-called "problem of time" in general relativity. A generally accepted calculational framework to account for this constraint has yet to be found A plausible candidate for the quantum Hamiltonian constraint is the operator introduced by Thiemann.

LQG is formally background independent. The equations of LQG are not embedded in, or presuppose space and time (except for its invariant topology). Instead, they are expected to give rise to space and time at distances which are large compared to the Planck length. The issue of background independence in LQG still has some unresolved subtleties. For example, some derivations require a fixed choice of the topology, while any consistent quantum theory of gravity should include topology change as a dynamical process.

(Phys.org) Blog:

(1) Physicists Rodolfo Gambini and Jorge Pullin of University of the Republic in Montevideo, Uruguay, and Louisiana State University respectively, have applied the theory of Loop Quantum Gravity (LQG) **to (e&eb) a** simplified black hole.
(2) In so doing, as they describe in their paper published in the journal Physical Review Letters, they suggest that instead of a singularity existing at its center, **there (eb) is a** portal to another universe.
(3) For many years theoretical physicists have believed that the universe **came about (e) as a result of** a single Big bang event—Einstein's theories suggested it was so.
(4) The problem with this line of thinking however is that the theory of general relativity **can't (e)** describes what came before the singularity, which should exist at the point in time just before the Big Bang.
(5) Theory also suggests that a similar singularity **should (eb) exist at** the center of black holes
(6) General relativity **fails (e) to** describe them properly.
(7) Worse, there is the problem of the information loss paradox—if something falls into a black hole and is eventually squeezed to a singularity, what **happens(eb)** to the information it contained? Big Bang physicists can't say.
(8) To address these problems, Abhay Ashtekar and his team at Pennsylvania State University, back in 2006, came up with a theory known as loop quantum gravity. They suggested that instead of a singularity existing just before the Big Bang, there was the remains of a crunched down universe that had existed prior to the one that exists now. The universe didn't just Big Bang itself into

existence from nothing, they said, instead there is an infinite loop **where a universe(eb) shrinks down to** a very tiny spot, then explodes in a Big Bang, then shrinks down again, over and over again forever—hence the use of the term "loop" in the theory. Since that time, some in the field have begun to refer to the theory as the Big Bounce, to replace the name Big Bang.

(9) In this new effort Gambini and Pullin **applied (e&eb)** LQG to a simplified model of a black hole.

(10) Their experiment **showed(eb) that** everything that was pulled into the black hole didn't compress to a singularity after all—instead it was compressed to a certain small size, **then was spit(eb) out in** another part of the universe or into another universe entirely.

(11) Because their model worked so well, the two suggest that it would likely work with real black holes as well. If this new theory is correct, they note, **it would do (e) away with** the information loss paradox and open the door to the possibility of black holes being portals to other universes. **Via Synopsis Journal reference: Physical Review Letters Read more at: http://phys.org/news/2013-05-theorists-loop-quantum-gravity-theory.html#**

NOTATION

Module One

(1) LQG **is also a** theory of quantum space and quantum time, **because,** as discovered with general relativity, the geometry of spacetime is a manifestation of gravity.

(2) LQG is an attempt to merge and adapt standard quantum mechanics **and** standard general relativity.

(3) The main output of the theory **is a** physical picture of space where space is granular

(4) The granularity is a direct **consequence of the** quantization

G_{13} : Category one of Loop quantum gravity (LQG) ; geometry of spacetime is a manifestation of gravity; main output of the theory; quantization;

G_{14} : Category two of Loop quantum gravity (LQG); geometry of spacetime is a manifestation of gravity; main output of the theory; quantization;

G_{15} : Category three of Loop quantum gravity (LQG) ;geometry of spacetime is a manifestation of gravity; main output of the theory; quantization

T_{13} : Category one of quantum properties of gravity; LQG **is also a** theory of quantum space and quantum time; physical picture of space where space is granular; granularity;

T_{14} : Category two of quantum properties of gravity; LQG **is also a** theory of quantum space and quantum time; physical picture of space where space is granular; granularity;

T_{15} : Category three of quantum properties of gravity; LQG **is also a** theory of quantum space and quantum time; physical picture of space where space is granular; granularity;

Module Two

just fields on fields

G_{16} : Category one of fields

G_{17} : Category two of fields

G_{18} : Category three of fields

T_{16} : Category one of fields

T_{17} : Category two of fields

T_{18} : Category three of fields

Module three

No more fields on spacetime

G_{20} : Category one of space and time

G_{21} : Category two of space and time

G_{22} : Category three of space and time

T_{20} : Category one of no fields (non existence of any sort of field. Make a list of all fields and the concomitant nonexistence thereof)

T_{21} : Category two of non existence of any sort of field

T_{22} : Category three of non existence of any sort of field

Module four

Idea that the statistical entropy governing thermal interactions of a black hole with its exterior **is (e) determined by** the microstates of the hole having distinct effects on the exterior, and over which a hole in a given macroscopic configuration thermally fluctuates. For a (macroscopically) Schwarzschild black hole this ensemble **is (e) formed by** horizons with the same area. We compute the number of states in this ensemble from first principles using nonperturbative loop quantum gravity. Statistical entropy **proportional to** the area, as in the Bekenstein-Hawking formula.© 1996 The American Physical SocietyURL:http://link.aps.org/doi/10.1103/PhysRevLett.77.3288DOI: 10.1103/PhysRevLett.77.3288
Black Hole Entropy from Loop Quantum Gravity Carlo Rovelli

G_{24} : Category one of microstates of the hole having distinct effects on the exterior, and over which a hole in a given macroscopic configuration thermally fluctuates; horizons with the same area

G_{25} : Category two of microstates of the hole having distinct effects on the exterior, and over which a hole in a given macroscopic configuration thermally fluctuates; horizons with the same area

G_{26} : Category three of microstates of the hole having distinct effects on the exterior, and over which a hole in a given macroscopic configuration thermally fluctuates; horizons with the same area

T_{24} : Category one of statistical entropy governing thermal interactions of a black hole with its exterior; Schwarzschild black hole this ensemble;

T_{25} : Category two of statistical entropy governing thermal interactions of a black hole with its exterior; Schwarzschild black hole this ensemble;

T_{26} : Category three of statistical entropy governing thermal interactions of a black hole with its exterior; Schwarzschild black hole this ensemble;

Module five

In so doing, as they describe in their paper published in the journal Physical Review Letters, they suggest that instead of a singularity existing at its center, **there (eb) is a** portal to another universe

G_{28} : Category one of nonexistence of singularity at the centre

G_{29} : Category two of nonexistence of singularity at the centre

G_{30} : Category three of nonexistence of singularity at the centre

T_{28} : Category one of singularity is portal to another universe

T_{29} : Category two of singularity is portal to another universe

T_{30} : Category three of singularity is portal to another universe

Module six

(1) For many years theoretical physicists have believed that the universe **came about (e) as a result of** a single Big bang event—Einstein's theories suggested it was so.
(2) The problem with this line of thinking however is that the theory of general relativity **can't (e)** describes what came before the singularity, which should exist at the point in time just before the Big Bang.
(3) Theory also suggests that a similar singularity **should (eb) exist at** the center of black holes
(4) General relativity **fails (e) to** describe them properly.
(5) Worse, there is the problem of the information loss paradox—if something falls into a black hole and is eventually squeezed to a singularity, what **happens(eb)** to the information it contained? Big Bang physicists can't say

G_{32} : Category one of single Big bang event; theory of general relativity; Theory also suggests that a similar singularity; Big bang theory;

G_{33} : Category two of single Big bang event; theory of general relativity; Theory also suggests that a similar singularity; Big bang theory;

G_{34} : Category three of single Big bang event; theory of general relativity; Theory also suggests that a similar singularity; Big bang theory;

T_{32} : Category one of universe; **can't (e)** describes what came before the singularity, which should exist at the point in time just before the Big Bang; center of black holes; cannot expatiate or justify information paradox

T_{33} : Category two of universe; ; **can't (e)** describes what came before the singularity, which should exist at the point in time just before the Big Bang; center of black holes; cannot expatiate or justify information paradox

T_{34} : Category three of universe; ; **can't (e)** describes what came before the singularity, which should exist at the point in time just before the Big Bang; center of black holes; cannot expatiate or justify information paradox

Module seven

To address these problems, Abhay Ashtekar and his team at Pennsylvania State University, back in 2006, came up with a theory known as loop quantum gravity. They suggested that instead of a singularity existing just before the Big Bang, there was the remains of a crunched down universe that had existed prior to the one that exists now. The universe didn't just Big Bang itself into existence from nothing, they said, instead there is an infinite loop **where a universe(eb) shrinks down to** a very tiny spot, then explodes in a Big Bang, then shrinks down again, over and over again forever—hence the use of the term "loop" in the theory.

Since that time, some in the field have begun to refer to the theory as the Big Bounce, to replace the name Big Bang.

G_{36} : Category one of universe didn't just Big Bang itself into existence from nothing, they said, instead there is an infinite loop

G_{37} : Category two of universe didn't just Big Bang itself into existence from nothing, they said, instead there is an infinite loop

G_{38} : Category three of universe didn't just Big Bang itself into existence from nothing, they said, instead there is an infinite loop

T_{36} : Category one of **universe (eb) shrinks down to** a very tiny spot, then explodes in a Big Bang, and then shrinks down again, over and over again forever—hence the use of the term "loop" in the theory. Since that time, some in the field have begun to refer to the theory as the Big Bounce, to replace the name Big Bang.

T_{37} : Category two of **universe (eb) shrinks down to** a very tiny spot, then explodes in a Big Bang, and then shrinks down again, over and over again forever—hence the use of the term "loop" in the theory. Since that time, some in the field have begun to refer to the theory as the Big Bounce, to replace the name Big Bang.

T_{38} : Category three of **universe (eb) shrinks down to** a very tiny spot, then explodes in a Big Bang, and then shrinks down again, over and over again forever—hence the use of the term "loop" in the theory. Since that time, some in the field have begun to refer to the theory as the Big Bounce, to replace the name Big Bang.

Module eight

everything that was pulled into the black hole didn't compress to a singularity after all—instead it was compressed to a certain small size, **then was spit(eb) out in** another part of the universe or into another universe entirely

G_{40} : Category one of everything that was pulled into the black hole didn't compress to a singularity after all—instead it was compressed to a certain small size

G_{41} : Category two of everything that was pulled into the black hole didn't compress to a singularity after all—instead it was compressed to a certain small size

G_{42} : Category three of everything that was pulled into the black hole didn't compress to a singularity after all—instead it was compressed to a certain small size

T_{40} : Category one of **spit(eb) out in** another part of the universe or into another universe entirely

T_{41} : Category two of **spit(eb) out in** another part of the universe or into another universe entirely

T_{42} : Category three of **spit(eb) out in** another part of the universe or into another universe entirely

Module Nine

Loop quantum gravity can account for the Bekenstein-Hawking entropy of a black hole **provided** a free parameter is chosen appropriately. New choice of the Immirzi parameter could predict both black hole entropy and the frequencies of quasinormal modes in the large n limit, **but at the price of changing** the

gauge group of the theory. . **Alejandro Cori chi's** argument **uses** strongly the necessity of having fermions satisfying basic symmetry and conservation principles, and therefore supports SU(2) as the relevant gauge group of the theory.(© 2003 The American Physical SocietyURL: http://link.aps.org/doi/10.1103/PhysRevD.67.087502DOI: 10.1103/PhysRevD.67.087502 **Quasinormal modes, black hole entropy, and quantum geometry Alejandro Corichi)**

G_{44} : Category one of Loop quantum gravity can account for the Bekenstein-Hawking entropy of a black hole(systems that satisfy LQG are taken in to account); gauge group of the theory; strongly the necessity of having fermions satisfying basic symmetry and conservation principles, and therefore supports SU(2) as the relevant gauge group of the theory;

G_{45} : Category two of Loop quantum gravity can account for the Bekenstein-Hawking entropy of a black hole; gauge group of the theory; strongly the necessity of having fermions satisfying basic symmetry and conservation principles, and therefore supports SU(2) as the relevant gauge group of the theory;

G_{46} : Category three of Loop quantum gravity can account for the Bekenstein-Hawking entropy of a black hole; strongly the necessity of having fermions satisfying basic symmetry and conservation principles, and therefore supports SU(2) as the relevant gauge group of the theory;

T_{44} : Category one of free parameter is chosen appropriately; New choice of the Immirzi parameter could predict both black hole entropy and the frequencies of quasinormal modes in the large n limit; **Alejandro Cori chi's** argument;

T_{45} : Category two of free parameter is chosen appropriately; New choice of the Immirzi parameter could predict both black hole entropy and the frequencies of quasinormal modes in the large n limit; **Alejandro Cori chi's** argument;

T_{46} : Category three of free parameter is chosen appropriately; New choice of the Immirzi parameter could predict both black hole entropy and the frequencies of quasinormal modes in the large n limit; **Alejandro Cori chi's** argument;

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$

$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$
$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,

2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt}=(a_{16})^{(2)}G_{17}-\begin{bmatrix}(a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t)\\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t)\end{bmatrix}G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt}=(a_{17})^{(2)}G_{16}-\begin{bmatrix}(a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t)\\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t)\end{bmatrix}G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt}=(a_{18})^{(2)}G_{17}-\begin{bmatrix}(a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t)\\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t)\end{bmatrix}G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt}=(b_{16})^{(2)}T_{17}-\begin{bmatrix}(b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t)\\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t)\\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t)\end{bmatrix}T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt}=(b_{17})^{(2)}T_{16}-\begin{bmatrix}(b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t)\\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t)\\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t)\end{bmatrix}T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad 73$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad 74$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad 76$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)} & \boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad 78$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad 81$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8\ ,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\left[-(b''_{36})^{(7)}(G_{39},t)\right] & \left[-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\left[-(b''_{37})^{(7)}(G_{39},t)\right] & \left[-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\left[-(b''_{38})^{(7)}(G_{39},t)\right] & \left[-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{matrix} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{matrix} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{matrix} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\left[-(b''_{40})^{(8)}(G_{43},t)\right] & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\left[-(b''_{41})^{(8)}(G_{43},t)\right] & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\left[-(b''_{42})^{(8)}(G_{43},t)\right] & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$

$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)}$ 98

$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$

$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + \ (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$

$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + \ (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)}$ 102

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with (116)
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \quad (117)$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - \ T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32{,}33{,}34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32{,}33{,}34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(IIIIIIIII) $\qquad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36{,}37{,}38$ 131

(JJJJJJJJJ)The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(KKKKKKKKK) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(LLLLLLLLL)

$\lim_{G\to\infty}(b_i'')^{(7)}\big((G_{39}),t\big) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\big((G_{39}),t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(MMMMMMMMM) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(NNNNNNNNN) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad (169)$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that (170)

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that (172)

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that (174)

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(kk) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})($and\ not\ on\ \ t$) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b'_i)^{(7)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}),(\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}),(\widetilde{T_{43}})\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad (333)$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad (334)$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad (335)$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad (336)$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(kk) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SIXTY SIX

Light Harvesting And Quantum Computation: Noscitur A Sociis : Meaning Of A Word Is From The Context

Abstract: Light harvestment and quantum computation are so much entangled with each other that only disentanglement and decoherence can separate them! It gives a big bold hi-tech face, differential griller projector recessed fog lamps, all muzzle and swanky style and wedgy swoop of belt line to the theory of quantum computation Near-unity efficiency of energy transfer in photosynthesis makes photosynthetic light-harvesting complexes a promising avenue **for developing** new renewable energy technologies. Knowledge of the energy landscape of these complexes **is essential** in understanding their function, but its experimental determination has proven elusive. Here, the observation of quantum coherence **using** two-dimensional electronic spectroscopy is employed to directly measure the 14 lowest electronic energy levels in light-harvesting complex II (LHCII), the most abundant antenna complex in plants containing approximately 50% of the world's chlorophyll. Electronically excited states are relatively evenly distributed, highlighting an important **design principle** of photosynthetic complexes **that explains** the observed ultrafast intracomplex energy transfer in LHCII. (Phys**. Chem. B, 2009, 113 (51), pp 16291–16295DOI: 10.1021/jp908300c (Web): October 20, 2009 © 2009 American Chemical Society Quantum Coherence Enabled Determination of the Energy Landscape in Light-Harvesting Complex II)** Light-harvesting components of photosynthetic organisms are complex, coupled, many-body quantum systems, in which electronic coherence has recently been shown to survive for relatively, long timescales, **despite the** decohering effects of their environments. Entanglement in multichromophoric light-harvesting complexes, and establish methods for quantification of entanglement **is done (eb) by the** authors by describing necessary and sufficient conditions for entanglement and by deriving a measure of global entanglement. These methods are then applied to the Fenna–Matthews–Olson protein **to (eb) extract the** initial state and temperature dependencies of entanglement. **Mohan Sarovar et al** show that, although the Fenna–Matthews–Olson protein in natural conditions largely **contains** bipartite entanglement between dimerized chromophores, a small amount of long-range and multipartite entanglement should exist even at physiological temperatures. **This constitutes** the first rigorous quantification of entanglement in a biological system. Practical use of entanglement in densely packed molecular aggregates such as light-harvesting complexes **is emphasised. Quantum entanglement in photosynthetic light-harvesting complexes Mohan Sarovar et al Nature Physics 6, 462–467 (2010) doi: 10.1038/nphys1652**

INTRODUCTION—VARIABLES USED

Light harvesting:

A light-harvesting complex is a complex of subunit proteins that may be part of a larger super complex of a photo system, the functional unit in photosynthesis. It is used by plants and photosynthetic bacteria to collect more of the incoming light than would be captured by the photosynthetic reaction center alone. Light-harvesting complexes are found in a wide variety among the different photosynthetic species. The complexes consist of proteins and photosynthetic pigments and surround a photosynthetic reaction center to

focus energy, attained from photons absorbed by the pigment, toward the reaction center using Förster resonance energy transfer.

Chlorophylls and carotenoids are important in light-harvesting complexes present in plants. Chlorophyll b is almost identical to chlorophyll a except it has a formyl group in place of a methyl group. This small difference makes chlorophyll b absorb light with wavelengths between 400 and 500 nm more efficiently. Carotenoids are long linear organic molecules that have alternating single and double bonds along their length. Such molecules are called polyenes. Two examples of carotenoids are lycopene and β-carotene. These molecules also absorb light most efficiently in the 400 – 500 nm range. Due to their absorption region, carotenoids appear red and yellow and provide most of the red and yellow colours present in fruits and flowers. The carotenoid molecules also serve a safeguarding function. Carotenoid molecules suppress damaging photochemical reactions, in particular those including oxygen, which exposure to sunlight can cause. Plants that lack carotenoid molecules quickly die upon exposure to oxygen and light.

Quantum biology refers to applications of quantum mechanics to biological objects and problems. Usually, it is taken to refer to applications of the "non-trivial" quantum features such as superposition, nonlocality, entanglement and tunneling, as opposed to the "trivial" but ubiquitous quantum mechanical nature of chemical bonding, ionization, and other phenomena that are the basis of the fundamental biophysics and biochemistry of organisms. Austrian born physicist and theoretical biologist Erwin Schrödinger, one of the founders of quantum theory in physics, was also one of the first scientists to suggest a study of quantum biology in his 1944 book "What is Life?" Many biological processes involve the conversion of energy into forms that are usable for chemical transformations and are quantum mechanical in nature. Such processes involve chemical reactions, light absorption, formation of excited electronic states, transfer of excitation energy, and the transfer of electrons and protons (hydrogen ions) in chemical processes such as photosynthesis and cellular respiration. Quantum biology uses computation to model biological interactions in light of quantum mechanical effects. Some examples of the biological phenomena that have been studied in terms of quantum processes are the absorbance of frequency-specific radiation (i.e., photosynthesis and vision); the conversion of chemical energy into motion; magneto reception in animals, DNA mutation and Brownian motors in many cellular processes. Recent studies have identified quantum coherence and entanglement between the excited states of different pigments in the light-harvesting stage of photosynthesis. Although this stage of photosynthesis is highly efficient, it remains unclear exactly how or if these quantum effects are relevant biologically.

Substantive limits, ontological consonance, presuppositioal resemblances, singular pauses, general rests and logical attributes and necessity of substantives and expressional proposition of physically intuitive mechanisms for the effect of noise **on (e&eb)** excitation energy transfer (EET) in networks is provided in extenso. Using these mechanisms of dephasing-assisted transport (DAT) in a hybrid basis of both excitons and sites, authors **develop a** detailed picture of how noise enables energy transfer with efficiencies well above $90%$ across the Fenna-Matthew-Olson (FMO) complex, a type of light harvesting molecule. It is demonstrated in explicit terms and unmistakable words how noise **alters(e&eb) the** pathways of energy transfer across the complex, **suppressing(e)** ineffective pathways and facilitating direct ones to the reaction centre. Fundamental mechanisms underpinning DAT are expected to be robust **with (e&eb) respect to** the considered noise model. Concomitant ramification thereof of the fact that specific details of the exciton-phonon coupling, which remains largely unknown in these types of complexes, and in particular the impact of non-Markovian effects, **results (eb) in** variations of dynamical features that should be amenable to experimental verification within current or planned technology. Detailed understanding of DAT in natural compounds **should (eb) open up** a new paradigm of `noise-engineering' by which EET can be optimized in artificial light-harvesting structures. Version published in NJP Contains a comprehensive new section on non-Markovian dynamics and also efficient FMO transport **induced(e) by** local low frequency vibrations

New J. Phys. 12 065002 (2010) arXiv:0910.4153 [quant-ph] **Noise-assisted energy transfer in quantum networks and light-harvesting complexes Alex W. Chin et al**

Researchers explain how dye-based nanotubes can help harvest light's energy tiny cylinders help reveal how natural-light-harvesting antennae collect light with exceptional efficiency. David L. Chandler, MIT News Office

Green sulfur bacteria, whose exceptional light-harvesting capabilities inspired the artificial system analyzed by postdoc Dörthe Eisele and her co-workers, dominate this hot spring at Yellowstone National Park and give it its striking green color.

Companies that make commercial solar cells are happy if they can achieve 20 percent efficiency when converting sunlight to electricity; an improvement of even 1 percent is seen as major progress. But nature, which has had billions of years to fine-tune photosynthesis, can do much better: Microorganisms called green sulfur bacteria, which live deep in the ocean where there's hardly any light available, manage to harvest 98 percent of the energy in the light that reaches them.

(1) Researchers led by an MIT postdoc have analyzed an artificial system that models the light-capturing method **used (e) by** deep-sea bacteria. Further advances in understanding fundamental light-harvesting processes **may yield** entirely new approaches to capturing solar energy, the researchers say. Their results were reported July 1 in the journal Nature Chemistry.

(2) The artificial system, described in an earlier paper by postdoc Dörthe M. Eisele of MIT's Research Laboratory of Electronics and collaborators, consists of a self-assembling system of dye molecules **that (eb) form** perfectly uniform double-walled nanotubes.

(3) These tubes — only about 10 nanometers wide but thousands of times longer — are similar in size, shape and function to natural receptors **used(e) by** green sulfur bacteria that collect energy from the tiny amounts of sunlight that penetrate to the ocean's depths."It is one of nature's great secrets, how to harvest light so efficiently," Eisele says. Her co-authors include Moungi G. Bawendi and the late Robert J. Silbey, both MIT professors of chemistry, along with collaborators at Humboldt University of Berlin, the University of Texas at Austin and the University of Groningen in the Netherlands.

(4) Eisele says this particular type of nanotube is unlikely to find practical applications. Rather, she says, these experiments were designed to study underlying principles that could **then be (e) used** to find optimal materials for particular uses. "This system is so interesting because it is a beautiful model system," she says, for analyzing how such structures respond to light.

(5) Unlike typical self-assembling systems in which each structure can be a bit different, these double-walled tubes, made from a cyanine-based dye, form perfectly uniform shapes and sizes.

(6) That makes it a perfect model system, Eisele explains, because knowing that all the nanotubes in a solution are identical **makes it possible (eb) to study their** properties in bulk, rather than having to isolate the response of each individual tube.

(7) One fundamental question the team wanted to address was whether the two concentric cylinders of the double-walled tubes were working together as an integrated system **for(e) capturing** light's energy, or whether each cylinder was acting on its own.

(8) To answer that question, Eisele and her team devised a way of deactivating one of the two cylinders by oxidizing the outer wall's molecules. "The tubular structure is still intact, **but (e) it kills the** optical response of the outer wall, so what is left is the optical response of the inner wall," she says. "It's a very simple and elegant way to isolate the spectrum of the inner wall."

(9) By comparing optical responses when both cylinders are functioning and when only one is working,

it's possible to **determine how** much interaction is taking place between the two cylinders. "When you watch the dynamics of the redox reaction," Eisele says, "you see that these two cylinders can be seen as two separate systems."

Characterizing this simplified artificial structure **may allow researchers** to build more efficient light-harvesting devices. "Nature had millions of years to optimize" how organisms capture energy, Eisele says; understanding how it did so may lead to better man-made systems.

"We don't want to improve the efficiency of solar cells we have now," she says. "We want to learn from nature how to build entirely new light-harvesting devices."

(10) Gregory Scholes, the D.J. Leroy Distinguished Professor of Chemistry at the University of Toronto, who was not involved in this work, says, "The researchers used exquisite experiments to test how the nanoscale components of this system interact after photoexcitation." He adds that the work "**provides (eb) important** insights into the design of large assemblies of molecules for applications in 'light harvesting.'"

The research was supported by the Deutsche Forschungsgemeinschaft, the Integrative Research Institute for the Sciences in Berlin, the National Science Foundation, the Alexander von Humboldt Foundation, the Department of Energy Center for Excitonics, the Army Research Office and the Defense Advanced Research Projects Agency.

NOTATION

Module One

Near-unity efficiency of energy transfer in photosynthesis makes photosynthetic light-harvesting complexes a promising avenue **for developing** new renewable energy technologies. Knowledge of the energy landscape of these complexes **is essential** in understanding their function, but its experimental determination has proven elusive. Here, the observation of quantum coherence **using** two-dimensional electronic spectroscopy is employed to directly measure the 14 lowest electronic energy levels in light-harvesting complex II (LHCII), the most abundant antenna complex in plants containing approximately 50% of the world's chlorophyll. Electronically excited states are relatively evenly distributed, highlighting an important **design principle** of photosynthetic complexes **that explains** the observed ultrafast intracomplex energy transfer in LHCII. . **Phys. Chem. B, 2009, 113 (51), pp 16291–16295DOI: 10.1021/jp908300c (Web): October 20, 2009 © 2009 American Chemical Society Quantum Coherence Enabled Determination of the Energy Landscape in Light-Harvesting Complex II**

G_{13} : Category one of two-dimensional electronic spectroscopy is employed to directly measure the 14 lowest electronic energy levels in light-harvesting complex II (LHCII), the most abundant antenna complex in plants containing approximately 50% of the world's chlorophyll.

G_{14} : Category two of two-dimensional electronic spectroscopy is employed to directly measure the 14 lowest electronic energy levels in light-harvesting complex II (LHCII), the most abundant antenna complex in plants containing approximately 50% of the world's chlorophyll.

G_{15} : Category three of two-dimensional electronic spectroscopy is employed to directly measure the 14 lowest electronic energy levels in light-harvesting complex II (LHCII), the most abundant antenna complex in plants containing approximately 50% of the world's chlorophyll.

T_{13} : Category one of observation of quantum coherence

T_{14} : Category two of observation of quantum coherence

T_{15} : Category three of observation of quantum coherence

Module Two

Using mechanisms of dephasing-assisted transport (DAT) in a hybrid basis of both excitons and sites, authors **develop a** detailed picture of how noise enables energy transfer with efficiencies well above \$90%\$ across the Fenna-Matthew-Olson (FMO) complex, a type of light harvesting molecule. It is demonstrated in explicit terms and unmistakable words how noise **alters(e&eb) the** pathways of energy transfer across the complex, **suppressing(e)** ineffective pathways and facilitating direct ones to the reaction centre. Fundamental mechanisms underpinning DAT are expected to be robust **with (e&eb) respect to** the considered noise model. Concomitant ramification thereof of the fact that specific details of the exciton-phonon coupling, which remains largely unknown in these types of complexes, and in particular the impact of non-Markovian effects, **results (eb) in** variations of dynamical features that should be amenable to experimental verification within current or planned technology. Detailed understanding of DAT in natural compounds **should (eb) open up** a new paradigm of `noise-engineering' by which EET can be optimized in artificial light-harvesting structures. Version published in NJP Contains a comprehensive new section on non-Markovian dynamics and also efficient FMO transport **induced(e) by** local low frequency vibrations New J. Phys. 12 065002 (2010) arXiv:0910.4153 [quant-ph] **Noise-assisted energy transfer in quantum networks and light-harvesting complexes Alex W. Chin et al**

G_{16} : Category one of how noise enables energy transfer with efficiencies well above \$90%\$ across the Fenna-Matthew-Olson (FMO) complex, a type of light harvesting molecule; ineffective pathways and facilitating direct ones to the reaction centre; Fundamental mechanisms underpinning DAT are expected to be robust; exciton-phonon coupling, which remains largely unknown in these types of complexes, and in particular the impact of non-Markovian effects;

G_{17} : Category two of how noise enables energy transfer with efficiencies well above \$90%\$ across the Fenna-Matthew-Olson (FMO) complex, a type of light harvesting molecule; ineffective pathways and facilitating direct ones to the reaction centre; Fundamental mechanisms underpinning DAT are expected to be robust; exciton-phonon coupling, which remains largely unknown in these types of complexes, and in particular the impact of non-Markovian effects;

G_{18} : Category three of how noise enables energy transfer with efficiencies well above \$90%\$ across the Fenna-Matthew-Olson (FMO) complex, a type of light harvesting molecule; ineffective pathways and facilitating direct ones to the reaction centre; Fundamental mechanisms underpinning DAT are expected to be robust; exciton-phonon coupling, which remains largely unknown in these types of complexes, and in particular the impact of non-Markovian effects;

T_{16} : Category one of mechanisms of dephasing-assisted transport (DAT) in a hybrid basis of both excitons and sites; noise **alters(e&eb) the** pathways of energy transfer across the complex; considered noise model; variations of dynamical features that should be amenable to experimental verification within current or planned technology;

T_{17} : Category two of mechanisms of dephasing-assisted transport (DAT) in a hybrid basis of both excitons and sites; noise **alters(e&eb) the** pathways of energy transfer across the complex; considered noise model; variations of dynamical features that should be amenable to experimental verification within current or planned technology;

T_{18} : Category three of mechanisms of dephasing-assisted transport (DAT) in a hybrid basis of both excitons and sites; noise **alters(e&eb) the** pathways of energy transfer across the complex; considered noise model; variations of dynamical features that should be amenable to experimental verification within current or planned technology

Module three

Researchers led by an MIT postdoc have analyzed an artificial system that models the light-capturing method **used (e) by** deep-sea bacteria. Further advances in understanding fundamental light-harvesting

processes **may yield** entirely new approaches to capturing solar energy, the researchers say. Their results were reported July 1 in the journal Nature Chemistry

G_{20} : Category one of fundamental light-harvesting processes

G_{21} : Category two of fundamental light-harvesting processes

G_{22} : Category three of fundamental light-harvesting processes

T_{20} : Category one of entirely new approaches to capturing solar energy

T_{21} : Category two of entirely new approaches to capturing solar energy

T_{22} : Category three of entirely new approaches to capturing solar energy

Module four

Concomitant ramification thereof of the fact that specific details of the exciton-phonon coupling, which remains largely unknown in these types of complexes, and in particular the impact of non-Markovian effects, **results (eb) in** variations of dynamical features that should be amenable to experimental verification within current or planned technology. New J. Phys. 12 065002 (2010) arXiv:0910.4153 [quant-ph] **Noise-assisted energy transfer in quantum networks and light-harvesting complexes Alex W. Chin et al**

G_{24} : Category one of exciton-phonon coupling, which remains largely unknown in these types of complexes, and in particular the impact of non-Markovian effects

G_{25} : Category two of exciton-phonon coupling, which remains largely unknown in these types of complexes, and in particular the impact of non-Markovian effects

G_{26} : Category three of exciton-phonon coupling, which remains largely unknown in these types of complexes, and in particular the impact of non-Markovian effects

T_{24} : Category one of variations of dynamical features that should be amenable to experimental verification within current or planned technology

T_{25} : Category two of variations of dynamical features that should be amenable to experimental verification within current or planned technology

T_{26} : Category three of variations of dynamical features that should be amenable to experimental verification within current or planned technology

Module five

Mohan Sarovar et al show that, although the Fenna–Matthews–Olson protein in natural conditions largely **contains** bipartite entanglement between dimerized chromophores, a small amount of long-range and multipartite entanglement should exist even at physiological temperatures

G_{28} : Category one of Fenna–Matthews–Olson protein in natural conditions largely **contains** bipartite entanglement between dimerized chromophores, a small amount of long-range and multipartite entanglement

G_{29} : Category two of Fenna–Matthews–Olson protein in natural conditions largely **contains** bipartite entanglement between dimerized chromophores, a small amount of long-range and multipartite entanglement

G_{30} : Category three of Fenna–Matthews–Olson protein in natural conditions largely **contains** bipartite entanglement between dimerized chromophores, a small amount of long-range and multipartite entanglement

T_{28} : Category one of physiological temperatures

T_{29} : Category two of physiological temperatures

T_{30} : Category three of physiological temperatures

Module six

Eisele says this particular type of nanotube is unlikely to find practical applications. Rather, she says, these experiments were designed to study underlying principles that could **then be (e) used** to find optimal materials for particular uses. "This system is so interesting because it is a beautiful model system," she says, for analyzing how such structures respond to light.

G_{32} : Category one of optimal materials for particular uses.

G_{33} : Category two of optimal materials for particular uses.

G_{34} : Category three of optimal materials for particular uses.

T_{32} : Category one of experiments designed to study underlying principles

T_{33} : Category two of experiments designed to study underlying principles

T_{34} : Category three of experiments designed to study underlying principles

Module seven

One fundamental question the team wanted to address was whether the two concentric cylinders of the double-walled tubes were working together as an integrated system **for(e) capturing** light's energy, or whether each cylinder was acting on its own.

G_{36} : Category one of light's energy, or whether each cylinder was acting on its own.

G_{37} : Category two of light's energy, or whether each cylinder was acting on its own.

G_{38} : Category three of light's energy, or whether each cylinder was acting on its own.

T_{36} : Category one of two concentric cylinders of the double-walled tubes were working together as an integrated system

T_{37} : Category two of two concentric cylinders of the double-walled tubes were working together as an integrated system

T_{38} : Category three of two concentric cylinders of the double-walled tubes were working together as an integrated system

Module eight

To answer that question, Eisele and her team devised a way of deactivating one of the two cylinders by oxidizing the outer wall's molecules. "The tubular structure is still intact, **but (e) it kills the** optical response

of the outer wall, so what is left is the optical response of the inner wall," she says. "It's a very simple and elegant way to isolate the spectrum of the inner wall."

G_{40} : Category one of optical response of the outer wall, so what is left is the optical response of the inner wall

G_{41} : Category two of optical response of the outer wall, so what is left is the optical response of the inner wall

G_{42} : Category three of optical response of the outer wall, so what is left is the optical response of the inner wall

T_{40} : Category one of deactivating one of the two cylinders by oxidizing the outer wall's molecules. "The tubular structure is still intact

T_{41} : Category two of deactivating one of the two cylinders by oxidizing the outer wall's molecules. "The tubular structure is still intact

T_{42} : Category three of deactivating one of the two cylinders by oxidizing the outer wall's molecules. "The tubular structure is still intact

Module Nine

Gregory Scholes, the D.J. Leroy Distinguished Professor of Chemistry at the University of Toronto, who was not involved in this work, says, "The researchers used exquisite experiments to test how the nanoscale components of this system interact after photoexcitation." He adds that the work "**provides (eb) important** insights into the design of large assemblies of molecules for applications in 'light harvesting

G_{44} : Category one of researchers used exquisite experiments to test how the nanoscale components of this system interact after photoexcitation

G_{45} : Category two of researchers used exquisite experiments to test how the nanoscale components of this system interact after photoexcitation

G_{46} : Category three of researchers used exquisite experiments to test how the nanoscale components of this system interact after photoexcitation

T_{44} : Category one of design of large assemblies of molecules for applications in 'light harvesting

T_{45} : Category two of design of large assemblies of molecules for applications in 'light harvesting

T_{46} : Category three of design of large assemblies of molecules for applications in 'light harvesting

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27}),t\big)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27}),t\big)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27}),t\big)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\big((G_{27}),t\big) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\big((G_{31}),t\big)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\big((G_{31}),t\big)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\big((G_{31}),t\big)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\big((G_{31}),t\big) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$
$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$
$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**
$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{37}'')^{(7,7,)}(G_{39},t)$, $-(b_{36}'')^{(7,7,)}(G_{39},t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8)}(G_{43},t)$ $-(b_{41}'')^{(8,8)}(G_{43},t)$ $-(b_{42}'')^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17},t) + (a_{13}'')^{(1,1,)}(T_{14},t) + (a_{20}'')^{(3,3,3)}(T_{21},t) \\ + (a_{24}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{28}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{32}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{36}'')^{(7,7,7)}(T_{37},t) + (a_{40}'')^{(8,8,8)}(T_{41},t) + (a_{44}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17},t) + (a_{14}'')^{(1,1,)}(T_{14},t) + (a_{21}'')^{(3,3,3)}(T_{21},t) \\ + (a_{25}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{29}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{33}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{37}'')^{(7,7,7)}(T_{37},t) + (a_{41}'')^{(8,8,8)}(T_{41},t) + (a_{45}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17},t) + (a_{15}'')^{(1,1,)}(T_{14},t) + (a_{22}'')^{(3,3,3)}(T_{21},t) \\ + (a_{26}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{30}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{34}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{38}'')^{(7,7,7)}(T_{37},t) + (a_{42}'')^{(8,8,8)}(T_{41},t) + (a_{46}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad 63$$

Where $+(a_{16}'')^{(2)}(T_{17},t)$, $+(a_{17}'')^{(2)}(T_{17},t)$, $+(a_{18}'')^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for*

category 1, 2 and 3

$-(b_{13}'')^{(1,1,1,)}(G,t)$, $-(b_{14}'')^{(1,1,1,)}(G,t)$, $-(b_{15}'')^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b_{36}'')^{(7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7)}(G_{39},t)$ $-(b_{38}'')^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{46}'')^{(9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a_{24}')^{(4)}+(a_{24}'')^{(4)}(T_{25},t) & +(a_{28}'')^{(5,5,)}(T_{29},t) & +(a_{32}'')^{(6,6,)}(T_{33},t)\\ +(a_{13}'')^{(1,1,1,1)}(T_{14},t) & +(a_{16}'')^{(2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3)}(T_{21},t)\\ +(a_{36}'')^{(7,7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9)}(T_{45},t)\end{bmatrix}G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a_{25}')^{(4)}+(a_{25}'')^{(4)}(T_{25},t) & +(a_{29}'')^{(5,5,)}(T_{29},t) & +(a_{33}'')^{(6,6)}(T_{33},t)\\ +(a_{14}'')^{(1,1,1,1)}(T_{14},t) & +(a_{17}'')^{(2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3)}(T_{21},t)\\ +(a_{37}'')^{(7,7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9)}(T_{45},t)\end{bmatrix}G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a_{26}')^{(4)}+(a_{26}'')^{(4)}(T_{25},t) & +(a_{30}'')^{(5,5,)}(T_{29},t) & +(a_{34}'')^{(6,6,)}(T_{33},t)\\ +(a_{15}'')^{(1,1,1,1)}(T_{14},t) & +(a_{18}'')^{(2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3)}(T_{21},t)\\ +(a_{38}'')^{(7,7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9)}(T_{45},t)\end{bmatrix}G_{26} \quad (75)$$

$(a_{24}'')^{(4)}(T_{25},t)$, $(a_{25}'')^{(4)}(T_{25},t)$, $(a_{26}'')^{(4)}(T_{25},t)$ *are first augmentation coefficients category* $1,2\ 3$

$+(a_{28}'')^{(5,5,)}(T_{29},t)$, $+(a_{29}'')^{(5,5,)}(T_{29},t)$, $+(a_{30}'')^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* $1,2$ *and* 3

$+(a_{32}'')^{(6,6,)}(T_{33},t)$, $+(a_{33}'')^{(6,6,)}(T_{33},t)$, $+(a_{34}'')^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* $1,2$ *and* 3

$+(a_{13}'')^{(1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category $1,2$ *and* 3

$+(a_{16}'')^{(2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category $1,2$ *and* 3

$+(a_{20}'')^{(3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category $1,2$ *and* 3

$+(a_{36}'')^{(7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category $1,2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a_{32}')^{(6)} & +(a_{32}'')^{(6)}(T_{33},t) & +(a_{28}'')^{(5,5,5)}(T_{29},t) & +(a_{24}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a_{33}')^{(6)} & +(a_{33}'')^{(6)}(T_{33},t) & +(a_{29}'')^{(5,5,5)}(T_{29},t) & +(a_{25}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a_{34}')^{(6)} & +(a_{34}'')^{(6)}(T_{33},t) & +(a_{30}'')^{(5,5,5)}(T_{29},t) & +(a_{26}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a_{32}'')^{(6)}(T_{33},t)$, $+(a_{33}'')^{(6)}(T_{33},t)$, $+(a_{34}'')^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a_{28}'')^{(5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a_{24}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{c} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{c} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(OOOOOOOOO) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(PPPPPPPPP) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(QQQQQQQQQ) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(RRRRRRRRR)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(SSSSSSSSS) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(TTTTTTTTT) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\leq\ \left(\hat{P}_{13}\right)^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\leq\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\leq\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\leq\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(ll) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$

In the same way , one can obtain

$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad (208)$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad (209)$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself (210)

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric (211)

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t}\}$$

Indeed if we denote (212)

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results (213)

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad (214)$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. (215)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}\ and\ (\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$

In the same way , one can obtain

$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25}\ \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25}\ \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}\ , G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\ , \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})},\widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})},\widetilde{(T_{31})}\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}$, $if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)}$,

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}$, $if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}$, $if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}$,

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(\mathrm{T}_{17},t)+(a''_{17})^{(2)}(\mathrm{T}_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)}G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)}G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)}T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^{0}e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^{0}e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^{0}}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t}-e^{-(b'_{34})^{(6)}t}\right]+T_{34}^{0}e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^{0}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T_{34}^{0}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)}+(C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $v^{(4)}$:- $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

For $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)}=\frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(ll) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523
A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SIXTY SEVEN

Quantum Computation In Neurobiological Framework: Nosce Te Ipsum :Know Thyself

Abstract: When we get to the topic and understanding of the Transcendental Logic, the brilliant and baffling Transcendental Deduction (TD) comes to the fore with all its scintillating and scarlet rocky razzmatazz and autumn sonata. Recall the two movements just discussed, the one from experience to its conditions and the one from the forms of valid inference to the concepts that we must use in all judging (the Categories). This duality **led Kant** to his famous question of right (quid juris) (A84=B116): with what right do we apply the Categories, which are not acquired from experience, to the contents of experience? (A85=B117). Kant's problem here is not as arcane as it might seem. It reflects an important question: How is it that the world as we experience it conforms to our logic? In briefest form, Kant thought that the trick to showing how it is possible for the Categories to apply to experience is to show that it is necessary that they apply (A97).TD has two sides, though Kant never treats them separately John Marlado, accounts, attributes, and ascribes the philosophy of Nishida KitarÅ is lucid, insightful, and deeply informative despite the highly questionable argument that gives the book its structure. In comparison with other, technically more accurate accounts of Nishida's philosophy, it is broader in scope than most but more single-minded in its approach. The analysis is for the most part limited to four books that have been translated into English, out of approximately 16 volumes of philosophical essays in Japanese, but it aims to capture the contours and most important details of Nishida's entire philosophy. To appreciate the achievement of this account, one would need only to read, in English translation or in the Japanese original, any of the volumes with their streams of repetitive, puzzling expressions and circuitous, often dead- end argumentation. Wilkinson's elucidation of details in the passages he samples makes it easier to understand why Nishida is considered Japan's greatest academic philosopher. A critical look at the frame of the book and some of its own claims reveals where, to my mind, it has gone astray. The argument that frames this book is repeated throughout: Nishida's philosophy is an **attempt to articulate, in a characteristically Western philosophical manner, the Zen experience he had as a** young man. Nishida first made the attempt by using the conceptual frameworks of

Western philosophers, and after they failed him he developed his own, largely by recasting Buddhist insights. In the end, a philosophical articulation of the Zen worldview in Western terms or even in Eastern terms recast by conceptual, Western philosophical methods approved impossible. What Nishida attempted is incommensurable with Western philosophy. While the **author first offers these statements as hypotheses (p. 2), one premise of his argument that there is such a thing as "the Zen world-view" or "the Zen experience "remains unquestioned; and another that Nishida's pivotal experience** was the Zen experienceâis asserted as "beyond question" (p. 151; see also pp. 28, 48, 10, 114).Wilkinson's presentation of Zen in the first chapter is drawn largely from D. T. Suzuki, with help from Suzuki's portage Abe Masao; some comments on time, birth, and death in; and a few other sources. The result is a caricature that imagines Zen as an invariable, unitary experience. In fact, Dagan's teachings on practice as the manifestation of enlightenment do not sit well with D. T. Suzuki's emphasis on a satori experience, and Suzuki does not enlighten us about what Zen in its various practices has historically been. The author would have benefited from acquaintance with other presentations by Zen teachers, such as that of ShunryÅ Suzuki'sZen Mind Beginner's Mind, with historical accounts of Zen traditions, and with contrasts between DÅgen's view of time and Sam Â· sÄraand that of other Buddhist philosophers. A little research in contemporary Zen scholarship would have shown how implausible is the talk of "nirvana or awareness of mu" (p. 24), "the basic Zen position [that] never varies" (p. 3), "the Zen vision" (pp. 5, 7), "Zen epistemology" , "Zen experience" (p. 59), "the conception of time involved in Zen" (p. 15), and "the fundamental Zen assertion that there is no absolute time" (p. 16). Even if DÅgen's notion that time and being are nondual (p. 16) epitomized most of Zen, Nishida's early notion of "a transcendent, unchanging reality apart from time" (p. 51) would contrast sharply with it. Equally questionable is Wilkinson's contention that Nishida's "pure experience" is equivalent to Zen satori. The notion of pure **experience Nishida developed in his first major work may be pivotal for his entire philosophy, but he never claims or i**mplies that any part of his philosophy presupposes an exceptional experience like satori that grounds his convictions and renders them unverifiable to those who have not had the experience, as Wilkinson intimates (pp. 55, 159, and 161). **(Source: Wikipedia, Stanford Encyclopedia, Philosopher's pages etc,)**

INTRODUCTION—VARIABLES USED

The Philosophy of Mysticism: Perennialism and Constructivism

Randolph T Dible I Study **the experiencer or the observer** and the transcendental subjectivity itself. Pure objectivity is essentially a deconstruction, apophatic and negativa The encountering of the experiencer or observer—transcendental subjectivity itself—at the foundation of the world leads inevitably to the recognition of pure objectivity as ultimate reality (which can be taken as its ultimate deconstruction, analogous to the apophatic or via negativa), from which objects derive their value, weight, significance, meaning or objectivity. In this way, pure objectivity can be seen as the supra-self-evident Axiological Axiom, so to speak, even Unconditional Love, in romantic terms. This axiology (value theory) has a structure inverse to the relationship between transcendental subjectivity as the radical unity of pure self-reference and on the other hand, the world of forms, as mere traces (representations, indications) of the unique, original "first distinction" Spencer-Brown speaks of at the foundation of his calculus. That is, all forms (i.e., distinctions, differences) would reduce to being the first distinction, also known as the marked state, which can be called penultimate reality (pure self-reference or transcendental subjectivity: the Spirit which animates us), except that forms are complimentary to their content, which is their objectivity or value, which would reduce to the unmarked state or ultimate reality. It is the incongruity of form (thoughts; Whitehead's "negative prehensions") and value (feelings; Whitehead's "positive prehensions," or my notion of objectivity, meaning and qualia; in short, the non-formal aspects of experience) that holds forms open and keeps them from absolute reduction. This accounts for the brute, concrete persistence of the "functional illusion"-- to use a term from Dzogchen Buddhism-- of the world. Thus this system has an axiology of

metaphysical objectivity grounded on the ideal of pure objectivity as the source of all value, meaning and significance, itself the very fecundity of profundity, which is the motive of drawing the distinction in the first place.(Wikipedia, Stanford encyclopedia and other Author's blogs)

Shifting Paradigms

The wisdom and understanding of our own divine essence is like a beaconing begorra, bureaucratic burdock, befuddling binary and digital digamma, that invites us inward toward ever deeper connection with all around us. Yet rather than look within, most in this world prefer to look outward for divine wisdom, thus sustaining a dependence upon a vast hierarchy that stretches between the individual and the Divine. In all our wanderings away from the Divine, humankind has obscured, obfuscated, obliterated, adjured, abnegated and abjured its most compelling features through a persistent belief systems in limitations arising out of the controlling mechanisms of this hierarchy of sub ordinates, superordinates and coordinates. Has my consciousness changed? Why that doubt? Twist and shake, ponderous novelty themes, pathos, poignancy in writing in different styles, exciting optimism of youth, power obmbudsman,pulsations and throbbing of pop, hard luck stories of people, modernity and mischief, grimalkins of rurality, wobbling boats of lovers, saxophone in mouths all point to newer value orientations.

The Divine dances and the **"Grand design "**is perpetrated outside of the confines of any hierarchical structures, be it feudal, organisational. It is complete within itself, and has a singular purpose of demonstrating the collective potential of all life within the universe. It is the archetype of perfection. It is the standard bearer of each soul's innate design and ultimate destiny. The essence of the Divine is far beyond mental conception, yet humanity's tendency is to resort to the limiting language of the hierarchical paradigm to define and understand it. Searching for Connection and Wholeness calls for not lumping elements like me made of clay feet and weak knees, owing allegiance to the fringe, without courante and courage of conviction and will for vindication, visual restlessness and exaggerated narratives of self to its own,exciting potentialities available, terrestrial passion, and skew of scams all bear an infallible demonstration for new order of consciousness. Old consciousness will not do. Newer one has to be imbibed and cultivated.

When people are unaware of their inner **wisdom and wholeness**, they tend to search for order and security outside themselves. That is my personal experience. There is always search for sanctity of emotion, aesthetic expression, pursuit of harmony, unraveling realms, syncopated ending and stillness of mind and shiver of heart at the mirror broken somewhere are the strains of newer wisdom if it can be called one. Same questions shall be asked and same shall be answer in the similar vein. Uncertain of their place within the hierarchical order of the world, they define themselves based upon their insecurities. Individuals thus become only pieces of their wholeness and like shards of glass from a beautiful vase; they bear little resemblance to their aggregate beauty. Within the hierarchy, many in high positions of power have taken advantage of our collective insecurities in attempting to guide the development of all humankind. They have obscured the direct connection between the individual and the Divine through a variety of means designed to intercede between our inner essence and our divine source.

Each individual can come to know them to be free of all forms of institutional and organisational and societal hierarchical control. This is not to imply that we should not trust others or join together in bonds of friendship and community. It is simply a reminder that relative truth is constantly shifting in the hands of those who desire to control. And even when the motive for their controlling behaviors may be of good will, it is still a form of control. When the revealers of "truth" within the hierarchy withhold and suppress information, they are usually positioning themselves to maintain or acquire more power rather than to disseminate empowerment to all. The desire for connection and wholeness is a fuel that drives us to seek out and explore the hierarchical paradigm. This inner longing **provides us with** the motivation to seek help and guidance from a specific group within the hierarchy, and in so doing, cultivate a sense of belonging and

connection. Furthermore, the hierarchical paradigm is a stage whereby we develop a sense of connection to some grand, encompassing vision. This is why the hierarchy nurtures prophets who point toward a greater vision. Spiritual leaders are able to peer deeply beneath the surface reality of life and experience how intricately connected every life form is, and how the composite of all life is intelligent far beyond measure. Scientists can interpret reality through their personal abilities to perceive and express life's dimensional depth and limitless intelligence. Yet no one is able to articulate life's full dimensional depth and breadth with the tools of language. They can only, at best, describe their interpretation or their impressions.

In actuality, all of us are able to peer beneath the surface reality of life and perceive a unique vision of the universe. We require only time and intention to develop our own interpretations. And this is precisely what many great spiritual leaders have taught. Life's deeper essence is not an absolute to be experienced by the chosen few, but an evolving, dynamic intelligence that wears as many faces as there are life forms. No group or people have the exclusive portal into the universe by which the Divine expresses itself in all its majesty. The portal is open and available to all, because the Divine is within all things. Those recognized as great prophets each produced a vision of the universe beyond what was currently defined by the hierarchy. Because their interpretations were articulated with authority and depth of insight, they became a target of debate among various groups in the hierarchy. This debate then created a polarity of belief. A sympathetic constituency emerged to defend and embellish their leader's interpretation, while established groups held it in contempt of previously held beliefs. Invariably, the leader's vision became confined and shaped into dogma by followers who desired to create a new religion or sect. Thus, this infusion of fresh insight quietly receded into the hands of the hierarchy, where its deeper meaning was obscured by the very fact that it was incorporated into a rigid structure that both protected and promoted it.

Paradigm Shift:

A new paradigm is emerging that promotes a clear connection of individual consciousness to the compelling features of the Divine without the intervention of a hierarchy. This is when the fables and myths of history step into the light and become known as they were originally intended. This is the time when language will be transformed into a new form of communication that breaks down all barriers of control. Personal transformation, through the awakening of inner wisdom and understanding, is the pathway into wholeness. The transformation paradigm is initiated simply by the recognition that rather than the dependency-inducing ways of the hierarchical paradigm, there are accelerated, independent pathways that bypass the hierarchy and lead to self-mastery. These new pathways lead to the divine wisdom and understanding that is present within all of us. This wisdom can be accessed through the practice of three principles of transformation: seeing the Divine in all, nurturance of life, and gratitude. The application of these life principles **disengages** individuals from the controlling elements of the hierarchy, **thus initiating the** transformation experience. As there are relative truths, there are relative freedoms. As individuals evolve through the hierarchical process, an ever-increasing sense of freedom is gained, yet external forces continue to exert control through limiting language, confining belief systems, and more. These controlling influences lead to continued reliance upon the hierarchy as it unceasingly attempts to impose a sense of inequality between us. The underlying equation of the hierarchical paradigm is: **individual + hierarchy = God connection**. In the case of personal transformation it is: **individual + inner wisdom and understanding divine equality with all**. The time has come to integrate the dominant paradigm of the hierarchy with the liberating transformation paradigm. That shall redeem all promises made to oneself. There shall be a **Matrix force or balancing force** that shall remain un satisfies until conservativeness is reached. This integration occurs naturally once we have fully explored the two paradigms and develop a synthesis paradigm whereby transformation is attained by searching for connection and wholeness through our own inner wisdom and understanding. It is this combination of release from dependency on the hierarchy and transformation from within that initiates the synthesis paradigm. Once we take responsibility for our own transformation and integration, it does not

mean that the hierarchy is to be shunned or avoided. The hierarchy is quite benign as a manipulative force and merely represents one important stage in the journey toward wholeness. What is being set in motion now is the initial preparation for these paradigm shifts. More specifically, these paradigms will be simultaneously played out over the coming years. As always, it will be the choice of each one of us as to which paradigm we embrace in our journey. All of the highest imaginings of humanity are yet unaware of our deepest foundation. We have sought the upper reaches of the building, yet remain unaware of the foundation's design. It is here, at the very core of existence that the Divine is bursting forth with its creative energy while simultaneously reintegrating with its invitation to wholeness. It is here that equality is realized, not in the lofty places of relative truth lodged in the hierarchy, but rather in the deepest part of the foundational plan of life's original source and ultimate destiny. The origin and destiny of existence is the tone of equality in life. Listen for this tone – this frequency of vibration – and follow it back into the very foundation from whence all things arise and return.

Consider these words as symbols only. Feel the truth that stands behind these words, and tap into this empowering energy force that reaches out for you. Know it as a tone or vibration – a resonance that waits for you around every corner in which your life will turn. It is a beacon of the Divine gathering itself into the form of language in order to usher you to a place from which you can experience the formless tone of equality – the bypass of limitation. It is the primal language of our divine source that bestows to you the freedom to generate your own deepest beauty in the expression of your highest truth. (**REFERENCE: WIKEPEDIA)**

PURE CONSCIOUSNESS

Subjective reports and physiological correlates, phenomenological attributes, predicational anteority, character constitution, primordial frontier of hypostatized signification and diurnal dynamics, anamensial alienisms, anchorite aperitif, Arcadian Atticism, of the experience of "consciousness itself" - self awareness isolated from the processes and objects of experience during Transcendental Meditation practice. Subjectively, this state is characterized by the absence of the very framework (time, space, and body sense) and content (qualities of inner and outer perception) that define waking experiences. Physiologically, this state is distinguished by the presence of apneustic breathing, autonomic orienting at the onset of breath changes, and increases in the frequency of peak EEG power. A model, called the junction point model, is presented that integrates pure consciousness with waking, dreaming, or sleeping. It could provide a structure to generate a coherent program of research to test the full range of consciousness and so enable us to understand what it means to be fully human. Some commentators believe that Kant's views on the mind are dependent on his idealism (he called it transcendental idealism). For the most part, that is not so. At worst, most of what he said about the mind and consciousness can be detached from his idealism. Though often viewed as a quintessentially German philosopher, Kant is said to have been one-quarter Scottish. Some philosophers (often Scottish) hold that 'Kant' is a Germanization of the Scottish name 'Candt', though many scholars now reject the idea. It is noteworthy, however, that his work on epistemology, which led him to his ideas about the mind, was a response to Hume as much as to any other philosopher. In general structure, Kant's model of the mind was the dominant model in the empirical psychology that flowed from his work and then again, after a hiatus during which behaviorism reigned supreme (roughly 1910 to 1965), toward the end of the 20th century, especially in cognitive science. Central elements of the models of the mind of thinkers otherwise as different as Sigmund Freud and Jerry Fodor are broadly Kantian, for example. Three ideas define the basic shape ('cognitive architecture') of Kant's model and one its dominant method. They have all become part of the foundation of cognitive science. The mind is complex set of abilities (functions). (As Meerbote 1989 and many others have observed, Kant held a functionalist view of the mind almost 200 years before functionalism was officially articulated in the 1960s by Hilary Putnam and others.) The functions crucial for mental, knowledge-generating activity are spatio-temporal processing of, and

application of concepts to, sensory inputs. Cognition requires concepts as well as percepts. These functions are forms of what Kant called synthesis. Synthesis (and the unity in consciousness required for synthesis) is central to cognition. These three ideas are fundamental to most thinking about cognition now. Kant's most important method, the transcendental method, is also at the heart of contemporary cognitive science. To study the mind, infer the conditions necessary for experience. Arguments having this structure are called transcendental arguments. Translated into contemporary terms, the core of this method is inference to the best explanation, the method of postulating unobservable mental mechanisms in order to explain observed behaviour. To be sure, Kant thought that he could get more out of his transcendental arguments than just 'best explanations'. He thought that he could get a priori (experience independent) knowledge out of them. Kant had a tripartite doctrine of the a priori. He held that some features of the mind and its knowledge had a priori origins, i.e., must be in the mind prior to experience (because using them is necessary to have experience). That mind and knowledge have these features are a priori truths, i.e., necessary and universal (B3/4). And we can come to know these truths, or that they are a priori at any rate, only by using a priori methods, i.e., we cannot learn these things from experience (B3) (Brook 1993). Kant thought that transcendental arguments were a priori or yielded the a priori in all three ways. Nonetheless, at the heart of this method is inference to the best explanation. When introspection fell out of favour about 100 years ago, the alternative approach adopted was exactly this approach. Its nonempirical roots in Kant notwithstanding, it are now the major method used by experimental cognitive scientists.

Other things equally central to Kant's approach to the mind have not been taken up by cognitive science, as we will see near the end, a key part of his doctrine of synthesis and most of what he had to say about consciousness of self in particular. Far from his model having been superseded by cognitive science, some important things have not even been assimilated by it.

The major works so far as Kant's views on the mind are concerned are the monumental Critique of Pure Reason (CPR) and his little, late Anthropology from a Pragmatic Point of View, first published in 1798 only six years before his death. Since the Anthropology was worked up from notes for popular lectures, it is often superficial compared to CPR. Kant's view of the mind arose from his general philosophical project in CPR the following way. Kant aimed among other things to, justify our conviction that physics, like mathematics, is a body of necessary and universal truth. Insulate religion, including belief in immortality, and free will from the corrosive effects of this very same science. Kant accepted without reservation that "God, freedom and immortality" (1781/7, Bxxx) exist but feared that, if science were relevant to their existence at all, it would provide reasons to doubt that they exist. As he saw it and very fortunately, science cannot touch these questions. "I have found it necessary to deny knowledge, in order to make room for faith." (Bxxx, his italics). Laying the foundation for pursuit of the first aim, which as he saw it was no less than the aim of showing why physics is a science, was what, led Kant to his views about how the mind works. He approached the grounding of physics by asking: What are the necessary conditions of experience (A96)? Put simply, he held that for our experience, and therefore our minds, to be as they are, the way that our experience is tied together must reflect the way that physics says that objects in the world must be tied together. Seeing this connection also tells us a lot about what our minds must be like.

His pursuit of the second aim, and specifically his critique of some arguments of his predecessors that entailed that we can know more about the mind's consciousness of itself than Kant could also, led him to some extraordinarily penetrating ideas about our consciousness of ourselves. In CPR, Kant discussed the mind only in connection with his main projects, never in its own right, so his treatment is remarkably scattered and sketchy. As he put it, "Enquiry … [into] the pure understanding itself, its possibility and the cognitive faculties upon which it rests … is of great importance for my chief purpose, … [but] does not form an essential part of it" (Axvii). Indeed, Kant offers no sustained, focused discussion of the mind anywhere in his work except the popular Anthropology, which, as we just said, is quite superficial**.(Wikipedia and other**

Philosophers' especially Kant's page)

If the nature is nonlinear, and thinking is linear, speaking is linear, and then the very act of speaking and interaction must create a bias as a participant. The concept of witness consciousness is destroyed here. Does Klossowski's thought mean to say that speaking prevents us from thinking nasty things? No! It cannot be! What he says is in fact that pure knowledge that produces an impure silence is a provocation of mind by the body; similarly, the impure language which produces a pure silence is the revocation of the body by mind. As Sade's heroes say it is not the bodies which are present that excite the libertine, but the great idea that is not there. Language in its wide ranging manifestations becomes forces giving thereby to the disintegrated body and the dissolved self access to a silence which is that of innocence's it means there is purity of responsibility and purity of innocence. Le Baphiomet says: either the words are recalled but their senses become obscure; or the sense appears when the memory of the words disappears.

In the profound, the nature of dilemma is still theological Octave is a Professor of Theology. Le Baphiomet is in its entirety is a theological model, which opposes the system of God, and is therefore Anti Brahman. So these terms are of fundamental disjunction. In the order of Brahman, in the order of existence, bodies give to minds or impose on them, catapult them to two phenomenological properties: identity and immortality, personality and resurrect ability, incommunicability and integrity. What is incommunicability? It is the principle according to which individual would not be attributable to several individuals and being of the individual to be a self identical person (Antoine)

The order of Brahman includes the following elements:

The identity of Brahman as the ultimate terra firma

The identities of human bodies or in general bodies as the bastion

The identity of the world as the principal frontier of bipolar counter actualities and variant and virtuous action.

The identity of bodies as the sentinel of depth-length, theme –potentialities, profiles and unity of perspectives, fringe-centre, text-context, the tic and non the tic

But this order of Brahman is constructed against the order of Anti Brahman, an order, this order of Anti Brahman subsists in the Brahman (See the Model) and weakens him, dissipates him little by little. We shall here do not make any assumptions of co creations which some Vedas and scriptures do. It is at this point that the story of Baphiomet begins: in the service of Brahman, the great master of Templars has as its mission, the sorting out of spirits and preventing their mixing together while waiting for the day of resurrection. The return of God, Christ, Brahman. We will not further elaborate upon the Theories of Knowledge, especially that of Beingness and Nothingness which has been vehemently and trenchantly criticized and has on the other hand received accolades and plaudits; There exits transitive states and substantive sub states of determinate orientation. Thus we shall refer Brahman and Anti Brahman are the two sides of the same coin. What was virtue yesterday is vice today; and what is vice today is virtue tomorrow; the cycle never ends……..

Ervin Laszlo Systems philosopher, integral theorist, concert pianist Posted: March 8, 2010 11:19 AMBIOBecome a Fan Get Email Alerts Bloggers' Index Why Your Brain Is A Quantum Computer emphasis mine . I have removed some sentences due to spatial constraints. Notwithstanding the essence and point of view that the article conveys remains the same.

(1) Scientists have believed that the brain operates **as a** biochemical and bioelectric system.

(2) Individual brain cells, so-called neurons, fire in complex coordinated patterns, and their chemical and electrical discharges **make (e) up a** network that processes information.
(3) Somehow, this information (or some part of it) **gets (e&eb) translated** into conscious "mind-events": shapes and colors, sounds, and the other "data" of your senses.

However, there is mounting evidence that your brain doesn't operate merely by biochemical and bioelectric information processing. What your brain does is far more complex, fast, and sophisticated than could be accomplished by this standard kind of information-processing. Thousands of chemical reactions take place every second in every cell of your body, and your brain and nervous system ensure that they are sufficiently coherent and coordinated so that your body can maintain itself in the complex and physically highly improbable state we call living.

(4) Your brain is the command center that directs the flow of the precise, highly coordinated information crucial to your body's genetic, chemical, and physiological processes. These processes **are based on** the interaction of molecules and atoms, and subatomic particles.
(5) Though in large part biochemical, these processes have a significant nonlocal component as well. By "nonlocal," I refer to the finding in physics that particles once connected **retain (e&eb)** their connection and influence upon each other even at a distance--they are said to remain "entangled."
(6) Particles and atoms in body are "entangled" with each other: they **receive and (e&eb) transmit information** not just by biochemical means, but by the remarkable process known in quantum physics as "phase-conjugate quantum resonance." Phase-conjugate quantum resonance is a term used by physicists that means that the particles are "non-locally entangled."
(7) What's revealed at the leading edge of quantum physics and quantum biology is that your body is not just a biochemical system: it's also a "macroscopic quantum system." Quantum systems were believed to exist only at the submicroscopic level, where quanta are in the state known as "coherent," **which means that** they are able to get into synch with each other.
(8) Interactions at ordinary dimensions and temperatures were believed **to destroy** their coherence: they render macroscale bodies "decoherent."
(9) In other words, it was believed that once these quanta assembled themselves into particles, atoms, molecules, cells, and ultimately complex systems like the human body, the relationships that existed between these building blocks **ceased to operate** according to the same laws of physics or to function in the same manner as they did on the submicroscopic level. Instead of continuing their non-local dance of coherence, **they ceased to be** in this astonishingly close kind of relationship and went largely their separate ways.
(10) But this is not entirely the case. Some theories claim that specifically organized networks of quanta--for example, networks where the particles are "woven" or "braided"--**are sufficiently (eb) robust to maintain** quantum coherence at macroscopic dimensions and ordinary temperatures. In other words, the dance continues even when the particles are organized into larger systems like a human brain or body.
(11) This makes sense because living systems **exhibit highly** and until recently inexplicably coherent behavior.
(12) Their cells and organs resonate in phase, and the entire living organism **seems(eb) to obey one** encompassing "macroscopic wave-function."
(13) In other words, instead of functioning like a bunch of cells and chemicals each doing their own thing unaware of each other, all the biochemical and bioelectric **dance in superb (e&eb) coordination** acts like a giant wave which moves and flows as one, despite the many individual droplets that are within it. It is like the individual debits and credits of the bank . Individually they tally with each other and after disorganization and reorganization at the General ledger, they balance each other with the general theory of the General ledger stating Assets= Liabilities,

irrespective and regardless of the transactions or the volume and the centres of origination or destination. **This however is my analogy**

(14) This means that your body is not just a biochemical system: **it's also a** macroscopic quantum system.

(15) Your brain is not just a bioelectric and biochemical computer, **but also** a quantum computer.

(16) The cells of your body, and the neurons and networks of neurons of your brain, **are entangled** with each other.

(17) This is why your brain can perform functions that are entire dimensions beyond the capacity of any conceivable biochemical system. If and when quantum computers are developed, we may for the first time see man-made systems that come close to matching your brain's information-processing powers.

Scientists map process by which brain cells form long-term memories June 9, 2013 in Neuroscience

(1) How connections in the brain must change **to form** memories could help **to(eb) develop** artificial cognitive computers November 7, 2012 in Neuroscience A schematic diagram depicting the recall of a sequence of memory items when the network containing the pool of memory items is triggered **by (e) a** stimulus. Credit: 2012 A*STAR Institute for Infocomm Research Exactly how memories are stored and accessed in the brain is unclear. **We give a Sundry Deposit Model of Bank wherein the amount is credited and accessed no sooner than the account is found (Model by me).**

(2) Neuroscientists, however, do know that a primitive structure buried in the center of the brain, called the hippocampus, is a pivotal region **of (e)** memory formation.

(3) Here, changes in the strengths of connections between neurons, which are called synapses, **are (e) the basis** for memory formation.

(4) Networks of neurons linking up in the hippocampus are likely **to (e) encode** specific memories.

(5) Since direct tests cannot be performed in the brain, experimental evidence for this process of memory formation is difficult to obtain but mathematical and computational models can provide insight. To this end, Eng Yeow Cheu and co-workers at the A*STAR Institute for Infocomm Research, Singapore, have developed a model that **sheds light** on the exact synaptic conditions required in memory formation. Their work builds on a previously proposed model of auto-associative memory, a process **whereby (e) a** memory is retrieved or completed after partial activation of its constituent neural network.

(6) The earlier model proposed that neural networks encoding short-term memories are **activated (eb)** at specific points during oscillations of brain activity.

(7) Changes in the strengths of synapses, and therefore the abilities of neurons in the network to activate each other, **lead to** an auto-associative long-term memory.

NOTATION

Module One

Quantum mechanics in the brain Does the enormous computing power of neurons mean consciousness can be explained within a purely neurobiological framework, or is there scope for quantum computation in the brain? Christof Koch and Klaus Hepp

G_{13} : Category one of coherent states ;neurons either spikes **(**fasten, impale, lance, make fast, nail, pin, prick, skewer, spear, spit, stick, transfix)at a particular point in time or it does not

G_{14} : Category two of coherent states ;neurons either spikes **(**fasten, impale, lance, make fast, nail, pin, prick, skewer, spear, spit, stick, transfix)at a particular point in time or it does not

G_{15} : Category three of coherent states ;neurons either spikes **(**fasten, impale, lance, make fast, nail, pin, prick, skewer, spear, spit, stick, transfix)at a particular point in time or it does not

T_{13} : Category one of chemical transmission across the synaptic cleft and generation of action potentials which involve thousands of ions coupled by diffusion or by membrane potential that extends across tens of micrometers; neurons does not exist in spike and nonspike states

T_{14} : Category two of chemical transmission across the synaptic cleft and generation of action potentials which involve thousands of ions coupled by diffusion or by membrane potential that extends across tens of micrometers; neurons does not exist in spike and nonspike states

T_{15} : Category three of chemical transmission across the synaptic cleft and generation of action potentials which involve thousands of ions coupled by diffusion or by membrane potential that extends across tens of micrometers; neurons does not exist in spike and nonspike states

Module Two

Everything=Nothing

G_{16} : Category one of Nothing (Nothing exists in everything. In all walks of life that is the final conclusion we arrive at. There is nothing in what is being done. Only another job)

G_{17} : Category two of Nothing (Nothing exists in everything. In all walks of life that is the final conclusion we arrive at. There is nothing in what is being done. Only another job)

G_{18} : Category three of Nothing (Nothing exists in everything. In all walks of life that is the final conclusion we arrive at. There is nothing in what is being done. Only another job)

T_{16} : Category one of Everything (it is based on perception and the classification on the idiosyncrasies' and propensities of the investigating systems that are expositional in its thematic and discursive form)

T_{17} : Category two of Everything (it is based on perception and the classification on the idiosyncrasies' and propensities of the investigating systems that are expositional in its thematic and discursive form)

T_{18} : Category three of Everything (it is based on perception and the classification on the idiosyncrasies' and propensities of the investigating systems that are expositional in its thematic and discursive form)

Module three (Important module)

Ervin Laszlo Systems philosopher, integral theorist, concert pianist Posted: March 8, 2010 11:19 AMBIOBecome a Fan Get Email Alerts Bloggers' Index Why Your Brain Is A Quantum Computer

Individual brain cells, so-called neurons, fire in complex coordinated patterns, and their chemical and electrical discharges **make (e) up a** network that processes information.

Somehow, this information (or some part of it) **gets (e&eb) translated** into conscious "mind-events": shapes and colors, sounds, and the other "data" of your senses.

Your brain is the command center that directs the flow of the precise, highly coordinated information crucial to your body's genetic, chemical, and physiological processes. These processes **are based on** the interaction of molecules and atoms, and subatomic particles.

Though in large part biochemical, these processes have a significant nonlocal component as well. By "nonlocal," reference is made to the finding in physics that particles once connected **retain (e&eb)** their

connection and influence upon each other even at a distance--they are said to remain "entangled."

Particles and atoms in body are "entangled" with each other: they **receive and (e&eb) transmit information** not just by biochemical means, but by the remarkable process known in quantum physics as "phase-conjugate quantum resonance." Phase-conjugate quantum resonance is a term used by physicists that means that the particles are "non-locally entangled."

What's revealed at the leading edge of quantum physics and quantum biology is that your body is not just a biochemical system: it's also a "macroscopic quantum system." Quantum systems were believed to exist only at the submicroscopic level, where quanta are in the state known as "coherent," **which means that** they are able to get into synch with each other.

Interactions at ordinary dimensions and temperatures were believed **to destroy** their coherence: they render macroscale bodies "decoherent."

In other words, it was believed that once these quanta assembled themselves into particles, atoms, molecules, cells, and ultimately complex systems like the human body, the relationships that existed between these building blocks **ceased to operate** according to the same laws of physics or to function in the same manner as they did on the submicroscopic level. Instead of continuing their non-local dance of coherence, **they ceased to be** in this astonishingly close kind of relationship and went largely their separate ways.

But this is not entirely the case. Some theories claim that specifically organized networks of quanta--for example, networks where the particles are "woven" or "braided"--**are sufficiently (eb) robust to maintain** quantum coherence at macroscopic dimensions and ordinary temperatures. In other words, the dance continues even when the particles are organized into larger systems like a human brain or body.

This makes sense because living systems **exhibit highly** and until recently inexplicably coherent behavior. Their cells and organs resonate in phase, and the entire living organism **seems (eb) to obey one** encompassing "macroscopic wave-function."

In other words, instead of functioning like a bunch of cells and chemicals each doing their own thing unaware of each other, all the biochemical and bioelectric **dance in superb (e&eb) coordination** acts like a giant wave which moves and flows as one, despite the many individual droplets that are within it. It is like the individual debits and credits of the bank. Individually they tally with each other and after disorganization and reorganization at the General ledger, they balance each other with the general theory of the General ledger stating Assets= Liabilities, irrespective and regardless of the transactions or the volume and the centres of origination or destination. **This however is my analogy**

This means that your body is not just a biochemical system: **it's also a** macroscopic quantum system.

Your brain is not just a bioelectric and biochemical computer, **but also** a quantum computer.

The cells of your body, and the neurons and networks of neurons of your brain, **are entangled** with each other.

This is why your brain can perform functions that are entire dimensions beyond the capacity of any conceivable biochemical system. If and when quantum computers are developed, we may for the first time see man-made systems that come close to matching your brain's information-processing powers.

Note: This module contains 19 models, attributed and ascribed to spatial constraints. Reader is advised to

identify the parameters correctly before reading the holistic and consummate model.

G_{20} : Category one of Individual brain cells, so-called neurons, fire in complex coordinated patterns, and their chemical and electrical discharges; this information (or some part of it); interaction of molecules and atoms, and subatomic particles; Though in large part biochemical, these processes have a significant nonlocal component as well. By "nonlocal," reference is made to the finding in physics that particles once connected; Particles and atoms in body are "entangled" with each other: they **receive and (e&eb) transmit information** ; quantum mechanical system; according to the same laws of physics or to function in the same manner as they did on the submicroscopic level. Instead of continuing their non-local dance of coherence, **they ceased to be** in this astonishingly close kind of relationship and went largely their separate ways; This makes sense because living systems **exhibit highly** and until recently inexplicably coherent behavior. Their cells and organs resonate in phase, and the entire living organism;

G_{21} : Category two of Individual brain cells, so-called neurons, fire in complex coordinated patterns, and their chemical and electrical discharges; this information (or some part of it); interaction of molecules and atoms, and subatomic particles; Though in large part biochemical, these processes have a significant nonlocal component as well. By "nonlocal," reference is made to the finding in physics that particles once connected; Particles and atoms in body are "entangled" with each other: they **receive and (e&eb) transmit information** ; quantum mechanical system; according to the same laws of physics or to function in the same manner as they did on the submicroscopic level. Instead of continuing their non-local dance of coherence, **they ceased to be** in this astonishingly close kind of relationship and went largely their separate ways; This makes sense because living systems **exhibit highly** and until recently inexplicably coherent behavior. Their cells and organs resonate in phase, and the entire living organism;

G_{22} : Category three of Individual brain cells, so-called neurons, fire in complex coordinated patterns, and their chemical and electrical discharges; this information (or some part of it); interaction of molecules and atoms, and subatomic particles; Though in large part biochemical, these processes have a significant nonlocal component as well. By "nonlocal," reference is made to the finding in physics that particles once connected; Particles and atoms in body are "entangled" with each other: they **receive and (e&eb) transmit information** ;quantum mechanical system; according to the same laws of physics or to function in the same manner as they did on the submicroscopic level. Instead of continuing their non-local dance of coherence, **they ceased to be** in this astonishingly close kind of relationship and went largely their separate ways; This makes sense because living systems **exhibit highly** and until recently inexplicably coherent behavior. Their cells and organs resonate in phase, and the entire living organism;

T_{20} : Category one of network that processes information; conscious "mind-events": shapes and colors, sounds, and the other "data" of your senses; brain is the command center that directs the flow of the precise, highly coordinated information crucial to your body's genetic, chemical, and physiological processes; connection and influence upon each other even at a distance--they are said to remain "entangled."; not just by biochemical means, but by the remarkable process known in quantum physics as "phase-conjugate quantum resonance." Phase-conjugate quantum resonance is a term used by physicists that means that the particles are "non-locally entangled.";body; once these quanta assembled themselves into particles, atoms, molecules, cells, and ultimately complex systems like the human body, the relationships that existed between these building blocks; encompassing "macroscopic wave-function." ;

T_{21} : Category two of network that processes information; conscious "mind-events": shapes and colors, sounds, and the other "data" of your senses; brain is the command center that directs the flow of the precise, highly coordinated information crucial to your body's genetic, chemical, and physiological processes; connection and influence upon each other even at a distance--they are said to remain "entangled."; not just by biochemical means, but by the remarkable process known in quantum physics as "phase-conjugate

quantum resonance." Phase-conjugate quantum resonance is a term used by physicists that means that the particles are "non-locally entangled.";body; once these quanta assembled themselves into particles, atoms, molecules, cells, and ultimately complex systems like the human body, the relationships that existed between these building blocks; encompassing "macroscopic wave-function." ;

T_{22} : Category three of network that processes information; conscious "mind-events": shapes and colors, sounds, and the other "data" of your senses; brain is the command center that directs the flow of the precise, highly coordinated information crucial to your body's genetic, chemical, and physiological processes; connection and influence upon each other even at a distance--they are said to remain "entangled."; not just by biochemical means, but by the remarkable process known in quantum physics as "phase-conjugate quantum resonance." Phase-conjugate quantum resonance is a term used by physicists that means that the particles are "non-locally entangled.";body; once these quanta assembled themselves into particles, atoms, molecules, cells, and ultimately complex systems like the human body, the relationships that existed between these building blocks; encompassing "macroscopic wave-function." ;

Module four

How connections in the brain must change **to form** memories could help **to (eb) develop** artificial cognitive computers November 7, 2012 in Neuroscience A schematic diagram depicting the recall of a sequence of memory items when the network containing the pool of memory items is triggered **by (e) a** stimulus. Credit: 2012 A*STAR Institute for Infocomm Research Exactly how memories are stored and accessed in the brain is unclear. **We give a Sundry Deposit Model of Bank wherein the amount is credited and accessed no sooner than the account is found (Model by me).**

G_{24} : Category one of connections in the brain must change

G_{25} : Category two of connections in the brain must change

G_{26} : Category three of connections in the brain must change

T_{24} : Category one of memories could help **to (eb) develop** artificial cognitive computers

T_{25} : Category two of memories could help **to (eb) develop** artificial cognitive computers

T_{26} : Category three of memories could help **to (eb) develop** artificial cognitive computers

Module five

Here, changes in the strengths of connections between neurons, which are called synapses, **are (e) the basis** for memory formation

G_{28} : Category one of memory formation

G_{29} : Category two of memory formation

G_{30} : Category three of memory formation

T_{28} : Category one of changes in the strengths of connections between neurons, which are called synapses

T_{29} : Category two of changes in the strengths of connections between neurons, which are called synapses

T_{30} : Category three of changes in the strengths of connections between neurons, which are called synapses

Module six

Networks of neurons linking up in the hippocampus are likely **to (e) encode** specific memories

G_{32} : Category one of **encode** specific memories

G_{33} : Category two of **encode** specific memories

G_{34} : Category three of **encode** specific memories

T_{32} : Category one of Networks of neurons linking up in the hippocampus

T_{33} : Category two of Networks of neurons linking up in the hippocampus

T_{34} : Category three of Networks of neurons linking up in the hippocampus

Module seven

Since direct tests cannot be performed in the brain, experimental evidence for this process of memory formation is difficult to obtain but mathematical and computational models can provide insight. To this end, Eng Yeow Cheu and co-workers at the A*STAR Institute for Infocomm Research, Singapore, have developed a model that **sheds light** on the exact synaptic conditions required in memory formation. Their work builds on a previously proposed model of auto-associative memory, a process **whereby (e) a** memory is retrieved or completed after partial activation of its constituent neural network.

G_{36} : Category one of memory is retrieved or completed after partial activation of its constituent neural network.

G_{37} : Category two of memory is retrieved or completed after partial activation of its constituent neural network.

G_{38} : Category three of memory is retrieved or completed after partial activation of its constituent neural network.

T_{36} : Category one of exact synaptic conditions required in memory formation, auto-associative memory

T_{37} : Category two of exact synaptic conditions required in memory formation, auto-associative memory

T_{38} : Category three of exact synaptic conditions required in memory formation, auto-associative memory

Module eight

The earlier model proposed that neural networks encoding short-term memories are **activated (eb)** at specific points during oscillations of brain activity.

G_{40} : Category one of neural networks encoding short-term memories

G_{41} : Category two of neural networks encoding short-term memories

G_{42} : Category three of neural networks encoding short-term memories

T_{40} : Category one of specific points during oscillations of brain activity

T_{41} : Category two of specific points during oscillations of brain activity

T_{42} : Category three of specific points during oscillations of brain activity

Module Nine

Changes in the strengths of synapses, and therefore the abilities of neurons in the network to activate each other, **lead to** an auto-associative long-term memory

G_{44} : Category one of Changes in the strengths of synapses, and therefore the abilities of neurons in the network to activate each other,

G_{45} : Category two of Changes in the strengths of synapses, and therefore the abilities of neurons in the network to activate each other

G_{46} : Category three of Changes in the strengths of synapses, and therefore the abilities of neurons in the network to activate each other

T_{44} : Category one of auto-associative long-term memory

T_{45} : Category two of auto-associative long-term memory

T_{46} : Category three of auto-associative long-term memory

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$

$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$
$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$
$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$
$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} \quad \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{ccc} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} \quad \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} \quad \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation

coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\left[-(b''_{13})^{(1)}(G,t)\right] & \left[-(b''_{16})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\left[-(b''_{14})^{(1)}(G,t)\right] & \left[-(b''_{17})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\left[-(b''_{15})^{(1)}(G,t)\right] & \left[-(b''_{18})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad 73$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad 74$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad 76$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)} & \boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category $1, 2$ *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{matrix} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)}\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{matrix} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)}\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{matrix} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)}\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32{,}33{,}34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32{,}33{,}34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(UUUUUUUUU) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36{,}37{,}38$ 131

(VVVVVVVVV) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$

132

(WWWWWWWWW) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(XXXXXXXXX)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(YYYYYYYYY) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(ZZZZZZZZZ) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(mm) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \le (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \le \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}ds_{(16)} +$

$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t} \; and \; (\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2 \; and \; \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37} \; and \; G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \le \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \le \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \ge (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \ge \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \ge \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$
Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \qquad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le \frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \qquad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \qquad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le -(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le -(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}\ (v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}\ (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq v^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(mm) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

(586 A)

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SIXTY EIGHT

Cognitive Informatics: Mutantur Omnia Nos Et Mutamur In Illis :All Things Change, And We Change With Them


Abstract: Notwithstanding the imperative compatibilities, structural variabilities, inner differentiation, comparative dependability and structural morphology, and motivational value orientation institutionalization of pattern variables, Cognitive science is the interdisciplinary scientific study of the mind and its processes. It examines what cognition is, what it does and how it works. It includes research on intelligence and behavior, especially focusing on how information is represented, processed, and transformed (in faculties such as perception, language, memory, reasoning, and emotion) within nervous systems (human or other animal) and machines (e.g. computers). Cognitive science **consists of** multiple research disciplines, including psychology, artificial intelligence, philosophy, neuroscience, linguistics, and anthropology. It spans many levels of analysis, from low-level learning and decision mechanisms to high-level logic and planning; from neural circuitry to modular brain organization. The fundamental concept of cognitive science is "that thinking can best be understood in terms of representational structures in the mind and computational procedures that operate on those structures." A central tenet of cognitive science **is that a** complete understanding of the mind/brain cannot be attained by studying only a single level. An example would be the problem of remembering a phone number and recalling it later. One approach to understanding this process would be to study behavior through direct observation. A person could be presented with a phone number, asked to recall it after some delay. Then the accuracy of the response **could be measured**. Another approach would be to study the firings of individual neurons while a person is trying to remember the phone number. Neither of these experiments on its own would fully explain how the process of remembering a phone number works. Even if the technology to map out every neuron in the brain in real-time was available, and it was known when each neuron was firing, it would still be impossible to know how a particular firing of neurons translates into the observed behavior. Thus an understanding of how these two **levels relate to each other** is needed. **The Embodied Mind: Cognitive Science and Human Experience say**s "the new sciences of the mind need to enlarge their horizon to encompass both lived human experience and the possibilities for transformation inherent in human experience." This can be provided by a functional level account of the process. Studying a particular phenomenon from multiple levels **creates a** better understanding of the processes that occur in the brain to give rise to a particular behavior. Marr gave a famous description of three levels of analysis: the computational theory, specifying the goals of the computation; representation and algorithm, giving a representation of the input and output and the algorithm **which transforms** one into the other; and the hardware implementation, how algorithm and representation may be physically realized. The study of language processing in cognitive science is closely tied to the field of linguistics. Linguistics was traditionally studied as a part of the humanities, including studies of history, art and literature. In the last fifty years or so, more and more researchers have studied knowledge **and use of language** as a cognitive

phenomenon, the main problems being how knowledge of language can be acquired and used, and what precisely it consists of. Linguists have found that, while humans form sentences in ways apparently governed by very complex systems, they are remarkably unaware of the rules **that govern their** own speech. Thus linguists must resort to indirect methods to determine what those rules might be, if indeed rules as such exist. In any event, if speech is indeed governed by rules, they appear to be opaque to any conscious consideration. Mention may be made of the fact that in sociology emphasis is laid on the fact that the optimization of the facilities systems **leads to the** dynamical improvement of the instrumental efficacies, with necessary qualitative gradient and ascribed diffuse solidarity abstraction. Attention is drawn to the study of these factors with mathematical model. Kloowski's ideas are briefly stated in the introduction.

INTRODUCTION—VARIABLES USED

Introduction:

Information is the third essence in modeling the natural world with the internal differentiation, structural morphology and comparative dynamics, and phenomenological coextensive representations acting as the bridge between different cross section of the universe. Wang, Yingxu makes an attempt to explore an emerging discipline know as cognitive informatics. Cognitive Informatics **is a** profound interdisciplinary research area that tackles the common root problems of modern informatics, computation, software engineering, artificial intelligence (AI), neural psychology, and cognitive science. Cognitive informatics studies the internal information processing mechanisms **and** natural intelligence of the brain. This paper describes historical development of informatics from the classical information theory, contemporary informatics, to cognitive informatics. The domain of cognitive informatics and its interdisciplinary nature are explored. Foundations of cognitive informatics, particularly the brain vs. the mind, the acquired life functions vs. the inherited ones, and generic relationships between information, matter and energy are investigated. The potential engineering applications of cognitive informatics and perspectives on future research are discussed. It is expected that the investigation into cognitive informatics **will result in** fundamental findings towards the development of next generation IT and software technologies, such as neural computers, bioinformatics, quantum information processing, new software development approaches, and new architectures of information systems. **Cognitive Informatics, 2002. Proceedings. First IEEE International Conference on Date of Conference: 2002 INSPEC Accession Number: 7456030 Digital Object Identifier: 10.1109/COGINF.2002.103928On cognitive informatics Wang, Yingxu .**Quantum artificial neural network architectures and components Ajit Narayanan et al http://dx.doi.org/10.1016/S0020-0255 (00)00055-4 concentrate on the primary focus and locus on the homologues receptive of quantum architecture in the association of artificial cognitive memory and differentiated system of expressly oriented actions. This is against the backdrop of paradigm of relational context. It is shown by classical simulation and experimentation that quantum artificial neural networks (QUANNs) are more efficient and in some cases more powerful **than** classical artificial neural networks (CLANNs) for a variety of experimental tasks. This effect is particularly noticeable with larger and more complex domains. The gain in efficiency is achieved with no generalisation loss in most cases. QUANNs are also more powerful than CLANNs, again for some of the tasks examined, in terms of what the network can learn. What is more, it appears that not all components of QUANN architecture **need to be quantum** for these advantages to surface. It is demonstrated that a fully quantum neural network **has no advantage over** a partly quantum network and in **fact produce** worse results. Overall, this work **provides a first insight into** the expected behaviour of individual components of QUANNs, if and when quantum hardware is ever built, and raises questions about the interface between quantum and classical components of future QUANNs.



Alexandr A. Ezhov1 and Dan Ventura2 outline the research, development and perspectives of quantum neural networks – a burgeoning new field which integrates classical Neurocomputing with quantum computation. It is argued that the study of quantum neural networks may give us both new understanding of brain function as well as unprecedented possibilities in creating new systems for information processing, including solving classically intractable problems, associative memory with exponential capacity and possibly **overcoming the** limitations posed by the Church-Turing thesis. The Penrose-Hameroff orchestrated objective reduction (orch. OR) model **assigns a** cognitive role to quantum computations in microtubules within the neurons of the brain. Despite an apparently "warm, wet, and noisy" intracellular milieu, the proposal **suggests that** microtubules avoid environmental decoherence long enough to reach threshold for "self-collapse" (objective reduction) by a quantum gravity mechanism put forth by Penrose. The model has been criticized as regards the issue of environmental decoherence, and a recent report by Tegmark finds that microtubules can maintain quantum coherence for only 10-13 s, far too short to be neurophysiologically relevant. Critical examination of the common value patterns of manifested actions and the decoherence mechanisms likely to dominate in a biological setting **and find that** (1) Tegmark's commentary is not aimed at an existing model in the literature but rather at a hybrid that replaces the superposed protein conformations of the orch. OR theory with a soliton in superposition along the microtubule; (2) recalculation after correcting for differences between the model on which Tegmark bases his calculations and the orch. OR model (superposition separation, charge vs dipole, dielectric constant) **lengthens the** decoherence time to 10-5–10-4 s; (3) decoherence times on this order invalidate the assumptions of the derivation and determine the approximation regime considered by Tegmark to be **inappropriate to** the orch. OR superposition; (4) Tegmark's formulation **yields** decoherence times that increase with temperature contrary to well-established physical intuitions and the observed behavior of quantum coherent states; (5) incoherent metabolic energy supplied to the collective dynamics ordering water in the vicinity of microtubules at a rate exceeding that of decoherence **can counter** decoherence effects (in the same way that lasers avoid decoherence at room temperature); (6) microtubules are surrounded by a Debye layer of counterions, which can screen thermal fluctuations, and by an actin gel that **might enhance the** ordering of water in bundles of microtubules, further increasing the decoherence-free zone by an order of magnitude and, if the dependence on the distance between environmental ion and superposed state is accurately reflected in Tegmark's calculation, extending decoherence times by three orders of magnitude; (7) topological quantum computation in microtubules may be error **correcting, resistant** to decoherence; and (8) the decohering effect of radiative scatterers on microtubule quantum states **is negligible**. These considerations bring microtubule decoherence **into a regime in** which quantum gravity could **interact** with neurophysiology (. © 2002 The American Physical SocietyURL: http://link.aps.org/doi/10.1103/PhysRevE.65.061901DOI:10.1103/PhysRevE.65.061901 **Quantum computation in brain microtubules: Decoherence and biological feasibility S. Hagan1, S. R.**

Hameroff2, and J. A. Tuszyński3 emphasis mine) Supplementary to matter and energy, information is the third essence for modeling the natural world. An emerging discipline known as cognitive informatics (CI) is developed recently that forms a profound interdisciplinary study of cognitive and information sciences, and tackles the common root problems sharing by informatics, computing, software engineering, artificial intelligence, cognitive science, neuropsychology, philosophy, linguistics, and life science. CI focuses on internal information processing mechanisms and the natural intelligence of the brain. This paper describes the historical development of informatics from the classical information theory and contemporary informatics, to CI. The domain of CI and its interdisciplinary nature are explored. Foundations of CI, particularly the brain versus the mind, the acquired life functions versus the inherited ones, and generic relationships between information, matter, and energy are investigated. The potential engineering applications of CI and perspectives on future research are discussed. It is expected that the investigation into CI will result in fundamental findings towards the development of next generation IT and software technologies, and new architectures of computing systems. **Brain and Mind August 2003, Volume 4, Issue 2, pp 151-167 On Cognitive Informatics Yingxu Wang**

Following is excerpted from The Struggle for Your Mind: Conscious Evolution and the Battle to Control How We Think, available from Inner Traditions Kingsley L Dennis

The human body **is a** constant flux of thousands of chemical/biological interreactions and processes connecting molecules, cells, organs, and fluids, throughout the brain, body, and nervous system. Up until recently it was thought that all these interactions operated in a linear sequence, passing on information much like a runner passing the baton to the next runner. However, the latest findings in quantum biology and biophysics have discovered that there is in fact a tremendous degree of coherence within all living systems.

Extensive scientific investigation has found that a form of quantum coherence operates within living biological systems through what is known as biological excitations and biophoton emission. What this means is that metabolic energy is stored **as a form of** electromechanical and electromagnetic excitations. These coherent excitations are considered responsible for **generating and maintaining** long-range order via the transformation of energy and very weak electromagnetic signals. After nearly twenty years of experimental research, Fritz-Albert Popp put forward the hypothesis that biophotons are emitted from a coherent electrodynamic field within the living system.2

What this means is that each living cell is giving off, or resonating, a biophoton field of coherent energy. If each cell is emitting this field, then the whole living system is, in effect, a resonating field-a ubiquitous nonlocal field. And since biophotons **are the** entities through which the living system communicates, there is near-instantaneous intercommunication throughout. And this, claims Popp, is the basis for coherent biological organization **-- referred to as** quantum coherence. This discovery led Popp to state that the capacity for evolution rests not on aggressive struggle and rivalry but on the capacity for communication and cooperation. In this sense the built-in capacity for species evolution is not based on the individual but rather living systems that are interlinked within a coherent whole: Living systems are thus neither the subjects alone, nor objects isolated, but both subjects and objects in a mutually communicating universe of meaning. . . . Just as the cells in an organism take on different tasks for the whole, different populations enfold information not only for themselves, but for all other organisms, expanding the consciousness of the whole, while at the same time becoming more and more aware of this collective consciousness.3

Biophysicist Mae-Wan Ho describes how the living organism, including the human body, is coordinated throughout and is "coherent beyond our wildest dreams." It appears that every part of our body is "in communication with every other part through a dynamic, tuneable, responsive, liquid crystalline medium that pervades the whole body, from organs and tissues to the interior of every cell."4

What this tells us is that the medium of our bodies is a form of liquid crystal, an ideal transmitter of communication, resonance, and coherence. These relatively new developments in biophysics have discovered that all biological organisms are constituted of a liquid crystalline medium. Further, DNA is a liquid-crystal, lattice-type structure (which some refer to as a liquid crystal gel), whereby body cells are involved in a holographic instantaneous communication via the emitting of biophotons (a source based on light). This implies that all living biological organisms continuously emit radiations of light that form a field of coherence and communication. Moreover, biophysics has discovered that living organisms are permeated by quantum wave forms. Ho informs us that

. . . the visible body just happens to be where the wave function of the organism is most dense. Invisible quantum waves are spreading out from each of us and permeating into all other organisms. At the same time, each of us has the waves of every other organism entangled within our own make-up. . . . We are participants in the creation drama that is constantly unfolding. We are constantly co-creating and re-creating ourselves and other organisms in the universe. . . .5

This incredible new information actually positions each living being within a nonlocal quantum field consisting of wave interferences (where bodies meet). The liquid crystalline structure within living systems is also responsible for the direct current (DC) electrodynamic field that permeates the entire body of all animals. It has also been noted that the DC field has a mode of semiconduction that is much faster than the nervous system.6 If biological living systems are operating within a nonlocal interwoven field of resonating energy, then perhaps it is possible to see this manifesting in physical behavior?

Mae-Wan Ho describes how coherent excitations in living systems operate in much the same way as a boat race, where the oars (people) must row in step so as to create a phase transition. This indicates that there is an inherent tendency in Nature and in living systems to resonate in sync as a way of maintaining order and coherency. This type of behavior serves to reinforce the relationship between the individual and the collective that before had been thought random. This discovery is important in that it lends validity to the emerging paradigm of the global brain and of the growth of a planetary empathy-the third revolution (as discussed in chapter 6). Each person is thus not only in an empathic relationship with others but also entangled. This view has recently been corroborated by neuroscience with its finding of mirror neurons.

A mirror neuron is a brain neuron that is activated (fires) when a living being (such as a human, other primates, or mammal) observes the action of another. In other words, if an individual watches another person eat an apple, then the exact same brain neurons will fire in the person observing the action as if he were performing the act. Such neuron behavior has been found in humans to operate in the premotor and inferior parietal cortex. This phenomenon of mirror neurons was first discovered by a research team in Italy in the 1990s when studying the neuronal activity of macaques. This discovery has led to many notable neuroscientists to declare that mirror neurons are important for learning processes (imitation) as well as language acquisition. In more modern general terms we might also say that this capacity is what ties a person in sympathy and empathy to another's situation. It may also explain why people become so emotionally attached to events on television, and even cry in response to watching someone crying on the screen. In this way we are emotionally entangled through a mirroring of brain neuronal firing. When we also consider that our bodies are entangled through a quantum field of electrical bio-photon resonance, it explains how we are affected by and from others -- via wave/field interference. This information is significant when considering a shift toward heightened empathy between people both near and at a distance (via digital communications) as well as the potential for catalyzing future abilities for telepathic communication between individuals.

Neuroscience, quantum biology, and quantum physics are all beginning to converge to reveal that our bodies are not only biochemical systems but also a sophisticated resonating quantum system. This helps us to

understand how the body can be efficiently coherent, as well as explaining how we feel drawn to others, especially when we use terms such as good vibes, good energies, and we just seem to click. Our bodies, then, as well as our brains appear to function like receivers/decoders within a constantly in-flux information energy field. This explains how the human brain is able to store a lifetime of memories and experiences* as a wealth of data may well be stored within the informational field that encompasses the brain, and indeed the whole body. (Eminent mathematician John von Neumann calculated that during an average lifetime of seventy years we accumulate some 280 trillion bits of information.)

This new understanding of the quantum human informational field also gives credibility to the existence of extrasensory perceptions (ESP) and related abilities. Human consciousness is not only empathic, in a "wave-interference" relationship with other mind fields, but also is constantly transmitting and receiving information. If this is indeed an inherent aspect of human functionality then we can see why the power hierarchies maintained by a minority have been active in suppressing its operation. As children we are told/conditioned from very young age to dismiss our fantasies-to grow out of and grow up from such illusions and get with the "real world" (whose world?). Early educational and social-peer conditioning serves to wire our brain neurons into a particular set: a fixed pattern of receiving and interpreting the world. Thus we are literally hardwired into a specific reality paradigm and social operating system. Within this paradigm any thought of extrasensory perceptions are sneered at as childish nonsense (manufactured social peer pressure). Many of our early expressions of intuition are thus suppressed and stifled and replaced with "normal" thoughts and perceptions. Imaginative insights and visions are usually left to the eccentric artists, mystics, and fringe creative innovators. Much of our modern minds have been denied their left-right brain full function and pulled into a tight left-brain rational functioning that operates as mechanical, linear, competitive, and narrow.

The Modernity Project has fashioned a mind-set that is a highly focused and logical narrow-band receiver. This arrangement has been further strengthened by modern social institutions in order to suppress visionary and creative insights and our intuitive capacities. The abstract right brain, with its magical world of creative visionary thinking, has been sidelined. Much of this right-brain activity was the source for indigenous wisdom, shamanic practices, and similar traditions that modern materialism has mercilessly eliminated over the years. We have been conditioned to think of such "magical practices" as primitive, barbaric, and worthy of little more than Western colonialism and imperial rule.

The social institutions in our modern materialistic age act to influence us to reject anything extrasensory as a load of nonsense, wishful thinking, or New Age delusion. Thus with our left-hemisphere-dominated brain we live in the everyday world of matter: of material objects and external attractions. We are shown to exist as separate forces, as islands in a chaotic sea of physical and natural impacts, and at the whim of random neutral influences. Yet we now know that this is not the case.

To recap, quantum biology has shown that the body displays an incredible degree of quantum coherence, and that a quantum consciousness field exists throughout the human DNA and thus the human nervous system. Our biochemical structure is composed of a confluence of energies in complete entanglement and that operate as a nonlocal field within and outside the human body. Further DNA is a liquid-crystal, lattice-type structure that emits biophotons, which are light based. What this leads to is a new understanding that human DNA operates also as a quantum field. In other words, we can begin referring to DNA as quantum DNA. Therefore the 97 percent of human DNA that is not involved in protein building is active within a quantum state. It may well be that a future manifestation of quantum consciousness will come from part activation of the 97 percent quantum DNA that so far has baffled our scientists with its function. This quantum DNA activation may likely be related to the state of human consciousness and has remained dormant in response to human consciousness not being sufficiently prepared, or made ready, for its

manifestation. This field "life force" may be similar to the pervasive pranic energy that, as Gopi Krishna states, forms the impulse for evolutionary growth in the human nervous system:

. . . . An ever-present possibility, existing in all human beings by virtue of the evolutionary process still at work in the race, tending to create a condition of the brain and nervous system that can enable one to transcend the existing boundaries of the mind and acquire a state of consciousness far above that which is the normal heritage of mankind at present.7

This transcendental stage of consciousness that is depicted above as being a part of our natural evolutionary heritage is connected with the human brain and nervous system. We now know that we have a DNA quantum field activated within our bodies. Some biophysicists are already discussing whether quantum behavior may not be a common denominator for all living processes. As such a quantum informational field throughout the human body will determine the coherence of our light (biophoton) resonance as a vibratory rate. If human consciousness begins to shift its vibratory rate then there is every likelihood that DNA -- as a quantum field -- will likewise undergo a resonance shift, bringing into activation parts of its 97 percent hitherto "inactive" capacities. This may or may not be linked to the increase in electromagnetic frequencies now impacting our solar system from the galactic core. Is there a possibility that a phase step in the "engine of evolutionary energies" is under way?

The Russian biophysicist and molecular biologist Pjotr Garjajev, who has studied human DNA with his research team in Moscow, found that the 97 percent "inactive" DNA actually has complex properties. Garjajev discovered that the DNA, which is not used for protein synthesis, is instead actually used for communication -- more exactly for hyper communication. In their terms, hyper communication is a data exchange on a DNA level. Garjajev and his group analyzed the vibration response of the DNA and concluded that it can function much like networked intelligence, and that it allows for hyper communication of information among all sentient beings.

For example, the Moscow research group proved that damaged chromosomes (such as those harmed by x-rays) can be repaired. Their method was to capture the information patterns of particular DNA and then transmit these patterns, using focused light frequencies, onto another genome as a way of reprogramming the cells. In this way they successfully transformed frog embryos to salamander embryos simply by transmitting the DNA information patterns. Garjajev's research shows that certain frequency patterns can be "beamed" (such as with a laser) to transfer genetic information. This shows how DNA operates through resonance and vibratory frequencies. It also shows that human DNA can be modified -- or altered -- through the impact of external frequencies. This may also help to go some way toward validating the existence of such phenomena as remote acts of healing and other psychic attributes. It also suggests that DNA is a living, fluid, and dynamic "language" that as a quantum informational field is responsive not only to laser waves (as in the above experiment) but also EM waves and sound, given that the correct frequencies are applied.

The knowledge that human DNA can be influenced and modulated by frequencies (sound, light, language, and thought) is likely to have been known to various spiritual traditions, mystics, and teachers over the ages. This is perhaps why a variety of exercises have existed that utilize thought focus (prayer), sounds (music, chanting, singing), light (both natural light and produced light, such as in stained glass), and language (specific recitations such as a mantra and zikr). DNA appears to function not only as a protein builder (the minority function) but also as a medium for the storage, receiving, and communication of information.

Somewhat more controversially, Garjajev and his Russian colleagues also found examples where DNA could cause disturbing patterns in a vacuum, resulting in the production of what seemed to be magnetized wormholes. (For more information see the work of Grazyna Fosar and Franz Bludorf.) These wormholes appeared to function as connections outside our normal fields of time and space (which hints at

interdimensional communication). This phenomenon is indeed worthy of further analysis and experimentation. Yet it does seem probable that DNA is involved with various forms of hyper communication of which, at present, we know very little about. However, there are examples of hyper communication at work in Nature. For example, the organization of ant colonies appears to make use of this distributed form of communication. When a queen ant is separated from her colony, the worker ants continue to build and construct the colony as if following some form of blueprint. Yet if the queen ant is killed, then all work in the colony ceases, as if the blueprint had suddenly been taken offline. This suggests that the queen ant need not be in physical contact to continue to transmit the blueprint, yet upon her death the group consciousness ceases to operate within a hypercommunicative informational field. We can thus refer to these forms of hyper communication as quantum-field consciousness, or simply as quantum consciousness (since quantum implies non-local field effect).

At-a-distance human phenomena such as remote healing, remote sensing, and telepathy may work along comparable lines. On a more basic level we could say that many of us experience this as the sense of intuition and moments of inspiration. We may even be receiving these forms of hyper communication when we are asleep. There are countless examples of people, artists, and designers who gained inspiration for their work in their dreams. One example is the Italian composer Giuseppe Tartini who one night dreamt that a devil sat beside his bed playing the violin. The next morning Tartini wrote down the piece from memory and called it the Devil's Trill sonata. Not only do these experiences seem to be increasing (or perhaps people are more open to speaking of them?), but also newer generations of children are manifesting a higher level of clairvoyance and other extrasensory capacities. In recent times they have been referred to as indigo children, or the "new children." These developments may indicate that a higher form of group consciousness is emerging within humanity and that these abilities are now finding greater expression. This does not, however, deny the presence of negative influences against collective quantum consciousness since stress, fear, and similar impacts (see Armageddon meme) all serve to disrupt awareness and manifestation of quantum states. In this context we would do well to return to those practices recommended for centuries by spiritual traditions and teachers, that is, mediation, reflection, watchfulness, and mindfulness. Einstein was famous as a daydreamer throughout his life, and he often claimed that his greatest inspiration came to him when in such states. Enhanced connectivity between humanity may thus be served by each of us paying more attention to our inner states and striving for harmony and balance in our lives.

Scientists map process by which brain cells form long-term memories June 9, 2013 in Neuroscience

(1) Cheu and his team then adapted a mathematical model that describes the activity of a single neuron to **incorporate (e)** specific characteristics of cells in the hippocampus, including their inhibitory activity.
(2) This mathematical model that describes the activity of a single neuron to **incorporate (e)** specific characteristics of cells in the hippocampus, including their inhibitory activity **allowed (eb) them** to model neural networks in the hippocampus that encode short-term memories.
(3) They showed that for successful formation of auto-associative memories, the strength of synapses needs to be within a certain range: if synapses become too strong, the associated neurons **are (eb) activated** at the wrong time and networks become muddled, destroying the memories.
(4) If they are not strong enough, however, activation of some neurons in the network **is not (e) enough** to activate the rest, and memory retrieval fails.
(5) As well as providing insight into how memories may be stored and retrieved in the brain, Cheu thinks this work also has practical applications. "This study has **significant (eb) implications** in the construction of artificial cognitive computers in the future," he says.
(6) "It helps with developing artificial cognitive memory, in which memory sequences **can (e) be retrieved** by the presentation of a partial query." According to Cheu, one can compare it to a single

image being used to retrieve a sequence of images from a video clip. More information: Cheu, E. Y., Yu, J., Tan, C. H. & Tang, H. Synaptic conditions for auto-associative memory storage and pattern completion in Jensen et al.'s model of hippocampal area CA3. Journal of Computational Neuroscience advance online publication, 30 May 2012 (doi: 10.1007/s10827-012-0394-8). http://link.springer.com/article/10.1007%2Fs10827-012-0394-8 .Provided by Agency for Science, Technology and Research (A*STAR), Singapore Read more at: http://medicalxpress.com/news/2012-11-brain-memories-artificial-cognitive.html#jCp

(7) Transcranial magnetic stimulation (TMS) **is (=) a** noninvasive method to cause depolarization or Hyperpolarization in the neurons of the brain. TMS uses electromagnetic induction to induce weak electric currents using a rapidly changing magnetic field; http://en.wikipedia.org/wiki/Transcr...ic_stimulation

(8) You can induce action potential trains artificially **using** TMS. As far as natural communication between cells, we usually model these as electrochemical interactions. So, although magnetism always goes along with the territory when speaking of these actions, we typically do not invoke it a causal agent in natural brain function.

NOTATION

Module One

Nirguna Brahman and Saguna Brahman. Nirguna Brahman the characterless one, or for that matter spacelessness and timelessness should not mistaken for something beyond the subterranean realm and ceratoid dualism of space time. Infact it is the constitution, consummation and composition of all the characterstics, spaces and times (negative time is extant and existentialist that is what is understood), all combine to make it the Universal constant Zero. Like a general ledger in a bank is representative of the contrast between the transfer scrolls and the transfer and cash slips that flown in finally leading to the proposition Assets =liabilities. Remarks hereof in done in the context of gratification deprivation complex which is considered by the author as something that is individually non conservative but holistically nonconservative. This module is of utmost importance which is hoped for further development both by more erudite researchers and the author. In essence and sum and substance what is said is the dualities of debit and credit, of space and time, of character and propertilessness, energy mass all cease in this cauldron cram of brick and brac the melting pot of nature's general ledger. There are three different models in this module, due to spatial constraints. Freewill and destiny is also dual at one stage and at the General ledger stage closes down to zero. What then is the role of compunction, contrition and confession? It helps increase gratification (total memory at all three stages including anagrammatic acting mind leading to dreams, phantasmagoria, and even pathology; ASCII characters and repetitive record in the brain. It is true that after 20 years of experience "mind would be somewhere" and "mechanical operation' takes place. At this stage, write up still would be there in the information but the influence is minimal for human beings do their work "mechanically")

G_{13} : Category one of Saguna Brahman; space; time

G_{14} : Category two of Saguna Brahman; space; time

G_{15} : Category three of Saguna Brahman; space; time

T_{13} : Category one of Nirguna Brahman ;spacelessness; timelessness

T_{14} : Category two of Nirguna Brahman ; spacelessness; timelessness

T_{15} : Category three of Nirguna Brahman ; spacelessness; timelessness

Module Two

Cheu and his team then adapted a mathematical model that describes the activity of a single neuron to **incorporate (e)** specific characteristics of cells in the hippocampus, including their inhibitory activity.

G_{16} : Category one of specific characteristics of cells in the hippocampus, including their inhibitory activity

G_{17} : Category two of specific characteristics of cells in the hippocampus, including their inhibitory activity

G_{18} : Category three of specific characteristics of cells in the hippocampus, including their inhibitory activity

T_{16} : Category one of activity of a single neuron

T_{17} : Category two of activity of a single neuron

T_{18} : Category three of activity of a single neuron

Module three

This mathematical model that describes the activity of a single neuron to **incorporate (e)** specific characteristics of cells in the hippocampus, including their inhibitory activity **allowed (eb) them** to model neural networks in the hippocampus that encode short-term memories.

G_{20} : Category one of activity of a single neuron to **incorporate (e)** specific characteristics of cells in the hippocampus, including their inhibitory activity

G_{21} : Category two of activity of a single neuron to **incorporate (e)** specific characteristics of cells in the hippocampus, including their inhibitory activity

G_{22} : Category three of activity of a single neuron to **incorporate (e)** specific characteristics of cells in the hippocampus, including their inhibitory activity

T_{20} : Category one of neural networks in the hippocampus that encode short-term memories models

T_{21} : Category two of neural networks in the hippocampus that encode short-term memories models

T_{22} : Category three of neural networks in the hippocampus that encode short-term memories models

Module four

They showed that for successful formation of auto-associative memories, the strength of synapses needs to be within a certain range: if synapses become too strong, the associated neurons **are (eb) activated** at the wrong time and networks become muddled, destroying the memories.

G_{24} : Category one of successful formation of auto-associative memories, the strength of synapses needs to be within a certain range: if synapses become too strong, the associated neurons

G_{25} : Category two of successful formation of auto-associative memories, the strength of synapses needs to be within a certain range: if synapses become too strong, the associated neurons

G_{26} : Category three of successful formation of auto-associative memories, the strength of synapses needs to be within a certain range: if synapses become too strong, the associated neurons

T_{24} : Category one of wrong time and networks become muddled, destroying the memories

T_{25} : Category two of wrong time and networks become muddled, destroying the memories

T_{26} : Category three of wrong time and networks become muddled, destroying the memories

Module five

As well as providing insight into how memories may be stored and retrieved in the brain, Cheu thinks this work also has practical applications. "This study has **significant (eb) implications** in the construction of artificial cognitive computers in the future," he says.

G_{28} : Category one of successful formation of auto-associative memories, the strength of synapses needs to be within a certain range: if synapses become too strong, the associated neurons **are (eb) activated** at the wrong time and networks become muddled, destroying the memories

G_{29} : Category two of successful formation of auto-associative memories, the strength of synapses needs to be within a certain range: if synapses become too strong, the associated neurons **are (eb) activated** at the wrong time and networks become muddled, destroying the memories

G_{30} : Category three of successful formation of auto-associative memories, the strength of synapses needs to be within a certain range: if synapses become too strong, the associated neurons **are (eb) activated** at the wrong time and networks become muddled, destroying the memories

T_{28} : Category one of construction of artificial cognitive computers

T_{29} : Category two of construction of artificial cognitive computers

T_{30} : Category three of construction of artificial cognitive computers

Module six

"It helps with developing artificial cognitive memory, in which memory sequences **can (e) be retrieved** by the presentation of a partial query." According to Cheu, one can compare it to a single image being used to retrieve a sequence of images from a video clip. More information: Cheu, E. Y., Yu, J., Tan, C. H. & Tang, H. Synaptic conditions for auto-associative memory storage and pattern completion in Jensen et al.'s model of hippocampal area

G_{32} : Category one of **retrieved** by the presentation of a partial query which are the conditions for auto-associative memory storage and pattern completion in Jensen et al.'s model of hippocampal area

G_{33} : Category two of **retrieved** by the presentation of a partial query which are conditions for auto-associative memory storage and pattern completion in Jensen et al.'s model of hippocampal area

G_{34} : Category three of **retrieved** by the presentation of a partial query which are conditions for auto-associative memory storage and pattern completion in Jensen et al.'s model of hippocampal area

T_{32} : Category one of artificial cognitive memory, in which memory sequences

T_{33} : Category two of artificial cognitive memory, in which memory sequences

T_{34} : Category three of artificial cognitive memory, in which memory sequences

Module seven

Transcranial magnetic stimulation (TMS) **is a** noninvasive method to cause depolarization or Hyperpolarization in the neurons of the brain. TMS uses electromagnetic induction to induce weak electric currents using a rapidly changing magnetic field

G_{36} : Category one of noninvasive method to cause depolarization or Hyperpolarization in the neurons of the brain. TMS uses electromagnetic induction to induce weak electric currents using a rapidly changing magnetic field

G_{37} : Category two of noninvasive method to cause depolarization or Hyperpolarization in the neurons of the brain. TMS uses electromagnetic induction to induce weak electric currents using a rapidly changing magnetic field

G_{38} : Category three of noninvasive method to cause depolarization or Hyperpolarization in the neurons of the brain. TMS uses electromagnetic induction to induce weak electric currents using a rapidly changing magnetic field

T_{36} : Category one of Transcranial magnetic stimulation (TMS)

T_{37} : Category two of Transcranial magnetic stimulation (TMS)

T_{38} : Category three of Transcranial magnetic stimulation (TMS)

Module eight

You can induce action potential trains artificially **using** TMS. As far as natural communication between cells, we usually model these as electrochemical interactions. So, although magnetism always goes along with the territory when speaking of these actions, we typically do not invoke it a causal agent in natural brain function.

G_{40} : Category one of TMS. Minds are different and classification is based on the pen chance, proclivities, predilections and propensities of the brain, rajas, tamas, and sattva

G_{41} : Category two of TMS

G_{42} : Category three of TMS

T_{40} : Category one of induction of action potential trains artificially

T_{41} : Category two of induction of action potential trains artificially

T_{42} : Category three of induction of action potential trains artificially

Module Nine

G_{44} : Category one of electrochemical interactions

G_{45} : Category two of electrochemical interactions

G_{46} : Category three of electrochemical interactions

T_{44} : Category one of natural communication between cells

T_{45} : Category two of natural communication between cells

T_{46} : Category three of natural communication between cells

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}$ $(a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$ $(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$ $(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$

$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)\right]T_{18}$ 12

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}((G_{19}), t) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20}$ 13

$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21}$ 14

$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22}$ 15

$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20}$ 16

$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21}$ 17

$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22}$ 18

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24}$ 19

$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25}$ 20

$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26}$ 21

$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)\right]T_{24}$ 22

$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)\right]T_{25}$ 23

$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)\right]T_{26}$ 24

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}), t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28}$ 25

$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29}$ 26

$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30}$ 27

$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)\right]T_{28}$ 28

$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)\right]T_{29}$ 29

$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)\right]T_{30}$ 30

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}), t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}), t\right)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}), t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}), t\right)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}), t\right)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}), t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45}, t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}), t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad (55)$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad (56)$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad (57)$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $-(b_{13}'')^{(1)}(G,t)$, $-(b_{14}'')^{(1)}(G,t)$, $-(b_{15}'')^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b_{16}'')^{(2,2,)}(G_{19},t)$, $-(b_{17}'')^{(2,2,)}(G_{19},t)$, $-(b_{18}'')^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{37}'')^{(7,7,)}(G_{39},t)$, $-(b_{36}'')^{(7,7,)}(G_{39},t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8)}(G_{43},t)$ $-(b_{41}'')^{(8,8)}(G_{43},t)$ $-(b_{42}'')^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{ccc} (a_{16}')^{(2)} & +(a_{16}'')^{(2)}(T_{17},t) & +(a_{13}'')^{(1,1,)}(T_{14},t) \quad +(a_{20}'')^{(3,3,3)}(T_{21},t) \\ +(a_{24}'')^{(4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{36}'')^{(7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{ccc} (a_{17}')^{(2)} & +(a_{17}'')^{(2)}(T_{17},t) & +(a_{14}'')^{(1,1,)}(T_{14},t) \quad +(a_{21}'')^{(3,3,3)}(T_{21},t) \\ +(a_{25}'')^{(4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{37}'')^{(7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a_{18}')^{(2)} & +(a_{18}'')^{(2)}(T_{17},t) & +(a_{15}'')^{(1,1,)}(T_{14},t) \quad +(a_{22}'')^{(3,3,3)}(T_{21},t) \\ +(a_{26}'')^{(4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{38}'')^{(7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad (63)$$

Where $+(a_{16}'')^{(2)}(T_{17},t)$, $+(a_{17}'')^{(2)}(T_{17},t)$, $+(a_{18}'')^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)}\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b_{20}'')^{(3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{16}'')^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{13}'')^{(1,1,1,)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,)}(G,t)}$, $\boxed{-(b_{15}'')^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{36}'')^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b_{38}'')^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a_{24}')^{(4)} & \boxed{+(a_{24}'')^{(4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,)}(T_{29},t)} \quad \boxed{+(a_{32}'')^{(6,6,)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a_{16}'')^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{36}'')^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9)}(T_{45},t)} \end{array}\right] G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a_{25}')^{(4)} & \boxed{+(a_{25}'')^{(4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,)}(T_{29},t)} \quad \boxed{+(a_{33}'')^{(6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a_{17}'')^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{37}'')^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9)}(T_{45},t)} \end{array}\right] G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a_{26}')^{(4)} & \boxed{+(a_{26}'')^{(4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,)}(T_{29},t)} \quad \boxed{+(a_{34}'')^{(6,6,)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a_{18}'')^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{38}'')^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9)}(T_{45},t)} \end{array}\right] G_{26} \quad (75)$$

$\boxed{(a_{24}'')^{(4)}(T_{25},t)}$, $\boxed{(a_{25}'')^{(4)}(T_{25},t)}$, $\boxed{(a_{26}'')^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a_{28}'')^{(5,5,)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a_{32}'')^{(6,6,)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a_{13}'')^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{14}'')^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a_{16}'')^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a_{20}'')^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3)}(T_{21},t)}$

are sixth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$

are seventh augmentation coefficients for category 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt}=(b_{26})^{(4)}T_{25}-\begin{bmatrix}(b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

are fourth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$

are fifth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$

are sixth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$

are seventh detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$

are eighth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt}=(a_{28})^{(5)}G_{29}-\begin{bmatrix}(a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad 81$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8\ ,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad 84$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ *are second detrition coefficients*

$for\ category$ 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}$ $are\ \ third\ \ detrition\ coefficients$ $for\ category$ 1,2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{34} \quad (87)$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6)}(T_{33},t)}$ $are\ \ first\ \ augmentation\ coefficients$ $for\ category$ 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ $are\ second\ augmentation$ $coefficients\ for\ category$ 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ $are\ third\ augmentation$ $coefficients\ for\ category$ 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$
seventh augmentation coefficients
$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$
Eighth augmentation coefficients
$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt}=(b_{32})^{(6)}T_{33}-\begin{bmatrix}(b'_{32})^{(6)}\,-(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t)\\ -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)\\ -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)\end{bmatrix}T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt}=(b_{33})^{(6)}T_{32}-\begin{bmatrix}(b'_{33})^{(6)}\,-(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t)\\ -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)\\ -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)\end{bmatrix}T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt}=(b_{34})^{(6)}T_{33}-\begin{bmatrix}(b'_{34})^{(6)}\,-(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t)\\ -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)\\ -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)\end{bmatrix}T_{34} \quad (90)$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3
$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3
$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3
$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3
$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2, and 3
$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$ *95*

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\boxed{-(b_{44}'')^{(9)}(G_{47},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\boxed{-(b_{45}'')^{(9)}(G_{47},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\boxed{-(b_{46}'')^{(9)}(G_{47},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \le (r_i)^{(1)} \le (b_i')^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \le (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} +\ (\hat{A}_{20})^{(3)} +\ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} +\ (\hat{B}_{20})^{(3)} +\ (\hat{Q}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le\ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) =\ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\ ,\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ \ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(AAAAAAAAAA) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(BBBBBBBBBB) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le \ (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(CCCCCCCCCC) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(DDDDDDDDDD)

$\lim_{G \to \infty} (b_i'')^{(7)}\big((G_{39}), t\big) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - \ T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\big((G_{39}), t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(EEEEEEEEEE) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(FFFFFFFFFF) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$
satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\leq\ \left(\hat{P}_{13}\right)^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\leq\ (\hat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\leq\ (\hat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\leq\ (\hat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(nn) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$

In the same way , one can obtain

$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18}. indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16}, G_{18} and G_{16}, G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18}. The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2:

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

208

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)}$$

209

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)}$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),\left((G_{23})^{(2)},(T_{23})^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

214

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$

In the same way , one can obtain

$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\,((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\hat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\hat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)}G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)}G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)}T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, ***if*** $(u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(\nu_1)^{(6)}, (\nu_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, \nu^{(6)}, u^{(6)}$: 350

By $(\nu_1)^{(6)} > 0$, $(\nu_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)},, (\bar{\nu}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, $\boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)},, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)}+(C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21}, t)\right) - (a_{21}'')^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $v^{(4)}$:- $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

For $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)}=\frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)}(T_{45}, t)\right) - (a_{45}'')^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(nn) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523

A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial (b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SIXTY NINE

Mahavairocana (Dainichi-Nyorai 大日如来) Mandala: Constituents of Dependent Arising

Abstract: It is to be noted at the outset that the appellation is only Buddhist and has nothing to do with the contents. Integrative function, motivational adequacy, latency of collective orientations of Lie Algebras, interactional role specifications of production group functions and specificity –diffusivity component of the Theory is what all expatiated and delineated here towards the end of constitution and consummation of the model."An Exceptionally Simple Theory of Everything" is a physics preprint proposing a basis for a unified field theory, very often referred to as "E8 Theory," which attempts **to describe all** known fundamental interactions in physics and to stand as a possible theory of everything. The paper was posted to the physics arXiv by Antony Garrett Lisi on November 6, 2007, and was not submitted to a peer-reviewed scientific journal The title is a pun on the algebra used, the Lie algebra of the largest "simple," "exceptional" Lie group, E8. The paper's goal is to describe how the combined structure and dynamics of all gravitational and Standard Model particle fields, including fermions, are part of the E8 Lie algebra. In the paper, Lisi states that all three generations of fermions do not directly embed in E8 with correct quantum numbers and spins, but that they might be described via atriality transformation, noting that the theory is incomplete and that a correct description of the relationship between triality and generations, if it exists, awaits a better understanding. In a follow-up paper, Lee Smolin proposes a **spontaneous symmetry breaking** mechanism **for obtaining the classical action** in Lisi's model, and speculates on the path to its quantization. In July 2009 Jacques Distler and Skip Garibaldi published a critical paper in Communications in Mathematical Physics called "There is no 'Theory of Everything' inside E8," arguing that Lisi's theory, and a large class of related models, cannot work. They offer a direct proof that it is impossible to embed all three generations of fermions in E8, or to obtain even the one-generation Standard Model without the presence of an antigeneration. In response to Distler and Garibaldi's paper, Lisi argues, in a news paper "An Explicit Embedding of Gravity and the Standard Model in E8," peer reviewed and published in a conference proceedings, that some assumptions about fermion embeddings are unnecessary and that the antigeneration is not per se a problem sufficient to rule out the one-generation Standard Model. In December 2010 and May 2011 Lisi wrote in the popular magazine Scientific American a feature article on the E8 Theory of Everything and an entry in the blog section of the magazine addressing some of the criticism of his theory

and how it has progressed, noting that the theory is still incomplete and makes only tenuous predictions, with the three generation issue remaining as a significant problem. N. E. Bonesteel is one author who has studied the internal differentiated evaluative interactional orientations as components of institutional integration and role differentiation in quantum fluids (Chiral spin liquids and quantum error-correcting codes N. E. Bonesteel) and the concomitant quantum correcting codes. Possibility of **using the** twofold topological degeneracy of spin-1/2 chiral spin liquid states on the torus to construct quantum error-correcting codes is investigated. It is shown that codes constructed using these states on finite periodic lattices **do not** meet the necessary and sufficient conditions for correcting even a single qubit error with perfect fidelity. However, for large enough lattice sizes, these conditions are approximately satisfied, **and the resulting** codes may therefore be viewed as approximate quantum error-correcting codes. © 2000 The American Physical SocietyURL: http://link.aps.org/doi/10.1103/PhysRevA.62.062310DOI: 10.1103/PhysRevA.62.062310

(Source: Wikipedia, Buddhist Sutras)

INTRODUCTION—VARIABLES USED

Nature 460, 1106-1109 (27 August 2009) | doi: 10.1038/nature08308; Received 13 May 2009; Accepted 21 July 2009; Published online 9 August 2009 Pedram Roushan et al

(1) Topological insulators are a new class of insulators in which a bulk gap for electronic excitations is generated **because (e) of** the strong spin–orbit coupling1, 2, 3, 4, 5 inherent to these systems.
(2) These materials are distinguished from ordinary insulators **by (eb) the presence** of gapless metallic surface states, resembling chiral edge modes in quantum Hall systems, but with unconventional spin textures.
(3) A key predicted feature of such spin-textured boundary states **is (=) their** insensitivity to spin-independent scattering, which is thought to protect them from backscattering and localization.
(4) Recently, experimental and theoretical efforts have **provided (eb) s**trong evidence for the existence of both two- and three-dimensional classes of such topological insulator materials in semiconductor quantum well structures and several bismuth-based compounds but so far experiments have not probed the sensitivity of these chiral states to scattering.
(5) Here authors use scanning tunnelling spectroscopy and angle-resolved photoemission spectroscopy to visualize the gapless surface states in the three-dimensional topological insulator Bi1-xSbx, and examine in detail the influence of scattering from disorder **caused(e) by** random alloying in this compound.
(6) Authors show that, despite strong atomic scale disorder, backscattering between states of opposite momentum and opposite spin **is** absent.
(7) Observations demonstrate that the chiral nature of these states **protects (e)** the spin of the carriers.
(8) These chiral states are therefore potentially **useful for** spin-based electronics, in which long spin coherence is critical14, and also for quantum computing applications, where topological protection **can(eb) enable** fault-tolerant information processing.

POSTED BY ROB BRYANTON AT 4:09 A.M LABELS: E8, MANY WORLDS, AND PHASE SPACE


(1) As we continue to add dimensions, the orthogonal hypercube projections that we're looking at here with each new dimension **become (=)** increasingly ornate, and (dare I say it) more mandala-like.
(2) Another eight-dimensional shape that we've looked at with this project **is** Garrett Lisi's E8 Rotation. Lisi created a huge stir in the physics world with a new proposal for what he calls An Exceptionally Simple Theory of Everything **that uses** "E8" - a complex, eight-dimensional pattern with 248

points.

(3) As visually appealing as this shape (pictured below) might be, its implications are startling: Lisi has demonstrated that there is a way to place the various forces and elementary particles (including their possible quantum spin values) **on (e: space)** E8's 248 points.

(4) Rotating Lisi's model in various ways reveals the explanation for a variety of interactions, some of which (like the clustering of quarks into families of three) **are (=)** natural outcomes from this structure.

(5) Some points in his model are currently **occupied (eb) by** particles which have been theorized but which have yet to be seen, and there is hope that the Large Hadron Collider in Switzerland may some day reveal some of those particles.



(6) Whether Garrett Lisi's theory proves the existence of higher dimensions or not is open to interpretation - Lisi himself says this geometric pattern, although **it is based(E) upon** an 8 dimensional construct, could be fully realized within our 4D space-time without requiring additional dimensions.

(7) On the other hand, in a New Scientist magazine article, string theorist Sabine Hossenfelder (of the Perimeter Institute for Theoretical Physics) points out that this could be **complimentary(EB)** to string theory, which she says also uses E8 to describe the Calabi-Yau manifold, the extra-dimensional shape that string theorists says our universe **is derived from**.

(8) Universe is **derived from** Calabi Yau Manifold

Lisi also acknowledges that this is not a finished theory but a work in progress, and that there are still some deficiencies in his model that need further refinement. For me, this dimensional connection is fascinating because with this project I've insisted that you can't have any physical expressions of matter in anything beyond the 8th dimension.

Why Do We Need the 8th Dimension?

(9) A commonly asked question about this project is, "if our universe and every other universe **can be (=) thought of** as "points", or perhaps "positions" within a multiverse landscape, and in this way of visualizing the dimensions we get to that landscape by the time we're at the seventh dimension, then why do we need to think about any more dimensions beyond that?

NOTATION

Module One

as we continue to add dimensions, the orthogonal hypercube projections that we're looking at here with each new dimension **become** increasingly ornate, and (dare I say it) more mandala-like

G_{13} : Category one of increasingly ornate, and (dare I say it) more mandala-like

G_{14} : Category two of increasingly ornate, and (dare I say it) more mandala-like

G_{15} : Category three of increasingly ornate, and (dare I say it) more mandala-like

T_{13} : Category one of as we continue to add dimensions, the orthogonal hypercube projections that we're looking at here with each new dimension

T_{14} : Category two of as we continue to add dimensions, the orthogonal hypercube projections that we're looking at here with each new dimension

T_{15} : Category three of as we continue to add dimensions, the orthogonal hypercube projections that we're looking at here with each new dimension

Module Two

Another eight-dimensional shape that we've looked at with this project **is** Garrett Lisi's E8 Rotation. Lisi created a huge stir in the physics world with a new proposal for what he calls An Exceptionally Simple Theory of Everything **that uses** "E8" - a complex, eight-dimensional pattern with 248 points.

G_{16} : Category one of "E8" - a complex, eight-dimensional pattern with 248 points.

G_{17} : Category two of "E8" - a complex, eight-dimensional pattern with 248 points.

G_{18} : Category three of "E8" - a complex, eight-dimensional pattern with 248 points.

T_{16} : Category one of Simple Theory of Everything. I repeat that this does not mean that there are three theories. What is meant is that investigating systems could be applied with the theory for study. Like Newton's law can be applied to various systems and one can think of "total gravity"

T_{17} : Category two of Simple Theory of Everything

T_{18} : Category three of Simple Theory of Everything

Module three

As visually appealing as this shape (pictured below) might be, its implications are startling: Lisi has demonstrated that there is a way to place the various forces and elementary particles (including their possible quantum spin values) **on (e: space)** E8's 248 points.

G_{20} : Category one of various forces and elementary particles (including their possible quantum spin values)

G_{21} : Category two of various forces and elementary particles (including their possible quantum spin values)

G_{22} : Category three of various forces and elementary particles (including their possible quantum spin values)

T_{20} : Category one of E8's 248 points

T_{21} : Category two of E8's 248 points

T_{22} : Category three of E8's 248 points

Module four

Rotating Lisi's model in various ways reveals the explanation for a variety of interactions, some of which (like the clustering of quarks into families of three) **are** natural outcomes from this structure

G_{24} : Category one of natural outcomes from this structure

G_{25} : Category two of natural outcomes from this structure

G_{26} : Category three of natural outcomes from this structure

T_{24} : Category one of Rotating Lisi's model in various ways reveals the explanation for a variety of interactions, some of which (like the clustering of quarks into families of three)

T_{25} : Category two of Rotating Lisi's model in various ways reveals the explanation for a variety of interactions, some of which (like the clustering of quarks into families of three)

T_{26} : Category three of Rotating Lisi's model in various ways reveals the explanation for a variety of interactions, some of which (like the clustering of quarks into families of three)

Module five

Some points in his model are currently **occupied (eb) by** particles which have been theorized but which have yet to be seen, and there is hope that the Large Hadron Collider in Switzerland may some day reveal some of those particles.

G_{28} : Category one of Some points in his model are currently

G_{29} : Category two of Some points in his model are currently

G_{30} : Category three of Some points in his model are currently

T_{28} : Category one of particles which have been theorized but which has yet to be seen, and there is hope that the Large Hadron Collider in Switzerland may some day reveal some of those particles.

T_{29} : Category two of particles which have been theorized but which have yet to be seen, and there is hope that the Large Hadron Collider in Switzerland may some day reveal some of those particles.

T_{30} : Category three of particles which have been theorized but which have yet to be seen, and there is hope that the Large Hadron Collider in Switzerland may some day reveal some of those particles.

Module six

Whether Garrett Lisi's theory proves the existence of higher dimensions or not is open to interpretation - Lisi himself says this geometric pattern, although **it is based(E) upon** an 8 dimensional construct, could be fully realized within our 4D space-time without requiring additional dimensions.

G_{32} : Category one of 8 dimensional construct, could be fully realized within our 4D space-time without requiring additional dimensions

G_{33} : Category two of 8 dimensional construct, could be fully realized within our 4D space-time without requiring additional dimensions

G_{34} : Category three of 8 dimensional construct, could be fully realized within our 4D space-time without requiring additional dimensions

T_{32} : Category one of geometric pattern

T_{33} : Category two of geometric pattern

T_{34} : Category three of geometric pattern

Module seven

On the other hand, in a New Scientist magazine article, string theorist Sabine Hossenfelder (of the Perimeter Institute for Theoretical Physics) points out that this could be **complimentary(EB)** to string theory, which she says also uses E8 to describe the Calabi-Yau manifold, the extra-dimensional shape that string theorists says our universe **is derived from**

G_{36} : Category one of Rotating Lisi's model in various ways reveals the explanation for a variety of interactions, some of which (like the clustering of quarks into families of three) **are** natural outcomes from this structure

G_{37} : Category two of Rotating Lisi's model in various ways reveals the explanation for a variety of interactions, some of which (like the clustering of quarks into families of three) **are** natural outcomes from this structure

G_{38} : Category three of Rotating Lisi's model in various ways reveals the explanation for a variety of interactions, some of which (like the clustering of quarks into families of three) **are** natural outcomes from this structure

T_{36} : Category one of string theory, which she says also uses E8 to describe the Calabi-Yau manifold, the extra-dimensional shape that string theorists says our universe **is derived from**

T_{37} : Category two of string theory, which she says also uses E8 to describe the Calabi-Yau manifold, the extra-dimensional shape that string theorists says our universe **is derived from**

T_{38} : Category three of string theory, which she says also uses E8 to describe the Calabi-Yau manifold, the extra-dimensional shape that string theorists says our universe **is derived from**

Module eight

Universe is **derived from** Calabi Yau Manifold

G_{40} : Category one of Calabi Yau Manifold. There are millions of them. Model predicts, projects, prognosticates the nature of correspondence of the universe vis a vis the Calabi Yau manifold

G_{41} : Category two of Calabi Yau Manifold

G_{42} : Category three of Calabi Yau Manifold

T_{40} : Category one of Universe (that part of the universe which is under investigation. Note the inflationary universe makes it impossible to perceive the entire gamut of universe, and interiors of black holes still lay beyond the horizons of human perception)

T_{41} : Category two of Universe (that part of the universe which is under investigation. Note the inflationary universe makes it impossible to perceive the entire gamut of universe, and interiors of black holes still lay beyond the horizons of human perception)

T_{42} : Category three of Universe (that part of the universe which is under investigation. Note the inflationary universe makes it impossible to perceive the entire gamut of universe, and interiors of black holes still lay beyond the horizons of human perception)

Module Nine

A commonly asked question about this project is, "if our universe and every other universe **can be thought of** as "points", or perhaps "positions" within a multiverse landscape, and in this way of visualizing the dimensions we get to that landscape by the time we're at the seventh dimension, then why do we need to think about any more dimensions beyond that?

G_{44} : Category one of "points", or perhaps "positions" within a multiverse landscape, and in this way of visualizing the dimensions we get to that landscape by the time we're at the seventh dimension, then why do we need to think about any more dimensions beyond that?

G_{45} : Category two of "points", or perhaps "positions" within a multiverse landscape, and in this way of visualizing the dimensions we get to that landscape by the time we're at the seventh dimension, then why do we need to think about any more dimensions beyond that?

G_{46} : Category three of "points", or perhaps "positions" within a multiverse landscape, and in this way of visualizing the dimensions we get to that landscape by the time we're at the seventh dimension, then why do we need to think about any more dimensions beyond that?

T_{44} : Category one of universe and every other universe

T_{45} : Category two of universe and every other universe

T_{46} : Category three of universe and every other universe

.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$

$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation

coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\left[-(b''_{13})^{(1)}(G,t)\right] & \left[-(b''_{16})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\left[-(b''_{14})^{(1)}(G,t)\right] & \left[-(b''_{17})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\left[-(b''_{15})^{(1)}(G,t)\right] & \left[-(b''_{18})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad 73$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad 74$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad 76$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)} & \boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category $1, 2$ *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8\ ,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b_{36}')^{(7)}\boxed{-(b_{36}'')^{(7)}(G_{39},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b_{37}')^{(7)}\boxed{-(b_{37}'')^{(7)}(G_{39},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b_{38}')^{(7)}\boxed{-(b_{38}'')^{(7)}(G_{39},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)}\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)}\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$

$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)}$ 98

$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$

$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$

$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)}$ 102

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(GGGGGGGGGG) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(HHHHHHHHHH) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$

132

(IIIIIIIII) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(JJJJJJJJJ)

$\lim_{G\to\infty}(b_i'')^{(7)}((G_{39}),t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}((G_{39}),t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(KKKKKKKKKK) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(LLLLLLLLLL) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad 146\ A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad (169)$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that (170)

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that (172)

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that (174)

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(oo) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right| \; e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$$
$$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$$

$$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$$

In the same way , one can obtain

$$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ and\ (\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$

$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$

$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$

$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq$ 248

$\frac{1}{(\widehat{M}_{32})^{(6)}} \left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(6)}$ and $(b''_i)^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 250

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(6)} - (a''_i)^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b'_i)^{(6)} t\right)} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(7)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)}$$ 267

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)}$$ 268

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}),(\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}),(\widetilde{T_{43}})\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

406

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case,

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

407

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(oo) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)} T_{45}^{*}}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)} T_{45}^{*}}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad (525)$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad (526)$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad (527)$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \quad (528)$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \quad (529)$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \quad (530)$$

ASYMPTOTIC STABILITY ANALYSIS (531)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \quad (532)$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad (533)$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad (534)$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad (535)$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad (536)$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \quad (537)$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\Big)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\Big)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SEVENTY

Vajrasattva (Kongō-Satta 金剛薩埵) Mandala: Adamantine Being

Abstract: Lisi's model **is a** variant and extension of a Grand Unification Theory (a "GUT," describing electromagnetism, the weak interaction and the strong interaction) to include gravitation, aHiggs boson and fermions in an attempt to describe all fields of the Standard Model and gravity as different parts of one field over four dimensional spacetime. More specifically, Lisi **combines t**he left-right symmetric Pati-Salam GUT **with a** MacDowell-Mansouri description of gravity, **using the** spin connection and gravitational frame combined with a Higgs boson, necessitating a cosmological constant. The model is formulated as a gauge theory, using a modified BF action, with E8 as the Lie group. Mathematically, this is an E8 **principal bundle**, with connection, over a four dimensional base manifold. Lisi's **embedding of the** Standard Model gauge group in E8 **leads him to predict the existence** of 22 new bosonic particles at an undetermined mass scale. Lisi states that the fermions enter **via an** unconventional use of the BRST technique, as Grassmann number fields valued in part of the E8 Lie algebra. The bosons are combined with these fermions as one-form and Grassmann number parts of a kind of superconnection, each valued in separate parts of the E8 Lie algebra. The curvature of this superconnection is calculated, **producing the** Riemann curvature, gauge field curvature, gravitational torsion, covariant derivative of the Higgs, and the covariant Dirac derivative of the fermions. This curvature is used **to build the** modified BF action by hand, in an attempt to match the dynamics of the Standard Model and gravity. In the paper, Lisi describes several deficiencies in this model. The most important deficiency is noted as an incorrect, or "poorly understood," inclusion of the second and third generations of fermions in E8, **relying on** triality. This deficiency, and the incomplete nature of the model, **precludes the** prediction of masses for new or existing particles. Also, Lisi notes the use of explicit symmetry breaking in building his action, rather than offering a more desirable spontaneous symmetry breaking mechanism. And, **no attempt is made to provide** a quantum description of the theory—this being left for future work. About it, Lee Smolin proposed a spontaneous symmetry breaking mechanism for obtaining the action in Lisi's model, and speculates on the path to its quantization as a spin foam (Source; Wikipedia, Stanford Encyclopedia, emphasis mine)

INTRODUCTION—VARIABLES USED

Chirality ,Lisi's Model and Quantum Computation:

Local conserved charges in principal chiral models in 1+1 dimensions and there is a classically conserved local charge for each totally symmetric invariant tensor of the underlying group. These local charges **are shown** to be in involution **with the** non-local Yangian charges. For each Poisson bracket algebra of the local charges, an infinite set of local charges with spins **equal to the** exponents modulo the Coxeter number is constructed, and **it is shown that** these **commute with** one another. Comments on the evidence for, and implications of, survival of these charges in the quantum theory follow in the paper **(J.M. Evansa et al Local conserved charges in principal chiral models)**

POSTED BY ROB BRYANTON AT 4:09 A.M LABELS: E8, MANY WORLDS, AND PHASE SPACE

(1) To my way of thinking, you can express the same concern about every definition of every single dimension, and the arguments have to keep coming back to "what are we missing within this current dimension?" For instance, people say that the third spatial dimension is really all you need, **because (e)** no matter what other universe you imagine it should have a third dimensional expression.
(2) Or there are people who say all we need is space-time**, because(e)** by the time you have the third

dimension plus time, which gives the third dimension a way to change from state to state, then you can imagine every single possible universe strung end to end within infinite space-time,


(3) **Because (e) that** string of possible states is infinite you will eventually get to every possible universe.

(4) Or you can have people who say space-time plus its probabilistic outcomes (which Everett's Many Worlds Interpretation says are "orthogonal" or "at right angles" to space-time, **leading to(eb) my conclusion** that these branches are in the fifth dimension) gives you everything you need to imagine every possible universe.

(5) From there, I've proposed that the branches for our universe which no amount of chance or choice will allow us to select (like the branches where dinosaurs never became extinct or the ones where I died in a car accident last year) are in the sixth dimension, **and this(e) uses** the same logical reasoning that defines any spatial dimension - there always needs to be a "new degree of freedom" added for us to be able to call what we're talking about a new dimension, and the new dimension allows us to get to something that was unavailable from the previous dimensions

(6) Likewise, I've proposed that those other universes with different basic physical laws are located in other positions within the seventh dimension and above, and this is why our universe, **constrained(e) at its** position within that multiverse landscape never "wanders off" into one of those other universes.

(The rotating shapes below are known as the Platonic Solids. Pictured here are the tetrahedron, the cube or hexahedron, the octahedron, the dodecahedron, and the icosahedron. We'll talk about these shapes later on in this entry.)



(7) So, why isn't the seventh dimension as far as we need to go? We have to continue with the logic we've used from the outset. Imagine a point of infinite size in the sixth dimension, **and it's (=) much the same as** imagining the indeterminate point we started from with our thought experiment - the point encompasses the whole dimension.

(8) If we can think of a different point that is not subsumed by the first point, then the point-line-plane postulate **tells us we've found a (e) way to get to** the "next dimension up": those two points define a line in the seventh dimension.

(9) What if the second point we just imagined **represents (eb) a** universe where the strength of gravity is different from ours?

NOTATION

Module One

To my way of thinking, you can express the same concern about every definition of every single dimension, and the arguments have to keep coming back to "what are we missing within this current dimension?" For instance, people say that the third spatial dimension is really all you need, **because (e)** no matter what other universe you imagine it should have a third dimensional expression.

G_{13} : Category one of **no** matter what other universe you imagine it should have a third dimensional expression.

G_{14} : Category two of no matter what other universe you imagine it should have a third dimensional expression.

G_{15} : Category three of no matter what other universe you imagine it should have a third dimensional expression.

T_{13} : Category one of third spatial dimension is really all you need

T_{14} : Category two of third spatial dimension is really all you need

T_{15} : Category three of third spatial dimension is really all you need

Module Two

Or there are people who say all we need is space-time**, because(e)** by the time you have the third dimension plus time, which gives the third dimension a way to change from state to state, then you can imagine every single possible universe strung end to end within infinite space-time

G_{16} : Category one of by the time you have the third dimension plus time, which gives the third dimension a way to change from state to state, then you can imagine every single possible universe strung end to end within infinite space-time

G_{17} : Category two of by the time you have the third dimension plus time, which gives the third dimension a way to change from state to state, then you can imagine every single possible universe strung end to end within infinite space-time

G_{18} : Category three of by the time you have the third dimension plus time, which gives the third dimension a way to change from state to state, then you can imagine every single possible universe strung end to end within infinite space-time

T_{16} : Category one of all we need is space-time

T_{17} : Category two of all we need is space-time

T_{18} : Category three of all we need is space-time

Module three

Because (e) that string of possible states is infinite you will eventually get to every possible universe.

G_{20} : Category one of eventually get to every possible universe

G_{21} : Category two of eventually get to every possible universe

G_{22} : Category three of eventually get to every possible universe

T_{20} : Category one of string of possible states is infinite

T_{21} : Category two of string of possible states is infinite

T_{22} : Category three of string of possible states is infinite

Module four

Or you can have people who say space-time plus its probabilistic outcomes (which Everett's Many Worlds Interpretation says are "orthogonal" or "at right angles" to space-time, **leading to (eb) my conclusion** that

these branches are in the fifth dimension) gives you everything you need to imagine every possible universe.

G_{24} : Category one of space-time plus its probabilistic outcomes (which Everett's Many Worlds Interpretation says are "orthogonal" or "at right angles" to space-time

G_{25} : Category two of space-time plus its probabilistic outcomes (which Everett's Many Worlds Interpretation says are "orthogonal" or "at right angles" to space-time

G_{26} : Category three of space-time plus its probabilistic outcomes (which Everett's Many Worlds Interpretation says are "orthogonal" or "at right angles" to space-time

T_{24} : Category one of these branches are in the fifth dimension) gives you everything you need to imagine every possible universe

T_{25} : Category two of these branches are in the fifth dimension) gives you everything you need to imagine every possible universe

T_{26} : Category three of these branches are in the fifth dimension) gives you everything you need to imagine every possible universe

Module five

From there, I've proposed that the branches for our universe which no amount of chance or choice will allow us to select (like the branches where dinosaurs never became extinct or the ones where I died in a car accident last year) are in the sixth dimension, **and this(e) uses** the same logical reasoning that defines any spatial dimension - there always needs to be a "new degree of freedom" added for us to be able to call what we're talking about a new dimension, and the new dimension allows us to get to something that was unavailable from the previous dimensions

G_{28} : Category one of same logical reasoning that defines any spatial dimension - there always needs to be a "new degree of freedom" added for us to be able to call what we're talking about a new dimension, and the new dimension allows us to get to something that was unavailable from the previous dimensions

G_{29} : Category two of same logical reasoning that defines any spatial dimension - there always needs to be a "new degree of freedom" added for us to be able to call what we're talking about a new dimension, and the new dimension allows us to get to something that was unavailable from the previous dimensions

G_{30} : Category three of same logical reasoning that defines any spatial dimension - there always needs to be a "new degree of freedom" added for us to be able to call what we're talking about a new dimension, and the new dimension allows us to get to something that was unavailable from the previous dimensions

T_{28} : Category one of branches for our universe which no amount of chance or choice will allow us to select (like the branches where dinosaurs never became extinct or the ones where I died in a car accident last year) are in the sixth dimension

T_{29} : Category two of branches for our universe which no amount of chance or choice will allow us to select (like the branches where dinosaurs never became extinct or the ones where I died in a car accident last year) are in the sixth dimension

T_{30} : Category three of branches for our universe which no amount of chance or choice will allow us to select (like the branches where dinosaurs never became extinct or the ones where I died in a car accident last

year) are in the sixth dimension

Module six

Likewise, I've proposed that those other universes with different basic physical laws are located in other positions within the seventh dimension and above, and this is why our universe, **constrained(e) at its** position within that multiverse landscape never "wanders off" into one of those other universes.

G_{32} : Category one of position within that multiverse landscape never "wanders off" into one of those other universes

G_{33} : Category two of position within that multiverse landscape never "wanders off" into one of those other universes

G_{34} : Category three of position within that multiverse landscape never "wanders off" into one of those other universes

T_{32} : Category one of those other universes with different basic physical laws are located in other positions within the seventh dimension and above, and this is why our universe

T_{33} : Category two of those other universes with different basic physical laws are located in other positions within the seventh dimension and above, and this is why our universe

T_{34} : Category three of those other universes with different basic physical laws are located in other positions within the seventh dimension and above, and this is why our universe

Module seven

So, why isn't the seventh dimension as far as we need to go? We have to continue with the logic we've used from the outset. Imagine a point of infinite size in the sixth dimension, **and it's much the same as** imagining the indeterminate point we started from with our thought experiment - the point encompasses the whole dimension.

G_{36} : Category one of imagining the indeterminate point we started from with our thought experiment - the point encompasses the whole dimension

G_{37} : Category two of imagining the indeterminate point we started from with our thought experiment - the point encompasses the whole dimension

G_{38} : Category three of imagining the indeterminate point we started from with our thought experiment - the point encompasses the whole dimension

T_{36} : Category one of point of infinite size in the sixth dimension

T_{37} : Category two of point of infinite size in the sixth dimension

T_{38} : Category three of point of infinite size in the sixth dimension

Module eight

If we can think of a different point that is not subsumed by the first point, then the point-line-plane postulate **tells us we've found a (eb) way to get to** the "next dimension up": those two points define a line in the

seventh dimension.

G_{40} : Category one of different points that is not subsumed by the first point, then the point-line-plane postulate

G_{41} : Category two of different points that is not subsumed by the first point, then the point-line-plane postulate

G_{42} : Category three of different points that is not subsumed by the first point, then the point-line-plane postulate

T_{40} : Category one of **get to** the "next dimension up": those two points define a line in the seventh dimension.

T_{41} : Category two of **get to** the "next dimension up": those two points define a line in the seventh dimension.

T_{42} : Category three of **get to** the "next dimension up": those two points define a line in the seventh dimension.

Module Nine

What if the second point we just imagined **represents (eb) a** universe where the strength of gravity is different from ours? We give a model to answer this question under the axiomatic predication being in the right. Note that we have different investigating systems that satisfy Lisi's Theory and the characterstics thereof is the basis, heart, root, mainstay, principle, fundamental, premise, starting point, principal element substructure, underpinning, groundwork, bedrock, justification, grounds, basis, warrant, reason of the classification scheme. **Note the model predicts the "second point' of various systems under investigation and corresponding universe where the strength of gravity is different from ours**

G_{44} : Category one of second point we just imagined

G_{45} : Category two of second point we just imagined

G_{46} : Category three of second point we just imagined

T_{44} : Category one of universe where the strength of gravity is different from ours

T_{45} : Category two of universe where the strength of gravity is different from ours

T_{46} : Category three of universe where the strength of gravity is different from ours

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}),t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) \quad +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)}\ \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{array}\right] G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{array}\right] G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{array}\right] G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for*

category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category $1, 2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a_{32}')^{(6)} & +(a_{32}'')^{(6)}(T_{33},t) & +(a_{28}'')^{(5,5,5)}(T_{29},t) & +(a_{24}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a_{33}')^{(6)} & +(a_{33}'')^{(6)}(T_{33},t) & +(a_{29}'')^{(5,5,5)}(T_{29},t) & +(a_{25}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a_{34}')^{(6)} & +(a_{34}'')^{(6)}(T_{33},t) & +(a_{30}'')^{(5,5,5)}(T_{29},t) & +(a_{26}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a_{32}'')^{(6)}(T_{33},t)$, $+(a_{33}'')^{(6)}(T_{33},t)$, $+(a_{34}'')^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a_{28}'')^{(5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a_{24}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{ccc} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(MMMMMMMMMM) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(NNNNNNNNNN) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(OOOOOOOOOO) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(PPPPPPPPPP) $\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(QQQQQQQQQ) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(RRRRRRRRRR) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \le (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and

$(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)},(\widehat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+(\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(b_i)^{(8)}+(b_i')^{(8)}+(\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146\,A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t} \ , \ \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t} \ , \ \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} \ , \ G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} \ , \ T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} \ , \ G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} \ , \ T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^{0} + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^{0} + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^{0} + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^{0} + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(pp) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18}. indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16}, G_{18} and G_{16}, G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18}. The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2:

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}\ and\ (\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$

In the same way , one can obtain

$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25}\ \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25}\ \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}\ , G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\ , \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}\left((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)}\right)}\left[e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}\left((S_1)^{(3)} - (a'_{22})^{(3)}\right)}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}\left((R_1)^{(3)} - (b'_{22})^{(3)}\right)}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}\left((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)}\right)}\left[e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right) - (b''_{25})^{(4)}\left((G_{27}), t\right) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t} \right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)} t} \right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t} \right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(\nu_1)^{(6)}, (\nu_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, \nu^{(6)}, u^{(6)}$: 350

By $(\nu_1)^{(6)} > 0$, $(\nu_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)},, (\bar{\nu}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad (354)$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad (355)$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad (356)$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)}+(C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)}(T_{45}, t)\right) - (a_{45}'')^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(pp) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523

A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\,(G_{47})^*\,) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SEVENTY ONE

Nagarjuna (Ryūju-Bosatsu 龍樹菩薩) Mandala: Proof of Convention

Abstract: Levels of magnification in the theory are: Macroscopic level – Matter 2. Molecular level 3. Atomic level - Protons, neutrons, and electrons 4. Subatomic level – Electron 5. Subatomic level – Quarks 6. Lie group geometrical representation level. In modern particle physics, the most common approach to describe elementary particles and their interactions is usually through a gauge theory based on a Lie group. A Lie group **is a** mathematical structure with much complex symmetry, which **can be described as** an object with a complex geometry. In the corresponding quantum field theory, there is a particle **associated with** each of these symmetries, and these particles can interact with each other according to the geometry of the group and how the particles **are related to** the group representation. In Lisi's model, the Lie group used is E8, a group with 248 parameters. In general, in each gauge theory **based on a** Yang-Mills action, the symmetries of the Lie group are associated to a specific kind of particles known as gauge bosons (like photons, W and Z bosons, and gluons in the Standard Model; in models involving supersymmetry this is a little more complicated). These gauge bosons **can interact** with each other and with fermions according to the geometry of the group and its fundamental representations. One of Lisi's challenges is that **his theory identifies** fermions with the symmetries that are usually associated only with gauge bosons. Generally this is considered not possible and this aspect of the theory still needs to be completed. Another aspect different from the common approaches is that Lisi's theory **includes also** gravity in the Lie Group E8. While this aspect has been proven to be possible in supersymmetric theories and impossible in a large class of theories (Coleman-Mandula theorem), it is sometimes attempted in other theories and models that do not strictly belong to those classes. **It is still not clear if this feature is achievable in Lisi's theory.** In general, a unified theory has a Lie group large enough to contain the Standard Model symmetries. There are many such theories, some of which have used E8 for a long time (like string theory). In Lisi's specific model, as introduced above, each of the 248 symmetries of E8 **corresponds to a** different elementary particle: Standard Model gauge bosons, gravitons and Standard Model fermions, which can all interact (as usual in this kind of theories) according to the geometry of the group, in this case E8. Lisi states that: "E8 **reproduce [es]** all known fields and dynamics through pure geometry." The complicated geometry of Lie groups, E8 amongst them, is described graphically **using** group representation theory. Using this mathematical description, each symmetry of a group—and so each kind of elementary particle—can be associated with a point in a weight diagram. The coordinates of these points are the quantum numbers—the charges—of elementary particles, which are conserved in interactions. In order to form a theory of everything, Lisi's model must eventually **predict the** exact number

of fundamental particles, all of their properties, masses, forces between them, the nature of spacetime, and the cosmological constant. Much of this work is still in the conceptual stage—in particular, quantization and predictions of particle masses have not been done and the model at the moment cannot **reproduce all the** known particles. Lisi himself acknowledges it as a work-in-progress: "The theory is very young, and still in development," and the three generation "issue remains the most significant problem, and until it is solved the theory is not complete and cannot be considered much more than a speculative proposal. Without fully describing how the three generations of fermions work, **the theory and all predictions from it remain tenuous**." (Wikipedia: emphasis mine

INTRODUCTION—VARIABLES USED

Lisi's model and quantum computation:

Nature 452, 970-974 (24 April 2008) | doi: 10.1038/nature06843; Received 25 November 2007; Accepted 14 February 2008 A topological Dirac insulator in a quantum spin Hall phase D. Hsieh et al

(1) When electrons are subject to a large external magnetic field, the conventional charge quantum Hall effect1, 2 **dictates that** an electronic excitation gap **is generated in** the sample bulk, but metallic conduction is permitted at the boundary.
(2) Recent theoretical models **suggest that** certain bulk insulators with large spin–orbit interactions may also naturally support conducting topological boundary states in the quantum limit3, 4, 5, **which(eb) opens up the** possibility for studying unusual quantum Hall-like phenomena in zero external magnetic fields6.
(3) Bulk Bi1-xSbx single crystals are predicted **to be prime candidates7**, 8 for one such unusual Hall phase of matter known as the topological insulator9, 10, 11.
(4) The hallmark of a topological insulator **is the** existence of metallic surface states that are higher-dimensional analogues of the edge states that characterize a quantum spin Hall insulator3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13.
(5) In addition to its interesting boundary states, the bulk of Bi1-xSbx is predicted **to exhibit** three-dimensional Dirac particles14, 15, 16, 17, another topic of heightened current interest following the new findings in two-dimensional graphene18, 19, 20 and charge quantum Hall fractionalization observed in pure bismuth21.
(6) However, despite numerous transport and magnetic measurements on the Bi1-xSbx family since the 1960s17, no direct evidence of either topological Hall states or bulk Dirac particles has been found. Here, **using** incident-photon-energy-modulated angle-resolved photoemission spectroscopy (IPEM-ARPES), authors report the direct observation of massive
(7) Dirac particles in the bulk of Bi0.9Sb0.1 locate the Kramers points at the sample's boundary **and provide** a comprehensive mapping of the Dirac insulator's gapless surface electron bands.
(8) These findings taken together **suggest that the** observed surface state on the boundary of the bulk insulator is a realization of the 'topological metal'9, 10, 11.
(9) They **also suggest that** this material has potential application in developing next-generation quantum computing devices that may incorporate 'light-like' bulk carriers and spin-textured surface currents


POSTED BY ROB BRYANTON AT 4:09 A.M LABELS: E8, MANY WORLDS, AND PHASE SPACE

(1) Then the line that passes through those two points **is (=) like** a one-dimensional line, and that line extends through all possible values for gravity.
(2) Different positions on that line would **include (e)** "impossible" universes which could not have come into existence **because (e) the** strength of gravity was above or below a certain threshold

needed **to allow a** physical universe to express itself, and such imaginary universes might be at many places along the line, interspersed with universes that were able to cohere into more organized states.

(3) What about a universe with a different value for the speed of light (or whatever physical constant you care to imagine)? That's not on the line we just drew. We could erase the other point **we drew,(e&eb,=)** then place a new point for a universe with a different speed of light, and the new line that passes through those two points would be all possible realities resulting from different values for the speed of light.

(4) But what if I want to consider both lines simultaneously? It can't be done without entering the next dimension up, the eighth dimension. So with the logic of the point-line-plane postulate we have a way of thinking of the eighth dimension as being like a plane and our universe **can be(=) viewed as** a point in the eighth dimension with those two lines or any other lines representing other different-initial-conditions universe passing through our point within that plane.



(5) Remember this: with the point-line-plane postulate, the point you start from is in dimension x, the line to a different point is in dimension x+1, and the plane defined by a third point not on that line **is (eb) in** dimension x+2.

(6) Having worked through three dimensions, nothing moves or changes about the first point, we are just adding dimensions **that (eb) allow us** to view that starting point from different perspectives.

(7) Thinking of our universe as a "largest possible point" encompassing the phase space of the sixth dimension, then, doesn't mean that we can't view that unchanging point from the widening perspective of the seventh or eighth dimension. Quite the contrary: this logic is the accepted approach to visualizing spatial dimensions. Likewise, the idealized shapes we've started each of these entries with (like the octeract at the start of this entry) have an underlying symmetry, where each of the points are equidistant to their adjacent ones **in the (=) same way** that the adjacent points of a 3D cube are equidistant when viewed within the third dimension.

(8) This means that the 8D octeract, like any of the other hypercube shapes, can be placed **within (eb) a** hypersphere with the same number of dimensions, and the outer points of the hypercube are pushed apart from the center of the hypersphere symmetrically:

(9) The point of indeterminate size **can (=) become** a sphere occupying as little or as much of the dimension it's placed within.

NOTATION

Module One

Then the line that passes through those two points **is like** a one-dimensional line, and that line extends through all possible values for gravity.

G_{13} : Category one of one-dimensional line and that line extends through all possible values for gravity.

G_{14} : Category two of one-dimensional line, and that line extends through all possible values for gravity.

G_{15} : Category three of one-dimensional line, and that line extends through all possible values for gravity.

T_{13} : Category one of line that passes through those two points

T_{14} : Category two of line that passes through those two points

T_{15} : Category three of line that passes through those two points

Module Two

Different positions on that line would **include (e)** "impossible" universes which could not have come into existence **because (e) the** strength of gravity was above or below a certain threshold needed **to allow a** physical universe to express itself, and such imaginary universes might be at many places along the line, interspersed with universes that were able to cohere into more organized states.

G_{16} : Category one of Different positions on that line would **include (e)** "impossible" universes which could not have come into existence **because (e) the** strength of gravity was above or below a certain threshold needed

G_{17} : Category two of Different positions on that line would **include (e)** "impossible" universes which could not have come into existence **because (e) the** strength of gravity was above or below a certain threshold needed

G_{18} : Category three of Different positions on that line would **include (e)** "impossible" universes which could not have come into existence **because (e) the** strength of gravity was above or below a certain threshold needed

T_{16} : Category one of physical universe to express itself, and such imaginary universes might be at many places along the line, interspersed with universes that were able to cohere into more organized states.

T_{17} : Category two of physical universe to express itself, and such imaginary universes might be at many places along the line, interspersed with universes that were able to cohere into more organized states.

T_{18} : Category three of physical universe to express itself, and such imaginary universes might be at many places along the line, interspersed with universes that were able to cohere into more organized states.

Module three(Both the cases are discussed. Care while reading the model)

What about a universe with a different value for the speed of light (or whatever physical constant you care to imagine)? That's not on the line we just drew. We could erase the other point **we drew,** then place a new point for a universe with a different speed of light, and the new line that passes through those two points would be all possible realities resulting from different values for the speed of light.

G_{20} : Category one of universe with a different value for the speed of light (or whatever physical constant you care to imagine) not on the line drawn, whence forth we could erase the point we drew; place a new point for a universe with a different speed of light, and the new line that passes through those two points would be all possible realities resulting from different values for the speed of light.

G_{21} : Category two of universe with a different value for the speed of light (or whatever physical constant you care to imagine) not on the line drawn, whence forth we could erase the point we drew; place a new point for a universe with a different speed of light, and the new line that passes through those two points would be all possible realities resulting from different values for the speed of light.

G_{22} : Category three of universe with a different value for the speed of light (or whatever physical constant you care to imagine) not on the line drawn, whence forth we could erase the point we drew; place a new point for a universe with a different speed of light, and the new line that passes through those two points would be all possible realities resulting from different values for the speed of light.

T_{20} : Category one of place a new point for a universe with a different speed of light, and the new line that passes through those two points would be all possible realities resulting from different values for the speed of light; universe with a different value for the speed of light (or whatever physical constant you care to imagine) not on the line drawn ,whence forth we could erase the point we drew

T_{21} : Category two of place a new point for a universe with a different speed of light, and the new line that passes through those two points would be all possible realities resulting from different values for the speed of light.; universe with a different value for the speed of light (or whatever physical constant you care to imagine) not on the line drawn ,whence forth we could erase the point we drew

T_{22} : Category three of place a new point for a universe with a different speed of light, and the new line that passes through those two points would be all possible realities resulting from different values for the speed of light; universe with a different value for the speed of light (or whatever physical constant you care to imagine) not on the line drawn ,whence forth we could erase the point we drew

Module four

But what if I want to consider both lines simultaneously? It can't be done without entering the next dimension up, the eighth dimension. So with the logic of the point-line-plane postulate we have a way of thinking of the eighth dimension as being like a plane and our universe **can be(=) viewed as** a point in the eighth dimension with those two lines or any other lines representing other different-initial-conditions universe passing through our point within that plane.

G_{24} : Category one of point in the eighth dimension with those two lines or any other lines representing other different-initial-conditions universe passing through our point within that plane.

G_{25} : Category two of point in the eighth dimension with those two lines or any other lines representing other different-initial-conditions universe passing through our point within that plane.

G_{26} : Category three of point in the eighth dimension with those two lines or any other lines representing other different-initial-conditions universe passing through our point within that plane.

T_{24} : Category one of consideration of both the lines simultaneously without entering the eighth dimension with the ramification of the logic of point line plane postulate we can think of eight dimension as being like a plane and our universe

T_{25} : Category two of consideration of both the lines simultaneously without entering the eighth dimension with the ramification of the logic of point line plane postulate we can think of eight dimension as being like a plane and our universe

T_{26} : Category three of consideration of both the lines simultaneously without entering the eighth dimension with the ramification of the logic of point line plane postulate we can think of eight dimension as being like a plane and our universe

Module five

Remember this: with the point-line-plane postulate, the point you start from is in dimension x, the line to a different point is in dimension x+1, and the plane defined by a third point not on that line **is (eb) in** dimension x+2.

G_{28} : Category one of point-line-plane postulate ramificatory, the point you start from is in dimension x, the line to a different point is in dimension x+1, and the plane defined by a third point not on that line

G_{29} : Category two of point-line-plane postulate ramificatory, the point you start from is in dimension x, the line to a different point is in dimension x+1, and the plane defined by a third point not on that line

G_{30} : Category three of point-line-plane postulate ramificatory, the point you start from is in dimension x, the line to a different point is in dimension x+1, and the plane defined by a third point not on that line

T_{28} : Category one of dimension x+2.

T_{29} : Category two of dimension x+2.

T_{30} : Category three of dimension x+2.

Module six

Having worked through three dimensions, nothing moves or changes about the first point, we are just adding dimensions **that (eb) allow us** to view that starting point from different perspectives.

G_{32} : Category one of nothing moves or changes about the first point having worked through three dimensions, there is addition of dimensions

G_{33} : Category two of nothing moves or changes about the first point having worked through three dimensions, there is addition of dimensions

G_{34} : Category three of nothing moves or changes about the first point having worked through three dimensions, there is addition of dimensions

T_{32} : Category one of view that starting point from different perspectives

T_{33} : Category two of view that starting point from different perspectives

T_{34} : Category three of view that starting point from different perspectives

Module seven

Thinking of our universe as a "largest possible point" encompassing the phase space of the sixth dimension, then, doesn't mean that we can't view that unchanging point from the widening perspective of the seventh or eighth dimension. Quite the contrary: this logic is the accepted approach to visualizing spatial dimensions. Likewise, the idealized shapes we've started each of these entries with (like the octeract at the start of this entry) have an underlying symmetry, where each of the points are equidistant to their adjacent ones **in the (=) same way** that the adjacent points of a 3D cube are equidistant when viewed within the third dimension.

G_{36} : Category one of adjacent points of a 3D cube are equidistant when viewed within the third dimension

G_{37} : Category two of adjacent points of a 3D cube are equidistant when viewed within the third dimension

G_{38} : Category three of adjacent points of a 3D cube are equidistant when viewed within the third dimension

T_{36} : Category one of idealized shapes we've started each of these entries with (like the octeract at the start of this entry) have an underlying symmetry, where each of the points are equidistant to their adjacent ones

T_{37} : Category two of idealized shapes we've started each of these entries with (like the octeract at the start of this entry) have an underlying symmetry, where each of the points are equidistant to their adjacent ones

T_{38} : Category three of idealized shapes we've started each of these entries with (like the octeract at the start of this entry) have an underlying symmetry, where each of the points are equidistant to their adjacent ones

Module eight

This means that the 8D octeract, like any of the other hypercube shapes, can be placed **within (eb) a** hypersphere with the same number of dimensions, and the outer points of the hypercube are pushed apart from the center of the hypersphere symmetrically

G_{40} : Category one of8D octeract, like any of the other hypercube shapes, can be placed

G_{41} : Category two of 8D octeract, like any of the other hypercube shapes, can be placed

G_{42} : Category three of 8D octeract, like any of the other hypercube shapes, can be placed

T_{40} : Category one of hypersphere with the same number of dimensions, and the outer points of the hypercube are pushed apart from the center of the hypersphere symmetrically

T_{41} : Category two of hypersphere with the same number of dimensions, and the outer points of the hypercube are pushed apart from the center of the hypersphere symmetrically

T_{42} : Category three of hypersphere with the same number of dimensions, and the outer points of the hypercube are pushed apart from the center of the hypersphere symmetrically

Module Nine

The point of indeterminate size **can become** a sphere occupying as little or as much of the dimension it's placed within.

G_{44} : Category one of sphere occupying as little or as much of the dimension it's placed within.

G_{45} : Category two of sphere occupying as little or as much of the dimension it's placed within.

G_{46} : Category three of sphere occupying as little or as much of the dimension it's placed within.

T_{44} : Category one of point of indeterminate size

T_{45} : Category two of point of indeterminate size

T_{46} : Category three of point of indeterminate size

SECTION SEVENTY TWO

Nagabodhi (Ryūchi-Bosatsu 龍智菩薩) Mandala: Hymn to the Absolute Reality

Abstract: By matching 226 known standard model particles to some of the 248 symmetries of E8, the theory is able **to predict the existence** and quantum numbers of 22 new particles.There of these are the same new $\mathrm{su}(2)_R$ and $\mathrm{u}(1)_{B-L}$ gauge bosons as predicted in the Pati-Salam model, the W' and Z' bosons. Another is a new $\mathrm{u}(1)$ gauge boson, with a corresponding new quantum number. And the remaining 18 new bosons predicted are new colored fields, interacting with the strong force. Lisi states that some of these 22 particles might be seen at the Large Hadron Collider. Since Lisi does not specify masses for these particles their prediction is not falsifiable by non-discovery in any given experiment**, because the** masses could exceed the experiment's reach. However, the discovery of new particles that do not fit in Lisi's classification, such as superpartners, would fall outside the model, and falsify Lisi's match to E8. Also, **because the** theory at the moment fails to predict all the known particles and the matching of the three fermion generations is tentative and problematic in the model, Lisi places a low confidence in these predictions. Three previous arXiv preprints by Lisi deal with mathematical physics related to the theory. "Clifford Geometrodynamics," in 2002, **endeavors to describe** fermions geometrically as BRST ghosts. "Clifford bundle formulation of BF gravity generalized to the standard model," in 2005**, describes** the algebra of gravitational and Standard Model fields acting on a generation of fermions, but does not mention E8. "Quantum mechanics from a universal action reservoir," in 2006, attempts to derive quantum mechanics using information theory. Before writing his 2007 paper, Lisi discussed his work on a Foundational Questions Institute (FQXi) forum, at an FQXi conference, and for an FQXi article. Lisi gave his first talk on E8 Theory at the Loops '07 conference in Morelia, Mexico soon followed by a talk at the Perimeter Institute. John Baez commented on Lisi's work in "This Week's Finds in Mathematical Physics (Week 253)", and Lisi was interviewed on Sabine Hossenfelder's "Backreaction" blog. Lisi's arXiv preprint, "An Exceptionally Simple Theory of Everything", appeared on November 6, 2007, and immediately attracted a great deal of attention. Lisi made a further presentation for the International Loop Quantum Gravity Seminar on November 13, 2007, and responded to press inquiries on an FQXi forum He presented his work at the TED Conference on February 28, 2008. Lisi's paper spawned a variety of reactions and debates across various physics blogs and online discussion groups. The first to comment was Sabine Hossenfelder, summarizing the paper and noting the lack of a dynamical symmetry breaking mechanism. Luboš Motl offered a colorful critique, **objecting to the addition** of bosons and fermions in Lisi's superconnection, and to the violation of the Coleman-Mandula theorem In the presentation "What's new at the arXiv?" on

May 20, 2008, Simeon Warner stated that Lisi's paper is the most downloaded article on the arXiv. Among the physicists early to comment on E8 Theory, Sabine Hossenfelder, Peter Woit and Lee Smolin were generally supportive, while Luboš Motl and Jacques Distler were critical. On his blog, Musings, Jacques Distler offered one of the strongest criticisms of Lisi's approach, **claiming to demonstrate that**, unlike in the Standard Model, Lisi's model **is** nonchiral — consisting of a generation and an anti-generation — and to prove that any alternative embedding in E8 must be similarly nonchiral. These arguments were distilled in a paper written jointly with Skip Garibaldi, "There is no 'Theory of Everything' inside E8," published in Communications in Mathematical Physics. In this paper, Distler and Garibaldi offer a proof that it is impossible **to embed all** three generations of fermions **in** E8, or to obtain even the one-generation Standard Model. In a press release from his university, "Rock climber takes on surfer's theory; Garibaldi states that his article with Distler is a rebuttal of Lisi's theory. In response, Lisi argues that Distler and Garibaldi made unnecessary assumptions about how the embedding needs to happen. Addressing the one generation case, in June 2010 Lisi posted a new paper on E8 Theory, "An Explicit Embedding of Gravity and the Standard Model in E8," peer reviewed and published in a conference proceedings, describing how the algebra of gravity and the Standard Model with one generation of fermions embeds in the E8 Lie algebra explicitly using matrix representations. When this embedding is done, Lisi agrees that there **is an** antigeneration of fermions (also known as "mirror fermions") remaining in E8; but while Distler and Garibaldi **state that these** mirror fermions make the theory nonchiral, Lisi states that these mirror fermions might have high masses, making the theory chiral, or that **they might be related to the** other generations. Addressing the three generation case, Lisi agrees that three generations of fermions cannot be directly embedded in E8, **but suggests that** a gauge transformation related to triality **might be used** to relate the 64 mirror fermions and 64 other E8 generators to two other generations of 64 fermions. Sixteen arXiv preprints have cited Lisi's work. Lee Smolin's "The Plebanski action extended to a unification of gravity **and** Yang-Mills theory," December 6, 2007, **proposes a** symmetry breaking mechanism to go from an E8 symmetric action to Lisi's action for the Standard Model and gravity. Roberto Percacci's "Mixing internal and spacetime transformations: some examples and counterexamples" addresses a general loophole in the Coleman-Mandula theorem also thought to work in E8 Theory Percacci and Fabrizio Nesti's "Chirality in unified theories of gravity" **confirms the** embedding of the algebra of gravitational and Standard Model forces acting on a generation of fermions in $so(3,11) \oplus 64$, mentioning that Lisi's "ambitious attempt to unify all known fields into a single representation of E8 **stumbled into** chirality issues. Mathematician Bertram Kostant discussed Lisi's work in a colloquium presentation at UC Riverside. In a joint paper with Lee Smolin and Simone Speziale, published in Journal of Physics A, Lisi proposes a new action and symmetry breaking mechanism. In "An Explicit Embedding of Gravity and the Standard Model in E8," Lisi describes E8 Theory using explicit matrix representations.(Wikipedia: emphasis mine)

INTRODUCTION—VARIABLES USED

Vyavahārasiddhi (Proof of Convention);Yuktiṣāṣṭika (Sixty Verses on Reasoning); Catuḥstava (Hymn to the Absolute Reality); Ratnāvalī (Precious Garland) ;Pratītyasamutpādahṛdayakārika (Constituents of Dependent Arising); Sūtrasamuccaya; Bodhicittavivaraṇa (Exposition of the Enlightened Mind) are other Buddhist texts that deal with the mandala studies. This is written as a reference. There is no connection between the E (8) Theory and the Buddhist mandala whatsoever.

LISI'S THEORY AND QUANTUM COMPUTATION:

Adiabatic quantum computation

Adiabatic quantum computation (AQC) relies on the adiabatic theorem to do calculations. First, a

complex Hamiltonian is found whose ground state describes the solution to the problem of interest. Next, a system with a simple Hamiltonian is prepared and initialized to the ground state. Finally, the simple Hamiltonian is adiabatically evolved to the complex Hamiltonian. By the adiabatic theorem, the system remains in the ground state, so at the end the state of the system describes the solution to the problem. AQC is a possible method to get around the problem of energy relaxation. Since the quantum system is in the ground state, interference with the outside world cannot make it move to a lower state. If the energy of the outside world (that is, the "temperature of the bath") is kept lower than the energy gap between the ground state and the next higher energy state, the system has a proportionally lower probability of going to a higher energy state. Thus the system can stay in a single system eigenstate as long as needed. Universality results in the adiabatic model are tied to quantum complexity and QMA-hard problems. The k-local Hamiltonian is QMA-complete for k ≥ 2. QMA-hardness results are known for physically realistic lattice models of qubits such as $H = \sum_i h_i Z_i + \sum_{i<j} J^{ij} Z_i Z_i + \sum_{i<j} K^{ij} X_i X_i$ where Z, X represent the Pauli matrices σ_z, σ_x. Such models are used for universal adiabatic quantum computation. The Hamiltonians for the QMA-complete problem can also be restricted to act on a two dimensional grid of qubits or a line of quantum particles with 12 states per particle. And if such models were found to be physically realisable, they too could be used to form the building blocks of a universal adiabatic quantum computer. In practice, there are problems during a computation. As the Hamiltonian is gradually changed, the interesting parts (quantum behaviour as opposed to classical) occur when multiple qubits are close to a tipping point. It is exactly at this point when the ground state (one set of qubit orientations) gets very close to a first energy state (a different arrangement of orientations). Adding a slight amount of energy (from the external bath or as a result of slowly changing the Hamiltonian) could take the system out of the ground state, and ruin the calculation. Trying to perform the calculation more quickly increases the external energy; scaling the number of qubits makes the energy gap at the tipping points smaller.

The Computer-simulated Universe: Divinity or Reality? Part 2: This article explores the search for a comprehensive creator and the possibility that our universe exists only within a highly advanced supercomputer. Posted by Andy Mac Nevin | Last updated: Mar 25, 2013 The E8 Lie Group and the Theory of Everything

(1) The man who may be on the brink of answering the most important question in science is a Physicist turned “beach bum” called Garrett Lisi, and his E8 Theory is (in his opinion) on its way to forming a Theory of Everything. The E8 Theory is essentially **based on** twisting a collection of identical circles around each other in different ways to create a mathematical shape known as a lie group and from this lie group the construction of our universe can be found. Large Hadron Collider, The Telegraph, Roger Highfield, 2007.

(2) It starts by wrapping one circle around another, keeping the second circle perpendicular to the first, **creating** a shape called a Torus, specifically a Ring Torus, which looks very similar to a ring donut. Then by wrapping a third circle around that, perpendicular to the first two circles you achieve a 3 dimensional shape. Garrett Lisi’s theory was discovered by continually repeating this process of twisting circles until he had 248 circles interlinked, **creating a** shape beyond standard human comprehension, almost impossible to visualise in 3 dimensions. He called this shape the E8 Lie group.

(3) Many may be wondering how this relates to the creation of our universe or our simulated universe or indeed fits into the Grand Unification theory. Garrett’s discovery becomes interesting when you begin to pull the E8 Lie Group shape apart and notice that the way in which these circles twist around each other **corresponds incredibly** closely to the way fundamental particles interact on a Quantum level.

(4) When dissected and mapped by computer programs, it can be seen that every circle **relates (E&EB) to a** different type of elementary particle, certain circles have been seen to directly relate to electrons, others to the strong force particles known as Kluons, others to Photons (the particles carrying Electromagnetism), and even to the weak force particles known as the W and Z Boson particles. TED Conference, G. Lisi, Feb 2008

(5) Then Garrett found a set of circles, which he believed **represented** the illusive graviton; the particles that make up gravity.

(6) This is a particle that has avoided detection since the birth of the human race. The route Garrett is going down is a very important one, **because(E) if** what he is proposing and working towards is even remotely close to being correct then it's a hypothesis that almost all have given up on and that is "a Unique unification within a beautiful mathematical structure" as put so eloquently by Dr lee Smolin, a theoretical physicist from the PITP (Perimeter Institute for Theoretical Physics). Life of the Cosmos, L. Smolin, Oxford University Press, 1997.

Computer Advancement: A Life in Pixels

(7) Garrett's theories highlight the huge importance shapes have in the lives of the human race. Whilst his discovery involves highly complex shapes and structures it is important not to disregard the simplest of shapes, for example the Pixel. It may be a tiny square but **it can (EB) provide** just as much insight into our existence as it's complex adversary the E8 Lie Group.

(8) As many will already know a computer-based image, no matter how realistic or complex, **can (E&EB) always be broke down** into pixels. These are the smallest addressable screen elements in a display device such as a computer screen or television.

(9) Each pixel **is (Eb) a part of** a larger image and it is given its coordinates on the display device by the computer system.

(10) These pixels vary in colour intensity **and by (E&EB) combining** many pixels of varying intensities of colour an image is created. The greater the number of pixels in a fixed boundary the sharper the image will be.

(11) The main way we discern between what's real and what isn't in our world **is (=)** pixel density.

No matter how good a graphical representation is the pixel density will always unveil its authenticity. To put in bluntly, zoom in and the quality decreases. This is where the idea of a computer-simulated universe begins to hold some weight, not much, but some. Most would assume that things in our universe don't break down into pixels but they would be wrong, zoom in and on a quantum level it is clear that all matter is made up of pixels.

(12) These are tiny fundamental particles, billions of times smaller than an atom. The theory that handles this far beyond microscopic world is called Quantum Mechanics and it applies to the whole universe. As Rich Terrell, a JPL (jet propulsion laboratory) scientist, states, space, time, energy and indeed our entire universe is quantised. It's all made up of tiny pixels**, which (EB) means** the universe has a finite number of components, which means a finite number of states, which means it's computable. Through the Wormhole, Discovery, R.Terrell, 2011

Many still believe that a computer will never possess the traits of a human or the ability **to create** self-aware programs and entire simulated worlds. Moore's law states that computational power doubles every 18 months to two years but over the last two decades it's been doubling roughly every 13 months. The current fastest computers on the planet are comparable and even exceeding the computational power of the human brain. To put that in perspective human brains can complete roughly 1 million billion operations per second. So following Moore's law and taking into consideration the increased exponential rate of technological advancement, that number of operations per second could double in 13 -18 months meaning that by 2012/2013 the fastest super computers on the planet will be twice as fast as the human brain. When you continue along the scale, that shows an increase by a factor of 500 over the next decade, meaning that by

2020 computers will be 500 times faster than our brains. If these figures hold true, a computer will be able to create a 100 percent photo-realistic recreation of human life and that is where the line between simulation and reality begins to blur. Rich Terrell claims that if you take a human brain and a laptop computer built 50 years in the future, both will be a similar size and volume and use roughly the same amount of power. If you then remove the element of physical appearance and begin to ask them questions and you can no longer differentiate between the human being answering and the computer answering then they are qualitatively equivalent. So in essence if you believe a human being to be conscious and self-aware then you must assume the same of the computer. Through the Wormhole, Discovery, R.Terrell, 2011

This whole concept of a computer simulated universe coupled **with** human desperation to discover the nature of our existence will no doubt lead to immense discoveries over the coming years, decades and centuries, but it's important never to discount any theory without first understanding it's fundamental laws. Our ability to understand and to analyse is one we have built up over millions of years of evolution. The big question is; did that evolution take place within a computer-simulated environment or within a truly organic planet and universe?

Spacetime May Have Fractal Properties on a Quantum Scale Mar 25, 2009: By Lisa Zyga

(13) As scale decreases, the number of dimensions of k-Minkowski spacetime (red line: figure deleted for spatial constraints), which is an example of a space **with (e&eb)** quantum group symmetry, decreases from four to three. In contrast, classical Minkowski spacetime (blue line) is …more

(14) (PhysOrg.com) -- Usually, we think of spacetime as being four-dimensional, with three dimensions of space and one dimension of time. However, this Euclidean perspective **is just (eb) one of** many possible multi-dimensional varieties of spacetime.

(15) For instance, string theory **predicts** the existence of extra dimensions - six, seven, even 20 or more. As physicists often explain, it's impossible to visualize these extra dimensions; they exist primarily to satisfy mathematical equations.

(16) As if extra dimensions weren't strange enough, new research has probed an even more mind-bending possibility: that spacetime has dimensions that change **depending on the** scale, and the dimensions could have fractal properties on small scales.

(17) In a recent study, Dario Benedetti, a physicist at the Perimeter Institute for Theoretical Physics in Waterloo, Ontario, has investigated two possible examples of spacetime with scale-dependent dimensions deviating from classical values at short scales. More than being just an interesting idea, this phenomenon might **provide insight** into a quantum theory of relativity, which also has been suggested to have scale-dependent dimensions. Benedetti's study is published in a recent issue of Physical Review Letters.

(18) "It is an old idea in quantum gravity that at short scales spacetime **might (=) appear** foamy, fuzzy, fractal or similar," Benedetti told PhysOrg.com.

(19) "In my work, I suggest that quantum groups **are (=) a** valid candidate for the description of such a quantum spacetime.

(20) Furthermore, computing the spectral dimension, I provide for the first time a link between quantum groups/noncommutative geometries **and** apparently unrelated approaches to quantum gravity, such as Causal Dynamical Triangulations and Exact Renormalization Group. And establishing links between different topics is often one of the best ways we have to understand such topics."

(21) In his study, Benedetti explains that a spacetime with quantum group symmetry **has (e) in general** a scale-dependent dimension.

(22) Unlike classical groups, which act on commutative spaces, quantum groups **act (e&eb) on** noncommutative spaces (e.g. where xy doesn't equal yx), which emerges through their unique curvature and quantum uncertainty.

(23) Here, Benedetti considers two types of spacetime with quantum group symmetry - a quantum sphere and k-Minkowski spacetime - and calculates their dimensions. In both spaces, the dimensions **have(e)** fractal properties at small scales, and only reach classical values at large scales

(24) "In simple words, the relation between quantum groups and noncommutative geometry is as follows," he explained. "Classically, we know that certain spaces **are (=)** invariant under the action of some classical groups; for example, Euclidean space is invariant under rotations and translations.

(25) A quantum group is a deformation of a given classical group, and is such that no classical space **can have it** as a symmetry group.

(26) The invariant space has to be as well a deformation of a classical space, a deformation **that (eb) makes it** noncommutative.

(27) No relation of all this to fractals is known, but in my work I've found that they do have a common property, **that is,** a non-integer dimensions (at some scale)."

(28) Compared to a Euclidean sphere, a quantum sphere's curvature and uncertainty **make it a** noncommutative space.

(29) When calculating the spectral dimension of the quantum sphere, Benedetti found that it closely **resembles a** standard sphere on large scales;

(30) However, as the scale decreases, the dimensions of the quantum sphere (=) deviate and go down to zero.

(31) He describes this phenomenon **as a (e) signature of the** fuzziness, or uncertainty, of the quantum sphere, and also as resulting from fractal behavior at small scales.

(32) In the second kind of space, k-Minkowski spacetime, the dimensions **also (e&eb) deviate** from the constant behavior of classical Minkowski spacetime. While the latter always has four dimensions, independent of the scale, the number of dimensions in the quantum version **decreases (e)** to three as a function of the scale. In both k-Minkowski spacetime and the quantum sphere, the dimensionality becomes non-integral, which is a typical signature of fractal geometry.

(33) Benedetti's results match previous approaches to quantum gravity, which also point to the **emergence of a** ground-scale spacetime with fractal properties.

(34) Together, these studies may help scientists **understand (e) the** unique Planck scale properties of spacetime, and possibly tie in to a quantum theory of gravity.

(35) For instance, as Benedetti explains, the fractal nature of quantum spacetime might **enable** gravity **to (e&eb) cure** its own ultraviolet behavior by dimensional reduction.

(36) The main problem with gravity is that apparently it cannot be quantized as other field theories; in jargon it **is said (=) to be** non-renormalizable," he said.

(37) "This problem **is specific (eb) to** four-dimensional spacetime.

(38) If spacetime had only two dimensions, then quantum gravity would be much simpler and treatable. The problem with a two-dimensional theory is that it is unphysical, as we see four dimensions at our scales. Things can be solved combining four **and (e&eb)** two dimensions at different scales.

(39) That is, if gravity itself provides a mechanism by which the dimension of spacetime depends on the scale at which we probe it (four at our and larger scales and two at very short scales), **then we (eb) could have** a physical theory (compatible with observations) that is free of quantum (short scale) troubles."**More information: Benedetti, Dario. "Fractal Properties of Quantum Spacetime." Physical Review Letters 102, 111303 (2009**)Emphasis mine

Fractal properties of quantum spacetime Dario Benedetti

Authors show that in general a spacetime having quantum group symmetry has also a scale dependent fractal dimension **which deviates** from its classical value at short scales, a phenomenon **that resembles** what observed in some approaches to quantum gravity. In particular we analyze the cases of a quantum sphere and of $\k$-Minkowski, the latter being **relevant in** the context of quantum gravity.Phys.Rev.Lett.102:111303,

2009DOI: 10.1103/PhysRevLett.102.111303 arXiv: 0811.1396 [hep-th]

Observation of a quarter of an electron charge at the ν = 5/2 quantum Hall state M. Dolev et al

(1) The fractional quantum Hall effect, where plateaus in the Hall resistance at values of h/νe2 **coexist with** zeros in the longitudinal resistance, **results (e) from** electron correlations in two dimensions under a strong magnetic field. (Here h is Planck's constant, ν the filling factor and e the electron charge.)

(2) Current flows along the sample edges and is carried by charged excitations (quasiparticles) whose charge **is a** fraction of the electron charge.

(3) Although earlier research concentrated on odd denominator fractional values of ν, the observation of the even denominator ν = 5/2 state sparked much interest. This state is conjectured to be **characterized (e)by** quasiparticles of charge e/4, whose statistics non-abelian'— in other words, are interchanging two quasiparticles **may(e&eb) modify** the state of the system into a different one, rather than just adding a phase as is the case for fermions or bosons.


(4) As such, these quasiparticles **may be (e) useful** for the construction of a topological quantum computer.

(5) Here authors report data on shot noise **generated (e) by** partitioning edge currents in the ν = 5/2 state, **consistent with** the charge of the quasiparticle being e/4, and **inconsistent** with other possible values, such as e/2 and e. Although this finding does not prove the non-abelian nature of the ν = 5/2 state, it is the first step towards a full understanding of these new fractional charges.

POSTED BY ROB BRYANTON AT 4:09 A.M LABELS: E8, MANY WORLDS, AND PHASE SPACE

(1) At its largest possible expression, this point **becomes (=) a** finite but unbounded hypersphere occupying the entire phase space of the dimension it's within.

(2) We can also look at the other Platonic Solids we've been looking at here with this same idea of symmetric points pushed out from a central position in mind, and this includes shapes like the nested tetrahedra we see below left, which can be used **to (eb)create** the dodecahedron at the right.



.

Speaking of phase space, here's a link to a recent New Scientist article about a new paper published by Lee Smolin and others at arxiv.org, proposing that all possible universes **could be (eb) contained within** an eight-dimensional phase space.

(3) To follow the logic we've been pursuing here, the six-dimensional phase space for our own universe (or any specific universe with locked-in physical laws) **would be a (eb) subset of** an eight-dimensional phase space that encompasses all possible universes.

(4) Likewise, it's worth noting that Everett didn't limit his Theory of the Universal Wavefunction to just our unique universe: so while the phase space of a unique different-initial-conditions universe such as ours could be encompassed by the sixth dimension, the logic we've followed here **shows (eb) us that** Everett's theory ultimately should be considered from the phase space of the eighth dimension

to include the wave function of all those other possible universes.

(5) So what's beyond the eighth dimension? The simplest way to take our thought experiment to the next dimension up now is to ask this: how would you jump from the universe with a different value for gravity **to (eb) the** universe with the different speed of light without passing through the intermediate possibilities?

(6) To get to that addition degree of freedom, you need the next dimension up. For our 2D flatlander ant, that was the third dimension. For our fly in the fifth-dimensional garden hose analogy, that freedom to flit from location to location was in the sixth dimension. And for all these different universes considered from the eighth dimension, where do you think we'd achieve this additional degree of freedom? **POSTED BY ROB BRYANTON AT 4:09 AM LABELS: E8, MANY WORLDS, AND PHASE SPACE**

The basic problem in many-worlds theories H P Stapp Canadian Journal of Physics, 2002, 80(9): 1043-1052, 10.1139/p02-068

(7) It is emphasized that a many-worlds interpretation of quantum theory **exists (EB)** only to the extent that the associated basis problem is solved. The core basis problem is that the robust enduring states specified by environmental decoherence effects **are (=)** essentially Gaussian wave packets that form continua of non-orthogonal states.

(8) environmental decoherence effects **are** essentially Gaussian wave packets that form continua of non-orthogonal states **are(E) not** a discrete set of orthogonal basis states to which finite probabilities can be assigned by the usual rules.

(9) The natural way to get an orthogonal basis without going outside the Schrödinger dynamics **is to use (E) the** eigenstates of the reduced density matrix, and this idea is the basis of some recent attempts by many-worlds proponents to solve the basis problem. But these eigenstates do not enjoy the locality and quasi-classicality properties of the states defined by environmental decoherence effects, and hence are not satisfactory preferred basis states. This core problem is obscured by approaches that treat the universe as a quantum computer, but it needs to be addressed and resolved before a many-worlds-type interpretation can be said to exist.

NOTATION

Module One

G_{13} : Category one of TOE e (8). This does not mean there are three theories of TOE. We are talking of applications of TOE to various investigating systems, and the corresponding concomitant characterization form the bastion, pillar, post, stylobate and sentinel of the stratifications schema.

G_{14} : Category two of TOE e(8)

G_{15} : Category three of TOE e(8)

T_{13} : Category one of so (7, 1), strong BL so (1, 7), fermions (3(8) X(8)). Again the propensities and characterstics of the investigating systems form the sole criterion.

T_{14} : Category two of so(7,1) ,strong BL so (1,7) ,fermions (3(8) X(8))

T_{15} : Category three of so(7,1) ,strong BL so (1,7) ,fermions (3(8) X(8))

Module Two

G_{16} : Category one of gravi (weak) so (7,1)

G_{17} : Category two of gravi (weak) so (7,1)

G_{18} : Category three of gravi (weak) so (7,1)

T_{16} : Category one of gravity so (3,1), frame higgs(4X2+2(bar)),electroweak$(su(2))_L$ +$(su(2))_R$

T_{17} : Category two of gravity so (3,1), frame higgs(4X2+2(bar)),electroweak$(su(2))_L$ +$(su(2))_R$

T_{18} : Category three of gravity so (3,1), frame higgs(4X2+2(bar)),electroweak$(su(2))_L$ +$(su(2))_R$

Module three

G_{20} : Category one of strong (B,L) so(1,7)

G_{21} : Category two of strong (B,L) so(1,7)

G_{22} : Category three of strong (B,L) so(1,7)

T_{20} : Category one of strong su(3), BL $u(1)_{B_L}$,new bosons u(1) + 3 X 6 ,

T_{21} : Category two of strong su(3), BL $u(1)_{B_L}$,new bosons u(1) + 3 X 6 ,

T_{22} : Category three of strong su(3), BL $u(1)_{B_L}$,new bosons u(1) + 3 X 6 ,

Module four

G_{24} : Category one of fermions (3X(8X8))

G_{25} : Category two of fermions (3X(8X8))

G_{26} : Category three of fermions (3X(8X8))

T_{24} : Category one of gen 1 8 s+ X 8s+ , gen 8_ X 8_{S-} , gen 3 8_V $X8_V$

T_{25} : Category two of gen 1 8 s+ X 8s+ , gen 8_ X 8_{S-} , gen 3 8_V $X8_V$

T_{26} : Category three of gen 1 8 s+ X 8s+ , gen 8_ X 8_{S-} , gen 3 8_V $X8_V$

Module five

At its largest possible expression, this point **becomes a** finite but unbounded hypersphere occupying the entire phase space of the dimension it's within

G_{28} : Category one of At its largest possible expression, this point

G_{29} : Category two of At its largest possible expression, this point

G_{30} : Category three of At its largest possible expression, this point

T_{28} : Category one of finite but unbounded hypersphere occupying the entire phase space of the dimension it's within

T_{29} : Category two of finite but unbounded hypersphere occupying the entire phase space of the dimension it's within

T_{30} : Category three of finite but unbounded hypersphere occupying the entire phase space of the dimension

it's within

Module six

We can also look at the other Platonic Solids we've been looking at here with this same idea of symmetric points pushed out from a central position in mind, and this includes shapes like the nested tetrahedra we see below left, which can be used **to (eb)create** the dodecahedron at the right

G_{32} : Category one of dodecahedron at the right

G_{33} : Category two of dodecahedron at the right

G_{34} : Category three of dodecahedron at the right

T_{32} : Category one of Platonic Solids we've been looking at here with this same idea of symmetric points pushed out from a central position in mind, and this includes shapes like the nested tetrahedra we see below left, which can be used

T_{33} : Category two of Platonic Solids we've been looking at here with this same idea of symmetric points pushed out from a central position in mind, and this includes shapes like the nested tetrahedra we see below left, which can be used

T_{34} : Category three of Platonic Solids we've been looking at here with this same idea of symmetric points pushed out from a central position in mind, and this includes shapes like the nested tetrahedra we see below left, which can be used

Module seven

To follow the logic we've been pursuing here, the six-dimensional phase space for our own universe (or any specific universe with locked-in physical laws) **would be a (eb) subset of** an eight-dimensional phase space that encompasses all possible universes

G_{36} : Category one of six-dimensional phase space for our own universe (or any specific universe with locked-in physical laws)

G_{37} : Category two of six-dimensional phase space for our own universe (or any specific universe with locked-in physical laws)

G_{38} : Category three of six-dimensional phase space for our own universe (or any specific universe with locked-in physical laws)

T_{36} : Category one of **subset of** an eight-dimensional phase space that encompasses all possible universes

T_{37} : Category two of **subset of** an eight-dimensional phase space that encompasses all possible universes

T_{38} : Category three of **subset of** an eight-dimensional phase space that encompasses all possible universes

Module eight

Likewise, it's worth noting that Everett didn't limit his Theory of the Universal Wavefunction to just our unique universe: so while the phase space of a unique different-initial-conditions universe such as ours could be encompassed by the sixth dimension, the logic we've followed here **shows (eb) us that** Everett's theory ultimately should be considered from the phase space of the eighth dimension to include the wave function

of all those other possible universes.

G_{40} : Category one of phase space of a unique different-initial-conditions universe such as ours could be encompassed by the sixth dimension

G_{41} : Category two of phase space of a unique different-initial-conditions universe such as ours could be encompassed by the sixth dimension

G_{42} : Category three of phase space of a unique different-initial-conditions universe such as ours could be encompassed by the sixth dimension

T_{40} : Category one of Everett's theories ultimately should be considered from the phase space of the eighth dimension to include the wave function of all those other possible universes.

T_{41} : Category two of Everett's theory ultimately should be considered from the phase space of the eighth dimension to include the wave function of all those other possible universes.

T_{42} : Category three of Everett's theory ultimately should be considered from the phase space of the eighth dimension to include the wave function of all those other possible universes.

Module Nine

So what's beyond the eighth dimension? The simplest way to take our thought experiment to the next dimension up now is to ask this: how would you jump from the universe with a different value for gravity **to (eb) the** universe with the different speed of light without passing through the intermediate possibilities?

environmental decoherence effects **are** essentially Gaussian wave packets that form continua of non-orthogonal states **are(E) not** a discrete set of orthogonal basis states to which finite probabilities can be assigned by the usual rules.

G_{44} : Category one of universe with a different value for gravity; discrete set of orthogonal basis states to which finite probabilities can be assigned by the usual rules.

G_{45} : Category two of universe with a different value for gravity; discrete set of orthogonal basis states to which finite probabilities can be assigned by the usual rules.

G_{46} : Category three of universe with a different value for gravity; discrete set of orthogonal basis states to which finite probabilities can be assigned by the usual rules.

T_{44} : Category one of universe with the different speed of light without passing through the intermediate possibilities; environmental decoherence effects **are** essentially Gaussian wave packets that form continua of non-orthogonal states;

T_{45} : Category two of universe with the different speed of light without passing through the intermediate possibilities; environmental decoherence effects **are** essentially Gaussian wave packets that form continua of non-orthogonal states;

T_{46} : Category three of universe with the different speed of light without passing through the intermediate possibilities; environmental decoherence effects **are** essentially Gaussian wave packets that form continua of non-orthogonal states;

SECTION SEVENTY THREE

Exotic matter: Rocky Razzmatazz

Abstract: Five requirements, plus two relating to the communication of quantum information, are extensively explored and related to the many schemes in atomic physics, quantum optics, nuclear and electron magnetic resonance spectroscopy, superconducting electronics, and quantum-dot physics, for achieving quantum computing are the essential functional prerequisites and the necessary elemental endowment of the exotic matter and quantum computation. The advent of quantum information processing, as an abstract concept, **has given birth to** a great deal of new thinking, of a very concrete form, about how to create physical computing devices that operate in the hitherto unexplored quantum mechanical regime. Scope proposed and underway, to the implementation of quantum hardware is remarkable, emerging from specialties in atomic physics; quantum optics; nuclear electron magnetic resonance spectroscopy; in superconducting device physics electron physics; mesoscopic and quantum dot research. Manipulation and transmission of information is today carried out by physical machines (computers, ∗Prepared for Fortschritte der Physik special issue, Experimental Proposals for Quantum Computation, eds. H.-K. Lo and S. Braunstein. Routers, scanners, etc.), in which the embodiment and transformations of this information can be described **using the** language of classical mechanics. But the final physical theory of the world is not Newtonian mechanics, and there is no reason to suppose that machines following the laws of quantum mechanics should have the same computational power as classical machines; indeed, since Newtonian mechanics emerges as a special limit of quantum mechanics, quantum machines can only have greater computational power than classical ones. The great pioneers and visionaries, who pointed the way towards quantum computers, Deutsch, Feynman, and others, were stimulated by such thoughts. Of course, by a similar line of reasoning, it may well be asked whether machines **embodying the** principles of other refine d descriptions of nature (perhaps general relativity or string theory) may have even more information processing capabilities; speculations exist about these **more exotic possibilities**, but they are beyond the scope of the present discussion. But computing with quantum mechanics really deserves a lot more attention than wormhole computing or quantum-gravity computing; quantum computing, while far in the future from the perspective of CMOS roadmaps and projections of chip fab advances, can certainly be seen as a real

prospect from the perspective of research studies in quantum physics. (**The Physical Implementation of Quantum Computation David P. DiVincenzo emphasis mine)** As the sizes of electronic components shrink, soon down to the size of single atoms or molecules, quantum interactions become ever more important, but conspicuous by its presence and perceptible by its intensity. Consequently, enhanced knowledge and exploitation of quantum effects is essential. Researchers have uncovered evidence for a long-sought-after quantum state of matter, a spin liquid. A quantum computer is a device for computation that makes direct use of quantum mechanical phenomena, such as superposition and entanglement, to perform operations on data. The basic principle behind quantum computation is that quantum properties can be used to represent data and perform operations on these data. Although quantum computing is still in its infancy, experiments have been carried out in which quantum computational operations **were executed** on a very small number of qubits (quantum binary digits). Quantum computing research helps development of quantum computers for both civilian and national security purposes, such as cryptanalysis. **Concept of spin** is at the heart of all magnetic phenomena. For instance, a refrigerator magnet, at the microscopic level, consists of trillions of trillions of iron atoms all lined up. Each of these atoms can be thought of loosely as a tiny spinning ball. The orientation of that spin **is what makes the** atom into a tiny magnet. The refrigerator magnet is an example of a ferromagnet, the ferro part coming from the Latin word for iron. In a ferromagnet, all the atomic spins are lined up in the same way, **producing** a large cooperative magnetic effect. Ferromagnets important though they may be aren't the only kind of material where magnetic interactions between spins are critical. In anti-ferromagnets, for instance, the neighboring spins are driven **to be** anti-aligned. That is, the orientations of the spins **alternate** up and down. Accumulative magnetic effect of all these up and down spins **are that the** material has no net magnetism. High-temperature superconducting materials discovered in the 1980s are an important example of an anti-ferromagnetic structure.(Physics.Org Blog) Stability analysis of exotic matter is studied by various authors such as Eric Poisson and Matt Visser, Matt Visser (Traversable wormholes from surgically modified Schwarzschild spacetimes) Eric Poisson and Matt Visser (Thin-shell wormholes: Linearization stability) Stability analysis of dynamic thin shells (Francisco S N Lobo and Paulo Crawford). Spherically symmetric thin-shell wormholes in the presence of a cosmological constant **are constructed** applying the cut-and-paste technique implemented by Visser. Darmois-Israel formalism **is used to find** the surface stresses, which are concentrated at the wormhole throat, are determined. This construction allows one to apply a dynamical analysis to the throat, considering linearized radial perturbations around static solutions. For a large positive cosmological constant, i.e., for the Schwarzschild Sitter solution, **there shall be significant increase** in the region of stability relatively to the null cosmological constant case,(Poisson and Visser). With a negative cosmological constant, i.e., the Schwarzschild-anti de Sitter solution, **there shall be** decrease in the region of stability is decreased. In particular, considering static solutions with a generic cosmological constant, the weak and dominant energy conditions are violated, while for $a0 \leq 3M$ the null and strong energy conditions are satisfied. The surface pressure of the static solution is strictly positive for the Schwarzschild and Schwarzschild-anti de Sitter spacetimes, **but takes** negative values, assuming a surface tension in the Schwarzschild-de Sitter solution, for high values of the cosmological constant and the wormhole throat radius. **(Linearized stability analysis of thin-shell wormholes with a cosmological constant Francisco S. N. Lobo and Paulo Crawford)**

INTRODUCTION—VARIABLES USED

Stanford physicists make new form of matter: The laser-cooled quantum gas opens exciting new realms of unconventional superconductivity. BY MAX MCCLURE (Some paragraphs are removed due to spatial constraints)

(1) Within the exotic world of macroscopic quantum effects, where fluids flow uphill, wires conduct without electrical resistance and magnets levitate, there is an even stranger family of "unconventional" phenomena. These effects often defy explanation by current theoretical physics,

but hold enormous promise for the development of such futuristic technologies as room-temperature superconductors, ultrasensitive microscopes and quantum computation. Much of the confusion surrounding the field is due to the sorts of materials that exhibit unconventional superconductivity. These substances **are (e) made up of** strongly interacting fermions, a class of particles that are often very difficult to understand on the quantum level.

(2) Last week's announcement by a Stanford team in Physical Review Letters that **it has created** the world's first dipolar quantum fermionic gas from the metal dysprosium – "an entirely new form of quantum matter," as Stanford applied physics Professor and lead author Benjamin Lev put it – **represents(eb) a** major step toward understanding the behavior of these systems of particles. And this understanding makes for a leap toward the supernatural-seeming applications that condensed-matter physics conjures.

(3) When the thermal energy of some substances drops below a certain critical point, it is often no longer possible to consider its component particles separately because the material **becomes** "strongly correlated" and its quantum effects become difficult to understand and study.

(4) Making the material out of a gas of atoms allows what is normally only observed on a nanometer scale to become visible. These quantum gases, the coldest objects known to man, **are where (e) researchers see** zero-viscosity fluids – superfluids – that are mathematical cousins of superconductors.

(5) Lev, as well as Stanford graduate students Mingwu Lu and Nathaniel Burdick, heated their particles in a crucible to around 1,300 degrees Celsius and shot them into a powerful vacuum. Using the world's most powerful continuous-wave blue laser, the particles were then cooled to within a thousandth of a degree of absolute zero. Subsequent lasers and an evaporative cooling process eventually **brought (eb)the gas down to the** experimental temperature of 64 nanokelvins – very, very cold.

(6) The initial step – trapping particles in the "optical molasses" **created by (e) the** high-powered lasers – works by exciting a particle and then allowing it to return to its initial state. The particle loses energy in this process, **producing (eb)** cooling dramatically. Typically, the substances that researchers cool in this way **only (=) have** two or three energy levels, making for a simple "optical loop." Dysprosium, on the other hand has more than 140.

(7) The evaporative cooling process the researchers hoped to use to bring the gas down to the nanokelvin range **depends** on collisions between particles. Collisions **produce (eb)** both dissipation of energy and knock high-energy particles out of the system. "But the lore in the field is that identical fermions never collide," Lev said.

(8) Fermions **obey (eb) the** Pauli Exclusion Principle, which states that no two fermions can occupy the same state simultaneously. This inviolable rule seems to disallow direct interactions between fermionic particles.

(9) Dysprosium's status as the most magnetic element was crucial **to (e) overcoming** this apparent impossibility. Extraordinarily strong magnetic dipolar interactions between the dysprosium atoms **lead (eb) the** particles to cool to below the critical temperature by colliding from afar, despite being quantum mechanically identical. Thus far, the result of the Lev lab's high-tech efforts is a tiny ball of ultracold quantum dipolar fluid. But the researchers have reason to believe that the humble substance **will exhibit the** seemingly contradictory characteristics of both crystals and superfluids. This combination **could lead to** quantum liquid crystals, or quantum-mechanical versions of the liquid crystals that make up most electronic displays. **Or it could yield** a supersolid – a hypothetical state of matter that would, in theory at least, be a solid with superfluid characteristics.

(10) The researchers have already begun developing a microscope **to make (e) use of** the dipolar quantum fluid's unique characteristics. Their "cryogenic atom chip microscope" is a magnetic probe that should measure magnetic fields with unprecedented sensitivity and resolution. This kind of probe may even allow for a more stable form of quantum computation **that (e) uses** exotic quantum

matter to process information, known as a topologically protected quantum computer.

X. G. WEN, Int. J. Mod. Phys. B, 04, 239 (1990). DOI: 10.1142/S0217979290000139 TOPOLOGICAL ORDERS IN RIGID STATES

(1) Topological order — in rigid states (the states with no local gapless excitations) concentrates on **characterization (eb) of** the different topological orders. Chiral spin states of 2 + 1 dimensional spin systems **are (e) described by** the topological Chern-Simons theories in the continuum limit.
(2) Topological orders **can be (e) characterized by** a non-Abelian gauge structure over the moduli space which parametrizes a family of the model Hamiltonians supporting topologically ordered ground states.
(3) In 2 + 1 dimensions, the non-Abelian gauge structure **determines** possible fractional statistics of the quasi-particle excitations over the topologically ordered ground states.
(4) The dynamics of the low lying global excitations **is shown (eb) to be** independent of random spatial dependent perturbations.
(5) The ground state degeneracy and the non-Abelian gauge structures discussed in this paper are very robust, **even (eb) against** those perturbations that break translation symmetry.
(6) Symmetry properties of the degenerate ground states of chiral spin states leads to repercussions of some degenerate ground states of chiral spin states **on (eb)** torus carry non-trivial quantum numbers of the 90° rotation.

Quantum Superfluid Could Be Akin To Exotic Matter Found In Quark Star Dec. 23, 2005 —

(1) In the bizarre and rule-bound world of quantum physics, every tiny spec of matter **has(e) something** called "spin" - an intrinsic trait like eye color - that cannot be changed and which dictates, very specifically, what other bits of matter the spec can share quantum space with.
(2) When fermions, the most antisocial type of quantum particle, do get together, they pair up in a wondrous dance **that enables** such things as superconductivity.
(3) For the first time, researchers at Rice University have succeeded in creating and observing an elusive and long-sought quantum state - a superfluid of fermions **with(e&eb)** mismatched numbers of dance partners. Despite more than 40 years of theoretical musings about what would occur in such a case, the result - a cluster of matched pairs surrounded by a cloud of would-be dance partners - was largely unexpected. Superconductivity and superfluidity are two factors that stand revealed therefore.
(4) Both phenomena **result (e) from a** change in the phase of matter. Anyone who has seen ice melt has seen matter change phases, and when electrons, atoms and other specs of matter change quantum phases, they behave just as differently as do ice and water in a glass.
(5) Superconducting and superfluid phases of matter **occur (eb) in** fermions - the antisocial particles that can't share quantum space - only when quantum forces become dominant.
(6) Because thermodynamic forces are typically so powerful that they overwhelm quantum interactions - like loud music overwhelms the whisper of someone nearby - superconductivity and superfluidity **only (eb) occur in** extreme cold.
(7) In the Rice experiment, when temperatures drop to within a few billionths of a degree of absolute zero, fermions with equal but opposite spin **become** (=) attracted to one another and behave, in some respects, like one particle.
(8) Like a couple on the dance floor, they don't technically share space, but they move in unison. In superconductors, these dancing pairs **allow(eb)** electrical current to flow through the material without any resistance at all, a property that engineers have long dreamed of harnessing to eliminate "leakage" in power cables, something that costs billions of dollars per year in the U.S. alone.

(9) The superconducting and superfluid phases are analogous except that superconductivity happens with particles carrying an electrical charge and superfluidity **occurs (eb) in** electrically neutral particles.
(10) In superfluids, fermionic pairing **leads (eb) to** a complete absence of viscosity - like a wave rippling back and forth in a swimming pool without ever diminishing.
(11) "Conventional theory tells us superconductivity or superfluidity **occurs (eb) only in** the presence of an equal number of spin-up and spin-down particles," said lead researcher Randy Hulet, the Fayez Sarofim Professor of Physics and Astronomy. "Physicists have speculated for almost 50 years about what would happen if this condition were not met.
(12) "**Because of the** pristine and controlled nature of our ultracold atoms, we're able to offer definitive evidence of what happens with mismatched numbers of spin-up and spin-down particles."
(13) Ultracold experiments at temperatures just a few billionths of a degree above absolute zero are Hulet's specialty. It's only been technically possible to chill atoms to these temperatures for the past 10 years, but in that time, this ability has proved remarkably **useful (e)** for testing the predictions of quantum mechanics and for exploring the properties of what physicists calls "many-body phenomena," including superconductivity and superfluidity.
(14) In the latest experiments, Hulet's team - which includes graduate students Guthrie Partridge, Ramsey Kamar and Yean-an Liao and postdoctoral researcher Wenhui Li - cooled a mixture of fermionic lithium-6 atoms to about 30-billionths of a degree above absolute zero. That's far colder than any temperature in nature - even in deepest interstellar space - and it's sufficient **to(e) quell** virtually all thermodynamic interaction in the atoms, leaving them subject to superfluid quantum pairing.
(15) Using radio waves, Hulet's team altered the ratio of spin-up and spin-down atoms in the cooled atoms with great precision. They found that the superfluid **was able to (e) tolerate** an excess of up to 10 percent unpaired fermions with no detrimental effects.
(16) Hulet's team found that increasing the ratio of spin-up to spin-down atoms eventually **caused (eb) a** phase change. When unpaired spin-up atoms rose above 10 percent of the total sample, the unpaired loners were suddenly expelled, leaving a core of superfluid pairs surrounded by a shell of excess spin-up atoms.
(17) "The gas behaves as if it is still perfectly paired, which is quite remarkable given the excess of spin-up atoms," Hulet said. "This was unexpected, **and it (eb) could signal a new**, exotic form of superfluidity that may be akin to the electron pairings in unconventional superconductors or to the quark soup that's predicted to exist at the heart of the densest neutron stars."
(18) In the largest neutron stars - known as "quark stars" - a mass about five times greater than the sun is pressed into a space smaller than the island of Manhattan. Some physics theorists believe gravity is so strong at the heart of these stars **that it creates** something called a "strange matter," a dense superfluid of up quarks, down quarks and strange quarks. The research is supported by the National Science Foundation, the Office of Naval Research, NASA and the R.A. Welch Foundation.

Rice University is consistently ranked one of America's best teaching and research universities. It is distinguished by its: size-2,850 undergraduates and 1,950 graduate students

Correlation phenomena observed by CMS may shed light on exotic state of matter Julia Velkovska and Wei Li 2013-05-31 (Figures and a few paragraphs are deleted attributable to spatial constraints)

(1) Using the same technique to analyse high-multiplicity pPb collisions **brings (eb) us** a step closer to deciphering the origin of the long-range correlations in smaller systems.
(2) In probing the ridge, CMS discovered a remarkable **similarity (=) between** the pPb and PbPb collision that produce the same number of particles in the final state.

(3) This observation provides more support to the conclusion that a "liquid" medium similar to those in PbPb collisions **might also be (e) produced in** pPb ones.

(4) CMS has advanced the correlation analysis beyond using just pairs of particles. Instead quartets of particles are studied for how these might be moving in a correlated manner. Indeed, whenever there are more than 50 particles produced in a collision**, CMS has(eb) found that** there are distinctive four-particle correlations occurring.

(5) The 50-particle threshold **corresponds t**o the multiplicity range where the ridge emerges in the pp, the pPb, and the PbPb collisions.

(6) These results **provide further (eb) evidence** of the collective nature of the effect and the similarity between different collisions systems.

(7) Several key observables **used to (e) characterise** nucleus-nucleus collisions are under investigation in the pPb system, in order to reach a consistent picture of the phenomena spanning various collisions systems.— by Julia Velkovska and Wei Li 2013-05-31, by Julia Velkovska

Loop Quantization of the Schwarzschild Black Hole, Phys. Rev. Lett. 110, 211301 (2013) prl.aps.org/abstract/PRL/v110/i21/e211301

(8) Authors quantize spherically symmetric vacuum gravity **without (e)** gauge fixing the diffeomorphism constraint.

(9) Through a rescaling, we make the algebra of Hamiltonian constraints Abelian, and therefore the constraint algebra is a true Lie algebra **which (eb) allows the** completion of the Dirac quantization procedure using loop quantum gravity techniques.

(10) They can construct explicitly the exact solutions of the physical Hilbert space annihilated by all constraints. New observables living in the bulk **appear(eb) at** the quantum level (analogous to spin in quantum mechanics) that are not present at the classical level and are associated with the discrete nature of the spin network states of loop quantum gravity. The resulting quantum space-times resolve the singularity present in the classical theory inside black holes

NOTATION

Module One

Using the same technique to analyse high-multiplicity pPb collisions **brings (eb) us** a step closer to deciphering the origin of the long-range correlations in smaller systems

G_{13} : Category one of same technique to analyse high-multiplicity pPb collisions

G_{14} : Category two of same technique to analyse high-multiplicity pPb collisions

G_{15} : Category three of same technique to analyse high-multiplicity pPb collisions

T_{13} : Category one of deciphering the origin of the long-range correlations in smaller systems

T_{14} : Category two of deciphering the origin of the long-range correlations in smaller systems

T_{15} : Category three of deciphering the origin of the long-range correlations in smaller systems

Module Two

Topological orders **can be (e) characterized by** a non-Abelian gauge structure over the moduli space which parametrizes a family of the model Hamiltonians supporting topologically ordered ground states

G_{16} : Category one of non-Abelian gauge structure over the moduli space which parametrizes a family of the model Hamiltonians supporting topologically ordered ground states

G_{17} : Category two of non-Abelian gauge structure over the moduli space which parametrizes a family of

the model Hamiltonians supporting topologically ordered ground states

G_{18} : Category three of non-Abelian gauge structure over the moduli space which parametrizes a family of the model Hamiltonians supporting topologically ordered ground states

T_{16} : Category one of Topological orders

T_{17} : Category two of Topological orders

T_{18} : Category three of Topological orders

Module three

In probing the ridge, CMS discovered a remarkable **similarity between** the pPb and PbPb collision that produce the same number of particles in the final state

G_{20} : Category one of pPb collision

G_{21} : Category two of pPb collision

G_{22} : Category three of pPb collision

T_{20} : Category one of PbPb collision

T_{21} : Category two of PbPb collision

T_{22} : Category three of PbPb collision

Module four

These results **provide further (eb) evidence** of the collective nature of the effect and the similarity between different collisions systems.

G_{24} : Category one of CMS discovered a remarkable **similarity between** the pPb and PbPb collision that produce the same number of particles in the final state

G_{25} : Category two of CMS discovered a remarkable **similarity between** the pPb and PbPb collision that produce the same number of particles in the final state

G_{26} : Category three of CMS discovered a remarkable **similarity between** the pPb and PbPb collision that produce the same number of particles in the final state

T_{24} : Category one of collective nature of the effect and the similarity between different collisions systems.

T_{25} : Category two of collective nature of the effect and the similarity between different collisions systems.

T_{26} : Category three of collective nature of the effect and the similarity between different collisions systems.

Module five

Through a rescaling, algebra of Hamiltonian constraints Abelian, and therefore the constraint algebra is a true Lie algebra **which (eb) allows the** completion of the Dirac quantization procedure using loop quantum gravity techniques.

G_{28} : Category one of through a rescaling, algebra of Hamiltonian constraints Abelian, and therefore the constraint algebra is a true Lie algebra

G_{29} : Category two of through a rescaling, algebra of Hamiltonian constraints Abelian, and therefore the constraint algebra is a true Lie algebra

G_{30} : Category three of through a rescaling, algebra of Hamiltonian constraints Abelian, and therefore the constraint algebra is a true Lie algebra

T_{28} : Category one of completion of the Dirac quantization procedure using loop quantum gravity techniques.

T_{29} : Category two of completion of the Dirac quantization procedure using loop quantum gravity techniques.

T_{30} : Category three of completion of the Dirac quantization procedure using loop quantum gravity techniques.

Module six

They can construct explicitly the exact solutions of the physical Hilbert space annihilated by all constraints. New observables living in the bulk **appear(eb) at** the quantum level (analogous to spin in quantum mechanics) that are not present at the classical level and are associated with the discrete nature of the spin network states of loop quantum gravity. The resulting quantum space-times resolve the singularity present in the classical theory inside black holes

G_{32} : Category one of New observables living in the bulk

G_{33} : Category two of New observables living in the bulk

G_{34} : Category three of New observables living in the bulk

T_{32} : Category one ofquantum level (analogous to spin in quantum mechanics) that are not present at the classical level and are associated with the discrete nature of the spin network states of loop quantum gravity. The resulting quantum space-times resolve the singularity present in the classical theory inside black holes

T_{33} : Category two of quantum level (analogous to spin in quantum mechanics) those are not present at the classical level and are associated with the discrete nature of the spin network states of loop quantum gravity. The resulting quantum space-times resolve the singularity present in the classical theory inside black holes

T_{34} : Category three of quantum level (analogous to spin in quantum mechanics) those are not present at the classical level and are associated with the discrete nature of the spin network states of loop quantum gravity. The resulting quantum space-times resolve the singularity present in the classical theory inside black holes

Module seven

The gas behaves as if it is still perfectly paired, which is quite remarkable given the excess of spin-up atoms," Hulet said. "This was unexpected, **and it (eb) could signal a new**, exotic form of superfluidity that may be akin to the electron pairings in unconventional superconductors or to the quark soup that's predicted to exist at the heart of the densest neutron stars."

G_{36} : Category one of gas behaves as if it is still perfectly paired, which is quite remarkable given the excess of spin-up atoms,

G_{37} : Category two of gas behaves as if it is still perfectly paired, which is quite remarkable given the excess

of spin-up atoms,

G_{38} : Category three of gas behaves as if it is still perfectly paired, which is quite remarkable given the excess of spin-up atoms,

T_{36} : Category one of exotic form of superfluidity that may be akin to the electron pairings in unconventional superconductors or to the quark soup that's predicted to exist at the heart of the densest neutron stars."

T_{37} : Category two of exotic form of superfluidity that may be akin to the electron pairings in unconventional superconductors or to the quark soup that's predicted to exist at the heart of the densest neutron stars."

T_{38} : Category three of exotic form of superfluidity that may be akin to the electron pairings in unconventional superconductors or to the quark soup that's predicted to exist at the heart of the densest neutron stars."

Module eight

In the largest neutron stars - known as "quark stars" - a mass about five times greater than the sun is pressed into a space smaller than the island of Manhattan. Some physics theorists believe gravity is so strong at the heart of these stars **that it creates** something called a "strange matter," a dense superfluid of up quarks, down quarks and strange quarks. The research is supported by the National Science Foundation, the Office of Naval Research, NASA and the R.A. Welch Foundation

G_{40} : Category one of largest neutron stars - known as "quark stars" - a mass about five times greater than the sun is pressed into a space smaller than the island of Manhattan. Some physics theorists believe gravity is so strong at the heart of these stars

G_{41} : Category two of largest neutron stars - known as "quark stars" - a mass about five times greater than the sun is pressed into a space smaller than the island of Manhattan. Some physics theorists believe gravity is so strong at the heart of these stars

G_{42} : Category three of largest neutron stars - known as "quark stars" - a mass about five times greater than the sun is pressed into a space smaller than the island of Manhattan. Some physics theorists believe gravity is so strong at the heart of these stars

T_{40} : Category one of something called a "strange matter," a dense superfluid of up quarks, down quarks and strange quarks

T_{41} : Category two of something called a "strange matter," a dense superfluid of up quarks, down quarks and strange quarks

T_{42} : Category three of something called a "strange matter," a dense superfluid of up quarks, down quarks and strange quarks

Module Nine

Dysprosium's status as the most magnetic element was crucial **to (e) overcoming** this apparent impossibility. Extraordinarily strong magnetic dipolar interactions between the dysprosium atoms **lead (eb) the** particles to cool to below the critical temperature by colliding from afar, despite being quantum mechanically identical. Thus far, the result of the Lev lab's high-tech efforts is a tiny ball of ultracold quantum dipolar fluid. But the

researchers have reason to believe that the humble substance **will exhibit the** seemingly contradictory characteristics of both crystals and superfluids which combination **could lead to** quantum liquid crystals, or quantum-mechanical versions of the liquid crystals that make up most electronic displays. **Or it could yield** a supersolid – a hypothetical state of matter that would, in theory at least, be a solid with superfluid characteristics

G_{44} : Category one of humble substance **will exhibit the** seemingly contradictory characteristics of both crystals and superfluids which combination

G_{45} : Category two of humble substance **will exhibit the** seemingly contradictory characteristics of both crystals and superfluids which combination

G_{46} : Category three of humble substance **will exhibit the** seemingly contradictory characteristics of both crystals and superfluids which combination

T_{44} : Category one of quantum liquid crystals, or quantum-mechanical versions of the liquid crystals that make up most electronic displays. **Or it could yield** a supersolid – a hypothetical state of matter that would, in theory at least, be a solid with superfluid characteristics

T_{45} : Category two of quantum liquid crystals, or quantum-mechanical versions of the liquid crystals that make up most electronic displays. **Or it could yield** a supersolid – a hypothetical state of matter that would, in theory at least, be a solid with superfluid characteristics

T_{46} : Category three of quantum liquid crystals, or quantum-mechanical versions of the liquid crystals that make up most electronic displays. **Or it could yield** a supersolid – a hypothetical state of matter that would, in theory at least, be a solid with superfluid characteristics

SECTION SEVENTY FOUR

Correlation Phenomenon: Omnia Causa Fiunt: Everything Happens For A Reason

Abstract: Classical phase transitions occur when a physical system reaches a state below a critical temperature characterized by a macroscopic order. Quantum phase transitions **occur at** absolute zero; they are induced by the change of an external parameter or coupling constant, and are driven by fluctuations. Examples include transitions in quantum Hall systems, localization in Si-MOSFETs (metal oxide silicon field-effect transistors; and the superconductor-insulator transition in two-dimensional systems. Both classical and quantum critical points **are governed by a** diverging correlation length, although quantum systems possess additional correlations that do not have a classical counterpart. This phenomenon, known as entanglement is the resource **that enables** quantum computation and communication. The role of the entanglement at a phase transition is not captured by statistical mechanics – a complete classification of the critical many-body state requires the introduction of concepts from quantum information theory. Theory of

critical phenomena **with** quantum information is correlated and consummated and consolidated **by exploring the** entangling resources of a system close to its quantum critical point. Class of one-dimensional magnetic systems, has entanglement **showing** scaling behaviour in the vicinity of the transition. Entanglement near a quantum phase transition by analyzing the properties of the concurrence for a class of exactly solvable models in one dimension to find that entanglement **can be (e&eb) classified** in the framework of scaling theory. Difference between classical correlations and the non-local quantum correlation, entanglement: the correlation length diverges at the phase transition, whereas entanglement in general **remains (eb)** short ranged. Nature 416, 608 (2002) DOI: 10.1038/416608a arXiv: quant-ph/0202029 **Scaling of Entanglement close to a Quantum Phase Transitions Osterloh** In 1968, Fröhlich showed that a driven set of oscillators **can condense** with nearly all of the supplied energy activating the vibrational mode of lowest frequency. This is a remarkable property usually compared with Bose–Einstein condensation, superconductivity, lasing, and other unique phenomena involving macroscopic quantum coherence. However, despite intense research, no unambiguous example has been documented. **Jeffrey R. Reimers et al** determine the most likely experimental signatures of Fröhlich condensation **and show that** they are significant features remote from the extraordinary properties normally envisaged. Fröhlich condensates are classified into 3 types: weak condensates in which profound effects on chemical kinetics are possible, strong condensates in which an extremely large amount of energy **is channeled** into 1 vibrational mode, and coherent condensates in which this energy is placed in a single quantum state. Coherent condensates **are shown to** involve extremely large energies, to not be produced by the Wu–Austin dynamical Hamiltonian **that provides** the simplest depiction of Fröhlich condensates formed using mechanically supplied energy, and to be extremely fragile. They are inaccessible in a biological environment. Hence the Penrose–Hameroff orchestrated objective-reduction model and related theories for cognitive function **that embody coherent** Fröhlich condensation as an essential element are untenable. Weak condensates, however, may have **profound effects** on chemical and enzyme kinetics, and may be produced from biochemical energy or from radio frequency, microwave, or terahertz radiation. Pokorný's observed 8.085-MHz microtubulin resonance is identified as a possible candidate, with microwave reactors (green chemistry) and terahertz medicine appearing as other feasible sources. **Proc Natl Acad Sci U S A. 2009 March 17; 106(11Published online 2009 February 26. doi: 10.1073/pnas.0806273106PMCID: PMC2657444 Chemistry, Biophysics and Computational Biology Weak, strong, and coherent regimes of Fröhlich condensation and their applications to terahertz medicine and quantum consciousness Jeffrey R. Reimers et al**

INTRODUCTION—VARIABLES USED

Introduction:

Quantum entanglement **is a** form of quantum superposition. When a measurement is made and **it causes** one member of such a pair to take on a definite value (e.g., clockwise spin), the other member of this entangled pair will at any subsequent time be found to have taken the appropriately correlated value (e.g., counterclockwise spin). Thus, there is a correlation between the results of measurements performed on entangled pairs, and this correlation is observed even though the entangled pair may have been separated by arbitrarily large distances. In quantum entanglement, part of the transfer happens instantaneously. Repeated experiments have verified that this works even when the measurements are performed more quickly than light could travel between the sites of measurement: there is no slower-than-light influence that can pass between the entangled particles. Recent experiments have shown that this transfer occurs at least 10,000 times faster than the speed of light, which does not remove the possibility of it being an instantaneous phenomenon, but only sets a lower limit. This behavior is consistent with quantum-mechanical theory, has been demonstrated experimentally, and it is an area of extremely active research by the physics community. However there is some heated debate about whether a possible classical underlying mechanism could

explain why this correlation occurs instantaneously even when the separation distance is large. The difference in opinion derives from espousal of various interpretations of quantum mechanics.

Correlation phenomena observed by CMS may shed light on exotic state of matter Julia Velkovska and Wei Li 2013-05-31 (Figures and a few paragraphs are deleted attributable to spatial constraints)

(1) The "ridge" was first observed in nucleus-nucleus collisions at the Relativistic Heavy Ion Collider and has a long history of competing theoretical explanations. Presently it is understood **to be the (e) consequence of the** explosion of a "droplet" of quark-gluon plasma (QGP) in which thousands of particles "flow" in a correlated fashion almost without any internal friction. However, a ridge was also observed in proton-proton (pp) and proton-nucleus collisions, where a QGP-like matter was not expected to be formed. In systems like pp and pPb the initial collision zone is about an order of magnitude **smaller (eb) in size** than in the lead-lead (PbPb) system and it is therefore difficult to reach a high-enough energy density required (e) to melt colliding nucleons into a soup of quarks and gluons.

(2) Yet, in rare cases, when the pp and pPb collisions produce a huge number of particles, these particles appear to be correlated. The question is, do these correlations in pp and pPb collisions share a common physical origin as in PbPb interactions, are **they(e) caused by** the internal gluon structure of the protons (as proposed by the "colour-glass condensate" model), or perhaps are they caused by both?

(3) In the 2013 run, on some occasions as many as 420 charged particles were observed coming (e) out of a single pPb collision in the CMS detector (see Image 1 for an event display of such a collision). This recorded data sample contained a relatively few, extraordinary cases where the multiplicity of particles produced is comparable (=) to that in an average PbPb collision.

(4) And sure enough, the two-particle correlations also look very similar. Direct comparison of two-particle correlations between PbPb and pPb collisions **by (e&eb) selecting** the same particle-multiplicity ranges.

(5) The new data set also makes it possible to analyse the long-range correlations in greater detail. A technique similar to the approach used to study the cosmic microwave background (CMB) spectrum was deployed to investigate the correlations.

(6) For the CMB data analysis, microscopic temperature fluctuations reveal the primordial inhomogeneity in the Universe immediately after the big bang, which eventually **gave rise to** galaxies.

(7) In each lead-lead collision a different microcosm is produced with its own set of fluctuations of hot spots in the collision zone that then propagate as ripples which can be uncovered by the CMS detector.

Entanglement in quantum critical phenomena G. Vidal et al

(1) Quantum phase transitions occur at zero temperature **and involve** the appearance of long-range correlations.

(2) These correlations are not due to thermal fluctuations but **due (e) to the** intricate structure of a strongly entangled ground state of the system. microscopic computation of the scaling properties of the ground-state entanglement in several 1D spin chain models both near and at the quantum critical regimes and quantify entanglement **by (e) using the** entropy of the ground state when the system is traced down to L spins.

(3) Entropy is seen to scale logarithmically with L, with a coefficient that corresponds to the central charge associated to the conformal theory **that describes** the universal properties of the quantum phase transition.

(4) Entanglement, a key concept of quantum information science, **obeys (eb)** universal scaling laws as dictated by the representations of the conformal group and its classification motivated by string theory.
(5) Entanglement, a key concept of quantum information science, **obeys (eb)** universal scaling laws as dictated by the representations of the conformal group and its classification motivated by string theory **unveils (eb) a** monotonicity law for ground-state entanglement along the renormalization group flow.
(6) Identification of majorization rule possibly associated to conformal invariance **and (e&eb) application to** present results to interpret the breakdown of density matrix renormalization group techniques near a critical point.**10.1103/PhysRevLett.90.227902arXiv:quant-ph/0211074**

Penrose-Hameroff orchestrated objective-reduction proposal for human consciousness is not biologically feasible Laura K. McKemmish et al

(1) Penrose and Hameroff have argued that the conventional models of a brain function **based on** neural networks alone cannot account for human consciousness, claiming that quantum-computation elements are also **required (e).**
(2) Specifically, in their Orchestrated Objective Reduction (Orch OR) model [R. Penrose and S. R. Hameroff J. Conscious. Stud. 2 99 (1995)], it is postulated that microtubules **act (e&eb) as** quantum processing units, with individual tubulin dimers forming the computational elements.
(3) Tubulin is able to switch between alternative conformational states in a coherent manner is an **essential functional requisite (e),** and that this process be rapid on the physiological time scale.
(4) Biological feasibility of the Orch OR proposal examined in light of recent experimental studies on microtubule assembly and dynamics **shows (eb) that the** tubulins do not possess essential properties required for the Orch OR proposal, as originally proposed, to hold. long-lived coherent motions in biological systems, is a feature critical to Orch OR, **which (eb) shows** that no reformation of the proposal based on known physical paradigms **could lead to** quantum computing within microtubules. Hence, the Orch OR model is not a feasible explanation of the origin of consciousness.© 2009 The American Physical SocietyURL:http://link.aps.org/doi/10.1103/PhysRevE.80.021912DOI: 10.1103/PhysRevE.80.021912

Quantum entanglement **is a** form of quantum superposition. When a measurement is made and **it causes** one member of such a pair to take on a definite value (e.g., clockwise spin), the other member of this entangled pair will at any subsequent time be found to have taken the appropriately correlated value (e.g., counterclockwise spin). Thus, there is a correlation between the results of measurements performed on entangled pairs, and this correlation is observed even though the entangled pair may have been separated by arbitrarily large distances. In quantum entanglement, part of the transfer happens instantaneously. Repeated experiments have verified that this works even when the measurements are performed more quickly than light could travel between the sites of measurement: there is no slower-than-light influence that can pass between the entangled particles. Recent experiments have shown that this transfer occurs at least 10,000 times faster than the speed of light, which does not remove the possibility of it being an instantaneous phenomenon, but only sets a lower limit. This behavior is consistent with quantum-mechanical theory, has been demonstrated experimentally, and it is an area of extremely active research by the physics community. However there is some heated debate about whether a possible classical underlying mechanism could explain why this correlation occurs instantaneously even when the separation distance is large. The difference in opinion derives from espousal of various interpretations of quantum mechanics.

he quantum mind or quantum consciousness hypothesis proposes that classical mechanics cannot explain

consciousness, while quantum mechanical phenomena, such as quantum entanglement and superposition, may play an important part in the brain's function, and could form the basis of an explanation of consciousness. It is not one theory, but a collection of distinct ideas described below.

A few theoretical physicists have argued that classical physics is intrinsically incapable of explaining the holistic aspects of consciousness, whereas quantum mechanics can. The idea that quantum theory has something to do with the workings of the mind goes back to Niels Bohr. Whereas Bohr **wrote in generalities on this matter**, Eugene Wigner assumed that the wave function **collapses due to its** interaction with consciousness. However, modern physicists and philosophers consider the arguments for an important role of quantum phenomena to be unconvincing. Physicist Victor Stenger characterized quantum consciousness as a "myth" having "no scientific basis" that "should take its place along with gods, unicorns and dragons." The philosopher David Chalmers has argued against quantum consciousness. He has discussed how quantum mechanics may relate to dualistic consciousness. Indeed, Chalmers is skeptical of the ability of any new physics to resolve the hard problem of consciousness.

De Broglie–Bohm theory is based on the following postulates:

There is a configuration q of the universe, described by coordinates q^k, which is an element of the configuration space Q. The configuration space is different for different versions of pilot wave theory. For example, this may be the space of positions $\mathbf{Q}_k$ of N particles, or, in case of field theory, the space of field configurations $\phi(x)$. The configuration evolves (for spin=0) according to the guiding equation

$$m_k \frac{dq^k}{dt}(t) = \hbar \nabla_k \operatorname{Im} \ln \psi(q,t) = \hbar \operatorname{Im} \left(\frac{\nabla_k \psi}{\psi} \right)(q,t)$$

.

Here, $\psi(q,t)$ is the standard complex-valued wavefunction known from quantum theory, which evolves according to Schrödinger's equation

$$i\hbar \frac{\partial}{\partial t}\psi(q,t) = -\sum_{i=1}^{N} \frac{\hbar^2}{2m_i} \nabla_i^2 \psi(q,t) + V(q)\psi(q,t)$$

NOTATION

Module One

De Broglie–Bohm theory is based on the following postulates

$$m_k \frac{dq^k}{dt}(t) = \hbar \nabla_k \operatorname{Im} \ln \psi(q,t) = \hbar \operatorname{Im} \left(\frac{\nabla_k \psi}{\psi} \right)(q,t)$$

G_{13} : Category one of LHS of the above equation

G_{14} : Category two of LHS of the above equation

G_{15} : Category three onLHS of the above equation

T_{13} : Category one of RHS of the above equation

T_{14} : Category two of RHS of the above equation

T_{15} : Category three of RHS of the above equation

Module Two

De Broglie–Bohm theory is based on the following postulates

$$i\hbar\frac{\partial}{\partial t}\psi(q,t) = -\sum_{i=1}^{N}\frac{\hbar^2}{2m_i}\nabla_i^2\psi(q,t) + V(q)\psi(q,t)$$

G_{16} : Category one of LHS of the above equation

G_{17} : Category two of LHS of the above equation

G_{18} : Category three of LHS of the above equation

T_{16} : Category one of RHS of the above equation

T_{17} : Category two of RHS of the above equation

T_{18} : Category three of RHS of the above equation

Module three

The "ridge" was first observed in nucleus-nucleus collisions at the Relativistic Heavy Ion Collider and has a long history of competing theoretical explanations with the existential and extant understanding to **be the (e) consequence of the** explosion of a "droplet" of quark-gluon plasma (QGP) in which thousands of particles "flow" in a correlated fashion almost without any internal friction.

However, a ridge was also observed in proton-proton (pp) and proton-nucleus collisions, where a QGP-like matter was not expected to be formed. In systems like pp and pPb the initial collision zone is about an order of magnitude **smaller (eb) in size** than in the lead-lead (PbPb) system and it is therefore difficult to reach a high-enough energy density required (e) to melt colliding nucleons into a soup of quarks and gluons.

G_{20} : Category one of explosion of a "droplet" of quark-gluon plasma (QGP) in which thousands of particles "flow" in a correlated fashion almost without any internal friction

G_{21} : Category two of explosion of a "droplet" of quark-gluon plasma (QGP) in which thousands of particles "flow" in a correlated fashion almost without any internal friction

G_{22} : Category three of explosion of a "droplet" of quark-gluon plasma (QGP) in which thousands of particles "flow" in a correlated fashion almost without any internal friction

T_{20} : Category one of "ridge" was first observed in nucleus-nucleus collisions at the Relativistic Heavy Ion Collider and has a long history of competing theoretical explanations with the existential and extant understanding

T_{21} : Category two of "ridge" was first observed in nucleus-nucleus collisions at the Relativistic Heavy Ion Collider and has a long history of competing theoretical explanations with the existential and extant

understanding

T_{22} : Category three of "ridge" was first observed in nucleus-nucleus collisions at the Relativistic Heavy Ion Collider and has a long history of competing theoretical explanations with the existential and extant understanding

Module four

in rare cases, when the pp and pPb collisions produce a huge number of particles, these particles appear to be correlated. The question is, do these correlations in pp and pPb collisions share a common physical origin as in PbPb interactions, are **they(e) caused by** the internal gluon structure of the protons (as proposed by the "colour-glass condensate" model), or perhaps are they caused by both?

G_{24} : Category one of internal gluon structure of the protons (as proposed by the "colour-glass condensate" model), or perhaps are they caused by both

G_{25} : Category two of internal gluon structure of the protons (as proposed by the "colour-glass condensate" model), or perhaps are they caused by both

G_{26} : Category three of internal gluon structure of the protons (as proposed by the "colour-glass condensate" model), or perhaps are they caused by both

T_{24} : Category one of correlations in pp and pPb collisions share a common physical origin as in PbPb interactions,

T_{25} : Category two of correlations in pp and pPb collisions share a common physical origin as in PbPb interactions,

T_{26} : Category three of correlations in pp and pPb collisions share a common physical origin as in PbPb interactions,

Module five

In the 2013 run, on some occasions as many as 420 charged particles were observed coming (e) out of a single pPb collision in the CMS detector (see Image 1 for an event display of such a collision),the recorded data sample contained a relatively few, extraordinary cases where the multiplicity of particles produced is comparable to that in an average PbPb collision

G_{28} : Category one of average PbPb collision

G_{29} : Category two of average PbPb collision

G_{30} : Category three of average PbPb collision

T_{28} : Category one of as many as 420 charged particles were observed coming (e) out of a single pPb collision in the CMS detector (see Image 1 for an event display of such a collision),the recorded data sample contained a relatively few, extraordinary cases where the multiplicity of particles produced is comparable

T_{29} : Category two of as many as 420 charged particles were observed coming (e) out of a single pPb collision in the CMS detector (see Image 1 for an event display of such a collision),the recorded data sample contained a relatively few, extraordinary cases where the multiplicity of particles produced is comparable

T_{30} : Category three of as many as 420 charged particles were observed coming (e) out of a single pPb collision in the CMS detector (see Image 1 for an event display of such a collision),the recorded data sample contained a relatively few, extraordinary cases where the multiplicity of particles produced is comparable

Module six

For the CMB data analysis, microscopic temperature fluctuations reveal the primordial inhomogeneity in the Universe immediately after the big bang, which eventually **gave rise to** galaxies.

G_{32} : Category one of CMB data analysis, microscopic temperature fluctuations reveal the primordial inhomogeneity in the Universe immediately after the big bang

G_{33} : Category two of CMB data analysis, microscopic temperature fluctuations reveal the primordial inhomogeneity in the Universe immediately after the big bang

G_{34} : Category three of CMB data analysis, microscopic temperature fluctuations reveal the primordial inhomogeneity in the Universe immediately after the big bang

T_{32} : Category one of galaxies.

T_{33} : Category two of galaxies.

T_{34} : Category three of galaxies.

Module seven

Biological feasibility of the Orch OR proposal examined in light of recent experimental studies on microtubule assembly and dynamics **shows (eb) that the** tubulins do not possess essential properties required for the Orch OR proposal, as originally proposed, to hold. long-lived coherent motions in biological systems, is a feature critical to Orch OR, **which (eb) shows** that no reformation of the proposal based on known physical paradigms **could lead to** quantum computing within microtubules. Hence, the Orch OR model is not a feasible explanation of the origin of consciousness.© 2009 The American Physical SocietyURL:http://link.aps.org/doi/10.1103/PhysRevE.80.021912DOI: 10.1103/PhysRevE.80.021912

G_{36} : Category one of Biological feasibility of the Orch OR proposal examined in light of recent experimental studies on microtubule assembly and dynamics **shows (eb) that the** tubulins do not possess essential properties required for the Orch OR proposal, as originally proposed, to hold. long-lived coherent motions in biological systems, is a feature critical to Orch OR

G_{37} : Category two of Biological feasibility of the Orch OR proposal examined in light of recent experimental studies on microtubule assembly and dynamics **shows (eb) that the** tubulins do not possess essential properties required for the Orch OR proposal, as originally proposed, to hold. long-lived coherent motions in biological systems, is a feature critical to Orch OR

G_{38} : Category three of Biological feasibility of the Orch OR proposal examined in light of recent experimental studies on microtubule assembly and dynamics **shows (eb) that the** tubulins do not possess essential properties required for the Orch OR proposal, as originally proposed, to hold. long-lived coherent motions in biological systems, is a feature critical to Orch OR

T_{36} : Category one of quantum computing within microtubules. Hence, the Orch OR model is not a feasible explanation of the origin of consciousness

T_{37} : Category two of quantum computing within microtubules. Hence, the Orch OR model is not a feasible explanation of the origin of consciousness

T_{38} : Category three of quantum computing within microtubules. Hence, the Orch OR model is not a feasible explanation of the origin of consciousness

Module eight

These correlations are not due to thermal fluctuations but **due (e) to the** intricate structure of a strongly entangled ground state of the system. microscopic computation of the scaling properties of the ground-state entanglement in several 1D spin chain models both near and at the quantum critical regimes and quantify entanglement **by (e) using the** entropy of the ground state when the system is traced down to L spins.

G_{40} : Category one of entropy of the ground state when the system is traced down to L spins

G_{41} : Category two of entropy of the ground state when the system is traced down to L spins

G_{42} : Category three of entropy of the ground state when the system is traced down to L spins

T_{40} : Category one of correlations are not due to thermal fluctuations but **due (e) to the** intricate structure of a strongly entangled ground state of the system. microscopic computation of the scaling properties of the ground-state entanglement in several 1D spin chain models both near and at the quantum critical regimes and quantify entanglement

T_{41} : Category two of correlations are not due to thermal fluctuations but **due (e) to the** intricate structure of a strongly entangled ground state of the system. microscopic computation of the scaling properties of the ground-state entanglement in several 1D spin chain models both near and at the quantum critical regimes and quantify entanglement

T_{42} : Category three of correlations are not due to thermal fluctuations but **due (e) to the** intricate structure of a strongly entangled ground state of the system. microscopic computation of the scaling properties of the ground-state entanglement in several 1D spin chain models both near and at the quantum critical regimes and quantify entanglement

Module Nine

Entanglement, a key concept of quantum information science, **obeys (eb)** universal scaling laws as dictated by the representations of the conformal group and its classification motivated by string theory **unveils (eb) a** monotonicity law for ground-state entanglement along the renormalization group flow.

G_{44} : Category one of Entanglement, a key concept of quantum information science

G_{45} : Category two of Entanglement, a key concept of quantum information science

G_{46} : Category three of Entanglement, a key concept of quantum information science

T_{44} : Category one of universal scaling laws as dictated by the representations of the conformal group and its classification motivated by string theory **unveils (eb) a** monotonicity law for ground-state entanglement along the renormalization group flow

T_{45} : Category two of universal scaling laws as dictated by the representations of the conformal group and its classification motivated by string theory **unveils (eb) a** monotonicity law for ground-state entanglement along the renormalization group flow

T_{46} : Category three of universal scaling laws as dictated by the representations of the conformal group and its classification motivated by string theory **unveils (eb) a** monotonicity law for ground-state entanglement along the renormalization group flow

SECTION SEVENTY FIVE

Quantum Phase transition: minima maxima sunt: The smallest things are most important

Abstract: Classical phase transitions occur when a physical system reaches a state below a critical temperature characterized by a macroscopic order. Quantum phase transitions **occur at** absolute zero; they are induced by the change of an external parameter or coupling constant, and are driven by fluctuations. Examples include transitions in quantum Hall systems, localization in Si-MOSFETs (metal oxide silicon

field-effect transistors; and the superconductor-insulator transition in two-dimensional systems. Both classical and quantum critical points **are governed by a** diverging correlation length, although quantum systems possess additional correlations that do not have a classical counterpart. This phenomenon, known as entanglement is the resource **that enables** quantum computation and communication. The role of the entanglement at a phase transition is not captured by statistical mechanics – a complete classification of the critical many-body state requires the introduction of concepts from quantum information theory. Theory of critical phenomena **with** quantum information is correlated and consummated and consolidated **by exploring the** entangling resources of a system close to its quantum critical point. Class of one-dimensional magnetic systems, has entanglement **showing** scaling behaviour in the vicinity of the transition. Entanglement near a quantum phase transition by analyzing the properties of the concurrence for a class of exactly solvable models in one dimension to find that entanglement **can be (e&eb) classified** in the framework of scaling theory. Difference between classical correlations and the non-local quantum correlation, entanglement: the correlation length diverges at the phase transition, whereas entanglement in general **remains (eb)** short ranged. Nature 416, 608 (2002) DOI: 10.1038/416608a arXiv: quant-ph/0202029 **Scaling of Entanglement close to a Quantum Phase Transitions Osterloh** In 1968, Fröhlich showed that a driven set of oscillators **can condense** with nearly all of the supplied energy activating the vibrational mode of lowest frequency. This is a remarkable property usually compared with Bose–Einstein condensation, superconductivity, lasing, and other unique phenomena involving macroscopic quantum coherence. However, despite intense research, no unambiguous example has been documented. **Jeffrey R. Reimers et al** determine the most likely experimental signatures of Fröhlich condensation **and show that** they are significant features remote from the extraordinary properties normally envisaged. Fröhlich condensates are classified into 3 types: weak condensates in which profound effects on chemical kinetics are possible, strong condensates in which an extremely large amount of energy **is channeled** into 1 vibrational mode, and coherent condensates in which this energy is placed in a single quantum state. Coherent condensates **are shown to** involve extremely large energies, to not be produced by the Wu–Austin dynamical Hamiltonian **that provides** the simplest depiction of Fröhlich condensates formed using mechanically supplied energy, and to be extremely fragile. They are inaccessible in a biological environment. Hence the Penrose–Hameroff orchestrated objective-reduction model and related theories for cognitive function **that embody coherent** Fröhlich condensation as an essential element are untenable. Weak condensates, however, may have **profound effects** on chemical and enzyme kinetics, and may be produced from biochemical energy or from radio frequency, microwave, or terahertz radiation. Pokorný's observed 8.085-MHz microtubulin resonance is identified as a possible candidate, with microwave reactors (green chemistry) and terahertz medicine appearing as other feasible sources. **Proc Natl Acad Sci U S A. 2009 March 17; 106(11Published online 2009 February 26. doi: 10.1073/pnas.0806273106PMCID: PMC2657444 Chemistry, Biophysics and Computational Biology Weak, strong, and coherent regimes of Fröhlich condensation and their applications to terahertz medicine and quantum consciousness Jeffrey R. Reimers et al**

INTRODUCTION—VARIABLES USED

Introduction:

Quantum entanglement **is a** form of quantum superposition. When a measurement is made and **it causes** one member of such a pair to take on a definite value (e.g., clockwise spin), the other member of this entangled pair will at any subsequent time be found to have taken the appropriately correlated value (e.g., counterclockwise spin). Thus, there is a correlation between the results of measurements performed on entangled pairs, and this correlation is observed even though the entangled pair may have been separated by arbitrarily large distances. In quantum entanglement, part of the transfer happens instantaneously. Repeated experiments have verified that this works even when the measurements are performed more quickly than

light could travel between the sites of measurement: there is no slower-than-light influence that can pass between the entangled particles. Recent experiments have shown that this transfer occurs at least 10,000 times faster than the speed of light, which does not remove the possibility of it being an instantaneous phenomenon, but only sets a lower limit. This behavior is consistent with quantum-mechanical theory, has been demonstrated experimentally, and it is an area of extremely active research by the physics community. However there is some heated debate about whether a possible classical underlying mechanism could explain why this correlation occurs instantaneously even when the separation distance is large. The difference in opinion derives from espousal of various interpretations of quantum mechanics.

Correlation phenomena observed by CMS may shed light on exotic state of matter Julia Velkovska and Wei Li 2013-05-31 (Figures and a few paragraphs are deleted attributable to spatial constraints)

(8) The "ridge" was first observed in nucleus-nucleus collisions at the Relativistic Heavy Ion Collider and has a long history of competing theoretical explanations. Presently it is understood **to be the (e) consequence of** the explosion of a "droplet" of quark-gluon plasma (QGP) in which thousands of particles "flow" in a correlated fashion almost without any internal friction.

(9) However, a ridge was also observed in proton-proton (pp) and proton-nucleus collisions, where a QGP-like matter was not expected to be formed. In systems like pp and pPb the initial collision zone is about an order of magnitude **smaller (eb)** in size than in the lead-lead (PbPb) system and it is therefore difficult to reach a high-enough energy density **required (e) to** melt colliding nucleons into a soup of quarks and gluons.

(10) Yet, in rare cases, when the pp and pPb collisions **produce a** huge number of particles, these particles appear to be correlated. The question is, do these correlations in pp and pPb collisions share a common physical origin as in PbPb interactions, **are they(e) caused by** the internal gluon structure of the protons (as proposed by the "colour-glass condensate" model), or perhaps are they caused by both?

(11) In the 2013 run, on some occasions as many as 420 charged particles were observed coming (e) out of a single pPb collision in the CMS detector (Image deleted due to spatial constraints). This recorded data sample contained a relatively few, extraordinary cases where the multiplicity of particles produced is **comparable to** that in an average PbPb collision.And sure enough, the two-particle correlations also look very similar. Image 2 shows a direct comparison of two-particle correlations between PbPb and pPb collisions **by (e&eb) selecting** the same particle-multiplicity ranges.

(12) The new data set also makes it possible to analyse the long-range correlations in greater detail. A technique similar to the approach used to study the cosmic microwave background (CMB) spectrum was deployed to investigate the correlations.

(13) For the CMB data analysis, microscopic temperature fluctuations **reveal the** primordial inhomogeneity in the Universe immediately after the big bang, which eventually gave rise to galaxies.

(14) In each lead-lead collision a different microcosm is produced with its own set of fluctuations of hot spots in the collision zone that then propagate as ripples which can be uncovered by the CMS detector.

Entanglement in quantum critical phenomena G. Vidal et al

(7) Quantum phase transitions occur at zero temperature **and involve** the appearance of long-range correlations. These correlations are not due to thermal fluctuations but **due (e) to the** intricate structure of a strongly entangled ground state of the system.

(8) microscopic computation of the scaling properties of the ground-state entanglement in several 1D

spin chain models both near and at the quantum critical regimes and quantification of entanglement is done **by (e) using the** entropy of the ground state when the system is traced down to L spins.

(9) Entropy is seen to scale logarithmically with L, with a coefficient that corresponds to the central charge associated to the conformal theory **that describes** the universal properties of the quantum phase transition.

(10) Entanglement, a key concept of quantum information science, **obeys (eb)** universal scaling laws as dictated by the representations of the conformal group and its classification motivated by string theory.

(11) Entanglement, a key concept of quantum information science, **obeys (eb)** universal scaling laws as dictated by the representations of the conformal group and its classification motivated by string theory **unveils (eb) a** monotonicity law for ground-state entanglement along the renormalization group flow.

(12) Identification of majorization rule possibly associated to conformal invariance **and (e&eb) application to** present results to interpret the breakdown of density matrix renormalization group techniques near a critical point.**10.1103/PhysRevLett.90.227902arXiv:quant-ph/0211074**

Penrose-Hameroff orchestrated objective-reduction proposal for human consciousness is not biologically feasible Laura K. McKemmish et al

(5) Penrose and Hameroff have argued that the conventional models of a brain function **based on** neural networks alone cannot account for human consciousness, claiming that quantum-computation elements are also **required (e).**

(6) Specifically, in their Orchestrated Objective Reduction (Orch OR) model [R. Penrose and S. R. Hameroff J. Conscious. Stud. 2 99 (1995)], it is postulated that microtubules **act (e&eb) as** quantum processing units, with individual tubulin dimers forming the computational elements.

(7) Tubulin is able to switch between alternative conformational states in a coherent manner is an **essential functional requisite (e),** and that this process be rapid on the physiological time scale.

(8) Biological feasibility of the Orch OR proposal examined in light of recent experimental studies on microtubule assembly and dynamics **shows (eb) that the** tubulins do not possess essential properties required for the Orch OR proposal, as originally proposed, to hold.

(9) long-lived coherent motions in biological systems, is a feature critical to Orch OR, **which (eb) shows** that no reformation of the proposal based on known physical paradigms **could lead to** quantum computing within microtubules. Hence, the Orch OR model is not a feasible explanation of the origin of consciousness.© 2009 The American Physical SocietyURL:http://link.aps.org/doi/10.1103/PhysRevE.80.021912DOI: 10.1103/PhysRevE.80.021912

Quantum entanglement **is a** form of quantum superposition. When a measurement is made and **it causes** one member of such a pair to take on a definite value (e.g., clockwise spin), the other member of this entangled pair will at any subsequent time be found to have taken the appropriately correlated value (e.g., counterclockwise spin). Thus, there is a correlation between the results of measurements performed on entangled pairs, and this correlation is observed even though the entangled pair may have been separated by arbitrarily large distances. In quantum entanglement, part of the transfer happens instantaneously. Repeated experiments have verified that this works even when the measurements are performed more quickly than light could travel between the sites of measurement: there is no slower-than-light influence that can pass between the entangled particles. Recent experiments have shown that this transfer occurs at least 10,000 times faster than the speed of light, which does not remove the possibility of it being an instantaneous phenomenon, but only sets a lower limit. This behavior is consistent with quantum-mechanical theory, has

been demonstrated experimentally, and it is an area of extremely active research by the physics community. However there is some heated debate about whether a possible classical underlying mechanism could explain why this correlation occurs instantaneously even when the separation distance is large. The difference in opinion derives from espousal of various interpretations of quantum mechanics.

The quantum mind or quantum consciousness hypothesis proposes that classical mechanics cannot explain consciousness, while quantum mechanical phenomena, such as quantum entanglement and superposition, may play an important part in the brain's function, and could form the basis of an explanation of consciousness. It is not one theory, but a collection of distinct ideas described below.

A few theoretical physicists have argued that classical physics is intrinsically incapable of explaining the holistic aspects of consciousness, whereas quantum mechanics can. The idea that quantum theory has something to do with the workings of the mind goes back to Niels Bohr. Whereas Bohr **wrote in generalities on this matter**, Eugene Wigner assumed that the wave function **collapses due to its** interaction with consciousness. However, modern physicists and philosophers consider the arguments for an important role of quantum phenomena to be unconvincing. Physicist Victor Stenger characterized quantum consciousness as a "myth" having "no scientific basis" that "should take its place along with gods, unicorns and dragons." The philosopher David Chalmers has argued against quantum consciousness. He has discussed how quantum mechanics may relate to dualistic consciousness. Indeed, Chalmers is skeptical of the ability of any new physics to resolve the hard problem of consciousness.

De Broglie–Bohm theory is based on the following postulates:

There is a configuration q of the universe, described by coordinates q^k, which is an element of the configuration space Q. The configuration space is different for different versions of pilot wave theory. For example, this may be the space of positions $\mathbf{Q}_k$ of N particles, or, in case of field theory, the space of field configurations $\phi(x)$. The configuration evolves (for spin=0) according to the guiding equation

$$m_k \frac{dq^k}{dt}(t) = \hbar \nabla_k \operatorname{Im} \ln \psi(q,t) = \hbar \operatorname{Im} \left(\frac{\nabla_k \psi}{\psi} \right)(q,t)$$

.

Here, $\psi(q,t)$ is the standard complex-valued wavefunction known from quantum theory, which evolves according to Schrödinger's equation

$$i\hbar \frac{\partial}{\partial t}\psi(q,t) = -\sum_{i=1}^{N} \frac{\hbar^2}{2m_i} \nabla_i^2 \psi(q,t) + V(q)\psi(q,t)$$

NOTATION

Module One

The “ridge” was first observed in nucleus-nucleus collisions at the Relativistic Heavy Ion Collider and has a long history of competing theoretical explanations understood **to be the (e) consequence of** the explosion of a “droplet” of quark-gluon plasma (QGP) in which thousands of particles “flow” in a correlated fashion almost without any internal friction

G_{13} : Category one of explosion of a "droplet" of quark-gluon plasma (QGP) in which thousands of particles "flow" in a correlated fashion almost without any internal friction

G_{14} : Category two of explosion of a "droplet" of quark-gluon plasma (QGP) in which thousands of particles "flow" in a correlated fashion almost without any internal friction

G_{15} : Category three of explosion of a "droplet" of quark-gluon plasma (QGP) in which thousands of particles "flow" in a correlated fashion almost without any internal friction

T_{13} : Category one of "ridge" was first observed in nucleus-nucleus collisions at the Relativistic Heavy Ion Collider and has a long history of competing theoretical explanations

T_{14} : Category two of "ridge" was first observed in nucleus-nucleus collisions at the Relativistic Heavy Ion Collider and has a long history of competing theoretical explanations

T_{15} : Category three of "ridge" was first observed in nucleus-nucleus collisions at the Relativistic Heavy Ion Collider and has a long history of competing theoretical explanations

Module two

However, a ridge was also observed in proton-proton (pp) and proton-nucleus collisions, where a QGP-like matter was not expected to be form and in systems like pp and pPb the initial collision zone is about an order of magnitude **smaller (eb)** in size than in the lead-lead (PbPb) system and it is therefore difficult to reach a high-enough energy density **required (e) to** melt colliding nucleons into a soup of quarks and gluons

G_{16} : Category one of melt colliding nucleons into a soup of quarks and gluons

G_{17} : Category two of melt colliding nucleons into a soup of quarks and gluons

G_{18} : Category three of melt colliding nucleons into a soup of quarks and gluons

T_{16} : Category one of ridge was also observed in proton-proton (pp) and proton-nucleus collisions, where a QGP-like matter was not expected to be form and in systems like pp and pPb the initial collision zone is about an order of magnitude **smaller (eb)** in size than in the lead-lead (PbPb) system and it is therefore difficult to reach a high-enough energy density

T_{17} : Category two of ridge was also observed in proton-proton (pp) and proton-nucleus collisions, where a QGP-like matter was not expected to be form and in systems like pp and pPb the initial collision zone is about an order of magnitude **smaller (eb)** in size than in the lead-lead (PbPb) system and it is therefore difficult to reach a high-enough energy density

T_{18} : Category three of ridge was also observed in proton-proton (pp) and proton-nucleus collisions, where a QGP-like matter was not expected to be form and in systems like pp and pPb the initial collision zone is about an order of magnitude **smaller (eb)** in size than in the lead-lead (PbPb) system and it is therefore difficult to reach a high-enough energy density

Module three(both the cases are discussed)

Yet, in rare cases, when the pp and pPb collisions **produce a** huge number of particles, these particles appear to be correlated, such correlations in pp and pPb collisions share a common physical origin as in PbPb interactions, **are they (e) caused by** the internal gluon structure of the protons (as proposed by the

"colour-glass condensate" model), or perhaps are they caused by both?

G_{20} : Category one of internal gluon structure of the protons (as proposed by the "colour-glass condensate" model); internal gluon structure of the protons and strongly correlated particles

G_{21} : Category two of internal gluon structure of the protons (as proposed by the "colour-glass condensate" model); internal gluon structure of the protons and strongly correlated particles

G_{22} : Category three of internal gluon structure of the protons (as proposed by the "colour-glass condensate" model); internal gluon structure of the protons and strongly correlated particles

T_{20} : Category one of pp and pPb collisions **produce a** huge number of particles, these particles appear to be correlated, such correlations in pp and pPb collisions share a common physical origin as in PbPb interactions,

T_{21} : Category two of pp and pPb collisions **produce a** huge number of particles, these particles appear to be correlated, such correlations in pp and pPb collisions share a common physical origin as in PbPb interactions,

T_{22} : Category three of pp and pPb collisions **produce a** huge number of particles, these particles appear to be correlated, such correlations in pp and pPb collisions share a common physical origin as in PbPb interactions,

Module four

In the 2013 run, on some occasions as many as 420 charged particles were observed coming (e) out of a single pPb collision in the CMS detector (Image deleted due to spatial constraints). This recorded data sample contained a relatively few, extraordinary cases where the multiplicity of particles produced is **comparable to** that in an average PbPb collision. And sure enough, the two-particle correlations also look very similar. Image 2 (deleted) shows a direct comparison of two-particle correlations between PbPb and pPb collisions **by (e&eb) selecting** the same particle-multiplicity ranges. For the CMB data analysis, microscopic temperature fluctuations **reveal the** primordial inhomogeneity in the Universe immediately after the big bang, which eventually gave rise to galaxies.

G_{24} : Category one of CMB data analysis, microscopic temperature fluctuations

G_{25} : Category two of CMB data analysis, microscopic temperature fluctuations

G_{26} : Category three of CMB data analysis, microscopic temperature fluctuations

T_{24} : Category one of primordial inhomogeneity in the Universe immediately after the big bang, which eventually gave rise to galaxies.

T_{25} : Category two of primordial inhomogeneity in the Universe immediately after the big bang, which eventually gave rise to galaxies.

T_{26} : Category three of primordial inhomogeneity in the Universe immediately after the big bang, which eventually gave rise to galaxies.

Module five

Quantum phase transitions occur at zero temperature **and involve** the appearance of long-range correlations

which are not due to thermal fluctuations but **due (e) to the** intricate structure of a strongly entangled ground state of the system

G_{28} : Category one of intricate structure of a strongly entangled ground state of the system

G_{29} : Category two of intricate structure of a strongly entangled ground state of the system

G_{30} : Category three of intricate structure of a strongly entangled ground state of the system

T_{28} : Category one of Quantum phase transitions occur at zero temperature **and involve** the appearance of long-range correlations which are not due to thermal fluctuations

T_{29} : Category two of Quantum phase transitions occur at zero temperature **and involve** the appearance of long-range correlations which are not due to thermal fluctuations

T_{30} : Category three of Quantum phase transitions occur at zero temperature **and involve** the appearance of long-range correlations which are not due to thermal fluctuations

Module six

microscopic computation of the scaling properties of the ground-state entanglement in several 1D spin chain models both near and at the quantum critical regimes and quantification of entanglement is done **by (e) using the** entropy of the ground state when the system is traced down to L spins.

G_{32} : Category one of entropy of the ground state when the system is traced down to L spins

G_{33} : Category two of entropy of the ground state when the system is traced down to L spins

G_{34} : Category three of entropy of the ground state when the system is traced down to L spins

T_{32} : Category one of microscopic computation of the scaling properties of the ground-state entanglement in several 1D spin chain models both near and at the quantum critical regimes and quantification of entanglement

T_{33} : Category two of microscopic computation of the scaling properties of the ground-state entanglement in several 1D spin chain models both near and at the quantum critical regimes and quantification of entanglement

T_{34} : Category three of microscopic computation of the scaling properties of the ground-state entanglement in several 1D spin chain models both near and at the quantum critical regimes and quantification of entanglement

Module seven

Entropy is seen to scale logarithmically with L, with a coefficient that corresponds to the central charge associated to the conformal theory **that describes** the universal properties of the quantum phase transition

G_{36} : Category one of Entropy is seen to scale logarithmically with L, with a coefficient that corresponds to the central charge associated to the conformal theory

G_{37} : Category two of Entropy is seen to scale logarithmically with L, with a coefficient that corresponds to the central charge associated to the conformal theory

G_{38} : Category three of Entropy is seen to scale logarithmically with L, with a coefficient that corresponds to the central charge associated to the conformal theory

T_{36} : Category one of universal properties of the quantum phase transition

T_{37} : Category two of universal properties of the quantum phase transition

T_{38} : Category three of universal properties of the quantum phase transition

Module eight

Entanglement, a key concept of quantum information science, **obeys (eb)** universal scaling laws as dictated by the representations of the conformal group and its classification motivated by string theory **unveils (eb) a** monotonicity law for ground-state entanglement along the renormalization group flow.

G_{40} : Category one of Entanglement, a key concept of quantum information science, **obeys (eb)** universal scaling laws as dictated by the representations of the conformal group and its classification motivated by string theory

G_{41} : Category two of Entanglement, a key concept of quantum information science, **obeys (eb)** universal scaling laws as dictated by the representations of the conformal group and its classification motivated by string theory

G_{42} : Category three of Entanglement, a key concept of quantum information science, **obeys (eb)** universal scaling laws as dictated by the representations of the conformal group and its classification motivated by string theory

T_{40} : Category one of monotonicity law for ground-state entanglement along the renormalization group flow

T_{41} : Category two of monotonicity law for ground-state entanglement along the renormalization group flow

T_{42} : Category three of monotonicity law for ground-state entanglement along the renormalization group flow

Module Nine

Identification of majorization rule possibly associated to conformal invariance **and (e&eb) application to** present results leads to interpretation that the breakdown of density matrix renormalization group techniques near a critical point

G_{44} : Category one of Identification of majorization rule possibly associated to conformal invariance; present results leads to interpretation that the breakdown of density matrix renormalization group techniques near a critical point

G_{45} : Category two of Identification of majorization rule possibly associated to conformal invariance; present results leads to interpretation that the breakdown of density matrix renormalization group techniques near a critical point

G_{46} : Category three of Identification of majorization rule possibly associated to conformal invariance; present results leads to interpretation that the breakdown of density matrix renormalization group techniques near a critical point

T_{44} : Category one of present results leads to interpretation that the breakdown of density matrix renormalization group techniques near a critical point; Identification of majorization rule possibly associated to conformal invariance

T_{45} : Category two of present results leads to interpretation that the breakdown of density matrix renormalization group techniques near a critical point; Identification of majorization rule possibly associated to conformal invariance

T_{46} : Category three of present results leads to interpretation that the breakdown of density matrix renormalization group techniques near a critical point; Identification of majorization rule possibly associated to conformal invariance

SECTION SEVENTY SIX

Quantum Floating Point: Autumn Sonata

Abstract: Tanaka et al propose a floating-point serial divider for a single-flux quantum (SFQ) logic implementation **based on** a hardware algorithm of division using subtract and shift operations. Fast serial calculations are performed by **utilizing the** signed-digit representation. The divider is implemented with a regularly aligned pipeline structure called a systolic array. The processing element of the systolic array is designed **using** several logic gates unique to the SFQ circuits, to reduce the number of pipeline stages. As a result, high throughput and low latency **is achieved** with a small circuit size. The throughput approaches the maximum throughput of serial processing. A 4-bit divider was fabricated **using** the 2.5-kA/cm2 niobium process technology, and successful operation up to 19 GHz was demonstrated. **A High-Throughput Single-Flux Quantum Floating-Point Serial Divider Using the Signed-Digit Representation Tanaka, M. SSN: 1051-8223 INSPEC Accession Number: 10781580 Digital Object Identifier: 10.1109/TASC.2009.2018537.** The floating point registers are created using an IEEE formula. Simple explanation is the two 16 bit registers are treated as one 32 bit register, where the first 12 bits represent the base number and the remaining 20 bits are the power of 10. It is not as if the wide range of amplitudinal ramifications and **cardinal** repercussions are restricted to the field of serial divider analysis and cellular automata. Quantum floating point studies are topical and thought provoking which has enclosed in its orbit with effervescence and chutzpah the Autopoiesis systems too. An exemplar and illustration of such a study is Integral **biomathics: A post-Newtonian view into the logos of Plamen L. Simeonov, http://dx.doi.org/10.1016/j.pbiomolbio.2010.01.005.** This work is an attempt for a state-of-the-art survey of natural and life sciences with the goal to define the scope and address the central questions of an original research program. It is focused on the phenomena of emergence, adaptive dynamics and evolution of self-assembling, self-organizing, self-maintaining and self-replicating biosynthetic systems viewed from a newly-arranged perspective and understanding of computation and communication **in the living nature**. The author regards this research **as an integral part of the** emerging discipline of nature-inspired or natural computation, i.e. **computation inspired by or occurring in nature**. Within this context, he is interested in studies which represent a significant departure from traditional theories appropriate, diagnostic, differentiating, discriminating, discriminative, and distinctive about complex systems and self-organization,

emergent phenomena and artificial biology. In particular, **these include** non-conventional approaches exploring the aggregation, collection, ensemble, entirety, accentuation, corroboration and consummation with concatenation composition, growth and development of physical forms and structures, Autopoiesis along with the associated abstract information structures and processes**.** This paper provides a critical review of the major assumptions which **guide the** development of modern computer science and engineering towards emulating biological systems. For this purpose, the author explores the potential and the virtues of biology **to reshape** contemporary science. The goal of this survey is to discuss the present state of natural and engineering sciences in the light of a necessary paradigm change in the structure and methodology of research and deliver some insights for developing a new kind of integral science based on the principles for dynamic interdependence of the constituting disciplines and on the evolving relationships among them**.(emphasis mine)**

INTRODUCTION—VARIABLES USED

Quantum computer might be the theoretician's dream, a figment of imagination, product of fecund prognostication and resultant orientationality the fertile thought with kinetic look and feel but as far as experimentalists are concerned; its realization is a nightmare a sordid and sardonic incubus, ordeal, and phantasm. The problem is that while some prototypes of the simplest elements needed to build a quantum computer have already been implemented in the laboratory, it is still an open question how **to combine these** elements into scalable systems. Superposition of qubits is what gives quantum computers their inherent parallelism. According to physicist David Deutsch, this parallelism allows a quantum computer to work on a million computations at once, while your desktop PC works on one. A 30-qubit quantum computer would equal the processing power of a conventional computer that could run at 10 teraflops (trillions of floating-point operations per second). Today's typical desktop computers run at speeds measured in gigaflops (billions of floating-point operations per second).(**How Quantum Computers Work Kevin Bonsor and Jonathan Strickland)**

Quantum computation and Shor's factoring algorithm Artur Ekert et al

(1) Current technology is beginning to allow us to **manipulate** rather than just observe individual quantum phenomena. This opens up the possibility of exploiting quantum effects **to (eb) perform** computations beyond the scope of any classical computer.Peter Shor discovered an efficient algorithm for factoring whole numbers, **which uses** characteristically quantum effects. The algorithm illustrates the potential power of quantum computation, as there is no known efficient classical method for solving this problem. The authors give an exposition of Shor's algorithm together with an introduction to quantum computation and complexity theory. They discuss experiments that may contribute to its practical implementation. [S0034-6861(96)00303-0]URL: http://link.aps.org/doi/10.1103/RevModPhys.68.733 DOI: 10.1103/RevModPhys.68.733

Quantum Computation over Continuous Variables Seth Lloyd et al

(2) This paper provides necessary and sufficient conditions for **constructing a** universal quantum computer over continuous variables. As an example, it is shown how a universal quantum computer for the amplitudes of the electromagnetic field might be constructed **using** simple linear devices such as beam splitters and phase shifters, together with squeezers and nonlinear devices such as Kerr-effect fibers and atoms in optical cavities.

(3) Such a device could in principle **perform** "quantum floating point" computations. Problems involving noise, finite precision, and error correction are discussed.URL: http://link.aps.org/doi/10.1103/PhysRevLett.82.1784DOI: 10.1103/PhysRevLett.82.1784

Characterization, applicability and defect analysis for tiles nanostructure of quantum dot cellular automata DOI:10.1080/08927022.2010.536543 Kunal Dasa et al

(4) Quantum dot cellular automata (QCA) are an emerging technology for nanoscale computing. In this paper, our contribution is to enhance the applicability of 3 × 3 tile nanostructure in QCA. We introduced few novel 3 × 3 active-type tiles, namely triple fan-output tiles or butterfly tile, five-input majority voter (MV) and fault-tolerant 3 × 3 tile-coupled majority–minority voter. Authors also explore the cascading effect **on (e&eb)** 3 × 3 tile.

(5) Triple fan-out tile is applicable to drive three outputs simultaneously. Five-input MV, which is another active-type tile we proposed, **is applicable for** designing five-variable or five addressing, advanced computations.

(6) In passive-type 3 × 3 tiles, they **introduce (eb)** symmetrical baseline tiles and asymmetrical fully populated fan-in tiles. QCA design was found to be having high error during fabrication of device. Authors make exhaustive simulation for analysing the defect that can occur during fabrication process.

(7) Defects such as extra cell deposition, missing cell deposition and displaced cell deposition for QCA are prone **to (eb) occur**. Fault analyses of the above-mentioned 3 × 3 tiles are performed. They also focused **on the effect** of extra dot or missing dot deposition, during the chemical synthesis phase.

NOTATION

Module One

Current technology is beginning to allow us to **manipulate** rather than just observe individual quantum phenomena. This opens up the possibility of exploiting quantum effects **to (eb) perform** computations beyond the scope of any classical computer.Peter Shor discovered an efficient algorithm for factoring whole numbers, **which uses** characteristically quantum effects. The algorithm illustrates the potential power of quantum computation, as there is no known efficient classical method for solving this problem. The authors give an exposition of Shor's algorithm together with an introduction to quantum computation and complexity theory. They discuss experiments that may contribute to its practical implementation. [S0034-6861(96)00303-0]URL: http://link.aps.org/doi/10.1103/RevModPhys.68.733 DOI: 10.1103/RevModPhys.68.733

G_{13} : Category one of **Shor's** algorithm. We are talking of the systems to which algorithm is applied and the differentiating, discriminating ,representative, singular characterstics thereof

G_{14} : Category two of Shor's algorithm

G_{15} : Category three of Shor's algorithm

T_{13} : Category one of quantum computation and complexity theory

T_{14} : Category two of quantum computation and complexity theory

T_{15} : Category three of quantum computation and complexity theory

Module Two

This paper provides necessary and sufficient conditions for **constructing a** universal quantum computer over continuous variables illustrated by the ramification how a universal quantum computer for the amplitudes of the electromagnetic field might be constructed **using** simple linear devices such as beam

splitters and phase shifters, together with squeezers and nonlinear devices such as Kerr-effect fibers and atoms in optical cavities

G_{16} : Category one of simple linear devices such as beam splitters and phase shifters, together with squeezers and nonlinear devices such as Kerr-effect fibers and atoms in optical cavities

G_{17} : Category two of simple linear devices such as beam splitters and phase shifters, together with squeezers and nonlinear devices such as Kerr-effect fibers and atoms in optical cavities

G_{18} : Category three of simple linear devices such as beam splitters and phase shifters, together with squeezers and nonlinear devices such as Kerr-effect fibers and atoms in optical cavities

T_{16} : Category one of necessary and sufficient conditions for **constructing a** universal quantum computer over continuous variables illustrated by the ramification how a universal quantum computer for the amplitudes of the electromagnetic field might be constructed

T_{17} : Category two of necessary and sufficient conditions for **constructing a** universal quantum computer over continuous variables illustrated by the ramification how a universal quantum computer for the amplitudes of the electromagnetic field might be constructed

T_{18} : Category three of necessary and sufficient conditions for **constructing a** universal quantum computer over continuous variables illustrated by the ramification how a universal quantum computer for the amplitudes of the electromagnetic field might be constructed

Module three

Such a device could in principle **perform** "quantum floating point" computations problems involving noise, finite precision, and error correction

G_{20} : Category one of necessary and sufficient conditions for **constructing a** universal quantum computer over continuous variables illustrated by the ramification how a universal quantum computer for the amplitudes of the electromagnetic field might be constructed **using** simple linear devices such as beam splitters and phase shifters, together with squeezers and nonlinear devices such as Kerr-effect fibers and atoms in optical cavities

G_{21} : Category two of necessary and sufficient conditions for **constructing a** universal quantum computer over continuous variables illustrated by the ramification how a universal quantum computer for the amplitudes of the electromagnetic field might be constructed **using** simple linear devices such as beam splitters and phase shifters, together with squeezers and nonlinear devices such as Kerr-effect fibers and atoms in optical cavities

G_{22} : Category three of necessary and sufficient conditions for **constructing a** universal quantum computer over continuous variables illustrated by the ramification how a universal quantum computer for the amplitudes of the electromagnetic field might be constructed **using** simple linear devices such as beam splitters and phase shifters, together with squeezers and nonlinear devices such as Kerr-effect fibers and atoms in optical cavities

T_{20} : Category one of "quantum floating point" computations problems involving noise, finite precision, and error correction

T_{21} : Category two of "quantum floating point" computations problems involving noise, finite precision, and error correction

T_{22} : Category three of "quantum floating point" computations problems involving noise, finite precision, and error correction

Module four

Quantum dot cellular automata (QCA) are an emerging technology for nanoscale computing and to enhance the applicability of 3 × 3 tile nanostructure in QCA is done **by introduction** few novel 3 × 3 active-type tiles, namely triple fan-output tiles or butterfly tile, five-input majority voter (MV) and fault-tolerant 3 × 3 tile-coupled majority–minority voter. Authors also explore the cascading effect **on (e&eb)** 3 × 3 tile.

G_{24} : Category one of Quantum dot cellular automata (QCA) are an emerging technology for nanoscale computing and to enhance the applicability of 3 × 3 tile nanostructure in QCA is done

G_{25} : Category two of Quantum dot cellular automata (QCA) are an emerging technology for nanoscale computing and to enhance the applicability of 3 × 3 tile nanostructure in QCA is done

G_{26} : Category three of Quantum dot cellular automata (QCA) are an emerging technology for nanoscale computing and to enhance the applicability of 3 × 3 tile nanostructure in QCA is done

T_{24} : Category one of few novel 3 × 3 active-type tiles, namely triple fan-output tiles or butterfly tile, five-input majority voter (MV) and fault-tolerant 3 × 3 tile-coupled majority–minority voter. Authors also explore the cascading effect **on (e&eb)** 3 × 3 tile.

T_{25} : Category two of few novel 3 × 3 active-type tiles, namely triple fan-output tiles or butterfly tile, five-input majority voter (MV) and fault-tolerant 3 × 3 tile-coupled majority–minority voter. Authors also explore the cascading effect **on (e&eb)** 3 × 3 tile.

T_{26} : Category three of few novel 3 × 3 active-type tiles, namely triple fan-output tiles or butterfly tile, five-input majority voter (MV) and fault-tolerant 3 × 3 tile-coupled majority–minority voter. Authors also explore the cascading effect **on (e&eb)** 3 × 3 tile.

Module five(Both cases are discussed)

Triple fan-out tile is applicable to drive three outputs simultaneously. Five-input MV, which is another active-type tile authors proposal, **is applicable for** designing five-variable or five addressing, advanced computations

G_{28} : Category one of Five-input MV, which is another active-type tile authors proposal; designing five-variable or five addressing, advanced computations

G_{29} : Category two of Five-input MV, which is another active-type tile authors proposal; designing five-variable or five addressing, advanced computations

G_{30} : Category three of Five-input MV, which is another active-type tile authors proposal; designing five-variable or five addressing, advanced computations

T_{28} : Category one of designing five-variable or five addressing, advanced computations; Five-input MV, which is another active-type tile authors proposal

T_{29} : Category two of designing five-variable or five addressing, advanced computations; Five-input MV, which is another active-type tile authors proposal

T_{30} : Category three of designing five-variable or five addressing, advanced computations; Five-input MV, which is another active-type tile authors proposal

Module six

In passive-type 3 × 3 tiles, they **introduce (eb)** symmetrical baseline tiles and asymmetrical fully populated fan-in tiles. QCA design was found to be having high error during fabrication of device. Authors make exhaustive simulation for analysing the defect that can occur during fabrication process

G_{32} : Category one of symmetrical baseline tiles and asymmetrical fully populated fan-in tiles

G_{33} : Category two of symmetrical baseline tiles and asymmetrical fully populated fan-in tiles

G_{34} : Category three of symmetrical baseline tiles and asymmetrical fully populated fan-in tiles

T_{32} : Category one of passive-type 3 × 3 tiles

T_{33} : Category two of passive-type 3 × 3 tiles

T_{34} : Category three of passive-type 3 × 3 tiles

Module seven

Defects such as extra cell deposition, missing cell deposition and displaced cell deposition for QCA are prone **to (eb) occur**. Fault analyses of the above-mentioned 3 × 3 tiles are performed. They also focused **on the effect** of extra dot or missing dot deposition, during the chemical synthesis phase.

G_{36} : Category one of Defects such as extra cell deposition, missing cell deposition and displaced cell deposition for QCA are prone **to (eb) occur**

G_{37} : Category two of Defects such as extra cell deposition, missing cell deposition and displaced cell deposition for QCA are prone **to (eb) occur**

G_{38} : Category three of Defects such as extra cell deposition, missing cell deposition and displaced cell deposition for QCA are prone **to (eb) occur**

T_{36} : Category one of fault analyses of the above-mentioned 3 × 3 tiles and implications and ramifications of extra dot or missing dot deposition, during the chemical synthesis phase

T_{37} : Category two of fault analyses of the above-mentioned 3 × 3 tiles and implications and ramifications of extra dot or missing dot deposition, during the chemical synthesis phase

T_{38} : Category three of fault analyses of the above-mentioned 3 × 3 tiles and implications and ramifications of extra dot or missing dot deposition, during the chemical synthesis phase

Module eight

Algorithm design is a highly complicated task, and in quantum computing it becomes even more complicated **due (e)** tothe attempts to harness quantum mechanical features to reduce the complexity of computational problems and to "speed-up" computation**. (Mystery, myth, Reality & Truth Sumit Raj IIIT Hyderabad)**

G_{40} : Category one of attempts to harness quantum mechanical features to reduce the complexity of computational problems and to "speed-up" computation

G_{41} : Category two of attempts to harness quantum mechanical features to reduce the complexity of computational problems and to "speed-up" computation

G_{42} : Category three of attempts to harness quantum mechanical features to reduce the complexity of computational problems and to "speed-up" computation

T_{40} : Category one of Algorithm design is a highly complicated task, and in quantum computing it becomes even more complicated

T_{41} : Category two of Algorithm design is a highly complicated task, and in quantum computing it becomes even more complicated

T_{42} : Category three of Algorithm design is a highly complicated task, and in quantum computing it becomes even more complicated

Module Nine (care: There are five models in this module)

(Mystery, myth, Reality & Truth Sumit Raj IIIT Hyderabad)

A 30-qubit quantum computer **would equal the** processing power of a conventional computer that could run at 10 teraflops (trillions of floating-point operations per second

Younis & knight theory

Younis & Knight: Power dissipation in conventional CMOS primarily occurs during device switching. One component of this dissipation **is due to** charging and discharging the gate capacitances through conducting, but slightly resistive, devices.

We note here that it is not the charging or the discharging of the gate that is necessarily dissipative, **but rather that a** small time is allocated to perform these operations. In conventionalCMOS, the time constant associated with charging the gate through a similar transistor is RC, where R is the ON resistance of the device and C its capacitance.

However, the cycle time can be, and usually is, much longer than RC. Obvious conclusion is that energy consumption **can be (e) reduced by** spreading the transitions over the whole cycle rather than "squeezing" it all inside one RC.

QC making can be achieved by Ion traps use optical or magnetic fields (or a combination of both) to trap ions. Electron spin. Quantum dots are made of semiconductor material and are used to contain and manipulate electrons. Optical traps use light waves to trap and control particles. Superconducting circuits allow electrons to flow with almost no resistance at very low temperatures. DNA computing is another method of making QC

Landauer theory

Landauer has posed the question of whether logical irreversibility is an unavoidable feature of useful computers, arguing that it is, and has demonstrated the physical and philosophical importance of this

question by showing that whenever a physical computer throws away information about its previous state **it must generate** a corresponding amount of entropy. Therefore, a computer must dissipate at leastkTln 2 of energy (about 3 X 10-21 joule at room temperature) for each bit of information it erases or otherwise throws away

Toffoli theory

At this time Tomasso Toffoli (1980) showed that there exists a reversible gate **which could play role** of a universal gate for reversible circuits. These two simultaneously **have lead to the** exploration in the field of Reversible Logic Gates and Circuits.

In recent years, reversible computing system design is attracting a lot of attention. Reversible computing is based on two concepts: logic reversibility and physical reversibility. A computational operation is said to be logically reversible if the logical state of the computational device before the operation of the device can be **determined (e) by** its state after the operation i.e., the input of the system can be retrieved from the output obtained from it.

G_{44} : Category one of processing power of a conventional computer that could run at 10 teraflops (trillions of floating-point operations per second; charging and discharging the gate capacitances through conducting, but slightly resistive, devices; dissipative; spreading the transitions over the whole cycle rather than "squeezing" it all inside one RC; logical irreversibility is an unavoidable feature of useful computers, arguing that it is, and has demonstrated the physical and philosophical importance of this question by showing that whenever a physical computer throws away information about its previous state; universal gate for reversible circuits; its state after the operation i.e., the input of the system can be retrieved from the output obtained from it.

G_{45} : Category two of processing power of a conventional computer that could run at 10 teraflops (trillions of floating-point operations per second; charging and discharging the gate capacitances through conducting, but slightly resistive, devices; dissipative; spreading the transitions over the whole cycle rather than "squeezing" it all inside one RC; logical irreversibility is an unavoidable feature of useful computers, arguing that it is, and has demonstrated the physical and philosophical importance of this question by showing that whenever a physical computer throws away information about its previous state; universal gate for reversible circuits; its state after the operation i.e., the input of the system can be retrieved from the output obtained from it.

G_{46} : Category three of processing power of a conventional computer that could run at 10 teraflops (trillions of floating-point operations per second; charging and discharging the gate capacitances through conducting, but slightly resistive, devices; dissipative; spreading the transitions over the whole cycle rather than "squeezing" it all inside one RC; logical irreversibility is an unavoidable feature of useful computers, arguing that it is, and has demonstrated the physical and philosophical importance of this question by showing that whenever a physical computer throws away information about its previous state; universal gate for reversible circuits; its state after the operation i.e., the input of the system can be retrieved from the output obtained from it.

T_{44} : Category one of30-qubit quantum computer; Power dissipation in conventional CMOS primarily occurs during device switching. One component of this dissipation; small time allocated to perform charging or the discharging of the gate; cycle time can be, and usually is, much longer than RC. Obvious conclusion is that energy consumption; corresponding amount of entropy; there exists a reversible gate; computational operation is said to be logically reversible if the logical state of the computational device before the

operation of the device;

T_{45} : Category two of 30-qubit quantum computer; Power dissipation in conventional CMOS primarily occurs during device switching. One component of this dissipation; small time allocated to perform charging or the discharging of the gate; cycle time can be, and usually is, much longer than RC. Obvious conclusion is that energy consumption; corresponding amount of entropy; there exists a reversible gate; computational operation is said to be logically reversible if the logical state of the computational device before the operation of the device;

T_{46} : Category three of 30-qubit quantum computer; Power dissipation in conventional CMOS primarily occurs during device switching. One component of this dissipation; small time allocated to perform charging or the discharging of the gate; cycle time can be, and usually is, much longer than RC. Obvious conclusion is that energy consumption; corresponding amount of entropy; there exists a reversible gate; computational operation is said to be logically reversible if the logical state of the computational device before the operation of the device;

SECTION SEVENTY SEVEN

Decoherence Engineering: Deux Ex machina

Abstract: There is new scorcher of scordatura and verkrampte reactionary on line. Grapevine is baffling and befuddling with newer stories. Currently a new dawn of template of rocky razzmatazz and scintillating and scarlet explorations have lead prospective to entering a new technological era in which we are able to build systems whose performance **is limited by** quantum physical effects and in which it may be possible to exploit non-classical phenomena in novel ways. To this end, there has been considerable recent interest in engineering quantum systems and at the heart of this is the development of a quantum control theory dedicated **to extending** classical control to the quantum domain. Examples already utilizing control of one sort or another **include** quantum electromechanical systems, **quantum dots**, cooper-pair boxes, superconducting interference devices, ion traps, as well as a large selection of optical devices. It is clear that a mathematical framework is essential for the future development of quantum control as an engineering discipline. The aim of the programme is to bring together experimentalists and theoreticians working in quantum engineering to identify the core mathematical issues and challenges ahead**.** These advances could enable the creation of immensely powerful computers as well as other applications, such as highly sensitive detectors capable of probing biological systems. "We are really excited about the possibilities of new semiconductor materials and new experimental systems that have become available in the last decade," said Jason Petta, one of the authors of the report and an associate professor of physics at Princeton University. **http://www.newton.ac.uk/programmes/QCE/. Quantum Control Engineering: Mathematical Principles and Applications** Method for scaling a neutral atom Rydberg gate quantum processor to a large number of qubits has been studied and investigated by **Saffman et al.,** Limits are derived **showing that** the number of qubits that can be directly connected by entangling gates with errors at the 10^{-3} level using long range Rydberg interactions between sites in an optical lattice, without mechanical motion or swap chains, is about 500 in two dimensions and 7500 in three dimensions. A scaling factor of 60 at a smaller number of sites can be obtained **using (e)** collective register encoding in the hyperfine ground states of the rare earth atom Holmium. Authors present a detailed analysis of operation of the 60 qubit register in

Holmium. Combining a lattice of multi-qubit ensembles **with** collective encoding results in a feasible design for a 1000 qubit fully connected quantum processor. Journal reference: Phys. Rev. A 78, 012336 (2008) DOI: 10.1103/PhysRevA.78.012336 arXiv: 0805.0440 [quant-ph] **Scaling the neutral atom Rydberg gate quantum computer by collective encoding in Holmium atoms M. Saffman et al** One other study linked decoherence engineering and adroit manipulation to EPR experiment. Quantum-state decoherence is the major **obstacle (e) to** the implementation of quantum computing and quantum information. Here authors present an experimental proof of principle for the quantum engineering of decohering interactions. An optically generated Einstein-Podolski-Rosen (EPR) entangled state was subjected **to severe (e) loss** by placing a 50% beam splitter in the path of each entangled light beam. The resulting severe degradation of the signature quantum correlations between the two beams was subsequently **alleviated (e)** when another, independent EPR state impinged upon the other input ports of the beam splitter, effectively **entangling the** losses inflicted to the initial EPR state. The additional EPR beam pair was classically coherent with the primary one but had no quantum correlations with it. This result may also find **application as a** "quantum tap" for entanglement. **Protecting an EPR state by quantum engineering of decoherence AIP Conf. Proc. 1363, pp. 233-238; doi: http://dx.doi.org/10.1063/1.3630186 (6 pages) QUANTUM COMMUNICATION, MEASUREMENT AND COMPUTING (QCMC): Russell Bloomer et al**

INTRODUCTION—VARIABLES USED

Source: NC University Page (emphasis mine)

Quantum Computing

(1) Although early demonstrations of quantum computing explored a wide variety of experimental methods, quantum computers **based on** solid-state systems, in spite of the potentially larger interactions limiting coherence, are believed to be worth serious investigation.
(2) Systems based on Josephson Junctions, electrons resident on the surface of superfluid Helium, compound semiconductors, electrons trapped in an acoustic wave, semiconductor quantum dots coupled via electromagnetic field, and solid-state nuclear magnetic resonance systems that have been proposed are described below.
(3) Previously we have investigated coherent coupling of electrons trapped at neighboring asymmetric quantum dots in a III-V pillar structure. However, we believe the best candidate **is based on** the coupled spin of electrons on trapped on adjacent quantum dots in a silicon environment currently being investigated in our research.

Quantum Josephson Junction Circuits

(4) A superconductor can be characterized by the absolute value (number of the quasiparticles) and the phase f of the macroscopic wavefunction, **representing (eb)** the order parameter.
(5) As for any quantum system, both of them cannot be determined simultaneously. If a Josephson junction is formed by the two massive superconductors **they (=) are** insensitive to the fluctuations of the quasiparticle number and the junction evolution can be described by the classical equations for the relative phase of the contacting superconductors.
(6) Small superconductors, however, have small capacitance; if the charging energy exceeds the Josephson energy, then the **(e) suppressed** charge fluctuations and the circuit demonstrates essentially quantum behavior.
(7) It has been demonstrated that in this case it is possible to create a quantum superposition of the two different charge states of the superconductor, **thus, forming** a qubit.
(8) Creation of a quantum superposition of the two different charge states of the superconductor, **thus,**

forming a qubit is possible **by (e) biasing** the additional gate contact or applying a microwave radiation.

(9) **As a (eb) result**, the Bloch oscillations of frequency between the two charge states are inspired.

(10) Addressing of the qubit can be accomplished by the **control (e&eb)of the** duration of the gate voltage pulse, while the two-qubit operation can be organized by coupling of the different circuits.

(11) The obvious advantage of this scheme is that **it (e) exploits** a quantum behavior of a macroscopic system that promises relatively easy control of the qubit.

(12) On the other hand, this concept is not free of some drawbacks. In particular, to inspire the Bloch oscillations, the applied voltage pulse **must (=) be** sharp on the time scale.

(13) Although the Josephson energy can be controlled by external magnetic flux, to avoid decoherence during the computation, it **should (=) be kept** at the level about 50 meV, as the London length and the coherence length of the Cooper which corresponds to about 10 GHz in the frequency domain.

(14) Obviously, it is a great challenge to achieve reliable operation at such high frequencies of massive circuits containing a large number of qubits. In addition, there are some restrictions in the possible miniaturization of the Josephson-junction circuits. Currently, the experiments accomplished where done with the superconductor having dimensions of the order of a micron that corresponds to the charging energy about 0.1 meV. Scaling down the junction dimensions **leads to production of (eb)** increase in the charging energy. As a result, a more precise adjustment of the gate voltage required for the resonance condition is necessary.

(15) What is more fundamental, the scaling of the junction dimensions **is (e) restricted** by such scales pairs.

Electrons Confined at the Surface of Superfluid Helium

(16) Close to the surface of the thin superfluid helium film an electron experiences Coulomb-like attraction **caused by the** electrostatic image forces.

(17) **As a result**, injected electrons are confined in the vicinity of the helium surface.

(18) Electron-electron interaction in such a system **causes (eb)** formation of a 2D Wigner crystal.

(19) The positions of the electrons in this crystal can be pinned **by (e) means of** an electrodes patterned at the bottom of the cell.

(20) The ground and excited states of an electron in the Coulomb-like potential are proposed to be used as basis states of a qubit. Addressing of the qubit can be accomplished **by (e) the** resonance microwave radiation causing Rabi oscillations.

(21) Adjustment of the resonance frequency of a particular qubit can be made by the electrodes **to(eb) allow** control of one-qubit rotations and two-qubit operation, **caused(e) by** the electron-electron coupling.

(22) Since the wavefunction of the ground state in the classically forbidden region **decays (e)** faster than that of the excited state, the tunneling of the electron **is sensitive** to the state of the qubit.

(23) Therefore, the electron tunneling current measurement by an additional electrode **can (eb) accomplishes** the qubit readout.

(24) Relatively weak electron interaction with the elementary excitations of helium, surface capillary waves and ripplons, **ensures (eb)** long decoherence times for the qubit and the possibility to implement the reliable quantum computations.

(25) However, it is unclear whether two-qubit coupling can be **properly (e&eb) controlled** for the quantum gate implementation.

(26) The possibility of the proposed readout procedure is also questionable. If a single electron is used as qubit, then its readout **is (e) complemented by** the electron removal. This complementation of and augmentation thereof of the single electron used as a qubit by the electron removal, obviously, **creates a** defect in the Wigner crystal and can strongly disturb the neighbor electrons. This

difficulty can be **overcome by (e) using** a cluster of electrons as a single qubit.

(27) In this case, however, it is more difficult to achieve the necessary decoherence level. One more serious disadvantage of this scheme is that even weak vibrations caused by the environment **strongly (e&eb) disturb** the superfluid helium surface giving rise additional electron decoherence or even complete destruction of the electron Wigner crystal. Obviously, this would establish very tough conditions for the mechanical isolation of such QC and restrict the area of its applications.

Spin-Split States in Compound Semiconductor Quantum Dots with Spin Injection

(28) This design of the QC has a lot in common with that proposed by us. The two main differences are as follows: (i) instead of silicon, a compound semiconductor is proposed as a quantum-dot material. In this case the two-level system is formed by the spin-split state, originated as a result of relatively strong spin-orbital coupling in compound semiconductors lacking inversion symmetry (Rashba effect). As a result, an effective voltage-control of the energy gap between the two states is possible with no magnetic field required; (ii) the initial spin-polarized state is created by the injection of a spin-polarized carrier from a magnetic material surrounding the quantum dot. Analogously, the readout of the qubit state is accomplished by measuring the spin-dependent current from the quantum dot to the magnetic material. Although this design could provide easier control of the energy splitting of the two states representing a qubit, it sacrifices such an advantage of silicon as very long spin decoherence time. In addition, injection of the spin-polarized carriers in semiconductors is a great challenge. Currently, it was observed in a few materials. In general, it seems to be very difficult to find a pair semiconductor-ferromagnetic material or semiconductor-Semimagnetic semiconductor with good-quality interface causing no strong spin decoherence.

Quantum Josephson junction Circuits

(29) This approach is based on a specific method to create 0D electron states and arrange their coupling. Recent experiments demonstrate that the piezoelectric potential of a gigahertz surface acoustic wave (SAW) can **effectively (e) trap** 2D electrons confined near the semiconductor surface.

(30) If a quasi-1D channel is formed in the 2D electron gas by gate electrodes or etching, the SAW traveling along the channel **drags (e) the** electrons.

(31) It is possible to calibrate the parameters of the SAW and 2D electrons **to (e) capture** a single electron from the channel in each potential minimum of the SAW.

(32) In this case the qubit is **represented (eb) by** the spin-split electron states in the external magnetic field.

(33) A two-qubit logic gate can be **achieved (eb)** by patterning a set of parallel 1D channel. In this case the electrons from the adjacent channels captured in the same potential well of the SAW are coupled **due (e) to the** exchange interaction.

(34) This approach allows control of the two-qubit logic gates by bringing the two adjacent channels closer for some portion of the channel, thus, controlling the exchange-induced two-qubit coupling **by (e&eb) the** geometry of the structure.

(35) However, it is questionable if it is possible to achieve a low enough level of electron spin decoherence for this design, since it assumes **using(e)** GaAs-based structures where the decoherence is much stronger than for silicon. Furthermore, some additional source of decoherence can be brought about by the SAW-generating circuit and the SAW itself.

Semiconductor Quantum Dots Coupled via Electromagnetic Field of a Microcavity

(36) In this approach it is proposed to implement in solid-state systems a method of quantum computation **using (e)** cavity QED.

(37) In this case a number of quantum dots are put into a common Microcavity **characterized (e) by** a high quality factor.

(38) The computation process utilizes the three lowest levels of a quantum dot; two of them represent a state of a qubit, while the third one **is (e) used** to arrange an electron interaction with the electromagnetic field.

(39) The two-qubit operation in this system is realized by the nonlinear interaction of the electron with the coherent electromagnetic radiation having the frequency different from the eigen-frequency of the cavity. The readout procedure **is (eb) accomplished** by means of the radiation detector.

(40) The advantage of this approach **is that (eb) it allows** entanglement of an arbitrary pair of qubits by tuning the energy levels of the individual quantum dots by the gate electrodes.

(41) However, the relatively high decoherence rate in such structure requires **use (e) of** the terahertz-range radiation; currently, it is a significant challenge **to create** high-quality microelectronic sources of terahertz radiation as well as terahertz detectors.

(42) In addition, it is necessary to mention that this approach **requires(e)** manufacturing a very sophisticated structure that integrates a set of individually controlled quantum dots, the cavity, laser, and detector, which comprises a significant technological challenge.

Solid-State NMR Quantum Computers

(43) Demonstration of the elements of the QC employing NMR in molecular systems inspired researches to investigate the possibility to use NMR in solids for the same purpose. Basically, the main principles of quantum computation based on nuclear and electron spins are similar. The main advantage of the nuclear spins is their weak coupling to the environment, **ensuring (eb) a** long decoherence time.

(44) On the other hand, the weak coupling makes the control of a qubit more difficult **and (e) reduces** the possible clock frequency of the QC.

(45) A number of propositions were generated **that (e) use** either electrical or optical methods to control the two-qubit rotations in NMR solid-state QC.

(46) Both of these schemes require the high-precision implantation of individual donor atoms in a silicon substrate, which is extremely hard to achieve, especially in many-qubit circuits. There is also an alternative NMR-based scheme for the quantum computation. Here an original method of the spin detection is proposed. It is based on the interaction of the nonuniform magnetic field of a ferromagnetic particle **and (e&eb) the** spins subject to the resonance radio frequency field. This interaction **causes the** mechanical oscillations of a micrometer cantilever, where the ferromagnetic particle is deposited.

(47) In turn, the mechanical oscillations are detected by the optical methods. Although this scheme **can provide** very weak decoherence, it would be extremely hard technologically to implement it for the many-qubit systems.

(48) More recently this scheme has been augmented by the suggestion that Si29 atoms can be preferentially deposited by MBE at the steps produced on a vicinal Si (111) 7´7 surface. With this structure placed in an extremely large magnetic field gradient to allow addressing, the readout **is (e) provided by** magnetic resonance force microscopy. It is suggested by initial state preparation could be aided by optical pumping and spin transfer from electrons. On the order of 105 identical systems are required to obtain adequate signal detection for ensemble measurement of the final state.

Engineering decoherence in Josephson persistent-current qubits Caspar H. van der Wal et al

(49) Authors discuss the relaxation and dephasing rates **that result from** the control and the measurement setup itself in experiments on Josephson persistent-current qubits.

(50) For control and measurement of the qubit state, the qubit is inductively **coupled to** electromagnetic circuitry.

(51) Authors show how this system can be mapped on the spin-boson model, and how the spectral density of the bosonic bath can be **derived from the** electromagnetic impedance that is coupled to the qubit.

(52) Part of the electromagnetic environment is a measurement apparatus (DC-SQUID) that is **permanently coupled to** the single quantum system that is studied.

(53) Since there is an obvious conflict between long coherence times and an efficient measurement scheme, the measurement process is analyzed in detail for different measurement schemes. They show that the coupling of the measurement apparatus to the qubit **can be (e&eb) controlled** in situ.

(54) Parameters that can be realized in experiments today are used for a quantitative evaluation, **and it is shown** that the relaxation and dephasing rates that are induced by the measurement setup can be made low enough for a time-resolved study of the quantum dynamics of Josephson persistent-current qubits. Our results can be generalized as engineering rules for the read-out of related qubit systems. ArXiv: cond-mat/0211664 (cond-matter)

Decoherence, Control, and Symmetry in Quantum Computers D. Bacon et al

(1) In this thesis we describe methods **for (e) avoiding the** detrimental effects of decoherence while at the same time still allowing for computation of the quantum information.

(2) The philosophy of the method discussed in the first part of this thesis is **to (e) use a** symmetry of the decoherence mechanism to find robust encodings of the quantum information.

(3) Stability, control, and methods **for (e) using** decoherence-free information in a quantum computer are presented with a specific emphasis on decoherence due to a collective coupling between the system and its environment.

(4) Universal quantum computation is **(eb) demonstrated** on such collective decoherence decoherence-free encodings.

(5) Rigorous definitions of control **and the use** of encoded universality in quantum computers are addressed. Explicit gate constructions for encoded universality on ion trap and exchange **based** quantum computers are given.

(6) In the second part of the thesis authors examine physical systems with error correcting properties. We examine systems that can store quantum information in their ground state such that decoherence processes **are (e) prohibited** via energetics.

(7) They present the theory of supercoherent systems whose ground states are quantum error detecting codes **and** describe a spin ladder whose ground state has both the error detecting and correcting properties. We conclude by discussing naturally fault-tolerant quantum computation. ArXiv:quant-ph/0305025

(8) On all levels accessible to experiment, quantum theory **generates** flawless predictions for every single one of matter's quirky operations. Faced with success after success the majority of physicists quietly decided that to give up reality in exchange for quantum theory's immense predictive power was not such a bad deal after all.

NOTATION

Module One

Demonstration of the elements of the QC employing NMR in molecular systems inspired researches to investigate the possibility to use NMR in solids for the same purpose. Basically, the main principles of quantum computation based on nuclear and electron spins are similar. The main advantage of the nuclear spins is their weak coupling to the environment, **ensuring (eb) a** long decoherence time.

G_{13} : Category one of main advantage of the nuclear spins is their weak coupling to the environment

G_{14} : Category two of main advantage of the nuclear spins is their weak coupling to the environment

G_{15} : Category three of main advantage of the nuclear spins is their weak coupling to the environment

T_{13} : Category one of long decoherence time.

T_{14} : Category two of long decoherence time.

T_{15} : Category three of long decoherence time.

Module Two

The philosophy of the method discussed in the first part of this thesis is **to (e) use a** symmetry of the decoherence mechanism to find robust encodings of the quantum information

G_{16} : Category one of symmetry of the decoherence mechanism to find robust encodings of the quantum information

G_{17} : Category two of symmetry of the decoherence mechanism to find robust encodings of the quantum information

G_{18} : Category three of symmetry of the decoherence mechanism to find robust encodings of the quantum information

T_{16} : Category one of philosophy of the method, nay the modus operandi

T_{17} : Category two of philosophy of the method, nay the modus operandi

T_{18} : Category three of philosophy of the method, nay the modus operandi

Module three

Stability, control, and methods **for (e) using** decoherence-free information in a quantum computer are presented with a specific emphasis on decoherence due to a collective coupling between the system and its environment

G_{20} : Category one of decoherence-free information in a quantum computer are presented with a specific emphasis on decoherence due to a collective coupling between the system and its environment

G_{21} : Category two of decoherence-free information in a quantum computer are presented with a specific emphasis on decoherence due to a collective coupling between the system and its environment

G_{22} : Category three of decoherence-free information in a quantum computer are presented with a specific emphasis on decoherence due to a collective coupling between the system and its environment

T_{20} : Category one of Stability, control, and methods

T_{21} : Category two of Stability, control, and methods

T_{22} : Category three of Stability, control, and methods

Module four

Universal quantum computation is **(eb) demonstrated** on such collective decoherence decoherence-free encodings

G_{24} : Category one of Universal quantum computation

G_{25} : Category two of Universal quantum computation

G_{26} : Category three of Universal quantum computation

T_{24} : Category one of collective decoherence decoherence-free encodings

T_{25} : Category two of collective decoherence decoherence-free encodings

T_{26} : Category three of collective decoherence decoherence-free encodings

Module five

Rigorous definitions of control **and the use** of encoded universality in quantum computers are addressed. Explicit gate constructions for encoded universality on ion trap and exchange **based** quantum computers are given

G_{28} : Category one of Universal quantum computation is **(eb) demonstrated** on such collective decoherence decoherence-free encodings

G_{29} : Category two of Universal quantum computation is **(eb) demonstrated** on such collective decoherence decoherence-free encodings

G_{30} : Category three of Universal quantum computation is **(eb) demonstrated** on such collective decoherence decoherence-free encodings

T_{28} : Category one of Explicit gate constructions for encoded universality on ion trap and exchange

T_{29} : Category two of Explicit gate constructions for encoded universality on ion trap and exchange

T_{30} : Category three of Explicit gate constructions for encoded universality on ion trap and exchange

Module six

They present the theory of supercoherent systems whose ground states are quantum error detecting codes **and** describe a spin ladder whose ground state has both the error detecting and correcting properties. We conclude by discussing naturally fault-tolerant quantum computation. ArXiv:quant-ph/0305025

G_{32} : Category one of supercoherent systems whose ground states are quantum error detecting codes

G_{33} : Category two of supercoherent systems whose ground states are quantum error detecting codes

G_{34} : Category three of supercoherent systems whose ground states are quantum error detecting codes

T_{32} : Category one of spin ladder whose ground state has both the error detecting and correcting properties and fault-tolerant quantum computation

T_{33} : Category two of spin ladder whose ground state has both the error detecting and correcting properties and fault-tolerant quantum computation

T_{34} : Category three of spin ladder whose ground state has both the error detecting and correcting properties and fault-tolerant quantum computation

Module seven

Parameters that can be realized in experiments today are used for a quantitative evaluation, **and it is shown** that the relaxation and dephasing rates that are induced by the measurement setup can be made low enough for a time-resolved study of the quantum dynamics of Josephson persistent-current qubits. Our results can be generalized as engineering rules for the read-out of related qubit systems. ArXiv: cond-mat/0211664 (cond-matter)

G_{36} : Category one of Parameters that can be realized in experiments today are used for a quantitative evaluation,

G_{37} : Category two of Parameters that can be realized in experiments today are used for a quantitative evaluation,

G_{38} : Category three of Parameters that can be realized in experiments today are used for a quantitative evaluation,

T_{36} : Category one of relaxation and dephasing rates that are induced by the measurement setup can be made low enough for a time-resolved study of the quantum dynamics of Josephson persistent-current qubits. Our results can be generalized as engineering rules for the read-out of related qubit systems

T_{37} : Category two of relaxation and dephasing rates that are induced by the measurement setup can be made low enough for a time-resolved study of the quantum dynamics of Josephson persistent-current qubits. Our results can be generalized as engineering rules for the read-out of related qubit systems

T_{38} : Category three of relaxation and dephasing rates that are induced by the measurement setup can be made low enough for a time-resolved study of the quantum dynamics of Josephson persistent-current qubits. Our results can be generalized as engineering rules for the read-out of related qubit systems

Module eight

Both of these schemes require the high-precision implantation of individual donor atoms in a silicon substrate, which is extremely hard to achieve, especially in many-qubit circuits notwithstanding the extant is also an alternative NMR-based scheme for the quantum computation. Here an original method of the spin detection is proposed based on the interaction of the nonuniform magnetic field of a ferromagnetic particle **and (e&eb) the** spins subject to the resonance radio frequency field, the interaction thereof **causes the** mechanical oscillations of a micrometer cantilever, where the ferromagnetic particle is deposited. (see serial 48)

G_{40} : Category one of high-precision implantation of individual donor atoms in a silicon substrate, which is extremely hard to achieve, especially in many-qubit circuits notwithstanding the extant is also an alternative NMR-based scheme for the quantum computation. Here an original method of the spin detection is proposed based on the interaction of the nonuniform magnetic field of a ferromagnetic particle **and (e&eb) the** spins subject to the resonance radio frequency field, the interaction thereof

G_{41} : Category two of high-precision implantation of individual donor atoms in a silicon substrate, which is extremely hard to achieve, especially in many-qubit circuits notwithstanding the extant is also an alternative NMR-based scheme for the quantum computation. Here an original method of the spin detection is proposed based on the interaction of the nonuniform magnetic field of a ferromagnetic particle **and (e&eb) the** spins subject to the resonance radio frequency field, the interaction thereof

G_{42} : Category three of high-precision implantation of individual donor atoms in a silicon substrate, which is extremely hard to achieve, especially in many-qubit circuits notwithstanding the extant is also an alternative NMR-based scheme for the quantum computation. Here an original method of the spin detection is proposed based on the interaction of the nonuniform magnetic field of a ferromagnetic particle **and (e&eb) the** spins subject to the resonance radio frequency field, the interaction thereof

T_{40} : Category one of spins subject to the resonance radio frequency field, the interaction thereof causes **the** mechanical oscillations of a micrometer cantilever, where the ferromagnetic particle is deposited. (see serial 48)

T_{41} : Category two of spins subject to the resonance radio frequency field, the interaction thereof causes **the** mechanical oscillations of a micrometer cantilever, where the ferromagnetic particle is deposited. (see serial 48)

T_{42} : Category three of spins subject to the resonance radio frequency field, the interaction thereof causes **the** mechanical oscillations of a micrometer cantilever, where the ferromagnetic particle is deposited. (see serial 48)

Module Nine

The possibility of the proposed readout procedure is also questionable. If a single electron is used as qubit, then its readout **is (e) complemented by** the electron removal. This complementation of and augmentation thereof of the single electron used as a qubit by the electron removal, obviously, **creates a** defect in the Wigner crystal and can strongly disturb the neighbor electrons which can be **overcome by (e) using** a cluster of electrons as a single qubit

G_{44} : Category one of possibilities of the proposed readout procedure is also questionable. If a single electron is used as qubit, then its readout **is (e) complemented by** the electron removal. This complementation of and augmentation thereof of the single electron used as a qubit by the electron removal, obviously

G_{45} : Category two of possibilities of the proposed readout procedure is also questionable. If a single electron is used as qubit, then its readout **is (e) complemented by** the electron removal. This complementation of and augmentation thereof of the single electron used as a qubit by the electron removal, obviously

G_{46} : Category three of possibilities of the proposed readout procedure is also questionable. If a single electron is used as qubit, then its readout **is (e) complemented by** the electron removal. This complementation of and augmentation thereof of the single electron used as a qubit by the electron removal, obviously

T_{44} : Category one of defect in the Wigner crystal and can strongly disturb the neighbor electrons which can be **overcome by (e) using** a cluster of electrons as a single qubit

T_{45} : Category two of defect in the Wigner crystal and can strongly disturb the neighbor electrons which can be **overcome by (e) using** a cluster of electrons as a single qubit

T_{46} : Category three of defect in the Wigner crystal and can strongly disturb the neighbor electrons which can be **overcome by (e) using** a cluster of electrons as a single qubit

SECTION SEVENTY EIGHT

Perception: Nosce Te Ipsum :Know Thyself

Abstract: Many epithets and alliterations and logomachies' have been used to describe the perception by poets and authors. There is an ensorcelled freneticness and nonlinear frenzy, an ergodic erythrism when it comes to deliberation, debate and discussion of the perception and it's wide ranging ramifications and repercussions on the human mind and hence upon the physiology. Example of Beta movement often confused with phi phenomenon, in which a succession of still images **gives the** illusion of a moving ball. Two or more stimuli that are switched on and off in alternation **can produce** two different motion percepts. The first, demonstrated in the figure to the right is "Beta movement", **often used in** billboard displays, in which an object is perceived as moving when, in fact, a series of stationary images is being presented. This is also termed "apparent motion" and **is the basis of** movies and television. However, at faster alternation rates, and if the distance between the stimuli is just right, an illusory "object" the same colour as the background **is seen** moving between the two stimuli and alternately occluding them. This is called the phi phenomenon and is an example of "pure" motion detection uncontaminated, as in Beta movement, by form cues. This pure motion perception is referred to as "first-order" motion perception **and is mediated** by relatively simple "motion sensors" in the visual system, that have evolved to detect a change in luminance at one point on the retina **and correlate** it with a change in luminance at a neighbouring point on the retina after a short delay. Sensors that work this way have been referred to as either Hassenstein-Reichardt detectors after the scientists Bernhard Hassenstein and Werner Reichardt, who first modeled them, motion-energy sensors, or Elaborated Reichardt Detectors. These sensors detect motion by spatio-temporal correlation and are plausible models for how the visual system may detect motion. There is still considerable debate regarding the exact nature of this process. Second-order motion **is** motion in which the moving contour is defined by contrast, texture, flicker or some other quality that does not result in an increase in luminance or motion energy in the Fourier spectrum of the stimulus. There is much **evidence to suggest** that early processing of first- and second-order motion is carried out by separate pathways. Second-order mechanisms have poorer temporal resolution and are low-pass in terms of the range of spatial frequencies to

which they respond. Second-order motion **produces a** weaker motion aftereffect unless tested with dynamically flickering stimuli. First and second-order signals appear to be fully combined at the level of Area V5/MT of the visual system. The grating appears to be moving down and to the right, perpendicular to the orientation of the bars. But it could be moving in many other directions, such as only down, or only to the right. It is impossible to determine unless the ends of the bars become visible in the aperture. Each neuron in the visual system **is sensitive to** visual input in a small part of the visual field, as if each neuron is looking at the visual field through a small window or aperture. The motion direction of a contour is ambiguous, **because the** motion component parallel to the line cannot be inferred based on the visual input. This means that a variety of contours of different orientations moving at different speeds can cause identical responses in a motion sensitive neuron in the visual system. Individual neurons early in the visual system (V1) respond to motion **that occurs** locally within their receptive field. Because each local motion-detecting neuron will suffer from the aperture problem, the estimates from many neurons need to be integrated into a global motion estimate. This appears to occur in Area MT/V5 in the human visual cortex. Having extracted motion signals (first- or second-order) from the retinal image, the visual system must **integrate those** individual local motion signals at various parts of the visual field into a 2-dimensional or global representation of moving objects and surfaces. Further processing is required to disambiguate true "global motion" direction. (Wikipedia) visual perception of motion patterns characteristic of living organisms in locomotion and motion patterns in animals and men are termed here as biological motion which are **characterized by** a far higher degree of complexity than the patterns of simple mechanical motions usually studied in our laboratories. More often than not, visual information from biological motion and from the corresponding figurative contour patterns (the shape of the body) **is intermingled**. A method for studying information from the motion pattern per se **without** interference with the form aspect was devised. In short, the motion of the living body was **represented by** a few bright spots describing the motions of the main joints. It is found that 10–12 such elements in adequate motion combinations in proximal stimulus **evoke a** compelling impression of human walking, running, dancing, etc. The kinetic-geometric model for visual vector analysis originally developed in the study of perception of motion combinations of the mechanical type **was applied to** these biological motion patterns. The validity of this model in the present context was experimentally tested and the results turned out to be highly positive. **(June 1973, Volume 14, Issue 2, Visual perception of biological motion and a model for its analysis Gunnar Johansson)**

INTRODUCTION—VARIABLES USED

Default perception of high-speed motion Mark Wexler et al

When human observers are exposed to even slight motion signals followed by brief visual transients—stimuli containing no detectable coherent motion signals—they perceive large and salient illusory jumps. This visually striking effect, which we call “high phi,” challenges well-entrenched assumptions about the perception of motion, namely the minimal-motion principle and the breakdown of coherent motion perception with steps above an upper limit called dmax. Our experiments with transients, such as texture randomization or contrast reversal, show that the magnitude of the jump depends on spatial frequency and transient duration—but not on the speed of the inducing motion signals—and the direction of the jump depends on the duration of the inducer. Jump magnitude is robust across jump directions and different types of transient. In addition, when a texture is actually displaced by a large step beyond the upper step size limit of dmax, a breakdown of coherent motion perception is expected; however, in the presence of an inducer, observers again perceive coherent displacements at or just above dmax. In summary, across a large variety of stimuli, we find that when incoherent motion noise is preceded by a small bias, instead of perceiving little or no motion—as suggested by the minimal-motion principle—observers perceive jumps whose amplitude closely follows their own dmax limits

Motion perception revisited: High Phi effect challenges established motion perception assumptions

April 23, 2013 by Stuart Mason Dambrot in Neuroscience Enlarge A schematic illustration of the basic high-phi stimuli. Animated versions may be seen in demonstrations 1–4.

(1) During the first phase, the inducer, a random texture is slowly rotated, very briefly or for a longer duration. This inducer is followed by the transient, which may be a replacement of the texture by another, uncorrelated random texture (once or several times); a sudden jump of the original texture by an amount large enough that it cannot be detected; or an inversion of the texture contrast. (Other transients also lead to the effect: see text.) The **transient leads** to the illusory perception of a very rapid rotation, either forward (with respect to the direction of the inducer)—provided the inducer is very brief—or backward—for longer inducers. Copyright © PNAS, doi:10.1073/pnas.1213997110 (Medical Xpress)

(2) Optical illusions abound in human visual perception, **as (e) demonstrated by** the following well-known examples. Although many are static illusions, motion illusions also occur. Recently, scientists at Université Paris Descartes and Centre National de la Recherche Scientifique, Paris, University of Reading, United Kingdom, and Kyushu University, Japan discovered and investigated a new illusory motion effect, termed high phi by the authors, in which **we (e) perceive** conspicuous large illusory jumps when presentation of motion signals are followed by brief visual stimuli free of detectable motion signals.

(3) The researchers found that the size of the illusory jump does not depend on the speed of the motion signals presented, but rather **depend on (=)** on spatial frequency and transient duration while jump duration depends on motion signal duration.

(4) The study's authors conclude that their findings demonstrate that existing explanations for this illusion – namely, **the (e) loss of** coherent motion perception above an upper limit and the preference for minimal motion – are incomplete at best.

(5) Lead researcher Mark Wexler describes some of the challenges he and his colleagues – Andrew Glennerster, Patrick Cavanagh, Hiroyuki Ito, and Takeharu Seno – encountered in conducting their study. "We had the idea that these illusory jumps are related to dmax, the supposed upper speed limit on the steps **that(eb) leads to** motion perception, varies between individuals, and must be measured using random textures," Wexler tells Medical Xpress.

(6) "For displacements **below (eb)** dmax you're supposed to see the motion more or less correctly," he explains, "while for displacement above dmax you're just supposed to see noise – and the latter also turns out to be false." (These illusory jumps are demonstrated in an online supplement to the paper.1)

(7) Interestingly, the researchers discovered the illusion **as (=) a** bug in a computer program whose purpose was to do something else. "The easy thing to do in those kinds of circumstances is to correct the bug and move on," Wexler comments, "and ignore how strange the effect of the bug actually is. Our key insight was not to move on."

(8) According to Wexler, the one finding in motion perception that everyone agrees with, for at least 100 years, **is (=) the** minimal-motion principle.

(9) "The minimal-motion principle **states that** whenever a stimulus is ambiguous and compatible with more than one motion – as it nearly always is – the brain is supposed to prefer the smallest, slowest motion, including stand-still, that is compatible with the stimulus," he explains. (In fact, he illustrates, many computer vision systems are built around this principle, and neuroscientists have verified it by recording signals from primate neurons.)

Temporal covariance model of human motion perception Jan P. H. van Santen and George Sperling JOSA A, Vol. 1, Issue 5, pp. 451-473 (1984) http://dx.doi.org/10.1364/JOSAA.1.000451

(1) Authors propose a model of **direction-sensitive units in human vision**. It is a modified and

elaborated version of a model by Reichardt [Z. Naturforsch. Teil B 12, 447 (1957)]. The model is **applied (e&eb) to** threshold experiments in which subjects view adjacent vertical bars with independently (typically sinusoidally), temporally modulated luminances.

(2) The subject must report whether the patterns moved to the left or to the right. According to the model, a basic motion-detecting unit **consists of** two subunits tuned to opposite directions. Each performs a spatial and temporal linear filtering of its input; outputs of the filters are multiplied, and the multiplied output **is (e&eb) integrated** (for a time that is long relative to the modulation period). The model's output consists of the difference between the subunit outputs. Direction of movement is indicated by the sign of the model output.

(3) Mathematical analysis of the model yielded several predictions that were confirmed experimentally. Specifically, authors find that (1) performance with complex patterns **can be predicted by** spatiotemporal Fourier analysis that results in the segregation and linear addition in the output for different temporal frequencies; (2) under special conditions, performance **depends on the** product of adjacent bar amplitudes, offering strong support for the multiplication principle; (3) performance is unaffected **by** addition of stationary patterns; and (4) addition of homogeneous flicker normally **produces** no effect but under special conditions reverses perceived direction. These and other results confirm our model and reject several other models, including Reichardt's original model.

NOTATION

Module One

Human mind is corrupt. There are dualities in nature. Nature is corrupt. God is corrupt in the sense both Brahman-Anti Brahman concept introduced. Human beings are always simulating situations thinking about the past or future. They are always want to achieve "ends" irrespective of the "means" There are dualities of space time, mass energy, perception reality, consciousness-unconsciousness,. So we suggest machine rising is the only way to control the corruption, widespread surveillance for the large scale cheating, garnering of money even in places like US where the 95 percent is the money is with 5 percent of people. Only way to keep things like murder, mayhem, plunder, pillage, nemesis, apocalypse, is to keep a watch on human mind and nature. Machine must control man. It must be able to know the perception of both the mind of man and God. Crimes continue unless the mind is watched and what the person is thinking. Differential of what is innocuous and the serious can be made out. How to know the mind of God. Nature's general ledger with quantum computation and maintenance of "progressive" of the general ledger both individual and that of nature seem would probably offer the solution.

G_{13} : Category one of mind of man and god(Brahman-AntiBrahman agency)

G_{14} : Category two of mind of man and god(Brahman-AntiBrahman agency)

G_{15} : Category three of mind of man and god(Brahman-AntiBrahman agency)

T_{13} : Category one of Quantum computers and creation of technical singularities

Vinge predicted four ways the singularity could occur:

(1) The development of computers that are "awake" and superhumanly intelligent
(2) Large computer networks (and their associated users) may "wake up" as a superhumanly intelligent entity
(3) Computer/human interfaces may become so intimate that users may reasonably be considered

superhumanly intelligent

(4) Biological science may find ways to improve upon the natural human intellect

T_{14} : Category two of Quantum computers and creation of technical singularities

T_{15} : Category three of Quantum computers and creation of technical singularities

Module Two

G_{16} : Category one of result of the module one

G_{17} : Category two of result of the module one

G_{18} : Category three of result of the module one

T_{16} : Category one of Dharma(axiomatic predications, postulation alcovishness, doctrines, dictums ,notions, nostrums as indoctrinated to every individual by peers: individual general ledger) and Karma (action to performed without any duress and compulsion .Freewill as is expatiated as the indiscretion of the discretionary vested and reposed in the individual)

T_{17} : Category two of Dharma(axiomatic predications, postulation alcovishness, doctrines, dictums ,notions, nostrums as indoctrinated to every individual by peers: individual general ledger) and Karma (action to performed without any duress and compulsion .Freewill as is expatiated as the indiscretion of the discretionary vested and reposed in the individual)

T_{18} : Category three of Dharma(axiomatic predications, postulation alcovishness, doctrines, dictums ,notions, nostrums as indoctrinated to every individual by reference groups and peers: individual general ledger) and Karma (action to performed without any duress and compulsion .Freewill as is expatiated as the indiscretion of the discretionary vested and reposed in the individual)

Module three

G_{20} : Category one of time

G_{21} : Category two of time

G_{22} : Category three of time

T_{20} : Category one of space

T_{21} : Category two of space

T_{22} : Category three of space

Module four

Three Stages and Two Systems of Visual Processing George Sperling Human Information Processing Laboratory, Department of Psychology, New York University, New York, NY 10003

Three stages of visual processing determination of how internal noise appears to an external observer. :Light adaptation; contrast gain control; Post sensory/decision stage.

(1) Dark noise occurs prior to adaptation, **determines** dark-adapted absolute thresholds, and mimics stationary external noise.
(2) Sensory noise occurs after dark adaptation, **determines** contrast thresholds for sine gratings and similar stimuli, and mimics external noise that increases with mean luminance.
(3) Post sensory noise **incorporates** perceptual, decision, and mnemonic processes.

(4) Post sensory noise **incorporates** perceptual, decision, and mnemonic processes occurs after contrast-gain control **and (=) mimics** external noise that increases with stimulus contrast (i.e., multiplicative noise).
(5) Dark noise and sensory noise are frequency specific and primarily affect weak signals. Only post sensory noise significantly **affects to** strong signals, and it has constant power over a wide spatial frequency range in which sensory noise varies enormously.
(6) Two parallel perceptual regimes jointly serve human object recognition and motion perception: a first-order linear (Fourier) regime that computes relations directly from luminances and a second-order nonlinear (non Fourier) rectifying regime **that uses the** absolute value of detector outputs.
(7) When objects or movements are defined by high spatial frequencies (i.e., texture carrier frequencies whose wavelengths are small compared to the object size), the responses of high-frequency receptors are demodulated by rectification **to (eb) facilitate** discrimination at the next hierarchical processing levels.
(8) Rectification **sacrifices (e) the** statistical efficiency (noise resistance) of the first-order regime for efficiency of connectivity and computation

CARE: THERE ARE EIGHT MODELS IN THIS MODULE DUE TO SPATIAL CONSTRAINTS. READER IS REQUESTED TO IDENTIFYTHE PARAMETERS BEFORE ARRIVING AT PARTICULAR CONCLUSION

G_{24} : Category one of Light adaptation; contrast gain control; Post sensory/decision stage; Dark noise occurs prior to adaptation; Sensory noise occurs after dark adaptation; perceptual, decision, and mnemonic processes; **mimics** external noise that increases with stimulus contrast (i.e., multiplicative noise); Dark noise and sensory noise are frequency specific and primarily affect weak signals. Only post sensory noise significantly; first-order linear (Fourier) regime that computes relations directly from luminances and a second-order nonlinear (non Fourier) rectifying regime **that uses the** absolute value of detector outputs; When objects or movements are defined by high spatial frequencies (i.e., texture carrier frequencies whose wavelengths are small compared to the object size), the responses of high-frequency receptors are demodulated by rectification;

G_{25} : Category two of Light adaptation; contrast gain control; Post sensory/decision stage; Dark noise occurs prior to adaptation; Sensory noise occurs after dark adaptation; perceptual, decision, and mnemonic processes; **mimics** external noise that increases with stimulus contrast (i.e., multiplicative noise); Dark noise and sensory noise are frequency specific and primarily affect weak signals. Only post sensory noise significantly; first-order linear (Fourier) regime that computes relations directly from luminances and a second-order nonlinear (non Fourier) rectifying regime **that uses the** absolute value of detector outputs; When objects or movements are defined by high spatial frequencies (i.e., texture carrier frequencies whose wavelengths are small compared to the object size), the responses of high-frequency receptors are demodulated by rectification;

G_{26} : Category three of Light adaptation; contrast gain control; Post sensory/decision stage; Dark noise occurs prior to adaptation; Sensory noise occurs after dark adaptation; perceptual, decision, and mnemonic processes; **mimics** external noise that increases with stimulus contrast (i.e., multiplicative noise); Dark noise and sensory noise are frequency specific and primarily affect weak signals. Only post sensory noise significantly; first-order linear (Fourier) regime that computes relations directly from luminances and a second-order nonlinear (non Fourier) rectifying regime **that uses the** absolute value of detector outputs; When objects or movements are defined by high spatial frequencies (i.e., texture carrier frequencies whose wavelengths are small compared to the object size), the responses of high-frequency receptors are

demodulated by rectification;

T_{24} : Category one of visual processing determination of how internal noise appears to an external observer; dark-adapted absolute thresholds, and mimics stationary external noise; contrast thresholds for sine gratings and similar stimuli, and mimics external noise that increases with mean luminance; Post sensory noise; Post sensory noise **incorporates** perceptual, decision, and mnemonic processes occurs after contrast-gain control; strong signals, and it has constant power over a wide spatial frequency range in which sensory noise varies enormously; Two parallel perceptual regimes jointly serve human object recognition and motion perception discrimination at the next hierarchical processing levels; Rectification;;

T_{25} : Category two of visual processing determination of how internal noise appears to an external observer; dark-adapted absolute thresholds, and mimics stationary external noise; contrast thresholds for sine gratings and similar stimuli, and mimics external noise that increases with mean luminance; Post sensory noise; Post sensory noise **incorporates** perceptual, decision, and mnemonic processes occurs after contrast-gain control; strong signals, and it has constant power over a wide spatial frequency range in which sensory noise varies enormously; Two parallel perceptual regimes jointly serve human object recognition and motion perception; discrimination at the next hierarchical processing levels; Rectification;

T_{26} : Category three of visual processing determination of how internal noise appears to an external observer; dark-adapted absolute thresholds, and mimics stationary external noise; contrast thresholds for sine gratings and similar stimuli, and mimics external noise that increases with mean luminance; Post sensory noise; Post sensory noise **incorporates** perceptual, decision, and mnemonic processes occurs after contrast-gain control; strong signals, and it has constant power over a wide spatial frequency range in which sensory noise varies enormously; Two parallel perceptual regimes jointly serve human object recognition and motion perception; discrimination at the next hierarchical processing levels; Rectification;

Module five

Optical illusions abound in human visual perception, **as (e) demonstrated by** the following well-known examples. Although many are static illusions, motion illusions also occur. Recently, scientists at Université Paris Descartes and Centre National de la Recherche Scientifique, Paris, University of Reading, United Kingdom, and Kyushu University, Japan discovered and investigated a new illusory motion effect, termed high phi by the authors, in which **we (e) perceive** conspicuous large illusory jumps when presentation of motion signals are followed by brief visual stimuli free of detectable motion signals

G_{28} : Category one of **perceive** conspicuous large illusory jumps when presentation of motion signals are followed by brief visual stimuli free of detectable motion signals

G_{29} : Category two of **perceive** conspicuous large illusory jumps when presentation of motion signals are followed by brief visual stimuli free of detectable motion signals

G_{30} : Category three of **perceive** conspicuous large illusory jumps when presentation of motion signals are followed by brief visual stimuli free of detectable motion signals

T_{28} : Category one of Optical illusions abound in human visual perception, **as (e) demonstrated by** the following well-known examples. Although many are static illusions, motion illusions also occur. Recently, scientists at Université Paris Descartes and Centre National de la Recherche Scientifique, Paris, University of Reading, United Kingdom, and Kyushu University, Japan discovered and investigated a new illusory motion effect, termed high phi by the authors, in which

T_{29} : Category two of Optical illusions abound in human visual perception, **as (e) demonstrated by** the

following well-known examples. Although many are static illusions, motion illusions also occur. Recently, scientists at Université Paris Descartes and Centre National de la Recherche Scientifique, Paris, University of Reading, United Kingdom, and Kyushu University, Japan discovered and investigated a new illusory motion effect, termed high phi by the authors, in which

T_{30} : Category three of Optical illusions abound in human visual perception, **as (e) demonstrated by** the following well-known examples. Although many are static illusions, motion illusions also occur. Recently, scientists at Université Paris Descartes and Centre National de la Recherche Scientifique, Paris, University of Reading, United Kingdom, and Kyushu University, Japan discovered and investigated a new illusory motion effect, termed high phi by the authors, in which

Module six

The researchers found that the size of the illusory jump does not depend on the speed of the motion signals presented, but rather **depend on** spatial frequency and transient duration while jump duration depends on motion signal duration

G_{32} : Category one of the size of the illusory jump does not depend on the speed of the motion signals presented, but rather **depend on**

G_{33} : Category two of the size of the illusory jump does not depend on the speed of the motion signals presented, but rather **depend on**

G_{34} : Category three of the size of the illusory jump does not depend on the speed of the motion signals presented, but rather **depend on**

T_{32} : Category one of spatial frequency and transient duration while jump duration depends on motion signal duration

T_{33} : Category two of spatial frequency and transient duration while jump duration depends on motion signal duration

T_{34} : Category three of spatial frequency and transient duration while jump duration depends on motion signal duration

Module seven

Lead researcher Mark Wexler describes some of the challenges he and his colleagues – Andrew Glennerster, Patrick Cavanagh, Hiroyuki Ito, and Takeharu Seno – encountered in conducting their study. "We had the idea that these illusory jumps are related to dmax, the supposed upper speed limit on the steps **that(eb) leads to** motion perception, varies between individuals, and must be measured using random textures," Wexler tells Medical Xpress

G_{36} : Category one of illusory jumps are related to dmax, the supposed upper speed limit on the steps

G_{37} : Category two of illusory jumps are related to dmax, the supposed upper speed limit on the steps

G_{38} : Category three of illusory jumps are related to dmax, the supposed upper speed limit on the steps

T_{36} : Category one of motion perception, varies between individuals, and must be measured using random textures

T_{37} : Category two of motion perception, varies between individuals, and must be measured using random textures

T_{38} : Category three of motion perception, varies between individuals, and must be measured using random textures

Module eight

Interestingly, the researchers discovered the illusion **as a** bug in a computer program whose purpose was to do something else. "The easy thing to do in those kinds of circumstances is to correct the bug and move on," Wexler comments, "and ignore how strange the effect of the bug actually is. Our key insight was not to move on."

G_{40} : Category one of bug in a computer program whose purpose was to do something else. "The easy thing to do in those kinds of circumstances is to correct the bug and move on," Wexler comments, "and ignore how strange the effect of the bug actually is

G_{41} : Category two of bug in a computer program whose purpose was to do something else. "The easy thing to do in those kinds of circumstances is to correct the bug and move on," Wexler comments, "and ignore how strange the effect of the bug actually is

G_{42} : Category three of bug in a computer program whose purpose was to do something else. "The easy thing to do in those kinds of circumstances is to correct the bug and move on," Wexler comments, "and ignore how strange the effect of the bug actually is

T_{40} : Category one of illusion There are millions types of illusions and million human systems that are being investigated or could be investigated. Properties thereof are the bastion of the stratification doxa.

T_{41} : Category two of illusion

T_{42} : Category three of illusion

Module Nine

According to Wexler, the one finding in motion perception that everyone agrees with, for at least 100 years, **is the** minimal-motion principle. "The minimal-motion principle **states that** whenever a stimulus is ambiguous and compatible with more than one motion – as it nearly always is – the brain is supposed to prefer the smallest, slowest motion, including stand-still, that is compatible with the stimulus," he explains. (In fact, he illustrates, many computer vision systems are built around this principle, and neuroscientists have verified it by recording signals from primate neurons.)

G_{44} : Category one of minimal-motion principle. "The minimal-motion principle **states that** whenever a stimulus is ambiguous and compatible with more than one motion – as it nearly always is – the brain is supposed to prefer the smallest, slowest motion, including stand-still, that is compatible with the stimulus," he explains. (In fact, he illustrates, many computer vision systems are built around this principle, and neuroscientists have verified it by recording signals from primate neurons.)

G_{45} : Category two of minimal-motion principle. "The minimal-motion principle **states that** whenever a stimulus is ambiguous and compatible with more than one motion – as it nearly always is – the brain is supposed to prefer the smallest, slowest motion, including stand-still, that is compatible with the stimulus,"

he explains. (In fact, he illustrates, many computer vision systems are built around this principle, and neuroscientists have verified it by recording signals from primate neurons.)

G_{46} : Category three of minimal-motion principle. "The minimal-motion principle **states that** whenever a stimulus is ambiguous and compatible with more than one motion – as it nearly always is – the brain is supposed to prefer the smallest, slowest motion, including stand-still, that is compatible with the stimulus," he explains. (In fact, he illustrates, many computer vision systems are built around this principle, and neuroscientists have verified it by recording signals from primate neurons.)

T_{44} : Category one of motion perception that everyone agrees with

T_{45} : Category two of motion perception that everyone agrees with

T_{46} : Category three of motion perception that everyone agrees with

.

SECTION SEVENTY EIGHT

High Phi Effect Challenges :Rex Non Potest Peccare: The King Can Do No Wrong

Abstract: Predicational anteriorities, ontological consonance, primordial exactitude, discrimination jurisprudence, phenomenological correlates, apocryphal aneurism, atrophied asseverations, anamensial alienisms, anchorite aperitif, Arcadian Atticism ,functional topology, dispossession and saturnine flows, coruscation locomotion, reiterations, preservations and conservations ,hypostatized significations, substitutional designation localities, engendered propositions, presupposition resemblances, contiguous similarities, of Human consciousness, the human soul, the human mind, human subjective feelings have been a matter of concern, not only for philosophers and theologians, but recently also for neuroscientists, physicists and others. Our approach to understanding this problem is **based on the** fact that even the simplest brain functions **depend on the** activity of an enormous number of neurons, on their synaptic connections **and on** associated ionic and electrical events. The synaptic delay in each of those synapses is at least 0.5 ms and therefore the parallel and **serial interactions** between millions of neurons would take a very long time, too long for the individual's adequate interaction with the environment. Therefore, there must be some other mechanism **governing the interactions** of large numbers of neurons, located even in remote parts of the brain. However, the neuronal function - with a spreading of depolarization's, Hyperpolarization and repolarizations, graded and ungraded electrical potentials, ionic movements and small local electrical fields - **creates a** unique and very complicated system of the movements of subatomic particles. When moving from one position to another, each electron fills a large space and its precise position cannot be exactly determined. Thus brain function **depends on** the movements of an enormous number of electrons **which influence** each other at the subatomic level, even though their position cannot be determined. Therefore, beside relatively slow "classical" electrochemical interactions, rapid quantum interactions originating in functioning cell membranes **may participate** in many, perhaps in all, brain functions. Together, all the moving electrons **produce a** non-local system which we call the Real Human Soul, RHS, which is **created by the** functioning neurons and, at the same time, can also influence other neurons. Thus, **it creates a** connection between all functioning parts of the brain. The brain then functions **as a** unified system in which everything is interconnected and is able to interact. Viewed thus, the brain functions **as a** quantum computer. This system, the RHS, **is not identical** with consciousness. Only under certain

conditions does some area of the brain **create** subjective consciousness, which may be **one of the products** of the RHS. Subjective consciousness is probably **based o**n simple particle communication, their .proto-consciousness., but is much more complex**, due to** the neuronal analysis of sensory input and other cognitive functions of the CNS. There are several definitions of consciousness, which usually **depend on** the philosophical views of their authors. Let us use a simple definition of consciousness, as found on the Internet (http://www.selfknowledge.com/19677 .htm): .The state of being conscious; **knowledge of** one's own existence, sensations, mental operations, acts, etc. Consciousness is thus, on the one hand, is **the** recognition of the mind or .ego. **This is a modified English translation of a paper published in Czech in Ceskoslovenska fyziologie, 2005, vol.54, pp.26 - 31.A quantum hypothesis of brain function and consciousness Stanislav Reinis et al (emphasis and minor alterations mine)** We perceive the shapes and material properties of objects quickly and reliably despite the complexity and objective ambiguities of natural images. Typical images are highly complex **because they** consist of many objects embedded in background clutter. Moreover, the image features of an object are extremely variable and ambiguous **owing to the** effects of projection, occlusion, background clutter, and illumination. The very success of everyday vision **implies** neural mechanisms, yet to be understood, that discount irrelevant information and organize ambiguous or noisy local image features into objects and surfaces. Recent work in Bayesian theories of visual perception **has shown how** complexity may be managed and ambiguity resolved through the task-dependent, probabilistic integration of prior object knowledge with image features. **Object Perception as Bayesian Inference Annual Review of Psychology Vol. 55: 271-304 (Volume publication date February 2004) First posted online on October 6, 2003 DOI: 10.1146/annurev.psych.55.090902.142005 Daniel Kersten et al Giuseppe Castagnoli** provide an alternative way of seeing quantum computation with an idealized classical problem **solving** machine whose coordinates are submitted to a nonfunctional relation **representing** all the problem constraints; moving an input part, reversibly and nondeterministic ally **produces a** solution through a many body interaction. The machine can be considered **the** many body generalization of another perfect machine, the bouncing ball model **of** reversible computation. The mathematical description of the machine's motion, as it is, is applicable to quantum problem solving, an extension of the quantum algorithms **that comprises the** physical representation of the interdependence between the problem and the solution. The configuration space of the classical machine **is replaced** by the phase space of the quantum machine. The relation between the coordinates of the machine parts **now applies** to the populations of the reduced density operators of the parts of the computer register throughout state vector reduction. Thus, reduction **produces the** solution of the problem under a nonfunctional relation representing the problem-solution interdependence. At the light of this finding, the quantum speed up **turns out to be** "**precognition"** of the solution, namely the reduction of the initial ignorance of the solution due to backdating, to before running the algorithm, a part of the state vector reduction on the solution (a time-symmetric part in the case of unstructured problems); as such, **it is bounded by** state vector reduction through an entropic inequality. The computation mechanism under discussion might also explain the wholeness appearing in the introspective analysis of perception**. International Journal of Theoretical Physics August 2008, Volume 47, Issue 8, The Mechanism of Quantum Computation Giuseppe Castagnoli**

INTRODUCTION—VARIABLES USED

This is a modified English translation of a paper published in Czech in Ceskoslovenska fyziologie, 2005, vol.54, pp.26 - 31.A quantum hypothesis of brain function and consciousness Stanislav Reinis et al (emphasis and minor alterations mine)

(1) Describing the mechanisms which cause certain events in the brain to be subjectively perceived is the hard problem of neuroscience. Consciousness **cannot (e) b**e reduced to neuronal firing and neuronal interactions.

(2) On one hand, there are brain activities that can be objectively observed, recorded and measured by an external investigator. On the other hand, there exists our private, subjective perception of some of these events. Consciousness **is (=) our** primary reality; through it, we perceive ourselves and our environment; we plan and accomplish our actions, evaluate them, think about them, record them.
(3) Some believe that consciousness **is(e) an** emergent property of brain activity
(4) Others assume that there is a duality of matter **and** spirit, and that there exists an immaterial principle, a homunculus, **controlling** brain functions. But there might be some other possibilities as well.

The human brain is composed of billions of neurons and glia cells. There is an extracellular space between them, filled with fluid. This space is rather minimal, comprising about 5% of brain volume. The neurons communicate one with another through at least nine mechanisms:

(5) Classical synaptic transmission, when the nerve impulse **passes(e&eb) between** neurons through synapses, with a synaptic delay of at least 0.5 ms
(6) Diffusion of neurotransmitters and neuropeptide from nerve endings and from axonal varicosities into the extracellular space, where these substances **may influence** a larger number of neurons and glia cells at once
(7) Neuromodulation, where neuropeptide **influence (e&eb)** synaptic transmission generated by neurotransmitters
(8) Hormones produced elsewhere in the body, **which (e&eb) influence** target cells that possess the corresponding hormonal receptors
(9) Transneuronally **transmitted (e&eb)** proteins responsible for trophic interactions
(10) Ionic **changes (e&eb)** in the extracellular space
(11) Ephaptic transmission, **accomplished (e) by** the direct injection of electrical current from one cell into another;
(12) Spreading of small electrical fields **around (e&eb)** the excited neurons
(13) Automatic excitation of the pacemaker neurons **by (e)** metabolic processes.
(14) All these are local phenomena belonging in the area of classical physics and its ramifications. The result of neuronal activation **is (=) an** action potential generated by movements of electrons and ions, e.g. of potassium, calcium and sodium.
(15) A neuron producing an action potential usually **requires (e) a**t least ten synaptic inputs from other neurons to reach its firing level.
(16) Therefore, the function of the central nervous system **depends on** many serial and parallel interactions of masses of individual neurons.
(17) Each neuron is connected to hundreds and thousands of other neurons. In the brain, we may observe convergence and divergence, feedbacks and circulating nerve impulses. The reverberating neuronal circuits may be rather long, lasting up to one second (Reinis, 1997). All these events **slow (e) down the** functioning of neuronal networks containing millions of neurons, so such sequences of neuronal firing cannot accomplish the function of more complex neuronal systems that are expected to respond in a real, sufficiently short time. Despite these imperfections, the human brain is a uniquely complex system of electrochemical activities unlike anything in the known universe.
(18) For this reason, we must search for another, more rapid mechanism of neuronal interactions **to explain** the speed of some fast reactions in the nervous system. Synaptic transmission and axonal transfer of nerve impulses are too slow **to (eb) organize** coordinated activity in large areas of the central nervous system. Numerous observations confirm this view.
(19) The same organizing forces which **gave (eb) a form** to nature in all its forms are also responsible for the structure of our mind. (W**erner Heisenberg**)
(20) "It is very difficult for us to take stock of the fact that the localization of the personality, of the conscious mind, inside the body is only symbolic, **just (e) an aid** for practical use." **(Erwin**

Schroedinger)

(21) Schroedinger is probably not correct in his belief that consciousness is located outside the brain, somewhere in the universe. It is, we assume, the human (and perhaps some other as well) brain that produces it, and the contents of our conscious activities, thoughts, memories and intentions depend on sensory input and a large number of coordinated neuronal interactions. Consciousness cannot be reduced to neuronal activity, the functions of neurotransmitters and neuronal spikes. It is, however, **influenced (e&eb) by** mutual neuronal interactions mediated by synaptic and non-synaptic interactions.

(22) There are three advantages to this quantum approach: First, that the temporary connection of various systems might be sufficiently fast; second, that the connections may be quickly terminated; and third, that quantum interactions may also help to **e**xplain subjective consciousness.

(23) Submicroscopic particles may penetrate seemingly solid matter. They may pass at a supraliminal speed and their movement **may be (e&eb) subject to** non-locality as described by David Bohm (1951). This process may also take place in the brain.

(24) Each neuron is composed of the nerve cell body, perikaryon, with the attached axon with its branches, telodendria, and with the dendrites. Synapses at the end of telodendria **connect them with** dendrites, perikarya or axons of other neurons. Several synapses must usually be **activated to (eb) achieve** production of a nerve impulse in the axon hillock. When a nerve impulse reaches the nerve ending, calcium ions enter the synaptic knob and elicit the release of synaptic vesicles containing a neurotransmitter. The released neurotransmitter **activates t**he postsynaptic membrane and elicits the formation of EPSPs and IPSPs. These electrical waves spread over the surface membrane of the neuron decrementally. When they reach the axon hillock, they may produce a nerve impulse which moves along the axon, using circulating currents stimulating the axon toward its end, the synaptic knob. Each neuron is therefore a sufficient source of moving electrons which, as quantum particles fill the space, may interact with other electrons. The neuronal role as a generator of particles changes each nanosecond. There are at least ten billion neurons in the brain, all of them producing scores of particles. All electrical phenomena in the neuron must be considered, those taking place on the surface of the brain cells, in the cell membrane, but also inside, in microtubules and mitochondria, those involved in the conformation of protein molecules etc. All of them together **represent (eb) a** powerful source of subatomic particles contacting, on a quantum level, particles generated by other neurons.

(25) How is the brain protected against errors coming from its environment, both within the body and outside? How are the errors corrected? Is there any similarity in the quantum error correction proposed by researchers involved in quantum computer research and correction of errors in the brain? How the classical neuronal function is related to the quantum computation events, in particular in view of the fact that a quantum computer may be formed just by a single molecule (Chuang et al, 1995) Give the example of the general ledger of a bank wherein there is continuous recordial evidence of transactions' and they are tallied at the end. Brain also loses memory, and there may be what may be called "balancing error".

(26) Another open question remains: What is the difference between the conscious **and the** unconscious processes in the brain?

(27) Studies by Merikle et al. (e.g., Visser and Merikle, 1999) indicate that the difference between them is not very sharp; the transitions between the two are rapid, temporary **and dependent on** emotions. It is, therefore, possible that the difference between the conscious and the unconscious **is(=)** quantitative. The more synaptic and nonsynaptic events there are in a certain area of the brain, the higher the possibility that the event itself enters the consciousness. As an example, the visual pathway from the eye into the lateral geniculate **body is** unconscious, area V1 of the cortex may be conscious under certain circumstances, and events in areas V2 and higher are conscious. This probably **depends on** the number of acting elements. The human cerebral cortex **represents** about

75% of the brain mass, and most conscious events may take place within it. On the other hand, at a low level of firing and neuronal interactions, consciousness does not substantially exceed the level of protoconscious ness.

Optical illusions:

(1) Lead researcher Mark Wexler describes some of the challenges he and his colleagues – Andrew Glennerster, Patrick Cavanagh, Hiroyuki Ito, and Takeharu Seno – encountered in conducting their study. "We had the idea that these illusory jumps are related to dmax, the supposed upper speed limit on the steps **that leads to** motion perception, varies between individuals, and must be measured using random textures," Wexler tells Medical Xpress. "For displacements below dmax you're supposed **to (e) see the** motion more or less correctly," he explains, "while for displacement above dmax **you're just (e) supposed to see** noise – and the latter also turns out to be false." (These illusory jumps are demonstrated in an online supplement to the paper.)

(2) Interestingly, the researchers discovered the illusion **as a** bug in a computer program whose purpose was to do something else. "The easy thing to do in those kinds of circumstances is to correct the bug and move on," Wexler comments, "and ignore how strange the effect of the bug actually is. Our key insight was not to move on."

(3) According to Wexler, the one finding in motion perception that everyone agrees with, for at least 100 years, **is (=) the** minimal-motion principle. "The minimal-motion principle **states that** whenever a stimulus is ambiguous and compatible with more than one motion – as it nearly always is – the brain is supposed to prefer the smallest, slowest motion, including stand-still, that is compatible with the stimulus," he explains. (In fact, he illustrates, many computer vision systems are built around this principle, and neuroscientists have verified it by recording signals from primate neurons.)

(4) "However," Wexler points out, "one consequence of the high phi effect **is (=) the** minimal-motion principle can be violated!

(5) When the stimulus is incompatible with any globally coherent motion, and therefore equally compatible with any motion, **people (e) perceive** not only a large jump, but the largest possible jump that they can perceive. This maximum jump is the one that steps by dmax, which acts as the speed limit on motion perception."

(6) Another principle that seems to be violated by the high phi illusion, according to Wexler**, is (=)** dmax itself. "Below dmax, steps should be more or less seen as what they are – and as can be seen in demonstration three, this is what happens. On the other hand, says Wexler, "above dmax you're supposed **to (e) perceive** noise, not motion – but this is not what actually occurs." Rather, you perceive the high phi jump, as can also be seen in demonstration three. "In one of our experiments," Wexler adds, "we showed that the amplitude of the jump is very closely **correlated with the** dmax limit, so that people who have higher dmax limits also see a larger high phi jump."

(7) It's known, Wexler points out, that the dmax limit **depends on** spatial frequency: the lower the frequency (that is, the larger the features in the stimulus) the higher the limit. "And indeed," he notes, "we found that the magnitude of the illusory jump depends on spatial frequency in exactly the same way: the lower the frequency, the farther the jump." This can be verified, Wexler adds, by viewing high phi demonstration six and demonstration seven.

(8) Discussing the finding that the direction of the jump **depends on** the duration of the inducing motion signals, Wexler notes, "We think that the preceding – that is, inducing – motion acts like a seed. For brief inducers, the motion itself acts as the seed, and the jump is experienced forwards with respect to the inducer. For longer inducers, vision begins **to adapt (e&eb) to** the motion – a result known as the motion aftereffect."

(9) Also known as the waterfall illusion, the motion aftereffect occurs when, after viewing a moving object for an extended period of time, and that object then becomes stationary, the object appears to

slowly move in the opposite direction. “Many people initially think that what we’ve found **is a (e) consequence of** adaptation to motion or the motion aftereffect,” he says. “If so, then it’s the fastest motion aftereffect known. We’ve measured that the illusory motion is 10-100 times faster than the inducing motion! We think that for motion inducers, the adaptation acts as the seed of the fast, backward jump.” (Brief and long inducers can be compared directly in demonstration nine.)

(10) Wexler also describes how their findings **relate(e&eb) to the** activity of neurons in of the primary visual cortex that respond to lines of a certain angle moving in one direction, as first described by Hubel and Wiesel (1959). “In the brain, motion detectors are sensitive to motion in a particular place – the receptive field – a particular direction, and usually a particular speed,” Wexler notes. “When faced with our stimulus, there can be many accidental matches at the local level. In one image there is a dark spot, for example, and in the next, uncorrelated image **there (eb) happens to be** dark spot just next to it.

(11) In that case, a local motion detector will react to this false match – so our stimulus actually **activates many** local motion detectors, but incoherently, in that all of these motion detectors are signaling different motions. The main point is that in all this incoherent mess the brain **finally (eb) prefers the** largest possible motion.”

(12) Commenting on other areas of research that might benefit from their study, Wexler cites computer vision. “The minimal-motion principle is enshrined in a lot of algorithms **for (eb) extracting** motion,” he concludes. “Our study shows that this principle **can be (e) violated.** Can we find a different way to extract motion?”

(13) “The dependence on transient duration – which can be clearly seen in demonstration five – is, to be completely honest, a mystery, but a very interesting one,” Wexler continues. “The amplitude of the jump is a very linear function of transient duration, at least for small durations. If some perceptual process goes linearly farther for longer durations, **then(eb)** something in the brain must be effectively rotating at constant speed. I have no idea what that something may be, but it’s an interesting challenge for the future.”This is the html version of the file http://www.dtic.mil/dtic/tr/fulltext/u2/a212670.pdf.

Motion perception revisited: High Phi effect challenges established motion perception assumptions April 23, 2013 by Stuart Mason Dambrot in Neuroscience Enlarge A schematic illustration of the basic high-phi stimuli. Animated versions may be seen in demonstrations 1–4.

(1) "However," Wexler points out, "one **consequence of the** high phi effect **is (=) the** minimal-motion principle can be violated!

(2) When the stimulus is incompatible with any globally coherent motion, and therefore equally compatible with any motion, people **perceive (e)** not only a large jump, but the largest possible jump that they can perceive.

(3) This maximum jump is the one that steps by dmax, which **acts (=) as the** speed limit on motion perception."

(4) Another principle that seems to be violated by the high phi illusion, according to Wexler, **is (=)** dmax itself.

(5) "Below dmax, steps should be more or less seen as what they are – and as can be seen in demonstration three, this is what happens. On the other hand, says Wexler, "above dmax you're supposed to **perceive (e)** noise, not motion – but this is not what actually occurs."

(6) Rather, you **perceive (e)** the high phi jump, as can also be seen in demonstration three. "In one of our experiments," Wexler adds, "we showed that the amplitude of the jump is very closely correlated with the dmax limit, so that people who have higher dmax limits also see a larger high phi jump."

(7) It's known, Wexler points out, that the dmax limit **depends o**n spatial frequency: the lower the frequency (that is, the larger the features in the stimulus) the higher the limit.

(8) "And indeed," he notes, "we found that the magnitude of the illusory jump depends on spatial frequency in exactly the same way: the lower the frequency, formation shall be of the farther the jump." This can be verified, Wexler adds, by viewing high phi demonstration six and demonstration seven. Discussing the finding that the direction of the jump depends on the duration of the inducing motion signals, Wexler notes, "We think that the preceding – that is, inducing – motion acts like a seed. For brief inducers, the motion itself **acts (=) as** the seed, and the jump is experienced forwards with respect to the inducer.

Three Stages and Two Systems of Visual Processing George Sperling New York University. Most of the paragraphs are deleted due to spatial constraints emphasis mine. Only part of the article is taken

(1) Bandpass filtering refers to the processing of images or sounds so that they **contain(e) only** a narrow range--typically one or two octaves--of component frequencies. In addition, bandpass filtering is used to **create** stimuli that stimulate only a small portion of the basilar membrane.
(2) By studying psychophysical responses to stimuli filtered in different bands, information processing mediated by each portion of the basilar membrane can be studied. In vision, the aim of bandpass filtering **is to (eb) create** stimuli that stimulate only one or a small number of the visual channels that operate in parallel to process visual stimuli.
(3) Ideally, stimuli filtered in high frequency bands **would (eb) stimulate** only receptors (channels) with small receptive fields.
(4) Stimuli filtered in low frequency bands **would (eb) stimulate** only channels that have large receptive fields. (The term channel is used here to designate an information processing system characterized by receptors of a particular size.)
(5) As in audition, there is substantial interest not only in how stimuli that are confined to a single band are processed, but also in how information from stimuli in different bands is perceptually combined. With the advent of affordable graphics processors, bandpass filtering **has (=) become** an increasingly widespread stimulus manipulation in vision.
(6) They studied the **role of (E&EB)three factors** in the ability of subjects to identify these letters when they were embedded in noise: (1) the signal-to-noise ratio, (2) the object-relative spatial frequency band in which the letters-plus-noise were filtered, and (3) the viewing distance (which determined retinal spatial frequency). They found that identification accuracy was independent of viewing distance over a range of more than 30:1. In this wide range, retinal spatial frequency did not matter **in (EB) determining** recognition accuracy; only object spatial frequency mattered.

Additive and Multiplicative Noise

(7) The term additive noise is used here to denote a stationary noise source that is independent of the signal and is added to the signal. Additive noise **can be (e) overcome** by increasing signal strength until the effective signal-to-noise ratio is sufficient support the desired level of performance.
(8) Multiplicative noise cannot be responsible for detection or discrimination thresholds that are reached by reducing the strength of a signal. Because multiplicative noise declines proportionately with diminishing signal strength, weak signals **are (e) not** worse off than strong signals.

.

NOTATION

Module One

Describing the mechanisms which cause certain events in the brain to be subjectively perceived is the hard problem of neuroscience. Consciousness **cannot (e) b**e reduced to neuronal firing and neuronal interactions.

G_{13} : Category one of reduced to neuronal firing and neuronal interactions

G_{14} : Category two of reduced to neuronal firing and neuronal interactions

G_{15} : Category three of reduced to neuronal firing and neuronal interactions

T_{13} : Category one of mechanisms which cause certain events in the brain to be subjectively perceived is the hard problem of neuroscience. Consciousness

T_{14} : Category two of mechanisms which cause certain events in the brain to be subjectively perceived is the hard problem of neuroscience. Consciousness

T_{15} : Category three of mechanisms which cause certain events in the brain to be subjectively perceived is the hard problem of neuroscience. Consciousness

Module Two

"However," Wexler points out, "one **consequence of the** high phi effect **is the** minimal-motion principle can be violated!

G_{16} : Category one of minimal-motion principle can be violated!

G_{17} : Category two of minimal-motion principle can be violated!

G_{18} : Category three of minimal-motion principle can be violated!

T_{16} : Category one of **consequence of the** high phi effect

T_{17} : Category two of **consequence of the** high phi effect

T_{18} : Category three of **consequence of the** high phi effect

Module three

When the stimulus is incompatible with any globally coherent motion, and therefore equally compatible with any motion, people **perceive (e)** not only a large jump, but the largest possible jump that they can perceive.

G_{20} : Category one of not only a large jump, but the largest possible jump that they can perceive

G_{21} : Category two of not only a large jump, but the largest possible jump that they can perceive

G_{22} : Category three of not only a large jump, but the largest possible jump that they can perceive

T_{20} : Category one of stimulus is incompatible with any globally coherent motion, and therefore equally compatible with any motion, people

T_{21} : Category two of stimulus is incompatible with any globally coherent motion, and therefore equally compatible with any motion, people

T_{22} : Category three of stimulus is incompatible with any globally coherent motion, and therefore equally compatible with any motion, people

Module four

This maximum jump is the one that steps by dmax, which **acts as the** speed limit on motion perception.

G_{24} : Category one of speed limit on motion perception

G_{25} : Category two of speed limit on motion perception

G_{26} : Category three of speed limit on motion perception

T_{24} : Category one of maximum jump is the one that steps by dmax

T_{25} : Category two of maximum jump is the one that steps by dmax

T_{26} : Category three of maximum jump is the one that steps by dmax

Module five

Below dmax, steps should be more or less seen as what they are – and as can be seen in demonstration three, this is what happens. On the other hand, says Wexler, "above dmax you're supposed to **perceive (e)** noise, not motion – but this is not what actually occurs

G_{28} : Category one of noise, not motion – but this is not what actually occurs

G_{29} : Category two of noise, not motion – but this is not what actually occurs

G_{30} : Category three of noise, not motion – but this is not what actually occurs

T_{28} : Category one of Below dmax, steps should be more or less seen as what they are – and as can be seen in demonstration three, this is what happens. On the other hand, says Wexler, "above dmax you're supposed to

T_{29} : Category two of Below dmax, steps should be more or less seen as what they are – and as can be seen in demonstration three, this is what happens. On the other hand, says Wexler, "above dmax you're supposed to

T_{30} : Category three of Below dmax, steps should be more or less seen as what they are – and as can be seen in demonstration three, this is what happens. On the other hand, says Wexler, "above dmax you're supposed to

Module six

Rather, you **perceive (e)** the high phi jump, as can also be seen in demonstration three. "In one of our experiments," Wexler adds, "we showed that the amplitude of the jump is very closely **correlated with** the dmax limit, so that people who have higher dmax limits also see a larger high phi jump."

G_{32} : Category one of dmax limit, so that people who have higher dmax limits also see a larger high phi jump

G_{33} : Category two of dmax limit, so that people who have higher dmax limits also see a larger high phi jump

G_{34} : Category three of dmax limit, so that people who have higher dmax limits also see a larger high phi jump

T_{32} : Category one of perception of high phi jump ; amplitude of the jump is very closely

T_{33} : Category two of perception of high phi jump ; amplitude of the jump is very closely

T_{34} : Category three of perception of high phi jump ; amplitude of the jump is very closely

Module seven

"And indeed," he notes, "we found that the magnitude of the illusory jump depends on spatial frequency in exactly the same way: the lower the frequency, formation shall be of the farther the jump." This can be verified, Wexler adds, by viewing high phi demonstration six and demonstration seven. Discussing the finding that the direction of the jump depends on the duration of the inducing motion signals, Wexler notes, "We think that the preceding – that is, inducing – motion acts like a seed. For brief inducers, the motion itself **acts as** the seed, and the jump is experienced forwards with respect to the inducer

G_{36} : Category one of magnitude of the illusory jump; spatial frequency

G_{37} : Category two of magnitude of the illusory jump; spatial frequency

G_{38} : Category three of magnitude of the illusory jump; spatial frequency

T_{36} : Category one of spatial frequency ;magnitude of the illusory jump

T_{37} : Category two of spatial frequency ;magnitude of the illusory jump

T_{38} : Category three of spatial frequency; magnitude of the illusory jump

Module eight

G_{40} : Category one of seed, and the jump is experienced forwards with respect to the inducer

G_{41} : Category two of seed, and the jump is experienced forwards with respect to the inducer

G_{42} : Category three of seed, and the jump is experienced forwards with respect to the inducer

T_{40} : Category one of brief inducers, the motion itself

T_{41} : Category two of brief inducers, the motion itself

T_{42} : Category three of brief inducers, the motion itself

Module Nine

(Important module because of contential dynamics; six models due to spatial constraints)

Bandpass filtering refers to the processing of images or sounds so that they **contain(e) only** a narrow range--typically one or two octaves--of component frequencies. In addition, bandpass filtering is used to **create** stimuli that stimulate only a small portion of the basilar membrane

Thus we cannot escape the fact the world we know is constructed in order to **see(e)** itself ;But in order to do so evidently, it must **cut(e&eb) i**tself up in to atleast one state which sees and atleast one other state which is seen (G.Spencer Brown)

Consciousness is in its original nature, quiet, pure, and above the dualism of subject and object. But here appears the principle of particularization, and with the rise of this wind of action, the waves are **agitated(e&eb)** over the mind (D.T.Suzuki) (as quoted in Spectrum of Consciousness: Ken Wilber)

Mind is now **differentiated (e&eb)** and evolves in to eight levels (D.T.Suzuki) (as quoted in Spectrum of Consciousness: Ken Wilber)

There is thus multiplication of inexhaustible one (Ananda Coomaraswamy) (as quoted in Spectrum of

Consciousness: Ken Wilber)

Unification of indefinitely many (Ananda Coomaraswamy) (as quoted in Spectrum of Consciousness: Ken Wilber). Again it is the case of "organisation"-"disorganization"-"organisation". Individual debits and credits are entered in to transfer scrolls and they are accounted for, and sorted out based on some head of accounts, finally re tallied at the general ledger)

Such are the beginnings and endings of worlds and individual beings expanded from (eb) a point **without (e) position or dimensions and a now without** date or duration. This means position cancels off dimension ,and past=future=now(my interpretation) (Ananda Coomaraswamy) (as quoted in Spectrum of Consciousness: Ken Wilber)

G_{44} : Category one of narrow range--typically one or two octaves--of component frequencies; bandpass filtering is used; world we know; world we know; **i**tself up in to atleast one state which sees and atleast one other state which is seen (G.Spencer Brown); mind (anagrammatic field acting over the information storage in all the three levels and the concomitant ramification in the form of dreams, pathology, or normal thinking); Mind is now **differentiated;** one; Unification; many (Ananda Coomaraswamy) (as quoted in Spectrum of Consciousness: Ken Wilber). Again it is the case of "organisation"-"disorganization"-"organisation". Individual debits and credits are entered in to transfer scrolls and they are accounted for, and sorted out based on some head of accounts, finally re tallied at the general ledger); dimension; positions; past;

G_{45} : Category two of narrow range--typically one or two octaves--of component frequencies; bandpass filtering is used; world we know; world we know; **i**tself up in to atleast one state which sees and atleast one other state which is seen (G.Spencer Brown); mind (anagrammatic field acting over the information storage in all the three levels and the concomitant ramification in the form of dreams, pathology, or normal thinking); Mind is now **differentiated;** one(this means that microcosm is also macrocosm. Like the individual debits and credits in a bank tally before it is finally tallied at the General Ledger. There are infinite number of functionaries both celestial and terrestrial assigned with particular work, be it improving the empowerment of women, development of pecuniary position of the poor, and down trodden or to create havoc, or to fleece money at the wall street); Unification; many (Ananda Coomaraswamy) (as quoted in Spectrum of Consciousness: Ken Wilber). Again it is the case of "organisation"-"disorganization"-"organisation". Individual debits and credits are entered in to transfer scrolls and they are accounted for, and sorted out based on some head of accounts, finally re tallied at the general ledger); positions; past;

G_{46} : Category three of narrow range--typically one or two octaves--of component frequencies; bandpass filtering is used; world we know; world we know; **i**tself up in to atleast one state which sees and atleast one other state which is seen (G.Spencer Brown); mind (anagrammatic field acting over the information storage in all the three levels and the concomitant ramification in the form of dreams, pathology, or normal thinking); Mind is now **differentiated;** Unification; many (Ananda Coomaraswamy) (as quoted in Spectrum of Consciousness: Ken Wilber). Again it is the case of "organisation"-"disorganization"-"organisation". Individual debits and credits are entered in to transfer scrolls and they are accounted for, and sorted out based on some head of accounts, finally re tallied at the general ledger); positions; past;

T_{44} : Category one of Bandpass filtering refers to the processing of images or sounds; stimuli that stimulate only a small portion of the basilar membrane; world we know is constructed in order to; **i**tself up in to atleast one state which sees and atleast one other state which is seen (G.Spencer Brown); world we know; principle of particularization, and with the rise of this wind of action, the waves; evolves in to eight levels; many (Ananda Coomaraswamy) (as quoted in Spectrum of Consciousness: Ken Wilber). Again it is the case of "organisation"-"disorganization"-"organisation". Individual debits and credits are entered in to transfer

scrolls and they are accounted for, and sorted out based on some head of accounts, finally re tallied at the general ledger);Unification; dimension; future;

T_{45} : Category two of Bandpass filtering refers to the processing of images or sounds; stimuli that stimulate only a small portion of the basilar membrane; world we know is constructed in order to; **i**tself up in to atleast one state which sees and atleast one other state which is seen (G.Spencer Brown); world we know; principle of particularization, and with the rise of this wind of action, the waves; evolves in to eight levels; many (Ananda Coomaraswamy) (as quoted in Spectrum of Consciousness: Ken Wilber). Again it is the case of "organisation"-"disorganization"-"organisation". Individual debits and credits are entered in to transfer scrolls and they are accounted for, and sorted out based on some head of accounts, finally re tallied at the general ledger);Unification; dimension; future;

T_{46} : Category three of Bandpass filtering refers to the processing of images or sounds; stimuli that stimulate only a small portion of the basilar membrane; world we know is constructed in order to; **i**tself up in to atleast one state which sees and atleast one other state which is seen (G.Spencer Brown); world we know; principle of particularization, and with the rise of this wind of action, the waves; evolves in to eight levels; many (Ananda Coomaraswamy) (as quoted in Spectrum of Consciousness: Ken Wilber). Again it is the case of "organisation"-"disorganization"-"organisation". Individual debits and credits are entered in to transfer scrolls and they are accounted for, and sorted out based on some head of accounts, finally re tallied at the general ledger);Unification; dimension; future;

SECTION SEVENTY NINE

Motion Inducers: Isotropy-Antistrophe

INTRODUCTION—VARIABLES USED

Motion perception revisited: High Phi effect challenges established motion perception assumptions April 23, 2013 by Stuart Mason Dambrot in Neuroscience Enlarge A schematic illustration of the basic high-phi stimuli. Animated versions may be seen in demonstrations 1–4(not incorporated due to spatial constraints).

(1) For longer inducers, vision begins **to (e&eb) adapt** to the motion – a result known as the motion aftereffect." Also known as the waterfall illusion, the motion aftereffect occurs when, after viewing a moving object for an extended period of time, and that object then becomes stationary, **one (e) perceives** the object appears to slowly move in the opposite direction.

(2) "Many people initially think that what we've found **is (=) a** consequence of adaptation to motion or the motion aftereffect," he says. "If so, then it's the fastest motion aftereffect known. We've measured that the illusory motion is 10-100 times **faster (eb) than** the inducing motion!

(3) We think that for motion inducers, the adaptation **acts (=) as** the seed of the fast, backward jump." (Brief and long inducers can be compared directly in demonstration nine.)

(4) Wexler also describes how their findings for motion inducers, the adaptation **acts (=) as** the seed of the fast, backward jump." (Brief and long inducers can be compared directly in demonstration nine.) **relate(e&eb) to** the activity of neurons in of the primary visual cortex that respond to lines of a certain angle moving in one direction, as first described by Hubel and Wiesel (1959)2.

(5) "In the brain, motion detectors **are (e&eb) sensitive** to motion in a particular place – the receptive field – a particular direction, and usually a particular speed," Wexler notes.

(6) "When faced with our stimulus, there can be many accidental matches at the local level. In one image there is a dark spot, for example, and in the next, uncorrelated image **there (eb) happens** to be dark spot just next to it.

(7) In that case, a local motion detector **will (eb) react to** this false match – so our stimulus actually activates many local motion detectors, but incoherently, in that all of these motion detectors are signaling different motions.

(8) The main point is that in all this incoherent mess the brain finally **prefers (eb) the** largest possible motion."

(9) Commenting on other areas of research that might benefit from their study, Wexler cites computer vision. "The minimal-motion principle **is (e&eb) enshrined in** a lot of algorithms for extracting motion," he concludes.

NOTATION

Module One

For longer inducers, vision begins **to (e&eb) adapt** to the motion – a result known as the motion aftereffect." Also known as the waterfall illusion, the motion aftereffect occurs when, after viewing a moving object for an extended period of time, and that object then becomes stationary, **one (e) perceives** the object appears to slowly move in the opposite direction

G_{13} : Category one of longer inducers, vision begins

G_{14} : Category two of longer inducers, vision begins

G_{15} : Category three of longer inducers, vision begins

T_{13} : Category one of motion – a result known as the motion aftereffect." Also known as the waterfall illusion, the motion aftereffect occurs when, after viewing a moving object for an extended period of time, and that object then becomes stationary, **one (e) perceives** the object appears to slowly move in the opposite direction; longer inducers, vision begins

T_{14} : Category two of motion – a result known as the motion aftereffect." Also known as the waterfall illusion, the motion aftereffect occurs when, after viewing a moving object for an extended period of time, and that object then becomes stationary, **one (e) perceives** the object appears to slowly move in the opposite direction; longer inducers, vision begins

T_{15} : Category three of motion – a result known as the motion aftereffect." Also known as the waterfall illusion, the motion aftereffect occurs when, after viewing a moving object for an extended period of time, and that object then becomes stationary, **one (e) perceives** the object appears to slowly move in the opposite direction; longer inducers, vision begins

Module Two

"Many people initially think that what we've found **is a** consequence of adaptation to motion or the motion aftereffect," he says. "If so, then it's the fastest motion aftereffect known. We've measured that the illusory motion is 10-100 times **faster (eb) than** the inducing motion!

G_{16} : Category one of consequence of adaptation to motion or the motion aftereffect," he says. "If so, then it's the fastest motion aftereffect known. We've measured that the illusory motion is 10-100 times **faster (eb) than** the inducing motion!

G_{17} : Category two of consequence of adaptation to motion or the motion aftereffect," he says. "If so, then it's the fastest motion aftereffect known. We've measured that the illusory motion is 10-100 times **faster (eb) than** the inducing motion!

G_{18} : Category three of consequence of adaptation to motion or the motion aftereffect," he says. "If so, then it's the fastest motion aftereffect known. We've measured that the illusory motion is 10-100 times **faster (eb) than** the inducing motion!

T_{16} : Category one of vision begins **to (e&eb) adapt** to the motion – a result known as the motion aftereffect."

T_{17} : Category two of vision begins **to (e&eb) adapt** to the motion – a result known as the motion aftereffect."

T_{18} : Category three of vision begins **to (e&eb) adapt** to the motion – a result known as the motion aftereffect."

Module three

We think that for motion inducers, the adaptation **acts as** the seed of the fast, backward jump." (Brief and long inducers can be compared directly in demonstration nine.)

G_{20} : Category one of seed of the fast, backward jump." (Brief and long inducers can be compared directly in demonstration nine.)

G_{21} : Category two of seed of the fast, backward jump." (Brief and long inducers can be compared directly

in demonstration nine.)

G_{22} : Category three of seed of the fast, backward jump." (Brief and long inducers can be compared directly in demonstration nine.)

T_{20} : Category one of motion inducers, the adaptation

T_{21} : Category two of motion inducers, the adaptation

T_{22} : Category three of motion inducers, the adaptation

Module four(both cases discussed)

Wexler also describes how their findings for motion inducers, the adaptation **act as** the seed of the fast, backward jump." (Brief and long inducers can be compared directly in demonstration nine.) **relate(e&eb) to** the activity of neurons in of the primary visual cortex that respond to lines of a certain angle moving in one direction, as first described by Hubel and Wiesel (1959)2.

G_{24} : Category one of findings for motion inducers, the adaptation **act as** the seed of the fast, backward jump." (Brief and long inducers can be compared directly in demonstration nine.); activity of neurons in of the primary visual cortex that respond to lines of a certain angle moving in one direction, as first described by Hubel and Wiesel (1959)2.

G_{25} : Category two of findings for motion inducers, the adaptation **act as** the seed of the fast, backward jump." (Brief and long inducers can be compared directly in demonstration nine.); activity of neurons in of the primary visual cortex that respond to lines of a certain angle moving in one direction, as first described by Hubel and Wiesel (1959)2.

G_{26} : Category three of findings for motion inducers, the adaptation **act as** the seed of the fast, backward jump." (Brief and long inducers can be compared directly in demonstration nine.); activity of neurons in of the primary visual cortex that respond to lines of a certain angle moving in one direction, as first described by Hubel and Wiesel (1959)2.

T_{24} : Category one of activity of neurons in of the primary visual cortex that respond to lines of a certain angle moving in one direction, as first described by Hubel and Wiesel (1959)2 ;findings for motion inducers, the adaptation **act as** the seed of the fast, backward jump." (Brief and long inducers can be compared directly in demonstration nine.)

T_{25} : Category two of activity of neurons in of the primary visual cortex that respond to lines of a certain angle moving in one direction, as first described by Hubel and Wiesel (1959)2 ;findings for motion inducers, the adaptation **act as** the seed of the fast, backward jump." (Brief and long inducers can be compared directly in demonstration nine.)

T_{26} : Category three of activity of neurons in of the primary visual cortex that respond to lines of a certain angle moving in one direction, as first described by Hubel and Wiesel (1959)2 ; findings for motion inducers, the adaptation **act as** the seed of the fast, backward jump." (Brief and long inducers can be compared directly in demonstration nine.)

Module five (both the cases discussed)

"In the brain, motion detectors **are (e&eb) sensitive** to motion in a particular place – the receptive field – a particular direction, and usually a particular speed," Wexler notes.

G_{28} : Category one of motion detectors in the brain; motion in a particular place – the receptive field – a

particular direction, and usually a particular speed

G_{29} : Category two of motion detectors in the brain; motion in a particular place – the receptive field – a particular direction, and usually a particular speed

G_{30} : Category three of motion detectors in the brain; motion in a particular place – the receptive field – a particular direction, and usually a particular speed

T_{28} : Category one of motion in a particular place – the receptive field – a particular direction, and usually a particular speed ;motion detectors in the brain

T_{29} : Category two of motion in a particular place – the receptive field – a particular direction, and usually a particular speed; motion detectors in the brain

T_{30} : Category three of motion in a particular place – the receptive field – a particular direction, and usually a particular speed; motion detectors in the brain

Module six

"When faced with our stimulus, there can be many accidental matches at the local level. In one image there is a dark spot, for example, and in the next, uncorrelated image **there (eb) happens** to be dark spot just next to it.

G_{32} : Category one of faced with our stimulus, there can be many accidental matches at the local level. In one image there is a dark spot, for example, and in the next, uncorrelated image

G_{33} : Category two of faced with our stimulus; there can be many accidental matches at the local level. In one image there is a dark spot, for example, and in the next, uncorrelated image

G_{34} : Category three of faced with our stimulus; there can be many accidental matches at the local level. In one image there is a dark spot, for example, and in the next, uncorrelated image

T_{32} : Category one of dark spot just next to it.

T_{33} : Category two of dark spot just next to it.

T_{34} : Category three of dark spot just next to it.

Module seven

In that case, a local motion detector **will (eb) react to** this false match – so our stimulus actually activates many local motion detectors, but incoherently, in that all of these motion detectors are signaling different motions

G_{36} : Category one of local motion detector

G_{37} : Category two of local motion detector

G_{38} : Category three of local motion detector

T_{36} : Category one of false match – so our stimulus actually activates many local motion detectors, but incoherently, in that all of these motion detectors are signaling different motions

T_{37} : Category two of false match – so our stimulus actually activates many local motion detectors, but incoherently, in that all of these motion detectors are signaling different motions

T_{38} : Category three of false match – so our stimulus actually activates many local motion detectors, but incoherently, in that all of these motion detectors are signaling different motions

Module eight

The main point is that in all this incoherent mess the brain finally **prefers (eb) the** largest possible motion.
G_{40} : Category one of brain finally **prefers**

G_{41} : Category two of brain finally **prefers**

G_{42} : Category three of brain finally **prefers**

T_{40} : Category one of largest possible motion

T_{41} : Category two of largest possible motion

T_{42} : Category three of largest possible motion

Module Nine

Commenting on other areas of research that might benefit from their study, Wexler cites computer vision. "The minimal-motion principle **is (e&eb) enshrined in** a lot of algorithms for extracting motion," he concludes.
G_{44} : Category one of lot of algorithms for extracting motion

G_{45} : Category two of lot of algorithms for extracting motion

G_{46} : Category three of lot of algorithms for extracting motion

T_{44} : Category one of minimal-motion principle

T_{45} : Category two of minimal-motion principle

T_{46} : Category three of minimal-motion principle

SECTION EIGHTY ONE

Change Of Mind: Rocky Razzmatazz And Autumn Sonata

INTRODUCTION—VARIABLES USED

A change of mind: One protein appears to control neurons' ability to react to new experiences Mar 24, 2010 by Anne Trafton

(1) MIT neuroscientists have shown that the protein Arc **is (e) necessary** for neurons like this one to adjust their responses to new sensory stimuli. (The blue circle is the neuron's nucleus, and the red strands are actin filaments.) Image: Jason Shepherd Read more at: http://phys.org/news188666378.html#jCp. Image deleted due to spatial constraints. Please see the article

(2) MIT neuroscientists have now found that a single protein, known as Arc, appears **to (e&eb) control** neurons' ability to strengthen and weaken their synapses by regulating the number of neurotransmitter receptors on their surfaces. The finding could help researchers **identify new** drug targets for Fragile X and Angelman syndromes — inherited forms of mental retardation that have been linked to deficits of Arc.

(3) "The more we understand the chain of cellular events that Arc is involved in, **there(eb) shall be production** of the more we can identify particular targets where we could intervene," says

Mriganka Sur, head of MIT's Department of Brain and Cognitive Sciences (BCS). Sur and Mark Bear, the Picower Professor of Neuroscience, are senior authors of a paper on the work that appeared in the March 14 online edition of Nature Neuroscience. http://phys.org/news188666378.html#jCp

A surprising discovery

(4) Jason Shepherd, co-lead author of the paper and a postdoctoral associate in Bear's lab, began studying Arc as a grad student at Johns Hopkins University. He and his colleagues showed that Arc **weakens (e)** synapses by removing receptors for glutamate, a neurotransmitter that stimulates neuron activity, from neuron cell membranes.

(5) They also discovered that when the Arc gene is turned off in mice, they **lose (e)** their ability to form long-term memories.

(6) In the new study, Shepherd and co-lead author Cortina McCurry, a recent PhD recipient in BCS, did a series of experiments designed to pinpoint the role of Arc in the visual cortex of mice. They started with a classic experimental setup that involves sealing one eye for two days, depriving the eye of visual input. In normal mice, **this (eb) strengthens** synapses in the part of the cortex receiving input from the open eye, and weakens them in cortical cells wired to the closed eye.

(7) Shepherd and McCurry performed their experiments on mice with a mutation in the Arc gene **that (eb) renders** the protein ineffective.

(8) Because of Arc's known role in weakening synapses, they expected mice without Arc not to show any synaptic weakening from the closed eye. That prediction came true, but to the researchers' surprise, **they found (eb) that** the synaptic strengthening normally seen from the open eye also disappeared. Read more at: http://phys.org/news188666378.html#jCp

(9) "If you knock out the gene, you don't get either response," says Shepherd. "The brain is not responsive **at all to the** changes in sensory input."They observed the same lack of plasticity in studies of Arc-deficient mice that were repeatedly exposed to the same visual stimulus (for example, a horizontal bar), which normally **provokes** (eb) neurons to enhance their response to that particular stimulus.

NOTATION

Module One

MIT neuroscientists have shown that the protein Arc **is (e) necessary** for neurons like this one to adjust their responses to new sensory stimuli. (The blue circle is the neuron's nucleus, and the red strands are actin filaments.) Image: Jason Shepherd Read more at: http://phys.org/news188666378.html#jCp. Image deleted due to spatial constraints. Please see the article

G_{13} : Category one of neurons like this one to adjust their responses to new sensory stimuli.

G_{14} : Category two of neurons like this one to adjust their responses to new sensory stimuli.

G_{15} : Category three of neurons like this one to adjust their responses to new sensory stimuli.

T_{13} : Category one of protein Arc

T_{14} : Category two of protein Arc

T_{15} : Category three of protein Arc

Module Two(both cases studied)

MIT neuroscientists have now found that a single protein, known as Arc, appears **to (e&eb) control** neurons' ability to strengthen and weaken their synapses by regulating the number of neurotransmitter receptors on their surfaces. The finding could help researchers **identify new** drug targets for Fragile X and

Angelman syndromes — inherited forms of mental retardation that have been linked to deficits of Arc

G_{16} : Category one of single protein, known as Arc; neurons' ability to strengthen and weaken their synapses by regulating the number of neurotransmitter receptors on their surfaces. The finding could help researchers **identify new** drug targets for Fragile X and Angelman syndromes — inherited forms of mental retardation that have been linked to deficits of Arc

G_{17} : Category two of single protein, known as Arc; neurons' ability to strengthen and weaken their synapses by regulating the number of neurotransmitter receptors on their surfaces. The finding could help researchers **identify new** drug targets for Fragile X and Angelman syndromes — inherited forms of mental retardation that have been linked to deficits of Arc

G_{18} : Category three of single protein, known as Arc; neurons' ability to strengthen and weaken their synapses by regulating the number of neurotransmitter receptors on their surfaces. The finding could help researchers **identify new** drug targets for Fragile X and Angelman syndromes — inherited forms of mental retardation that have been linked to deficits of Arc

T_{16} : Category one of neurons' ability to strengthen and weaken their synapses by regulating the number of neurotransmitter receptors on their surfaces. The finding could help researchers **identify new** drug targets for Fragile X and Angelman syndromes — inherited forms of mental retardation that have been linked to deficits of Arc; single protein, known as Arc(there are various systems in which this Arc exists. Propensities and properties thereof bear testimony and infallible observatory for the classificational postulation alcovishness and axiomatic predications.

T_{17} : Category two of neurons' ability to strengthen and weaken their synapses by regulating the number of neurotransmitter receptors on their surfaces. The finding could help researchers **identify new** drug targets for Fragile X and Angelman syndromes — inherited forms of mental retardation that have been linked to deficits of Arc

T_{18} : Category three of neurons' ability to strengthen and weaken their synapses by regulating the number of neurotransmitter receptors on their surfaces. The finding could help researchers **identify new** drug targets for Fragile X and Angelman syndromes — inherited forms of mental retardation that have been linked to deficits of Arc

Module three

"The more we understand the chain of cellular events that Arc is involved in, **there(eb) shall be production** of the more we can identify particular targets where we could intervene," says Mriganka Sur, head of MIT's Department of Brain and Cognitive Sciences (BCS). Sur and Mark Bear, the Picower Professor of Neuroscience, are senior authors of a paper on the work that appeared in the March 14 online edition of Nature Neuroscience. http://phys.org/news188666378.html#jCp

G_{20} : Category one of chain of cellular events that Arc is involved in

G_{21} : Category two of chain of cellular events that Arc is involved in

G_{22} : Category three of chain of cellular events that Arc is involved in

T_{20} : Category one of more we can identify particular targets where we could intervene

T_{21} : Category two of more we can identify particular targets where we could intervene

T_{22} : Category three of more we can identify particular targets where we could intervene

Module four

Jason Shepherd, co-lead author of the paper and a postdoctoral associate in Bear's lab, began studying Arc as a grad student at Johns Hopkins University. He and his colleagues showed that Arc **weakens (e)** synapses

by removing receptors for glutamate, a neurotransmitter that stimulates neuron activity, from neuron cell membranes

G_{24} : Category one of synapses by removing receptors for glutamate, a neurotransmitter that stimulates neuron activity, from neuron cell membranes

G_{25} : Category two of synapses by removing receptors for glutamate, a neurotransmitter that stimulates neuron activity, from neuron cell membranes

G_{26} : Category three of synapses by removing receptors for glutamate, a neurotransmitter that stimulates neuron activity, from neuron cell membranes

T_{24} : Category one of Arc

T_{25} : Category two of Arc

T_{26} : Category three of Arc

Module five

They also discovered that when the Arc gene is turned off in mice, they **lose (e)** their ability to form long-term memories

G_{28} : Category one of ability to form long-term memories

G_{29} : Category two of ability to form long-term memories

G_{30} : Category three of ability to form long-term memories

T_{28} : Category one of Arc gene is turned off in mice

T_{29} : Category two of Arc gene is turned off in mice

T_{30} : Category three of Arc gene is turned off in mice

Module six

In the new study, Shepherd and co-lead author Cortina McCurry, a recent PhD recipient in BCS, did a series of experiments designed to pinpoint the role of Arc in the visual cortex of mice. They started with a classic experimental setup that involves sealing one eye for two days, depriving the eye of visual input. In normal mice, **this (eb) strengthens** synapses in the part of the cortex receiving input from the open eye, and weakens them in cortical cells wired to the closed eye.

G_{32} : Category one of sealing one eye for two days, depriving the eye of visual input

G_{33} : Category two of sealing one eye for two days, depriving the eye of visual input

G_{34} : Category three of sealing one eye for two days, depriving the eye of visual input

T_{32} : Category one of synapses in the part of the cortex receiving input from the open eye, and weakens them in cortical cells wired to the closed eye.

T_{33} : Category two of synapses in the part of the cortex receiving input from the open eye, and weakens

them in cortical cells wired to the closed eye.

T_{34} : Category three of synapses in the part of the cortex receiving input from the open eye, and weakens them in cortical cells wired to the closed eye.

Module seven

Shepherd and McCurry performed their experiments on mice with a mutation in the Arc gene **that (eb) renders** the protein ineffective.

G_{36} : Category one of experiments on mice with a mutation in the Arc gene

G_{37} : Category two of experiments on mice with a mutation in the Arc gene

G_{38} : Category three of experiments on mice with a mutation in the Arc gene

T_{36} : Category one of protein ineffective

T_{37} : Category two of protein ineffective

T_{38} : Category three of protein ineffective

Module eight

Because of Arc's known role in weakening synapses, they expected mice without Arc not to show any synaptic weakening from the closed eye. That prediction came true, but to the researchers' surprise, **they found (eb) that** the synaptic strengthening normally seen from the open eye also disappeared. Read more at: http://phys.org/news188666378.html#jCp

G_{40} : Category one of Arc's known role in weakening synapses, they expected mice without Arc not to show any synaptic weakening from the closed eye

G_{41} : Category two of Arc's known role in weakening synapses, they expected mice without Arc not to show any synaptic weakening from the closed eye

G_{42} : Category three of Arc's known role in weakening synapses, they expected mice without Arc not to show any synaptic weakening from the closed eye

T_{40} : Category one of synaptic strengthening normally seen from the open eye also disappeared

T_{41} : Category two of synaptic strengthening normally seen from the open eye also disappeared

T_{42} : Category three of synaptic strengthening normally seen from the open eye also disappeared

Module Nine

If you knock out the gene, you don't get either response," says Shepherd. "The brain is not responsive **at all to the** changes in sensory input."They observed the same lack of plasticity in studies of Arc-deficient mice that were repeatedly exposed to the same visual stimulus (for example, a horizontal bar), which normally **provokes** (eb) neurons to enhance their response to that particular stimulus.

G_{44} : Category one of lack of plasticity in studies of Arc-deficient mice that were repeatedly exposed to the same visual stimulus (for example, a horizontal bar), which normally

G_{45} : Category two of lack of plasticity in studies of Arc-deficient mice that were repeatedly exposed to the

same visual stimulus (for example, a horizontal bar), which normally

G_{46} : Category three of lack of plasticity in studies of Arc-deficient mice that were repeatedly exposed to the same visual stimulus (for example, a horizontal bar), which normally

T_{44} : Category one of neurons to enhance their response to that particular stimulus

T_{45} : Category two of neurons to enhance their response to that particular stimulus

T_{46} : Category three of neurons to enhance their response to that particular stimulus

SECTION EIGHTY TWO

Living On The Edge: Abyssus Abyssum Invocat: Deep Calls Unto Deep

INTRODUCTION—VARIABLES USED

Living on the edge: cosmology on the boundary of anti-de Sitter space Carlos Barceló http://dx.doi.org/10.1016/S0370-2693(00)00520-7

(1) We sketch a particularly simple and compelling version of D-brane cosmology. Inspired by the semi-phenomenological Randall–Sundrum models, and their cosmological generalizations, **authors (eb) develop** a variant that contains a single (3+1)-dimensional D-brane which is located on the boundary of a single bulk (4+1)-dimensional region.
(2) The D-brane boundary is itself to be interpreted as our visible universe, with ordinary matter (planets, stars, galaxies) being trapped on this D-brane **by (e)** string theory effects.
(3) The (4+1)-dimensional bulk **is,** in its simplest implementation, adS4+1, anti-de Sitter space.
(4) Authors demonstrate that a k=+1 closed FLRW universe **is (=) the** most natural option, though the scale factor could quite easily be so large as to make it operationally indistinguishable from a k=0 spatially flat universe. (With minor loss of elegance, spatially flat and hyperbolic FLRW cosmologies can also be accommodated.). Authors demonstrate how this model can be made consistent with standard cosmology, and suggest some possible observational tests.
(5) dark flow **implies (eb) that** another universe is travelling towards us
(6) All creation is subjective with human being as pawns and actors dictated and governed by B-AB agency leading to a predetermined destination and combinational denouement of pre determined results
(7) Number of people looking at the moon **produce (eb) the** number of moons.

(8) Entire life is a **process of (e&eb)** simulation and dissimulation as in a quantum computer

(9) Quantum world is also (=) nothing. It is again observation ruled one.

(10) Classical world too is an (=) observer dependent one.

(11) B-AB agency produce results and give the impression that you have **produced (eb)** the results

(12) CERN simulation and dissimulation **gormandizes(e)** the very truth

Note: Last seven statements are mine. Models are given based on the assumption of the correctness of the statement.

NOTATION

Module One

We sketch a particularly simple and compelling version of D-brane cosmology. Inspired by the semi-phenomenological Randall–Sundrum models, and their cosmological generalizations, **authors (eb) develop** a variant that contains a single (3+1)-dimensional D-brane which is located on the boundary of a single bulk (4+1)-dimensional region

G_{13} : Category one of semi-phenomenological Randall–Sundrum models, and their cosmological generalizations

G_{14} : Category two of semi-phenomenological Randall–Sundrum models, and their cosmological generalizations

G_{15} : Category three of semi-phenomenological Randall–Sundrum models, and their cosmological generalizations

T_{13} : Category one of variant that contains a single (3+1)-dimensional D-brane which is located on the boundary of a single bulk (4+1)-dimensional region

T_{14} : Category two of variant that contains a single (3+1)-dimensional D-brane which is located on the boundary of a single bulk (4+1)-dimensional region

T_{15} : Category three of variant that contains a single (3+1)-dimensional D-brane which is located on the boundary of a single bulk (4+1)-dimensional region

Module Two

The D-brane boundary is itself to be interpreted as our visible universe, with ordinary matter (planets, stars, galaxies) being trapped on this D-brane **by (e)** string theory effects.

G_{16} : Category one of string theory effects. Systems that satisfy string theory enunciations and conditionalities are taken in to consideration, which implies that the effects thereof are also incorporated.

G_{17} : Category two of string theory effects

G_{18} : Category three of string theory effects

T_{16} : Category one of D-brane boundary is itself to be interpreted as our visible universe, with ordinary matter (planets, stars, galaxies) being trapped on this D-brane

T_{17} : Category two of D-brane boundary is itself to be interpreted as our visible universe, with ordinary matter (planets, stars, galaxies) being trapped on this D-brane

T_{18} : Category three of D-brane boundary is itself to be interpreted as our visible universe, with ordinary matter (planets, stars, galaxies) being trapped on this D-brane

Module three

The (4+1)-dimensional bulk **is,** in its simplest implementation, adS4+1, anti-de Sitter space

G_{20} : Category one of adS4+1, anti-de Sitter space

G_{21} : Category two of adS4+1, anti-de Sitter space

G_{22} : Category three of adS4+1, anti-de Sitter space

T_{20} : Category one of (4+1)-dimensional bulk

T_{21} : Category two of(4+1)-dimensional bulk

T_{22} : Category three of (4+1)-dimensional bulk

Module four

Authors demonstrate that a k=+1 closed FLRW universe **is the** most natural option, though the scale factor could quite easily be so large as to make it operationally indistinguishable from a k=0 spatially flat universe. (With minor loss of elegance, spatially flat and hyperbolic FLRW cosmologies can also be accommodated.). Authors demonstrate how this model can be made consistent with standard cosmology, and suggest some possible observational tests

G_{24} : Category one of natural option, though the scale factor could quite easily be so large as to make it operationally indistinguishable from a k=0 spatially flat universe. (With minor loss of elegance, spatially flat and hyperbolic FLRW cosmologies can also be accommodated.).

G_{25} : Category two of natural option, though the scale factor could quite easily be so large as to make it operationally indistinguishable from a k=0 spatially flat universe. (With minor loss of elegance, spatially flat and hyperbolic FLRW cosmologies can also be accommodated.).

G_{26} : Category three of natural option, though the scale factor could quite easily be so large as to make it operationally indistinguishable from a k=0 spatially flat universe. (With minor loss of elegance, spatially flat and hyperbolic FLRW cosmologies can also be accommodated.).

T_{24} : Category one of k=+1 closed FLRW universe

T_{25} : Category two of k=+1 closed FLRW universe

T_{26} : Category three of k=+1 closed FLRW universe

Module five

dark flow **implies (eb) that** another universe is travelling towards us

G_{28} : Category one of dark flow

G_{29} : Category two of dark flow

G_{30} : Category three of dark flow

T_{28} : Category one of another universe is travelling towards us

T_{29} : Category two of another universe is travelling towards us

T_{30} : Category three of another universe is travelling towards us

Module six(Important)

All creation is subjective with human being as pawns and actors dictated and governed by B-AB agency **leading to** a predetermined destination and combinational denouement of pre determined results

G_{32} : Category one of creation is subjective with human being as pawns and actors dictated and governed by B-AB agency(Grand Design)

G_{33} : Category two of creation is subjective with human being as pawns and actors dictated and governed by B-AB agency

G_{34} : Category three of creation is subjective with human being as pawns and actors dictated and governed by B-AB agency

T_{32} : Category one of predetermined destination and combinational denouement of pre determined results

T_{33} : Category two of predetermined destination and combinational denouement of pre determined results

T_{34} : Category three of predetermined destination and combinational denouement of pre determined results

Module seven

Number of people looking at the moon **produce (eb) the** number of moons

G_{36} : Category one of Number of people looking at the moon

G_{37} : Category two of Number of people looking at the moon

G_{38} : Category three of Number of people looking at the moon

T_{36} : Category one of number of moons

T_{37} : Category two of number of moons

T_{38} : Category three of number of moons

Module eight

Entire life is a **process of (e&eb)** simulation and dissimulation as in a quantum computer
Quantum world is also nothing. It is again observation ruled one.
Classical world too is an observer dependent one.

G_{40} : Category one of simulation and dissimulation as in a quantum computer; nothing (general ledger of all the transactions which are finally tallied at the final point. What is the final point is to be decided by the experiment being done; observer dependent one

G_{41} : Category two of simulation and dissimulation as in a quantum computer; nothing; observer dependent one;

G_{42} : Category three of simulation and dissimulation as in a quantum computer; nothing; observer dependent one

T_{40} : Category one of Entire life; Quantum world; Classical world too;

T_{41} : Category two of Entire life; Quantum world; Classical world too;

T_{42} : Category three of Entire life; Quantum world; Classical world too;

Module Nine

B-AB agency produce results and give the impression that you have **produced (eb)** the results
CERN simulation and dissimulation **gormandizes(e)** the very truth

G_{44} : Category one of B-AB agency produces results and gives the impression that you have; truth which is equivalent to perception and consciousness. Accentuation coefficients and dissipation coefficients represent consciousness corresponding to perception
G_{45} : Category two of B-AB agency produce results and give the impression that you have; truth which is equivalent to perception and consciousness. Accentuation coefficients and dissipation coefficients represent consciousness corresponding to perception
G_{46} : Category three of B-AB agency produce results and give the impression that you have; truth which is equivalent to perception and consciousness. Accentuation coefficients and dissipation coefficients represent consciousness corresponding to perception
T_{44} : Category one of results ; simulation and dissimulation;

T_{45} : Category two of results ; simulation and dissimulation;

T_{46} : Category three of results ; simulation and dissimulation

SECTION EIGHTY THREE

Dasein: Cygnus Inter Anates: Swan Among Ducks

INTRODUCTION—VARIABLES USED

Ideas: General Introduction to Pure Phenomenology. Trans. W.R. Boyce Gibson. Collier/Macmillan. [first translation 1931] Original German text – 1913.

(1) Let's assume that Christensen is right about this. The supporter of the intermittent-subject view might still argue that although Heidegger holds that Dasein **sometimes (eb) emerges as a** subject whose access to the world is non-theoretical (plausibly, in certain cases of un-readiness-to-hand)
(2) There is other textual evidence, beyond that which indicates the non-theoretical character of hitch-free skilled activity, **to (eb) suggest** that readiness-to-hand must remain non-subject-object in form. Whether or not there is such evidence would then need to be settled.

Being-in-the-World

(3) What the existential analytic has given us so far **is (=) a** phenomenological description of Dasein's within-the-world encounters with entities.
(4) The next clarification concerns the notion of world and the associated within-ness of Dasein. Famously, Heidegger writes of Dasein **as (=)** Being-in-the-world.

(5) In effect, then, the notion of Being-in-the-world **provides us** with a reinterpretation of the activity of existing (Dreyfus 1990, 40), where existence is given the narrow reading (existence) identified earlier.
(6) Understood as a unitary phenomenon (as opposed to a contingent, additive, tripartite combination of Being, in-ness, and the world), Being-in-the-world is an essential **characteristic (eb) of** Dasein.
(7) As Heidegger explains: Being-in is not a 'property' which Dasein sometimes has and sometimes does not have, and without which it could just **be (=) just** as well as it could be with it.
(8) It is not the case that man 'is' and then has, by way of an extra, **a (e&eb) relationship**-of-Being towards the 'world'—a world with which he provides himself occasionally.
(9) Dasein is never 'proximally' an entity which is, so to speak, free from Being-in, but which sometimes has the inclination to take up a **'relationship' (e&eb)** towards the world.

NOTATION

Module One

Let's assume that Christensen is right about this. The supporter of the intermittent-subject view might still argue that although Heidegger holds that Dasein **sometimes (eb) emerges as a** subject whose access to the world is non-theoretical (plausibly, in certain cases of un-readiness-to-hand)

G_{13} : Category one of Dasein

G_{14} : Category two of Dasein

G_{15} : Category three of Dasein

T_{13} : Category one of **emerges as a** subject whose access to the world is non-theoretical (plausibly, in certain cases of un-readiness-to-hand)

T_{14} : Category two of **emerges as a** subject whose access to the world is non-theoretical (plausibly, in certain cases of un-readiness-to-hand)

T_{15} : Category three of **emerges as a** subject whose access to the world is non-theoretical (plausibly, in certain cases of un-readiness-to-hand)

Module Two

There is other textual evidence, beyond that which indicates the non-theoretical character of hitch-free skilled activity, **to (eb) suggest** that readiness-to-hand must remain non-subject-object in form. Whether or not there is such evidence would then need to be settled.

G_{16} : Category one of textual evidence, beyond that which indicates the non-theoretical character of hitch-free skilled activity,

G_{17} : Category two of textual evidence, beyond that which indicates the non-theoretical character of hitch-free skilled activity,

G_{18} : Category three of textual evidence, beyond that which indicates the non-theoretical character of hitch-free skilled activity,

T_{16} : Category one of readiness-to-hand must remain non-subject-object in form. Whether or not there is such evidence would then need to be settled.

T_{17} : Category two of readiness-to-hand must remain non-subject-object in form. Whether or not there is such evidence would then need to be settled.

T_{18} : Category three of readiness-to-hand must remain non-subject-object in form. Whether or not there is such evidence would then need to be settled.

Module three

What the existential analytic has given us so far **is a** phenomenological description of Dasein's within-the-world encounters with entities

G_{20} : Category one of phenomenological description of Dasein's within-the-world encounters with entities

G_{21} : Category two of phenomenological description of Dasein's within-the-world encounters with entities

G_{22} : Category three of phenomenological description of Dasein's within-the-world encounters with entities

T_{20} : Category one of existential analytic

T_{21} : Category two of existential analytic

T_{22} : Category three of existential analytic

Module four

The next clarification concerns the notion of world and the associated within-ness of Dasein. Famously, Heidegger writes of Dasein **as** Being-in-the-world

G_{24} : Category one of Being-in-the-world(storage of information about body and mind. Identity with body and mind)

G_{25} : Category two of Being-in-the-world

G_{26} : Category three of Being-in-the-world

T_{24} : Category one of notion of world and the associated within-ness of Dasein

T_{25} : Category two of notion of world and the associated within-ness of Dasein

T_{26} : Category three of notion of world and the associated within-ness of Dasein

Module five

In effect, then, the notion of Being-in-the-world **provides us** with a reinterpretation of the activity of existing (Dreyfus 1990, 40), where existence is given the narrow reading (existence) identified earlier.

G_{28} : Category one of notion of Being-in-the-world

G_{29} : Category two of notion of Being-in-the-world

G_{30} : Category three of notion of Being-in-the-world

T_{28} : Category one of reinterpretation of the activity of existing (Dreyfus 1990, 40), where existence is given the narrow reading (existence) identified earlier

T_{29} : Category two of reinterpretation of the activity of existing (Dreyfus 1990, 40), where existence is given the narrow reading (existence) identified earlier

T_{30} : Category three of reinterpretation of the activity of existing (Dreyfus 1990, 40), where existence is given the narrow reading (existence) identified earlier

Module six

Understood as a unitary phenomenon (as opposed to a contingent, additive, tripartite combination of Being, in-ness, and the world), Being-in-the-world is an essential **characteristic (eb) of** Dasein

G_{32} : Category one of unitary phenomenon (as opposed to a contingent, additive, tripartite combination of Being, in-ness, and the world), Being-in-the-world is an essential bastion

G_{33} : Category two of unitary phenomenon (as opposed to a contingent, additive, tripartite combination of Being, in-ness, and the world), Being-in-the-world is an essential bastion

G_{34} : Category three of unitary phenomenon (as opposed to a contingent, additive, tripartite combination of Being, in-ness, and the world), Being-in-the-world is an essential bastion

T_{32} : Category one of Dasein

T_{33} : Category two of Dasein

T_{34} : Category three of Dasein

Module seven

As Heidegger explains: Being-in is not a 'property' which Dasein sometimes has and sometimes does not have, and without which it could just **be just** as well as it could be with it.
G_{36} : Category one of **just** as well as it could be with it.

G_{37} : Category two of **just** as well as it could be with it.

G_{38} : Category three of **just** as well as it could be with it.

T_{36} : Category one of Being-in is not a 'property' which Dasein sometimes has and sometimes does not have, and without which it could just **be**

T_{37} : Category two of Being-in is not a 'property' which Dasein sometimes has and sometimes does not have, and without which it could just **be**

T_{38} : Category three of Being-in is not a 'property' which Dasein sometimes has and sometimes does not have, and without which it could just **be**

Module eight

It is not the case that man 'is' and then has, by way of an extra, **a (e&eb) relationship**-of-Being towards the 'world'—a world with which he provides himself occasionally

G_{40} : Category one of It is not the case that man 'is' and then has, by way of an extra; **relationship**-of-Being towards the 'world'—a world with which he provides himself occasionally

G_{41} : Category two of It is not the case that man 'is' and then has, by way of an extra; **relationship**-of-Being towards the 'world'—a world with which he provides himself occasionally

G_{42} : Category three of It is not the case that man 'is' and then has, by way of an extra; **relationship**-of-Being towards the 'world'—a world with which he provides himself occasionally

T_{40} : Category one of **relationship**-of-Being towards the 'world'—a world with which he provides himself occasionally; It is not the case that man 'is' and then has, by way of an extra

T_{41} : Category two of **relationship**-of-Being towards the 'world'—a world with which he provides himself occasionally; It is not the case that man 'is' and then has, by way of an extra

T_{42} : Category three of **relationship**-of-Being towards the 'world'—a world with which he provides himself occasionally; It is not the case that man 'is' and then has, by way of an extra

Module Nine(both cases discussed)

Dasein is never 'proximally' an entity which is, so to speak, free from Being-in, but which sometimes has the inclination to take up a **'relationship' (e&eb)** towards the world.

G_{44} : Category one of Dasein is never 'proximally' an entity which is, so to speak, free from Being-in, but which sometimes has the inclination to take up; towards the world

G_{45} : Category two of Dasein is never 'proximally' an entity which is, so to speak, free from Being-in, but which sometimes has the inclination to take up; towards the world

G_{46} : Category three of Dasein is never 'proximally' an entity which is, so to speak, free from Being-in, but which sometimes has the inclination to take up; towards the world

T_{44} : Category one of towards the world ;Dasein is never 'proximally' an entity which is, so to speak, free from Being-in, but which sometimes has the inclination to take up

T_{45} : Category two of towards the world ;Dasein is never 'proximally' an entity which is, so to speak, free from Being-in, but which sometimes has the inclination to take up

T_{46} : Category three of towards the world ;Dasein is never 'proximally' an entity which is, so to speak, free from Being-in, but which sometimes has the inclination to take up

SECTION EIGHTY FOUR

Pure Phenomenology: Adversus Solem Ne Loquitor: Donot Speak Against The Sun

INTRODUCTION—VARIABLES USED

Ideas: General Introduction to Pure Phenomenology. Trans. W.R. Boyce Gibson. Collier/Macmillan. [first translation 1931] Original German text – 1913.

(1) Taking up relationships towards the world is possible only because Dasein, as Being-in-the-world, is as it is. This state of Being does not arise just **because (e)** some entity is present-at-hand outside of Dasein and meets up with it.

(2) Such an entity can **'meet up with' (e&eb)** Dasein only in so far as it can, of its own accord, show itself within a world. (Being and Time 12: 84)

(3) As this passage makes clear, the Being-in dimension of Being-in-the-world cannot be **thought (=) of as** a merely spatial relation in some sense that might be determined by a GPS device, since Dasein is never just present-at-hand within the world in the way demanded by that sort of spatial in-ness.

(4) Heidegger sometimes uses the term dwelling to capture the distinctive manner in which Dasein is in the world. To dwell in a house is not merely to be inside it spatially in the sense just canvassed. Rather, it is to **belong (e&eb) there**, to have a familiar place there.

(5) It is in this sense that Dasein is (essentially) in the world. (Heidegger will later introduce an existential notion of spatiality that does help to illuminate the sense in which Dasein is in the world. More on that below.) So now, what is the world such that Dasein (essentially) dwells in it? To answer this question we need to spend some time unpacking the Heideggerian concept of an 'involvement' (Bewandtnis). The German term Bewandtnis is extremely difficult to translate in a way that captures all its native nuances (for discussion, see Tugendhat 1967; thanks to a reviewer for emphasizing this point). And things are made more complicated by the fact that, during his exposition, Heidegger freely employs a number of closely related notions, including '**assignment', 'indication' and 'reference'**. Nevertheless, what is clear is that Heidegger introduces the term that Macquarrie and Robinson **translate(e&eb) as** 'involvement' to express the roles that equipmental entities play—the ways in which they are involved—in Dasein's everyday patterns of activity. Crucially, for Heidegger, an involvement is not a stand-alone structure, but rather a link in a network of intelligibility that he calls a totality of involvements.

(6) Take the stock Heideggerian example: the hammer is involved in an act of hammering; that hammering is involved in making something fast; and that making something fast is involved in protecting the human agent against bad weather. Such totalities of involvements are the contexts of everyday equipmental practice. As such, they define equipmental entities, so the hammer is intelligible as what it is only with respect to the shelter and, indeed, all the other items of equipment to which it meaningfully relates in Dasein's everyday practices. This relational ontology **generates(eb)** what Brandom (1983, 391–3) calls Heidegger's 'strong systematicity condition', as given voice in Heidegger's striking claim that "[t]aken strictly, there 'is' no such thing as an equipment" (Being and Time, 15: 97).

(7) And this radical holism spreads, because once one begins to trace a path through a network of involvements, one will inevitably **traverse (e&eb)** vast regions of involvement-space.

(8) Thus links will be traced not only from hammers to hammering to making fast to protection against the weather, but also from hammers to pulling out nails to dismantling wardrobes to moving house. This behaviour will refer back to many other behaviours (packing, van-driving) and thus to many other items of equipment (large boxes, removal vans), and so on. **The (eb) result is a** large-scale holistic network of interconnected relational significance. Such networks constitute worlds, in one of Heidegger's key senses of the term—an ontical sense that he describes as having a pre-ontological signification (Being and Time 14: 93).

(9) Before a second key sense of the Heideggerian notion of world is revealed, some important detail can be added to the emerging picture. Heidegger points out that involvements **are (=)** non uniform structures.

NOTATION

Module One

Taking up relationships towards the world is possible only because Dasein, as Being-in-the-world, is as it is. This state of Being does not arise just **because (e)** some entity is present-at-hand outside of Dasein and meets up with it.

G_{13} : Category one of **some** entity is present-at-hand outside of Dasein and meets up with it.

G_{14} : Category two of some entity is present-at-hand outside of Dasein and meets up with it.

G_{15} : Category three of some entity is present-at-hand outside of Dasein and meets up with it.

T_{13} : Category one of relationships towards the world is possible only because Dasein, as Being-in-the-world, is as it is. This state of Being does not arise

T_{14} : Category two of relationships towards the world is possible only because Dasein, as Being-in-the-world, is as it is. This state of Being does not arise

T_{15} : Category three of relationships towards the world is possible only because Dasein, as Being-in-the-world, is as it is. This state of Being does not arise

Module Two

Such an entity can **'meet up with' (e&eb)** Dasein only in so far as it can, of its own accord, show itself within a world. (Being and Time 12: 84)

G_{16} : Category one of Dasein, as Being-in-the-world, is as it is. This state of Being does not arise just **because (e)** some entity is present-at-hand outside of Dasein and meets up with it; Dasein only in so far as it can, of its own accord, show itself within a world. (Being and Time 12: 84)

G_{17} : Category two of Dasein, as Being-in-the-world, is as it is. This state of Being does not arise just **because (e)** some entity is present-at-hand outside of Dasein and meets up with it; Dasein only in so far as it can, of its own accord, show itself within a world. (Being and Time 12: 84);

G_{18} : Category three of Dasein, as Being-in-the-world, is as it is. This state of Being does not arise just **because (e)** some entity is present-at-hand outside of Dasein and meets up with it; Dasein only in so far as it can, of its own accord, show itself within a world. (Being and Time 12: 84);

T_{16} : Category one of Dasein only in so far as it can, of its own accord, show itself within a world. (Being and Time 12: 84); Dasein, as Being-in-the-world, is as it is. This state of Being does not arise just **because (e)** some entity is present-at-hand outside of Dasein and meets up with it

T_{17} : Category two of Dasein only in so far as it can, of its own accord, show itself within a world. (Being and Time 12: 84); Dasein, as Being-in-the-world, is as it is. This state of Being does not arise just **because (e)** some entity is present-at-hand outside of Dasein and meets up with it

T_{18} : Category three of Dasein only in so far as it can, of its own accord, show itself within a world. (Being and Time 12: 84); Dasein, as Being-in-the-world, is as it is. This state of Being does not arise just **because (e)** some entity is present-at-hand outside of Dasein and meets up with it

Module three

As this passage makes clear, the Being-in dimension of Being-in-the-world cannot be **thought of as** a merely spatial relation in some sense that might be determined by a GPS device, since Dasein is never just present-at-hand within the world in the way demanded by that sort of spatial in-ness.

G_{20} : Category one of spatial relation in some sense that might be determined by a GPS device, since Dasein is never just present-at-hand within the world in the way demanded by that sort of spatial in-ness

G_{21} : Category two of spatial relation in some sense that might be determined by a GPS device, since Dasein is never just present-at-hand within the world in the way demanded by that sort of spatial in-ness

G_{22} : Category three of spatial relation in some sense that might be determined by a GPS device, since Dasein is never just present-at-hand within the world in the way demanded by that sort of spatial in-ness

T_{20} : Category one of Being-in dimension of Being-in-the-world cannot be

T_{21} : Category two of Being-in dimension of Being-in-the-world cannot be

T_{22} : Category three of Being-in dimension of Being-in-the-world cannot be

Module four

Heidegger sometimes uses the term dwelling to capture the distinctive manner in which Dasein is in the world. To dwell in a house is not merely to be inside it spatially in the sense just canvassed. Rather, it is to **belong (eb) there**, to have a familiar place there.

G_{24} : Category one of Heidegger sometimes uses the term dwelling to capture the distinctive manner in which Dasein is in the world. To dwell in a house is not merely to be inside it spatially in the sense just canvassed. Rather, it is to

G_{25} : Category two of Heidegger sometimes uses the term dwelling to capture the distinctive manner in which Dasein is in the world. To dwell in a house is not merely to be inside it spatially in the sense just canvassed. Rather, it is to

G_{26} : Category three of Heidegger sometimes uses the term dwelling to capture the distinctive manner in which Dasein is in the world. To dwell in a house is not merely to be inside it spatially in the sense just canvassed. Rather, it is to

T_{24} : Category one of **belong (eb) there**, to have a familiar place there

T_{25} : Category two of **belong (eb) there**, to have a familiar place there

T_{26} : Category three of **belong (eb) there**, to have a familiar place there

BELONG [BIH-LAWNG, -LONG]

accord, agree, **appertain, apply, associate, attach to, be a component**, be a constituent, be a part, be akin, be an adjunct of, be connected with, be fitting, be linked with, be related, be relevant, bear, bear upon, become, befit, **chime, concern, correlate, correspond, exist**, fit, go, go with, harmonize, have relationship to, have respect to, have to do with, inhere, match, permeate, pertain, refer, regard, reside, set, suit, touch, vest

Module five

It is in this sense that Dasein is (essentially) in the world. (Heidegger will later introduce an existential notion of spatiality that does help to illuminate the sense in which Dasein is in the world. More on that below.) So now, what is the world such that Dasein (essentially) dwells in it? To answer this question we need to spend some time unpacking the Heideggerian concept of an 'involvement' (Bewandtnis). The German term Bewandtnis is extremely difficult to translate in a way that captures all its native nuances (for discussion, see Tugendhat 1967; thanks to a reviewer for emphasizing this point). And things are made more complicated by the fact that, during his exposition, Heidegger freely employs a number of closely related notions, including '**assignment', 'indication' and 'reference'**. Nevertheless, what is clear is that Heidegger introduces the term that Macquarrie and Robinson **translate(e&eb) as** 'involvement' to express the roles that equipmental entities play—the ways in which they are involved—in Dasein's everyday patterns of activity. Crucially, for Heidegger, an involvement is not a stand-alone structure, but rather a link in a network of intelligibility that he calls a totality of involvements.

G_{28} : Category one of an involvement is not a stand-alone structure

G_{29} : Category two of an involvement is not a stand-alone structure

G_{30} : Category three of an involvement is not a stand-alone structure

T_{28} : Category one of network of intelligibility that he calls a totality of involvements

T_{29} : Category two of network of intelligibility that he calls a totality of involvements

T_{30} : Category three of network of intelligibility that he calls a totality of involvements

Module six

Take the stock Heideggerian example: the hammer is involved in an act of hammering; that hammering is involved in making something fast; and that making something fast is involved in protecting the human agent against bad weather. Such totalities of involvements are the contexts of everyday equipmental practice. As such, they define equipmental entities, so the hammer is intelligible as what it is only with respect to the shelter and, indeed, all the other items of equipment to which it meaningfully relates in Dasein's everyday practices. This relational ontology **generates(eb)** what Brandom (1983, 391–3) calls Heidegger's 'strong systematicity condition', as given voice in Heidegger's striking claim that "[t]aken strictly, there 'is' no such thing as an equipment" (Being and Time, 15: 97).

G_{32} : Category one of relational ontology

G_{33} : Category two of relational ontology

G_{34} : Category three of relational ontology

T_{32} : Category one of Brandom (1983, 391–3) calls Heidegger's 'strong systematicity condition', as given voice in Heidegger's striking claim that "[t]aken strictly, there 'is' no such thing as an equipment" (Being and Time, 15: 97).

T_{33} : Category two of Brandom (1983, 391–3) calls Heidegger's 'strong systematicity condition', as given voice in Heidegger's striking claim that "[t]aken strictly, there 'is' no such thing as an equipment" (Being and Time, 15: 97).

T_{34} : Category three of Brandom (1983, 391–3) calls Heidegger's 'strong systematicity condition', as given voice in Heidegger's striking claim that "[t]aken strictly, there 'is' no such thing as an equipment" (Being and Time, 15: 97).

Module seven

And this radical holism spreads, because once one begins to trace a path through a network of involvements, one will inevitably **traverse (e&eb)** vast regions of involvement-space

G_{36} : Category one of radical holism spreads, because once one begins to trace a path through a network of involvements, one will inevitably

G_{37} : Category two of radical holism spreads, because once one begins to trace a path through a network of involvements, one will inevitably

G_{38} : Category three of radical holism spreads, because once one begins to trace a path through a network of

involvements, one will inevitably

T_{36} : Category one of vast regions of involvement-space

T_{37} : Category two of vast regions of involvement-space

T_{38} : Category three of vast regions of involvement-space

Module eight

Thus links will be traced not only from hammers to hammering to making fast to protection against the weather, but also from hammers to pulling out nails to dismantling wardrobes to moving house. This behaviour will refer back to many other behaviours (packing, van-driving) and thus to many other items of equipment (large boxes, removal vans), and so on. **The (eb) result is a** large-scale holistic network of interconnected relational significance. Such networks constitute worlds, in one of Heidegger's key senses of the term—an ontical sense that he describes as having a pre-ontological signification (Being and Time 14: 93).

G_{40} : Category one of links will be traced not only from hammers to hammering to making fast to protection against the weather, but also from hammers to pulling out nails to dismantling wardrobes to moving house. This behaviour will refer back to many other behaviours (packing, van-driving) and thus to many other items of equipment (large boxes, removal vans), and so on

G_{41} : Category two of links will be traced not only from hammers to hammering to making fast to protection against the weather, but also from hammers to pulling out nails to dismantling wardrobes to moving house. This behaviour will refer back to many other behaviours (packing, van-driving) and thus to many other items of equipment (large boxes, removal vans), and so on

G_{42} : Category three of links will be traced not only from hammers to hammering to making fast to protection against the weather, but also from hammers to pulling out nails to dismantling wardrobes to moving house. This behaviour will refer back to many other behaviours (packing, van-driving) and thus to many other items of equipment (large boxes, removal vans), and so on

T_{40} : Category one of large-scale holistic network of interconnected relational significance. Such networks constitute worlds, in one of Heidegger's key senses of the term—an ontical sense that he describes as having a pre-ontological signification (Being and Time 14: 93).

T_{41} : Category two of large-scale holistic network of interconnected relational significance. Such networks constitute worlds, in one of Heidegger's key senses of the term—an ontical sense that he describes as having a pre-ontological signification (Being and Time 14: 93).

T_{42} : Category three of large-scale holistic network of interconnected relational significance. Such networks constitute worlds, in one of Heidegger's key senses of the term—an ontical sense that he describes as having a pre-ontological signification (Being and Time 14: 93).

Module Nine

Before a second key sense of the Heideggerian notion of world is revealed, some important detail can be added to the emerging picture. Heidegger points out those involvements **are** non uniform structures.

G_{44} : Category one of non uniform structures

G_{45} : Category two of non uniform structures

G_{46} : Category three of non uniform structures

T_{44} : Category one of second key sense of the Heideggerian notion of world is revealed, some important detail can be added to the emerging picture. Heidegger points out that involvements

T_{45} : Category two of second key sense of the Heideggerian notion of world is revealed, some important detail can be added to the emerging picture. Heidegger points out that involvements

T_{46} : Category three of second key sense of the Heideggerian notion of world is revealed, some important detail can be added to the emerging picture. Heidegger points out that involvements

SECTION EIGHTY FIVE

Being And Time: Agere Sequitur Esse: Action Follows Being

INTRODUCTION—VARIABLES USED

Ideas: General Introduction to Pure Phenomenology. Trans. W.R. Boyce Gibson. Collier/Macmillan. [first translation 1931] Original German text – 1913.

(1) Thus I am currently working with a computer (a with-which), in the practical context of my office (an in-which), in order to write this encyclopedia entry (an in-order-to), which is aimed towards presenting an introduction to Heidegger's philosophy (a towards-this), for the sake of my academic work, that is, for the sake of my being an academic (a for-the-sake-of-which). The final involvement here, the for-the-sake-of-which, is crucial, **because (e)** according to Heidegger all totalities of involvements have a link of this type at their base.

(2) This forges a **connection between** (i) the idea that each moment in Dasein's existence constitutes a branch-point at which it chooses a way to be, and (ii) the claim that Dasein's projects and possibilities are essentially bound up with the ways in which other entities may become intelligible.

(3) This is because every for-the-sake-of-which **is the (e) base** structure of an equipment-defining totality of involvements and reflects a possible way for Dasein to be (an academic, a carpenter, a parent, or whatever).

(4) Moreover, given that entities are intelligible only within contexts of activity that, so to speak, arrive with Dasein, **this helps (eb) to explain** Heidegger's claim (Being and Time 16: 107) that, in encounters with entities, the world is something with which Dasein is always already familiar.

(5) Finally, it puts further flesh on the phenomenological category of the un-ready-to-hand. Thus when I am absorbed in trouble-free typing, the computer and the role that it plays in my academic activity

are transparent aspects of my experience. But if the computer crashes, I become aware of it as an entity with which I was working in the practical context of my office, in order to write an encyclopedia entry aimed towards presenting an introduction to Heidegger's philosophy. And I become aware of the fact that my behaviour is being organized for the sake of my being an academic. So disturbances **have (e&eb) the effect of** exposing totalities of involvements and, therefore, worlds. (For a second way in which worlds are phenomenologically 'lit up', see Heidegger's analysis of signs (Being and Time 17:107–114); for discussion, see Dreyfus 1990, 100–2, Cappuccio and Wheeler 2010.)

(6) As already indicated, Heidegger sometimes uses the expression 'world' in a different key sense, **to(eb) designate** what he calls the "ontologico-existential concept of worldhood" (Being and Time 14: 93).

(7) At this point in the existential analytic, worldhood is usefully **identified (e&eb)** as the abstract network mode of organizational configuration that is shared by all concrete totalities of involvements.

(8) We shall see, however, that as the hermeneutic spiral of the text unfolds, the notion of worldhood is subject to a series of reinterpretations until, finally, and its deep structure **gets (e) played out** in terms of temporality.

The Critique of Cartesianism

(9) Having completed what we might think of as the first phase of the existential analytic, Heidegger uses its results to launch an attack on one of the front-line representatives of the tradition, namely Descartes. This is the only worked-through example in **Being and Time** itself of what Heidegger calls the destruction (Destruktion) of the Western philosophical tradition, a process that was supposed to be a prominent theme in the ultimately unwritten second part of the text. The aim is to show that although the tradition takes theoretical knowledge to be primary, such knowledge (the prioritization of which is an aspect of the 'onticization' of Being mentioned earlier) **presupposes(eb)** the more fundamental openness to Being that Heidegger has identified as an essential characteristic of Dasein.

NOTATION

Module One

Thus I am currently working with a computer (a with-which), in the practical context of my office (an in-which), in order to write this encyclopedia entry (an in-order-to), which is aimed towards presenting an introduction to Heidegger's philosophy (a towards-this), for the sake of my academic work, that is, for the sake of my being an academic (a for-the-sake-of-which). The final involvement here, the for-the-sake-of-which, is crucial, **because (e)** according to Heidegger all totalities of involvements have a link of this type at their base. Note the primary homologues receptiveness of differentially oriented activity

G_{13} : Category one of final involvement here, the for-the-sake-of-which, is crucial

G_{14} : Category two of final involvement here, the for-the-sake-of-which, is crucial

G_{15} : Category three of final involvement here, the for-the-sake-of-which, is crucial

T_{13} : Category one of according to Heidegger all totalities of involvements have a link of this type at their base. Note the primary homologues receptiveness of differentially oriented activity

T_{14} : Category two of according to Heidegger all totalities of involvements have a link of this type at their base. Note the primary homologues receptiveness of differentially oriented activity

T_{15} : Category three of according to Heidegger all totalities of involvements have a link of this type at their

base. Note the primary homologues receptiveness of differentially oriented activity

Module Two

This forges a **connection between** (i) the idea that each moment in Dasein's existence constitutes a branch-point at which it chooses a way to be, and (ii) the claim that Dasein's projects and possibilities are essentially bound up with the ways in which other entities may become intelligible

G_{16} : Category one of all totalities of involvements have a link of this type at their base

G_{17} : Category two of all totalities of involvements have a link of this type at their base

G_{18} : Category three of all totalities of involvements have a link of this type at their base

T_{16} : Category one of forge of (i) the idea that each moment in Dasein's existence constitutes a branch-point at which it chooses a way to be, and (ii) the claim that Dasein's projects and possibilities are essentially bound up with the ways in which other entities may become intelligible

T_{17} : Category two of forge of (i) the idea that each moment in Dasein's existence constitutes a branch-point at which it chooses a way to be, and (ii) the claim that Dasein's projects and possibilities are essentially bound up with the ways in which other entities may become intelligible

T_{18} : Category three of forge of (i) the idea that each moment in Dasein's existence constitutes a branch-point at which it chooses a way to be, and (ii) the claim that Dasein's projects and possibilities are essentially bound up with the ways in which other entities may become intelligible

Module three

This is because every for-the-sake-of-which **is the (e) base** structure of an equipment-defining totality of involvements and reflects a possible way for Dasein to be (an academic, a carpenter, a parent, or whatever).

G_{20} : Category one of structure of an equipment-defining totality of involvements and reflects a possible way for Dasein to be (an academic, a carpenter, a parent, or whatever).

G_{21} : Category two of structure of an equipment-defining totality of involvements and reflects a possible way for Dasein to be (an academic, a carpenter, a parent, or whatever).

G_{22} : Category three of structure of an equipment-defining totality of involvements and reflects a possible way for Dasein to be (an academic, a carpenter, a parent, or whatever).

T_{20} : Category one of every for-the-sake-of-which

T_{21} : Category two of every for-the-sake-of-which

T_{22} : Category three of every for-the-sake-of-which

Module four

Moreover, given that entities are intelligible only within contexts of activity that, so to speak, arrive with Dasein, **this helps (eb) to explain** Heidegger's claim (Being and Time 16: 107) that, in encounters with entities, the world is something with which Dasein is always already familiar.

G_{24} : Category one of given that entities are intelligible only within contexts of activity that, so to speak, arrive with Dasein,

G_{25} : Category two of given that entities are intelligible only within contexts of activity that, so to speak,

arrive with Dasein,

G_{26} : Category three of given that entities are intelligible only within contexts of activity that, so to speak, arrive with Dasein,

T_{24} : Category one of Heidegger's claim (Being and Time 16: 107) that, in encounters with entities, the world is something with which Dasein is always already familiar

T_{25} : Category two of Heidegger's claim (Being and Time 16: 107) that, in encounters with entities, the world is something with which Dasein is always already familiar

T_{26} : Category three of Heidegger's claim (Being and Time 16: 107) that, in encounters with entities, the world is something with which Dasein is always already familiar

Module five

Finally, it puts further flesh on the phenomenological category of the un-ready-to-hand. Thus when I am absorbed in trouble-free typing, the computer and the role that it plays in my academic activity are transparent aspects of my experience. But if the computer crashes, I become aware of it as an entity with which I was working in the practical context of my office, in order to write an encyclopedia entry aimed towards presenting an introduction to Heidegger's philosophy. And I become aware of the fact that my behaviour is being organized for the sake of my being an academic. So disturbances **have (e&eb) the effect of** exposing totalities of involvements and, therefore, worlds. (For a second way in which worlds are phenomenologically 'lit up', see Heidegger's analysis of signs (Being and Time 17:107–114); for discussion, see Dreyfus 1990, 100–2, Cappuccio and Wheeler 2010.)

G_{28} : Category one of disturbances

G_{29} : Category two of disturbances

G_{30} : Category three of disturbances

T_{28} : Category one of exposing totalities of involvements and, therefore, worlds. (For a second way in which worlds are phenomenologically 'lit up', see Heidegger's analysis of signs (Being and Time 17:107–114);

T_{29} : Category two of exposing totalities of involvements and, therefore, worlds. (For a second way in which worlds are phenomenologically 'lit up', see Heidegger's analysis of signs (Being and Time 17:107–114);

T_{30} : Category three of exposing totalities of involvements and, therefore, worlds. (For a second way in which worlds are phenomenologically 'lit up', see Heidegger's analysis of signs (Being and Time 17:107–114);

Module six

As already indicated, Heidegger sometimes uses the expression 'world' in a different key sense, **to(eb) designate** what he calls the "ontologico-existential concept of worldhood" (Being and Time 14: 93).

G_{32} : Category one of Heidegger sometimes uses the expression 'world' in a different key sense

G_{33} : Category two of Heidegger sometimes uses the expression 'world' in a different key sense

G_{34} : Category three of Heidegger sometimes uses the expression 'world' in a different key sense

T_{32} : Category one of "ontologico-existential concept of worldhood" (Being and Time 14: 93).

T_{33} : Category two of "ontologico-existential concept of worldhood" (Being and Time 14: 93).

T_{34} : Category three of "ontologico-existential concept of worldhood" (Being and Time 14: 93).

Module seven

At this point in the existential analytic, worldhood is usefully **identified (e&eb)** as the abstract network mode of organizational configuration that is shared by all concrete totalities of involvements.
G_{36} : Category one of point in the existential analytic, worldhood is usefully

G_{37} : Category two of point in the existential analytic, worldhood is usefully

G_{38} : Category three of point in the existential analytic, worldhood is usefully

T_{36} : Category one of abstract network mode of organizational configuration that is shared by all concrete totalities of involvements

T_{37} : Category two of abstract network mode of organizational configuration that is shared by all concrete totalities of involvements

T_{38} : Category three of abstract network mode of organizational configuration that is shared by all concrete totalities of involvements

Module eight

We shall see, however, that as the hermeneutic spiral of the text unfolds, the notion of worldhood is subject to a series of reinterpretations until, finally, and its deep structure **gets (e) played out** in terms of temporality.
G_{40} : Category one of terms of temporality.

G_{41} : Category two of terms of temporality.

G_{42} : Category three of terms of temporality.

T_{40} : Category one of hermeneutic spiral of the text unfolds, the notion of worldhood is subject to a series of reinterpretations until, finally, and its deep structure

T_{41} : Category two of hermeneutic spiral of the text unfolds, the notion of worldhood is subject to a series of reinterpretations until, finally, and its deep structure

T_{42} : Category three of hermeneutic spiral of the text unfolds, the notion of worldhood is subject to a series of reinterpretations until, finally, and its deep structure

Module Nine

Having completed what we might think of as the first phase of the existential analytic, Heidegger uses its results to launch an attack on one of the front-line representatives of the tradition, namely Descartes. This is the only worked-through example in **Being and Time** itself of what Heidegger calls the destruction (Destruktion) of the Western philosophical tradition, a process that was supposed to be a prominent theme in

the ultimately unwritten second part of the text. The aim is to show that although the tradition takes theoretical knowledge to be primary, such knowledge (the prioritization of which is an aspect of the 'onticization' of Being mentioned earlier) **presupposes(eb)** the more fundamental openness to Being that Heidegger has identified as an essential characteristic of Dasein

G_{44} : Category one of tradition takes theoretical knowledge to be primary, such knowledge (the prioritization of which is an aspect of the 'onticization' of Being mentioned earlier)

G_{45} : Category two of tradition takes theoretical knowledge to be primary, such knowledge (the prioritization of which is an aspect of the 'onticization' of Being mentioned earlier)

G_{46} : Category three of tradition takes theoretical knowledge to be primary, such knowledge (the prioritization of which is an aspect of the 'onticization' of Being mentioned earlier)

T_{44} : Category one of more fundamental openness to Being that Heidegger has identified as an essential characteristic of Dasein

T_{45} : Category two of more fundamental openness to Being that Heidegger has identified as an essential characteristic of Dasein

T_{46} : Category three of more fundamental openness to Being that Heidegger has identified as an essential characteristic of Dasein

SECTION EIGHTY SIX

"Ontologico-Existential Concept Of Worldhood": Welträtsel: Nature Of The Universe And Meaning Of Life

INTRODUCTION—VARIABLES USED

Ideas: General Introduction to Pure Phenomenology. Trans. W.R. Boyce Gibson. Collier/Macmillan. [first translation 1931] Original German text – 1913.

(1) According to Heidegger, Descartes presents the world to us "with its skin off" (Being and Time 20: 132), i.e., **as a (e&eb) collection** of present-at-hand entities to be encountered by subjects.
(2) The consequence of this prioritizing of the present-at-hand **is(eb) that** the subject needs to claw itself into a world of equipmental meaning by adding what Heidegger calls 'value-predicates' (context-dependent meanings) to the present-at-hand.
(3) In stark contrast, Heidegger's own view is that Dasein is in primary epistemic **contact(e&eb)** not with context-independent present-at-hand primitives (e.g., raw sense data, such as a 'pure' experience of a patch of red), to which context-dependent meaning would need to be added via value-predicates**, but rather with** equipment, the kind of entity whose mode of Being is readiness-to-hand and which therefore comes already laden with context-dependent significance. What is perhaps Heidegger's best statement of this opposition comes later in Being and Time.
(4) What we 'first' hear is never noises or complexes of sounds, but the creaking waggon, the motor-cycle. We hear the column on the march, the north wind, the woodpecker tapping, the fire crackling… It **requires (e) a** very artificial and complicated frame of mind to 'hear' a 'pure noise'.

(5) The fact that motor-cycles and wagons are what we proximally hear is the **phenomenal (eb) evidence** that in every case Dasein, as Being-in-the-world, already dwells alongside what is ready-to-hand within-the-world

(6) it certainly does not dwell proximally alongside 'sensations'; nor would it first have to give shape to the swirl of sensations **to(eb) provide a** springboard from which the subject leaps off and finally arrives at a 'world'. Dasein, as essentially understanding, is proximally alongside what is understood. (Being and Time 34: 207)

(7) For Heidegger, then, we start not with the present-at-hand, moving to the ready-to-hand by adding value-predicates, **but (e&eb) with** the ready-to-hand, moving to the present-at-hand by stripping away the holistic networks of everyday equipmental meaning.

(8) It seems clear, then, that our two positions are diametrically opposed to each other, but why should we favour Heidegger's framework over Descartes'? Heidegger's flagship argument here is that the systematic additions of value-predicates to present-at-hand primitives **cannot (e) transform** our encounters with those objects into encounters with equipment.

(9) It comes in the following brief but dense passage: "Adding on value-predicates **cannot tell us** anything at all new about the Being of goods, but would **merely (eb) presuppose** again that goods have pure presence-at-hand as their kind of Being.

NOTATION

Module One

According to Heidegger, Descartes presents the world to us "with its skin off" (Being and Time 20: 132), i.e., **as a (e&eb) collection** of present-at-hand entities to be encountered by subjects.

G_{13} : Category one of present-at-hand entities to be encountered by subjects

G_{14} : Category two of present-at-hand entities to be encountered by subjects

G_{15} : Category three of present-at-hand entities to be encountered by subjects

T_{13} : Category one of world to us "with its skin off" (Being and Time 20: 132),

T_{14} : Category two of world to us "with its skin off" (Being and Time 20: 132),

T_{15} : Category three of world to us "with its skin off" (Being and Time 20: 132),

Module Two

The consequence of this prioritizing of the present-at-hand **is(eb) that** the subject needs to claw itself into a world of equipmental meaning by adding what Heidegger calls 'value-predicates' (context-dependent meanings) to the present-at-hand

G_{16} : Category one of consequence of this prioritizing of the present-at-hand

G_{17} : Category two of consequence of this prioritizing of the present-at-hand

G_{18} : Category three of consequence of this prioritizing of the present-at-hand

T_{16} : Category one of subject needs to claw itself into a world of equipmental meaning by adding what Heidegger calls 'value-predicates' (context-dependent meanings) to the present-at-hand

T_{17} : Category two of subject needs to claw itself into a world of equipmental meaning by adding what Heidegger calls 'value-predicates' (context-dependent meanings) to the present-at-hand

T_{18} : Category three of subject needs to claw itself into a world of equipmental meaning by adding what

Heidegger calls 'value-predicates' (context-dependent meanings) to the present-at-hand

Module three

In stark contrast, Heidegger's own view is that Dasein is in primary epistemic **contact(e&eb)** not with context-independent present-at-hand primitives (e.g., raw sense data, such as a 'pure' experience of a patch of red), to which context-dependent meaning would need to be added via value-predicates**, but rather with** equipment, the kind of entity whose mode of Being is readiness-to-hand and which therefore comes already laden with context-dependent significance. What is perhaps Heidegger's best statement of this opposition comes later in Being and Time.

G_{20} : Category one of Heidegger's own view is that Dasein is in primary epistemic **contact(e&eb)** not with context-independent present-at-hand primitives (e.g., raw sense data, such as a 'pure' experience of a patch of red), to which context-dependent meaning would need to be added via value-predicates

G_{21} : Category two of Heidegger's own view is that Dasein is in primary epistemic **contact(e&eb)** not with context-independent present-at-hand primitives (e.g., raw sense data, such as a 'pure' experience of a patch of red), to which context-dependent meaning would need to be added via value-predicates

G_{22} : Category three of Heidegger's own view is that Dasein is in primary epistemic **contact(e&eb)** not with context-independent present-at-hand primitives (e.g., raw sense data, such as a 'pure' experience of a patch of red), to which context-dependent meaning would need to be added via value-predicates

T_{20} : Category one of equipment, the kind of entity whose mode of Being is readiness-to-hand and which therefore comes already laden with context-dependent significance. What is perhaps Heidegger's best statement of this opposition comes later in Being and Time.

T_{21} : Category two of equipment, the kind of entity whose mode of Being is readiness-to-hand and which therefore comes already laden with context-dependent significance. What is perhaps Heidegger's best statement of this opposition comes later in Being and Time.

T_{22} : Category three of equipment, the kind of entity whose mode of Being is readiness-to-hand and which therefore comes already laden with context-dependent significance. What is perhaps Heidegger's best statement of this opposition comes later in Being and Time.

Module four

What we 'first' hear is never noises or complexes of sounds, but the creaking waggon, the motor-cycle. We hear the column on the march, the north wind, the woodpecker tapping, the fire crackling… It **requires (e) a** very artificial and complicated frame of mind to 'hear' a 'pure noise'.

G_{24} : Category one of artificial and complicated frame of mind to 'hear' a 'pure noise'.

G_{25} : Category two of artificial and complicated frame of mind to 'hear' a 'pure noise'.

G_{26} : Category three of artificial and complicated frame of mind to 'hear' a 'pure noise'.

T_{24} : Category one of first' hear is never noises or complexes of sounds, but the creaking waggon, the motor-cycle. We hear the column on the march, the north wind, the woodpecker tapping, the fire crackling…

T_{25} : Category two of first' hear is never noises or complexes of sounds, but the creaking waggon, the motor-cycle. We hear the column on the march, the north wind, the woodpecker tapping, the fire crackling…

T_{26} : Category three of first' hear is never noises or complexes of sounds, but the creaking waggon, the motor-cycle. We hear the column on the march, the north wind, the woodpecker tapping, the fire crackling…

Module five

The fact that motor-cycles and wagons are what we proximally hear is the **phenomenal (eb) evidence** that in every case Dasein, as Being-in-the-world, already dwells alongside what is ready-to-hand within-the-world

G_{28} : Category one of motor-cycles and wagons are what we proximally hear

G_{29} : Category two of motor-cycles and wagons are what we proximally hear

G_{30} : Category three of motor-cycles and wagons are what we proximally hear

T_{28} : Category one of every case Dasein, as Being-in-the-world, already dwells alongside what is ready-to-hand within-the-world

T_{29} : Category two of every case Dasein, as Being-in-the-world, already dwells alongside what is ready-to-hand within-the-world

T_{30} : Category three of every case Dasein, as Being-in-the-world, already dwells alongside what is ready-to-hand within-the-world

Module six

it certainly does not dwell proximally alongside 'sensations'; nor would it first have to give shape to the swirl of sensations **to(eb) provide a** springboard from which the subject leaps off and finally arrives at a 'world'. Dasein, as essentially understanding, is proximally alongside what is understood. (Being and Time 34: 207)

G_{32} : Category one of it certainly does not dwell proximally alongside 'sensations'; nor would it first have to give shape to the swirl of sensations

G_{33} : Category two of it certainly does not dwell proximally alongside 'sensations'; nor would it first have to give shape to the swirl of sensations

G_{34} : Category three of it certainly does not dwell proximally alongside 'sensations'; nor would it first have to give shape to the swirl of sensations

T_{32} : Category one of springboard from which the subject leaps off and finally arrives at a 'world'. Dasein, as essentially understanding, is proximally alongside what is understood. (Being and Time 34: 207)

T_{33} : Category two of springboard from which the subject leaps off and finally arrives at a 'world'. Dasein, as essentially understanding, is proximally alongside what is understood. (Being and Time 34: 207)

T_{34} : Category three of springboard from which the subject leaps off and finally arrives at a 'world'. Dasein, as essentially understanding, is proximally alongside what is understood. (Being and Time 34: 207)

Module seven

It seems clear, then, that our two positions are diametrically opposed to each other, but why should we favour Heidegger's framework over Descartes'? Heidegger's flagship argument here is that the systematic additions of value-predicates to present-at-hand primitives **cannot (e) transform** our encounters with those objects into encounters with equipment

G_{36} : Category one of **transform** our encounters with those objects into encounters with equipment

G_{37} : Category two of **transform** our encounters with those objects into encounters with equipment

G_{38} : Category three of **transform** our encounters with those objects into encounters with equipment

T_{36} : Category one of flagship argument here is that the systematic additions of value-predicates to present-at-hand primitives

T_{37} : Category two of flagship argument here is that the systematic additions of value-predicates to present-at-hand primitives

T_{38} : Category three of flagship argument here is that the systematic additions of value-predicates to present-at-hand primitives

Module eight

It seems clear, then, that our two positions are diametrically opposed to each other, but why should we favour Heidegger's framework over Descartes'? Heidegger's flagship argument here is that the systematic additions of value-predicates to present-at-hand primitives **cannot (e) transform** our encounters with those objects into encounters with equipment.

G_{40} : Category one of

G_{41} : Category two of

G_{42} : Category three of

T_{40} : Category one of

T_{41} : Category two of

T_{42} : Category three of

Module Nine

It comes in the following brief but dense passage: "Adding on value-predicates **cannot tell us** anything at all new about the Being of goods, but would **merely (eb) presuppose** again that goods have pure presence-at-hand as their kind of Being.

G_{44} : Category one of Adding on value-predicates **cannot tell us** anything at all new about the Being of goods, but would

G_{45} : Category two of Adding on value-predicates **cannot tell us** anything at all new about the Being of goods, but would

G_{46} : Category three of Adding on value-predicates **cannot tell us** anything at all new about the Being of goods, but would

T_{44} : Category one of goods have pure presence-at-hand as their kind of Being.

T_{45} : Category two of goods have pure presence-at-hand as their kind of Being.

T_{46} : Category three of goods have pure presence-at-hand as their kind of Being.

SECTION EIGHTY SEVEN

Cartesianism: Adequatio Intellectus Nostri Cum Re: Conformity Of Minds To The Facts

INTRODUCTION—VARIABLES USED

Ideas: General Introduction to Pure Phenomenology. Trans. W.R. Boyce Gibson. Collier/Macmillan. [first translation 1931] Original German text – 1913.

(1) Values would then **be (eb) determinate** characteristics which a thing possesses, and they would be present-at-hand" (Being and Time 21: 132).
(2) In other words, once we have assumed that we begin with the present-at-hand; values must **take (eb) the form of** determinate features of objects, and therefore constitute nothing but more present-at-hand structures.
(3) And if you add more present-at-hand structures to some existing present-at-hand structures, what you end up with is not equipmental meaning (totalities of involvements) **but (eb) merely** a larger number of present-at-hand structures.
(4) Heidegger's argument here is (at best) incomplete (for discussion, see Dreyfus 1990, Wheeler 2005). The defender of Cartesianism might concede that present-at-hand entities have determinate properties, but wonder why the fact that an entity has determinate properties is necessarily **an (eb)**

indication of presence-at-hand.

(5) On this view, having determinate properties **is (e) necessary** but not sufficient for an entity to be present-at-hand.

(6) More specifically, she might wonder why involvements cannot be thought **of (=) as** determinate features that entities possess just when they are embedded in certain contexts of use.

(7) Consider for example the various involvements specified in the academic writing context described earlier. They **certainly (=) seem to be** determinate, albeit context-relative, properties of the computer.

(8) Of course, the massively holistic character of totalities of involvements would make the task of specifying the necessary value-predicates (say, as sets of internal representations) incredibly hard, but it is unclear that it makes that task impossible. So it seems as if Heidegger doesn't really develop his case in sufficient detail. However, Dreyfus (1990) pursues a response that Heidegger might have given, one that draws on the familiar philosophical distinction between knowing-how **and** knowing-that.

(9) It seems that value-predicates **constitute (e) a** form of knowing-that (i.e., knowing that an entity has a certain context-dependent property) whereas the circumspective knowledge of totalities of involvements (Dasein's skilled practical activity) constitutes a form of knowing-how (i.e., knowing how to use equipment in appropriate ways; see the characterization of readiness-to-hand given earlier). Given the plausible (although not universally held) assumption that knowing-how cannot be reduced to knowledge-that, this would explain why value-predicates are simply the wrong sort of structures to capture the phenomenon of world-embeddedness.

NOTATION

Module One

Values would then **be (eb) determinate** characteristics which a thing possesses, and they would be present-at-hand" (Being and Time 21: 132).

G_{13} : Category one of Values

G_{14} : Category two of Values

G_{15} : Category three of Values

T_{13} : Category one of characteristics which a thing possesses, and they would be present-at-hand" (Being and Time 21: 132).
T_{14} : Category two of characteristics which a thing possesses, and they would be present-at-hand" (Being and Time 21: 132).
T_{15} : Category three of characteristics which a thing possesses, and they would be present-at-hand" (Being and Time 21: 132).

Module Two

In other words, once we have assumed that we begin with the present-at-hand; values must **take (eb) the form of** determinate features of objects, and therefore constitute nothing but more present-at-hand structures.
G_{16} : Category one of once we have assumed that we begin with the present-at-hand; values

G_{17} : Category two of once we have assumed that we begin with the present-at-hand; values

G_{18} : Category three of once we have assumed that we begin with the present-at-hand; values

T_{16} : Category one of determinate features of objects, and therefore constitute nothing but more present-at-hand structures

T_{17} : Category two of determinate features of objects, and therefore constitute nothing but more present-at-hand structures

T_{18} : Category three of determinate features of objects, and therefore constitute nothing but more present-at-hand structures

Module three

And if you add more present-at-hand structures to some existing present-at-hand structures, what you end up with is not equipmental meaning (totalities of involvements) **but (eb) merely** a larger number of present-at-hand structures

G_{20} : Category one of present-at-hand structures to some existing present-at-hand structures, what you end up with is not equipmental meaning (totalities of involvements)

G_{21} : Category two of present-at-hand structures to some existing present-at-hand structures, what you end up with is not equipmental meaning (totalities of involvements)

G_{22} : Category three of present-at-hand structures to some existing present-at-hand structures, what you end up with is not equipmental meaning (totalities of involvements)

T_{20} : Category one of larger number of present-at-hand structures

T_{21} : Category two of larger number of present-at-hand structures

T_{22} : Category three of larger number of present-at-hand structures

Module four

Heidegger's argument here is (at best) incomplete (for discussion, see Dreyfus 1990, Wheeler 2005). The defender of Cartesianism might concede that present-at-hand entities have determinate properties, but wonder why the fact that an entity has determinate properties is necessarily **an (eb) indication of** presence-at-hand.

G_{24} : Category one of defender of Cartesianism might concede that present-at-hand entities have determinate properties, but wonder why the fact that an entity has determinate properties is necessarily

G_{25} : Category two of defender of Cartesianism might concede that present-at-hand entities have determinate properties, but wonder why the fact that an entity has determinate properties is necessarily

G_{26} : Category three of defender of Cartesianism might concede that present-at-hand entities have determinate properties, but wonder why the fact that an entity has determinate properties is necessarily

T_{24} : Category one of presence-at-hand.

T_{25} : Category two of presence-at-hand.

T_{26} : Category three of presence-at-hand.

Module five

On this view, having determinate properties **is (e) necessary** but not sufficient for an entity to be present-at-hand.

G_{28} : Category one of **necessary** but not sufficient for an entity to be present-at-hand.

G_{29} : Category two of **necessary** but not sufficient for an entity to be present-at-hand.

G_{30} : Category three of **necessary** but not sufficient for an entity to be present-at-hand.

T_{28} : Category one of determinate properties

T_{29} : Category two of determinate properties

T_{30} : Category three of determinate properties

Module six

More specifically, one might wonder why involvements cannot be thought **of as** determinate features that entities possess just when they are embedded in certain contexts of use.

G_{32} : Category one of determinate features that entities possess just when they are embedded in certain contexts of use.

G_{33} : Category two of determinate features that entities possess just when they are embedded in certain contexts of use.

G_{34} : Category three of determinate features that entities possess just when they are embedded in certain contexts of use.

T_{32} : Category one of involvements

T_{33} : Category two of involvements

T_{34} : Category three of involvements

Module seven

Consider for example the various involvements specified in the academic writing context described earlier. They **certainly seem to be** determinate, albeit context-relative, properties of the computer.

G_{36} : Category one of determinate, albeit context-relative, properties of the computer

G_{37} : Category two of determinate, albeit context-relative, properties of the computer

G_{38} : Category three of determinate, albeit context-relative, properties of the computer

T_{36} : Category one of various involvements specified in the academic writing context

T_{37} : Category two of various involvements specified in the academic writing context

T_{38} : Category three of various involvements specified in the academic writing context

Module eight

Of course, the massively holistic character of totalities of involvements would **make the** task of specifying the necessary value-predicates (say, as sets of internal representations) incredibly hard, but it is unclear that it makes that task impossible. So it seems as if Heidegger doesn't really develop his case in sufficient detail.

However, Dreyfus (1990) pursues a response that Heidegger might have given, one that draws on the familiar philosophical distinction between knowing-how **and** knowing-that

G_{40} : Category one of massively holistic character of totalities of involvements

G_{41} : Category two of massively holistic character of totalities of involvements

G_{42} : Category three of massively holistic character of totalities of involvements

T_{40} : Category one of knowing-how **and** knowing-that

T_{41} : Category two of knowing-how **and** knowing-that

T_{42} : Category three of knowing-how **and** knowing-that

Module Nine

It seems that value-predicates **constitute (e) a** form of knowing-that (i.e., knowing that an entity has a certain context-dependent property) whereas the circumspective knowledge of totalities of involvements (Dasein's skilled practical activity) constitutes a form of knowing-how (i.e., knowing how to use equipment in appropriate ways; see the characterization of readiness-to-hand given earlier). Given the plausible (although not universally held) assumption that knowing-how cannot be reduced to knowledge-that, this would explain why value-predicates are simply the wrong sort of structures to capture the phenomenon of world-embeddedness

G_{44} : Category one of form of knowing-that (i.e., knowing that an entity has a certain context-dependent property) whereas the circumspective knowledge of totalities of involvements (Dasein's skilled practical activity) constitutes a form of knowing-how (i.e., knowing how to use equipment in appropriate ways; see the characterization of readiness-to-hand given earlier).

G_{45} : Category two of form of knowing-that (i.e., knowing that an entity has a certain context-dependent property) whereas the circumspective knowledge of totalities of involvements (Dasein's skilled practical activity) constitutes a form of knowing-how (i.e., knowing how to use equipment in appropriate ways; see the characterization of readiness-to-hand given earlier).

G_{46} : Category three of form of knowing-that (i.e., knowing that an entity has a certain context-dependent property) whereas the circumspective knowledge of totalities of involvements (Dasein's skilled practical activity) constitutes a form of knowing-how (i.e., knowing how to use equipment in appropriate ways; see the characterization of readiness-to-hand given earlier).

T_{44} : Category one of value-predicates

T_{45} : Category two of value-predicates

T_{46} : Category three of value-predicates

SECTION EIGHTY EIGHT

Spatio Temporal Actualization: Wertfreiheit: Value Neutrality

INTRODUCTION—VARIABLES USED

Ideas: General Introduction to Pure Phenomenology. Trans. W.R. Boyce Gibson. Collier/Macmillan. [first translation 1931] Original German text – 1913.

Spatiality

(1) In the wake of his critique of Cartesianism, Heidegger turns his attention to spatiality. He argues that Dasein dwells in the world **in (=) a** spatial manner, but that the spatiality in question—Dasein's existential spatiality—cannot be a matter of Dasein being located at a particular co-ordinate in physical, Cartesian space.

(2) That would be to conceive of Dasein as present-at-hand, and presence-at-hand **is(=) a** mode of Being that can belong only to entities other than Dasein.

(3) According to Heidegger, the existential spatiality of Dasein is characterized most fundamentally **by (e)** what he calls de-severance, a bringing close.

(4) " 'De-severing' amounts to making the farness **vanish(e)**—that is, making the remoteness of something disappear, bringing it close" (Being and Time: 23: 139).

(5) This is of course not a bringing close in the sense of reducing physical distance, although it may involve that. Heidegger's proposal is that spatiality as de-severance is in some way (exactly how is a matter of subtle interpretation; see e.g., Malpas 2006) intimately **related (e&eb) to** the 'reach' of Dasein's skilled practical activity. For example, an entity is 'near by' if it is readily available for some such activity, and 'far away' if it is not, whatever physical distances may be involved.

(6) Given the Dasein-world relationship highlighted above, **the (eb) implication** (drawn explicitly by Heidegger, see Being and Time 22: 136) is that the spatiality distinctive of equipmental entities, and thus of the world, is not equivalent to physical, Cartesian space.

(7) Equipmental space is a matter of pragmatically determined regions of functional places, **defined(e) by** Dasein-centred totalities of involvements (e.g., an office with places for the computers, the

photocopier, and so on—places that are defined by the way in which they make these equipmental entities available in the right sort of way for skilled activity).

(8) For Heidegger, physical, Cartesian space is possible as something meaningful for Dasein **only(e) because** Dasein has de-severance as one of its existential characteristics.

(9) Given the intertwining of de-severance and equipmental space, **this (eb) licenses** the radical view (one that is consistent with Heidegger's prior treatment of Cartesianism) that physical, Cartesian space (as something that we can find intelligible) presupposes equipmental space; the former is the present-at-hand phenomenon that is revealed if we strip away the worldhood from the latter.

NOTATION

Module One

In the wake of his critique of Cartesianism, Heidegger turns his attention to spatiality. He argues that Dasein dwells in the world **in a** spatial manner, but that the spatiality in question—Dasein's existential spatiality—cannot be a matter of Dasein being located at a particular co-ordinate in physical, Cartesian space

G_{13} : Category one of spatial manner, but that the spatiality in question—Dasein's existential spatiality—cannot be a matter of Dasein being located at a particular co-ordinate in physical, Cartesian space

G_{14} : Category two of spatial manner, but that the spatiality in question—Dasein's existential spatiality—cannot be a matter of Dasein being located at a particular co-ordinate in physical, Cartesian space

G_{15} : Category three of spatial manner, but that the spatiality in question—Dasein's existential spatiality—cannot be a matter of Dasein being located at a particular co-ordinate in physical, Cartesian space

T_{13} : Category one of Dasein dwells in the world

T_{14} : Category two of Dasein dwells in the world

T_{15} : Category three of Dasein dwells in the world

Module Two

That would be to conceive of Dasein as present-at-hand, and presence-at-hand **is a** mode of Being that can belong only to entities other than Dasein.

G_{16} : Category one of mode of Being that can belong only to entities other than Dasein.

G_{17} : Category two of mode of Being that can belong only to entities other than Dasein.

G_{18} : Category three of mode of Being that can belong only to entities other than Dasein.

T_{16} : Category one of conceive of Dasein as present-at-hand, and presence-at-hand

T_{17} : Category two of conceive of Dasein as present-at-hand, and presence-at-hand

T_{18} : Category three of conceive of Dasein as present-at-hand, and presence-at-hand

Module three

According to Heidegger, the existential spatiality of Dasein is characterized most fundamentally **by (e)** what he calls de-severance, a bringing close.

G_{20} : Category one of de-severance, a bringing close

G_{21} : Category two of de-severance, a bringing close

G_{22} : Category three of de-severance, a bringing close

T_{20} : Category one of existential spatiality of Dasein is characterized most fundamentally

T_{21} : Category two of existential spatiality of Dasein is characterized most fundamentally

T_{22} : Category three of existential spatiality of Dasein is characterized most fundamentally

Module four

'De-severing' amounts to making the farness **vanish(e)**—that is, making the remoteness of something disappear, bringing it close" (Being and Time: 23: 139).

G_{24} : Category one of making the remoteness of something disappear, bringing it close" (Being and Time: 23: 139).

G_{25} : Category two of making the remoteness of something disappear, bringing it close" (Being and Time: 23: 139).

G_{26} : Category three of making the remoteness of something disappear, bringing it close" (Being and Time: 23: 139).

T_{24} : Category one of 'De-severing' amounts to making the farness

T_{25} : Category two of 'De-severing' amounts to making the farness

T_{26} : Category three of 'De-severing' amounts to making the farness

Module five(both cases discussed and deliberated)

This is of course not a bringing close in the sense of reducing physical distance, although it may involve that. Heidegger's proposal is that spatiality as de-severance is in some way (exactly how is a matter of subtle interpretation; see e.g., Malpas 2006) intimately **related (e&eb) to** the 'reach' of Dasein's skilled practical activity. For example, an entity is 'near by' if it is readily available for some such activity, and 'far away' if it is not, whatever physical distances may be involved

G_{28} : Category one of bringing close in the sense of reducing physical distance, although it may involve that. Heidegger's proposal is that spatiality as de-severance is in some way (exactly how is a matter of subtle interpretation; see e.g., Malpas 2006) intimately

G_{29} : Category two of bringing close in the sense of reducing physical distance, although it may involve that. Heidegger's proposal is that spatiality as de-severance is in some way (exactly how is a matter of subtle interpretation; see e.g., Malpas 2006) intimately

G_{30} : Category three of bringing closes in the sense of reducing physical distance, although it may involve that. Heidegger's proposal is that spatiality as de-severance is in some way (exactly how is a matter of subtle interpretation; see e.g., Malpas 2006) intimately

T_{28} : Category one of 'reach' of Dasein's skilled practical activity. For example, an entity is 'near by' if it is readily available for some such activity, and 'far away' if it is not, whatever physical distances may be involved; bringing close in the sense of reducing physical distance, although it may involve that. Heidegger's proposal is that spatiality as de-severance is in some way (exactly how is a matter of subtle interpretation; see e.g., Malpas 2006) intimately

T_{29} : Category two of 'reach' of Dasein's skilled practical activity. For example, an entity is 'near by' if it is readily available for some such activity, and 'far away' if it is not, whatever physical distances may be involved; bringing close in the sense of reducing physical distance, although it may involve that. Heidegger's

proposal is that spatiality as de-severance is in some way (exactly how is a matter of subtle interpretation; see e.g., Malpas 2006) intimately

T_{30} : Category three of 'reach' of Dasein's skilled practical activity. For example, an entity is 'near by' if it is readily available for some such activity, and 'far away' if it is not, whatever physical distances may be involved ;bringing close in the sense of reducing physical distance, although it may involve that. Heidegger's proposal is that spatiality as de-severance is in some way (exactly how is a matter of subtle interpretation; see e.g., Malpas 2006) intimately

Module six

Given the Dasein-world relationship highlighted above, **the (eb) implication** (drawn explicitly by Heidegger, see Being and Time 22: 136) is that the spatiality distinctive of equipmental entities, and thus of the world, is not equivalent to physical, Cartesian space

G_{32} : Category one of Dasein-world relationship highlighted above

G_{33} : Category two of Dasein-world relationship highlighted above

G_{34} : Category three of Dasein-world relationship highlighted above

T_{32} : Category one of(drawn explicitly by Heidegger, see Being and Time 22: 136) is that the spatiality distinctive of equipmental entities, and thus of the world, is not equivalent to physical, Cartesian space

T_{33} : Category two of(drawn explicitly by Heidegger, see Being and Time 22: 136) is that the spatiality distinctive of equipmental entities, and thus of the world, is not equivalent to physical, Cartesian space

T_{34} : Category three of (drawn explicitly by Heidegger, see Being and Time 22: 136) is that the spatiality distinctive of equipmental entities, and thus of the world, is not equivalent to physical, Cartesian space

Module seven

Equipmental space is a matter of pragmatically determined regions of functional places, **defined(e) by** Dasein-centred totalities of involvements (e.g., an office with places for the computers, the photocopier, and so on—places that are defined by the way in which they make these equipmental entities available in the right sort of way for skilled activity).

G_{36} : Category one of Dasein-centred totalities of involvements (e.g., an office with places for the computers, the photocopier, and so on—places that are defined by the way in which they make these equipmental entities available in the right sort of way for skilled activity).

G_{37} : Category two of Dasein-centred totalities of involvements (e.g., an office with places for the computers, the photocopier, and so on—places that are defined by the way in which they make these equipmental entities available in the right sort of way for skilled activity).

G_{38} : Category three of Dasein-centred totalities of involvements (e.g., an office with places for the computers, the photocopier, and so on—places that are defined by the way in which they make these equipmental entities available in the right sort of way for skilled activity).

T_{36} : Category one of Equipmental space is a matter of pragmatically determined regions of functional places

T_{37} : Category two of Equipmental space is a matter of pragmatically determined regions of functional

places

T_{38} : Category three of Equipmental space is a matter of pragmatically determined regions of functional places

Module eight

For Heidegger, physical, Cartesian space is possible as something meaningful for Dasein **only(e) because** Dasein has de-severance as one of its existential characteristics

G_{40} : Category one of Dasein has de-severance as one of its existential characteristics

G_{41} : Category two of Dasein has de-severance as one of its existential characteristics

G_{42} : Category three of Dasein has de-severance as one of its existential characteristics

T_{40} : Category one of physical, Cartesian space is possible as something meaningful for Dasein

T_{41} : Category two of physical, Cartesian space is possible as something meaningful for Dasein

T_{42} : Category three of physical, Cartesian space is possible as something meaningful for Dasein

Module Nine

Given the intertwining of de-severance and equipmental space, **this (eb) licenses** the radical view (one that is consistent with Heidegger's prior treatment of Cartesianism) that physical, Cartesian space (as something that we can find intelligible) presupposes equipmental space; the former is the present-at-hand phenomenon that is revealed if we strip away the worldhood from the latter

G_{44} : Category one of Given the intertwining of de-severance and equipmental space

G_{45} : Category two of Given the intertwining of de-severance and equipmental space

G_{46} : Category three of Given the intertwining of de-severance and equipmental space

T_{44} : Category one of **licenses** the radical view (one that is consistent with Heidegger's prior treatment of Cartesianism) that physical, Cartesian space (as something that we can find intelligible) presupposes equipmental space; the former is the present-at-hand phenomenon that is revealed if we strip away the worldhood from the latter

T_{45} : Category two of **licenses** the radical view (one that is consistent with Heidegger's prior treatment of Cartesianism) that physical, Cartesian space (as something that we can find intelligible) presupposes equipmental space; the former is the present-at-hand phenomenon that is revealed if we strip away the worldhood from the latter

T_{46} : Category three of **licenses** the radical view (one that is consistent with Heidegger's prior treatment of Cartesianism) that physical, Cartesian space (as something that we can find intelligible) presupposes equipmental space; the former is the present-at-hand phenomenon that is revealed if we strip away the worldhood from the latter

SECTION EIGHTY NINE

Objective, Intersubjective And Subjective Dimensions: Altagsgeschichte: Micro History

INTRODUCTION—VARIABLES USED

Ideas: General Introduction to Pure Phenomenology. Trans. W.R. Boyce Gibson. Collier/Macmillan. [first translation 1931] Original German text – 1913.

(1) Malpas (forthcoming) rejects the account of spatiality given in Being and Time. Drawing on Kant, he argues that "[any] agent, insofar as it is capable of action at all (that is, insofar as it is, indeed, an agent), acts in a space that is an objective space, in which other agents also act, and yet which is always immediately**(arrange, build, compose, construct, design, form, pattern, shape) configured(eb)** subjectively in terms of the agent's own oriented locatedness" (Malpas forthcoming, 14).

(2) According to Malpas, then, equipmental space (a space ordered in terms of practical activity and within which an agent acts) **presupposes (eb) a** more fundamental notion of space as a complex unity with objective, intersubjective and subjective dimensions.

(3) If this is right, then of course equipmental space cannot itself explain the spatial. A further problem, as Malpas also notes, is that the whole issue of spatiality **brings into (eb) sharp focus** the awkward relationship that Heidegger has with the body in Being and Time.

(4) In what is now a frequently quoted remark, Heidegger sets aside Dasein's embodiment, commenting that "this 'bodily nature' **hides(e)** a whole problematic of its own, though we shall not treat it here" (Being and Time 23: 143).

(5) Indeed, at times, Heidegger might be interpreted as linking embodiment **with (e&eb)** Thing hood.

(6) For example: "[as] Dasein goes along its ways, it does not measure off a stretch of space as a corporeal Thing which is present-at-hand" (Being and Time 23: 140). Here one might plausibly contain the spread of presence-at-hand by appealing to a distinction between material (present-at-hand) and lived (existential) ways in which Dasein is embodied. Unfortunately this distinction isn't

made in Being and Time (a point noted by Ricouer 1992, 327), although Heidegger does adopt it in the much later Seminar in Le Thor (see Malpas forthcoming, 5). What seems clear, however, is that while the Heidegger of Being and Time seems to hold that Dasein's embodiment **somehow(e&eb) depends on** its existential spatiality (see e.g., 23: 143), the more obvious thing to say is that Dasein's existential spatiality somehow depends on its embodiment.

(7) Before leaving this issue, it is worth noting briefly that space reappears later in Being and Time (70: 418–21), where Heidegger argues that existential space **is derived (e) from** temporality.

(8) This makes sense within Heidegger's overall project, because, as we shall see, the deep structure of totalities of involvements (and thus of equipmental space) is finally understood **in terms (e&eb) of** temporality.

(9) Nevertheless, and although the distinctive character of Heidegger's concept of temporality needs to be recognized, there is reason to think that the dependency here may well travel in the opposite direction. The worry, as Malpas (forthcoming, 26) again points out, has a Kantian origin. Kant (1781/1999) argued that the temporal character of inner sense is possible only **because (e) i**t is mediated by outer intuition whose form is space. If this is right, and if we can generalize appropriately, then the temporality that matters to Heidegger will be dependent on existential spatiality, and not the other way round. All in all, one is tempted to conclude that Heidegger's treatment of spatiality in Being and Time, and (relatedly) his treatment (or lack of it) of the body, face serious difficulties.

NOTATION

Module One

Malpas (forthcoming) rejects the account of spatiality given in Being and Time. Drawing on Kant, he argues that "[any] agent, insofar as it is capable of action at all (that is, insofar as it is, indeed, an agent), acts in a space that is an objective space, in which other agents also act, and yet which is always immediately**(arrange, build, compose, construct, design, form, pattern, shape) configured(eb)** subjectively in terms of the agent's own oriented locatedness" (Malpas forthcoming, 14).

G_{13} : Category one of the account of spatiality given in Being and Time. Drawing on Kant, he argues that "[any] agent, insofar as it is capable of action at all (that is, insofar as it is, indeed, an agent), acts in a space that is an objective space, in which other agents also act, and yet which is always immediately**(arrange, build, compose, construct, design, form, pattern, shape)**

G_{14} : Category two of the account of spatiality given in Being and Time. Drawing on Kant, he argues that "[any] agent, insofar as it is capable of action at all (that is, insofar as it is, indeed, an agent), acts in a space that is an objective space, in which other agents also act, and yet which is always immediately**(arrange, build, compose, construct, design, form, pattern, shape)**

G_{15} : Category three of the account of spatiality given in Being and Time. Drawing on Kant, he argues that "[any] agent, insofar as it is capable of action at all (that is, insofar as it is, indeed, an agent), acts in a space that is an objective space, in which other agents also act, and yet which is always immediately**(arrange, build, compose, construct, design, form, pattern, shape)**

T_{13} : Category one of subjectively in terms of the agent's own oriented locatedness"

T_{14} : Category two of subjectively in terms of the agent's own oriented locatedness"

T_{15} : Category three of subjectively in terms of the agent's own oriented locatedness"

Module Two

According to Malpas, then, equipmental space (a space ordered in terms of practical activity and within which an agent acts) **presupposes (eb) a** more fundamental notion of space as a complex unity with

objective, intersubjective and subjective dimensions

G_{16} : Category one of equipmental space (a space ordered in terms of practical activity and within which an agent acts)

G_{17} : Category two of equipmental space (a space ordered in terms of practical activity and within which an agent acts)

G_{18} : Category three of equipmental space (a space ordered in terms of practical activity and within which an agent acts)

T_{16} : Category one of more fundamental notion of space as a complex unity with objective, intersubjective and subjective dimensions

T_{17} : Category two of more fundamental notion of space as a complex unity with objective, intersubjective and subjective dimensions

T_{18} : Category three of more fundamental notion of space as a complex unity with objective, intersubjective and subjective dimensions

Module three

If this is right, then of course equipmental space cannot itself explain the spatial. A further problem, as Malpas also notes, is that the whole issue of spatiality **brings into (eb) sharp focus** the awkward relationship that Heidegger has with the body in Being and Time.

G_{20} : Category one of If this is right, and then of course equipmental space cannot itself explain the spatial. A further problem, as Malpas also notes, is that the whole issue of spatiality

G_{21} : Category two of If this is right, and then of course equipmental space cannot itself explain the spatial. A further problem, as Malpas also notes, is that the whole issue of spatiality

G_{22} : Category three of If this is right, and then of course equipmental space cannot itself explain the spatial. A further problem, as Malpas also notes, is that the whole issue of spatiality

T_{20} : Category one of **sharp focus** the awkward relationship that Heidegger has with the body in Being and Time.

T_{21} : Category two of **sharp focus** the awkward relationship that Heidegger has with the body in Being and Time.

T_{22} : Category three of **sharp focus** the awkward relationship that Heidegger has with the body in Being and Time.

Module four

In what is now a frequently quoted remark, Heidegger sets aside Dasein's embodiment, commenting that "this 'bodily nature' **hides(e)** a whole problematic of its own, though we shall not treat it here" (Being and Time 23: 143).

G_{24} : Category one of whole problematic of its own, though we shall not treat it here" (Being and Time 23: 143).

G_{25} : Category two of whole problematic of its own, though we shall not treat it here" (Being and Time 23: 143).

G_{26} : Category three of whole problematic of its own, though we shall not treat it here" (Being and Time 23: 143).

T_{24} : Category one of Heidegger sets aside Dasein's embodiment, commenting that "this 'bodily nature'

T_{25} : Category two of Heidegger sets aside Dasein's embodiment, commenting that “this ‘bodily nature’

T_{26} : Category three of Heidegger sets aside Dasein's embodiment, commenting that “this ‘bodily nature’

Module five

Indeed, at times, Heidegger might be interpreted as linking embodiment **with (e&eb)** Thing hood

G_{28} : Category one of embodiment; Thing hood

G_{29} : Category two of embodiment; Thing hood

G_{30} : Category three of embodiment; Thing hood

T_{28} : Category one of Thing hood; embodiment;

T_{29} : Category two of Thing hood; embodiment;

T_{30} : Category three of Thing hood; embodiment;

Module six

For example: “[as] Dasein goes along its ways, it does not measure off a stretch of space as a corporeal Thing which is present-at-hand” (Being and Time 23: 140). Here one might plausibly contain the spread of presence-at-hand by appealing to a distinction between material (present-at-hand) and lived (existential) ways in which Dasein is embodied. Unfortunately this distinction isn't made in Being and Time (a point noted by Ricouer 1992, 327), although Heidegger does adopt it in the much later Seminar in Le Thor (see Malpas forthcoming, 5). What seems clear, however, is that while the Heidegger of Being and Time seems to hold that Dasein's embodiment **somehow(e&eb) depends on** its existential spatiality (see e.g., 23: 143), the more obvious thing to say is that Dasein's existential spatiality somehow depends on its embodiment

G_{32} : Category one of What seems clear, however, is that while the Heidegger of Being and Time seems to hold that Dasein's embodiment; existential spatiality (see e.g., 23: 143), the more obvious thing to say is that Dasein's existential spatiality somehow depends on its embodiment

G_{33} : Category two of What seems clear, however, is that while the Heidegger of Being and Time seems to hold that Dasein's embodiment; existential spatiality (see e.g., 23: 143), the more obvious thing to say is that Dasein's existential spatiality somehow depends on its embodiment

G_{34} : Category three of What seems clear, however, is that while the Heidegger of Being and Time seems to hold that Dasein's embodiment; existential spatiality (see e.g., 23: 143), the more obvious thing to say is that Dasein's existential spatiality somehow depends on its embodiment

T_{32} : Category one of What seems clear, however, is that while the Heidegger of Being and Time seems to hold that Dasein's embodiment; existential spatiality (see e.g., 23: 143), the more obvious thing to say is that Dasein's existential spatiality somehow depends on its embodiment

T_{33} : Category two of What seems clear, however, is that while the Heidegger of Being and Time seems to hold that Dasein's embodiment; existential spatiality (see e.g., 23: 143), the more obvious thing to say is that Dasein's existential spatiality somehow depends on its embodiment

T_{34} : Category three of What seems clear, however, is that while the Heidegger of Being and Time seems to

hold that Dasein's embodiment; existential spatiality (see e.g., 23: 143), the more obvious thing to say is that Dasein's existential spatiality somehow depends on its embodiment

Module seven

Before leaving this issue, it is worth noting briefly that space reappears later in Being and Time (70: 418–21), where Heidegger argues that existential space **is derived (e) from** temporality

G_{36} : Category one of temporality

G_{37} : Category two of temporality

G_{38} : Category three of temporality

T_{36} : Category one of space reappears later in Being and Time (70: 418–21), where Heidegger argues that existential space

T_{37} : Category two of space reappears later in Being and Time (70: 418–21), where Heidegger argues that existential space

T_{38} : Category three of space reappears later in Being and Time (70: 418–21), where Heidegger argues that existential space

Module eight

This makes sense within Heidegger's overall project, because, as we shall see, the deep structure of totalities of involvements (and thus of equipmental space) is finally understood **in terms (e&eb) of** temporality.
G_{40} : Category one of deep structure of totalities of involvements (and thus of equipmental space) is finally understood

G_{41} : Category two of deep structure of totalities of involvements (and thus of equipmental space) is finally understood

G_{42} : Category three of deep structure of totalities of involvements (and thus of equipmental space) is finally understood

T_{40} : Category one of temporality; deep structure of totalities of involvements (and thus of equipmental space) is finally understood

T_{41} : Category two of temporality; deep structure of totalities of involvements (and thus of equipmental space) is finally understood

T_{42} : Category three of temporality; deep structure of totalities of involvements (and thus of equipmental space) is finally understood

Module Nine

Nevertheless, and although the distinctive character of Heidegger's concept of temporality needs to be recognized, there is reason to think that the dependency here may well travel in the opposite direction. The worry, as Malpas (forthcoming, 26) again points out, has a Kantian origin. Kant (1781/1999) argued that the temporal character of inner sense is possible only **because (e) i**t is mediated by outer intuition whose form is space. If this is right, and if we can generalize appropriately, then the temporality that matters to Heidegger

will be dependent on existential spatiality, and not the other way round. All in all, one is tempted to conclude that Heidegger's treatment of spatiality in Being and Time, and (relatedly) his treatment (or lack of it) of the body, face serious difficulties.

G_{44} : Category one of mediated by outer intuition whose form is space. If this is right, and if we can generalize appropriately, then the temporality that matters to Heidegger will be dependent on existential spatiality, and not the other way round. All in all, one is tempted to conclude that Heidegger's treatment of spatiality in Being and Time, and (relatedly) his treatment (or lack of it) of the body, face serious difficulties.

G_{45} : Category two of mediated by outer intuition whose form is space. If this is right, and if we can generalize appropriately, then the temporality that matters to Heidegger will be dependent on existential spatiality, and not the other way round. All in all, one is tempted to conclude that Heidegger's treatment of spatiality in Being and Time, and (relatedly) his treatment (or lack of it) of the body, face serious difficulties.

G_{46} : Category three of mediated by outer intuition whose form is space. If this is right, and if we can generalize appropriately, then the temporality that matters to Heidegger will be dependent on existential spatiality, and not the other way round. All in all, one is tempted to conclude that Heidegger's treatment of spatiality in Being and Time, and (relatedly) his treatment (or lack of it) of the body, face serious difficulties.

T_{44} : Category one of temporal character of inner sense is possible only

T_{45} : Category two of temporal character of inner sense is possible only

T_{46} : Category three of temporal character of inner sense is possible only

SECTION NINETY

Living Lives At The Limits Of Itself…..:Fingerspitzengefuhl: Intuitive-Instinctive

INTRODUCTION—VARIABLES USED

Ideas: General Introduction to Pure Phenomenology. Trans. W.R. Boyce Gibson. Collier/Macmillan. [first translation 1931] Original German text – 1913.

Being-with

(1) Heidegger turns next to the question of "who it is that Dasein is in its everydayness" (Being and Time, Introduction to IV: 149). He rejects the idea of Dasein as a Cartesian 'I-thing' (the Cartesian thinking thing conceived as a substance), since once again this **would (=) be to think** of Dasein as present-at-hand. In searching for an alternative answer, Heidegger observes that equipment is often revealed to us as being for the sake of (the lives and projects of) other Dasein.
(2) The boat anchored at the shore is assigned in its Being-in-itself to an acquaintance who undertakes voyages with it; but even if it is a 'boat which is strange to us', it still **is (eb) indicative** of Others.
(3) The Others who are thus 'encountered' in a ready-to-hand, environmental context of equipment, are

not somehow **added(eb) on** in thought to some Thing which is proximally just present-at-hand; such 'Things' are encountered from out of a world in which they are ready-to-hand for Others—a world which is always mine too in advance. (Being and Time 26: 154)

(4) On the basis of such observations, Heidegger argues that to be Dasein at all **means(=)** to Be-with:

(5) "So far as Dasein is at all, **it (e) has** Being-with-one-another as its kind of Being" (Being and Time 26: 163). One's immediate response to this might be that it is just false. After all, ordinary experience establishes that each of us is often alone. But of course Heidegger is thinking in an **ontological register**.

(6) Being-with (Mitsein) is thus the a priori transcendental condition **that makes (eb) it possible** that Dasein can discover equipment in this Other-related fashion.

(7) And it's because Dasein has Being-with as one of its essential modes of Being that everyday Dasein can experience being alone. Being-with **is thus (=) the** a priori transcendental condition for loneliness.

(8) It is important to understand what Heidegger means by 'Others', a term that he uses interchangeably with the more evocative 'the "they" ' (das Man). He explains: By 'Others' we do not mean everyone else **but (=)** me—those over against whom the 'I' stands out. They are rather those from whom, for the most part, one does not distinguish oneself—those among whom one is too…

(9) By reason of this with-like Being-in-the-world, the world is always the one that I **share (e&eb)** with others. (Being and Time 26: 154–5)

NOTATION

Module One

Heidegger turns next to the question of "who it is that Dasein is in its everydayness" (Being and Time, Introduction to IV: 149). He rejects the idea of Dasein as a Cartesian 'I-thing' (the Cartesian thinking thing conceived as a substance), since once again this **would (=) be to think** of Dasein as present-at-hand. In searching for an alternative answer, Heidegger observes that equipment is often revealed to us as being for the sake of (the lives and projects of) other Dasein

G_{13} : Category one of **Dasein** as present-at-hand. In searching for an alternative answer, Heidegger observes that equipment is often revealed to us as being for the sake of (the lives and projects of) other Dasein

G_{14} : Category two of Dasein as present-at-hand. In searching for an alternative answer, Heidegger observes that equipment is often revealed to us as being for the sake of (the lives and projects of) other Dasein

G_{15} : Category three of Dasein as present-at-hand. In searching for an alternative answer, Heidegger observes that equipment is often revealed to us as being for the sake of (the lives and projects of) other Dasein

T_{13} : Category one of Dasein as a Cartesian 'I-thing' (the Cartesian thinking thing conceived as a substance), since once again this

T_{14} : Category two of Dasein as a Cartesian 'I-thing' (the Cartesian thinking thing conceived as a substance), since once again this

T_{15} : Category three of Dasein as a Cartesian 'I-thing' (the Cartesian thinking thing conceived as a substance), since once again this

Module Two

The boat anchored at the shore is assigned in its Being-in-itself to an acquaintance who undertakes voyages with it; but even if it is a 'boat which is strange to us', it still **is (eb) indicative** of Others

G_{16} : Category one of boat anchored at the shore is assigned in its Being-in-itself to an acquaintance who undertakes voyages with it; but even if it is a 'boat which is strange to us', it still

G_{17} : Category two of boat anchored at the shore is assigned in its Being-in-itself to an acquaintance who undertakes voyages with it; but even if it is a 'boat which is strange to us', it still

G_{18} : Category three of boat anchored at the shore is assigned in its Being-in-itself to an acquaintance who undertakes voyages with it; but even if it is a 'boat which is strange to us', it still

T_{16} : Category one of **indicative** of Others

T_{17} : Category two of **indicative** of Others

T_{18} : Category three of **indicative** of Others

Module three

The Others who are thus 'encountered' in a ready-to-hand, environmental context of equipment, are not somehow **added(eb) on** in thought to some Thing which is proximally just present-at-hand; such 'Things' are encountered from out of a world in which they are ready-to-hand for Others—a world which is always mine too in advance. (Being and Time 26: 154)

G_{20} : Category one of Others who are thus 'encountered' in a ready-to-hand, environmental context of equipment, are not somehow

G_{21} : Category two of Others who are thus 'encountered' in a ready-to-hand, environmental context of equipment, are not somehow

G_{22} : Category three of Others who are thus 'encountered' in a ready-to-hand, environmental context of equipment, are not somehow

T_{20} : Category one of thought to some Thing which is proximally just present-at-hand; such 'Things' are encountered from out of a world in which they are ready-to-hand for Others—a world which is always mine too in advance. (Being and Time 26: 154)

T_{21} : Category two of thought to some Thing which is proximally just present-at-hand; such 'Things' are encountered from out of a world in which they are ready-to-hand for Others—a world which is always mine too in advance. (Being and Time 26: 154)

T_{22} : Category three of thought to some Thing which is proximally just present-at-hand; such 'Things' are encountered from out of a world in which they are ready-to-hand for Others—a world which is always mine too in advance. (Being and Time 26: 154)

Module four

On the basis of such observations, Heidegger argues that to be Dasein at all **means** to Be-with

G_{24} : Category one of to Be-with

G_{25} : Category two of to Be-with

G_{26} : Category three of to Be-with

T_{24} : Category one of Dasein

T_{25} : Category two of Dasein

T_{26} : Category three of Dasein

Module five

"So far as Dasein is at all, **it (e) has** Being-with-one-another as its kind of Being" (Being and Time 26: 163). One's immediate response to this might be that it is just false. After all, ordinary experience establishes that each of us is often alone. But of course Heidegger is thinking in an **ontological register**

G_{28} : Category one of Being-with-one-another as its kind of Being" (Being and Time 26: 163). One's immediate response to this might be that it is just false. After all, ordinary experience establishes that each of us is often alone. But of course Heidegger is thinking in an **ontological register**

G_{29} : Category two of Being-with-one-another as its kind of Being" (Being and Time 26: 163). One's immediate response to this might be that it is just false. After all, ordinary experience establishes that each of us is often alone. But of course Heidegger is thinking in an **ontological register**

G_{30} : Category three of Being-with-one-another as its kind of Being" (Being and Time 26: 163). One's immediate response to this might be that it is just false. After all, ordinary experience establishes that each of us is often alone. But of course Heidegger is thinking in an **ontological register**

T_{28} : Category one of Dasein is at all

T_{29} : Category two of Dasein is at all

T_{30} : Category three of Dasein is at all

Module six

Being-with (Mitsein) is thus the a priori transcendental condition **that makes (eb) it possible** that Dasein can discover equipment in this Other-related fashion

G_{32} : Category one of Being-with (Mitsein) is thus the a priori transcendental condition

G_{33} : Category two of Being-with (Mitsein) is thus the a priori transcendental condition

G_{34} : Category three of Being-with (Mitsein) is thus the a priori transcendental condition

T_{32} : Category one of Dasein can discover equipment in this Other-related fashion

T_{33} : Category two of Dasein can discover equipment in this Other-related fashion

T_{34} : Category three of Dasein can discover equipment in this Other-related fashion

Module seven

And it's because Dasein has Being-with as one of its essential modes of Being that everyday Dasein can experience being alone. Being-with **is thus the** a priori transcendental condition for loneliness.

G_{36} : Category one of a priori transcendental condition for loneliness

G_{37} : Category two of a priori transcendental condition for loneliness

G_{38} : Category three of a priori transcendental condition for loneliness

T_{36} : Category one of Dasein has Being-with as one of its essential modes of Being that everyday Dasein can experience being alone. Being-with

T_{37} : Category two of Dasein has Being-with as one of its essential modes of Being that everyday Dasein can experience being alone. Being-with

T_{38} : Category three of Dasein has Being-with as one of its essential modes of Being that everyday Dasein can experience being alone. Being-with

Module eight

It is important to understand what Heidegger means by 'Others', a term that he uses interchangeably with the more evocative 'the "they" ' (das Man). He explains: By 'Others' we do not mean everyone else **but** me—those over against whom the 'I' stands out. They are rather those from whom, for the most part, one does not distinguish oneself—those among whom one is too…

G_{40} : Category one of me—those over against whom the 'I' stands out. They are rather those from whom, for the most part, one does not distinguish oneself—those among whom one is too…

G_{41} : Category two of me—those over against whom the 'I' stands out. They are rather those from whom, for the most part, one does not distinguish oneself—those among whom one is too…

G_{42} : Category three of me—those over against whom the 'I' stands out. They are rather those from whom, for the most part, one does not distinguish oneself—those among whom one is too…

T_{40} : Category one of' Others' we do not mean everyone else

T_{41} : Category two of 'Others' we do not mean everyone else

T_{42} : Category three of 'Others' we do not mean everyone else

Module Nine

By reason of this with-like Being-in-the-world, the world is always the one that I **share (e&eb)** with others. (Being and Time 26: 154–5)

G_{44} : Category one of By reason of this with-like Being-in-the-world, the world is always the one that

G_{45} : Category two of By reason of this with-like Being-in-the-world, the world is always the one that

G_{46} : Category three of By reason of this with-like Being-in-the-world, the world is always the one that

T_{44} : Category one of with others. (Being and Time 26: 154–5); By reason of this with-like Being-in-the-world, the world is always the one that

T_{45} : Category two of with others. (Being and Time 26: 154–5); By reason of this with-like Being-in-the-world, the world is always the one that

T_{46} : Category three of with others. (Being and Time 26: 154–5);By reason of this with-like Being-in-the-world, the world is always the one that

SECTION NINETY ONE

Worm Holes And Time Travel: Freiwirtschaft: Natural Order

INTRODUCTION—VARIABLES USED

Wormholes and Time Travel Shortcut through Spacetime General Relativity: Wormholes and Time Travel Shortcut through Spacetime - If Traversable Wormholes Exist, They Could Allow Time Travel

(1) Physicists have also not found any natural process which would be predicted **to form a** wormhole naturally in the context of general relativity, although the quantum foam hypothesis **is sometimes used to (eb) suggest** that tiny wormholes might appear and disappear spontaneously at the Planck scale.
(2) It has also been proposed that if a tiny wormhole held open by a negative-mass cosmic string had appeared around the time of the Big Bang, it could have been **inflated (eb**) to macroscopic size by cosmic inflation.

The American theoretical physicist John Archibald Wheeler coined the term wormhole in 1957; however, in 1921, the German mathematician Hermann Weyl already had proposed the wormhole theory, in connection with mass analysis of electromagnetic field energy. Unknown X

Wormholes, Time Machines, and the Weak Energy Condition Michael S. Morris et al

(1) It is argued that, if the laws of physics permit an advanced civilization **to create and maintain** a wormhole in space for interstellar travel, then that wormhole can be converted into a time machine with which causality might be viola table. Whether wormholes can be created and maintained entails deep, ill-understood issues about cosmic censorship, quantum gravity, and quantum field theory, including the question of whether field theory enforces an averaged version of the weak energy condition.© 1988 The American Physical SocietyURL: http://link.aps.org/doi/10.1103/PhysRevLett.61.1446DOI: 10.1103/PhysRevLett.61.1446

Quantum mechanics of time travel through post-selected teleportation Seth Lloyd

(2) This paper discusses the quantum mechanics of closed-timelike curves (CTCs) and of other potential methods **for (e)** time travel.
(3) Authors analyze a specific proposal for such quantum time travel, the quantum description of CTCs **based on (e) post**-selected teleportation (P-CTCs).

(4) They compare the theory of P-CTCs to previously proposed quantum theories of time travel: the theory is **inequivalent to** Deutsch's theory of CTCs, but it is consistent with path-integral approaches (which are the best suited for analyzing quantum-field theory in curved space-time).

(5) They derive the dynamical equations that a chronology-respecting system **interacting with** a CTC will experience.

(6) Authors also discuss the possibility of time travel **in the (e)absence of** general-relativistic closed-timelike curves, and investigate the implications of P-CTCs for enhancing the power of computation.© 2011 American Physical SocietyURL:http://link.aps.org/doi/10.1103/PhysRevD.84.025007DOI:10.1103/PhysRevD.84.025007

NOTATION

Module One

Physicists have also not found any natural process which would be predicted **to form a** wormhole naturally in the context of general relativity, although the quantum foam hypothesis **is sometimes used to (eb) suggest** that tiny wormholes might appear and disappear spontaneously at the Planck scale.

G_{13} : Category one of quantum foam hypothesis **is sometimes used**

G_{14} : Category two of quantum foam hypothesis **is sometimes used**

G_{15} : Category three ofquantum foam hypothesis **is sometimes used**

T_{13} : Category one of tiny wormholes might appear and disappear spontaneously at the Planck scale.

T_{14} : Category two of tiny wormholes might appear and disappear spontaneously at the Planck scale.

T_{15} : Category three of tiny wormholes might appear and disappear spontaneously at the Planck scale.

Module Two

It has also been proposed that if a tiny wormhole held open by a negative-mass cosmic string had appeared around the time of the Big Bang, it could have been **inflated (eb**) to macroscopic size by cosmic inflation

G_{16} : Category one of tiny wormhole held open by a negative-mass cosmic string had appeared around the time of the Big Bang, it could have been

G_{17} : Category two of tiny wormhole held open by a negative-mass cosmic string had appeared around the time of the Big Bang, it could have been

G_{18} : Category three of tiny wormhole held open by a negative-mass cosmic string had appeared around the time of the Big Bang, it could have been

T_{16} : Category one of macroscopic size by cosmic inflation

T_{17} : Category two of macroscopic size by cosmic inflation

T_{18} : Category three of macroscopic size by cosmic inflation

Module three

It is argued that, if the laws of physics permit an advanced civilization **to create and maintain** a wormhole in space for interstellar travel, then that wormhole can be converted into a time machine with which

causality might be **viola table**. Whether wormholes can be created and maintained entails deep, ill-understood issues about cosmic censorship, quantum gravity, and quantum field theory, including the question of whether field theory enforces an averaged version of the weak energy condition.© 1988 The American Physical SocietyURL: http://link.aps.org/doi/10.1103/PhysRevLett.61.1446DOI: 10.1103/PhysRevLett.61.1446

G_{20} : Category one of principle of causality

G_{21} : Category two of principle of causality

G_{22} : Category three of principle of causality

T_{20} : Category one of **creation and maintenance** a wormhole in space for interstellar travel, then that wormhole can be converted into a time machine

T_{21} : Category two of **creation and maintenance** a wormhole in space for interstellar travel, then that wormhole can be converted into a time machine

T_{22} : Category three of **creation and maintenance** a wormhole in space for interstellar travel, then that wormhole can be converted into a time machine

Module four

This paper discusses the quantum mechanics of closed-timelike curves (CTCs) and of other potential methods **for (e)** time travel

G_{24} : Category one of time travel

G_{25} : Category two of time travel

G_{26} : Category three of time travel

T_{24} : Category one of quantum mechanics of closed-timelike curves (CTCs) and of other potential methods

T_{25} : Category two of quantum mechanics of closed-timelike curves (CTCs) and of other potential methods

T_{26} : Category three of quantum mechanics of closed-timelike curves (CTCs) and of other potential methods

Module five

Authors analyze a specific proposal for such quantum time travel, the quantum description of CTCs **based on (e) post**-selected teleportation (P-CTCs).

G_{28} : Category one of **post**-selected teleportation (P-CTCs).

G_{29} : Category two of **post**-selected teleportation (P-CTCs).

G_{30} : Category three of **post**-selected teleportation (P-CTCs).

T_{28} : Category one of specific proposal for such quantum time travel, the quantum description of CTCs

T_{29} : Category two of specific proposal for such quantum time travel, the quantum description of CTCs

T_{30} : Category three of specific proposal for such quantum time travel, the quantum description of CTCs

Module six

They compare the theory of P-CTCs to previously proposed quantum theories of time travel: the theory is **inequivalent to** Deutsch's theory of CTCs, but it is consistent with path-integral approaches (which are the best suited for analyzing quantum-field theory in curved space-time).

G_{32} : Category one of P-CTCs to previously proposed quantum theories of time travel

G_{33} : Category two of P-CTCs to previously proposed quantum theories of time travel

G_{34} : Category three of P-CTCs to previously proposed quantum theories of time travel

T_{32} : Category one of Deutsch's theory of CTCs, but it is consistent with path-integral approaches (which are the best suited for analyzing quantum-field theory in curved space-time).

T_{33} : Category two of Deutsch's theory of CTCs, but it is consistent with path-integral approaches (which are the best suited for analyzing quantum-field theory in curved space-time).

T_{34} : Category three of Deutsch's theory of CTCs, but it is consistent with path-integral approaches (which are the best suited for analyzing quantum-field theory in curved space-time).

Module seven

They derive the dynamical equations that a chronology-respecting system **interacting with** a CTC will experience

G_{36} : Category one of chronology-respecting system; CTC

G_{37} : Category two of chronology-respecting system; CTC

G_{38} : Category three of chronology-respecting system; CTC

T_{36} : Category one of CTC ;chronology-respecting system

T_{37} : Category two of CTC; chronology-respecting system

T_{38} : Category three of CTC; chronology-respecting system

Module eight

Authors also discuss the possibility of time travel **in the (e)absence of** general-relativistic closed-timelike curves, and investigate the implications of P-CTCs for enhancing the power of computation

G_{40} : Category one of general-relativistic closed-timelike curves

G_{41} : Category two of general-relativistic closed-timelike curves

G_{42} : Category three of general-relativistic closed-timelike curves

T_{40} : Category one of possibility of time travel

T_{41} : Category two of possibility of time travel

T_{42} : Category three of possibility of time travel

Module Nine

G_{44} : Category one of P-CTCs

G_{45} : Category two of P-CTCs

G_{46} : Category three of P-CTCs

T_{44} : Category one of power of computation

T_{45} : Category two of power of computation

T_{46} : Category three of power of computation

References

(1) ^ "Quantum Computing with Molecules" article in Scientific American by Neil Gershenfeld and Isaac L. Chuang
(2) ^ Manin, Yu. I. (1980). Vychislimoe i nevychislimoe [Computable and Noncomputable] (in Russian). Sov.Radio. pp. 13–15. Retrieved 4 March 2013.
(3) ^ Feynman, R. P. (1982). "Simulating physics with computers". International Journal of Theoretical Physics 21 (6): 467–488. doi:10.1007/BF02650179.
(4) ^ Deutsch, David (1992-01-06). "Quantum computation". Physics World.
(5) ^ Finkelstein, David (1969). "Space-Time Structure in High Energy Interactions". In Gudehus, T.; Kaiser, G. Fundamental Interactions at High Energy. New York: Gordon & Breach.
(6) ^ New qubit control bodes well for future of quantum computing
(7) ^ Quantum Information Science and Technology Roadmap for a sense of where the research is heading.
(8) ^ Simon, D.R. (1994). "On the power of quantum computation". Foundations of Computer Science, 1994 Proceedings., 35th Annual Symposium on: 116–123. doi:10.1109/SFCS.1994.365701. ISBN 0-8186-6580-7.
(9) ^ Nielsen, Michael A.; Chuang, Isaac L. Quantum Computation and Quantum Information. p. 202.
(10) ^ Nielsen, Michael A.; Chuang, Isaac L. Quantum Computation and Quantum Information. p. 17.
(11) ^ Waldner, Jean-Baptiste (2007). Nanocomputers and Swarm Intelligence. London: ISTE. p. 157. ISBN 2-7462-1516-0.
(12) ^ a b David P. DiVincenzo (1995). "Quantum Computation". Science 270 (5234): 255–261. Bibcode:1995Sci...270..255D. doi:10.1126/science.270.5234.255. (subscription required)
(13) ^ Arjen K. Lenstra (2000). "Integer Factoring". Designs, Codes and Cryptography 19 (2/3): 101–128. doi:10.1023/A:1008397921377.

(14)^ a b Daniel J. Bernstein, Introduction to Post-Quantum Cryptography. Introduction to Daniel J. Bernstein, Johannes Buchmann, Erik Dahmen (editors). Post-quantum cryptography. Springer, Berlin, 2009. ISBN 978-3-540-88701-0

(15)^ See also pqcrypto.org, a bibliography maintained by Daniel J. Bernstein and Tanja Lange on cryptography not known to be broken by quantum computing.

(16)^ Robert J. McEliece. "A public-key cryptosystem based on algebraic coding theory." Jet Propulsion Laboratory DSN Progress Report 42–44, 114–116.

(17)^ Kobayashi, H.; Gall, F.L. (2006). "Dihedral Hidden Subgroup Problem: A Survey". Information and Media Technologies 1 (1): 178–185.

(18)^ Bennett C.H., Bernstein E., Brassard G., Vazirani U., The strengths and weaknesses of quantum computation. SIAM Journal on Computing 26(5): 1510–1523 (1997).

(19)^ Quantum Algorithm Zoo – Stephen Jordan's Homepage

(20)^ The Father of Quantum Computing By Quinn Norton 02.15.2007, Wired.com

(21)^ David P. DiVincenzo, IBM (2000-04-13). "The Physical Implementation of Quantum Computation". arXiv:quant-ph/0002077 [quant-ph].

(22)^ M. I. Dyakonov, Université Montpellier (2006-10-14). "Is Fault-Tolerant Quantum Computation Really Possible?". In: Future Trends in Microelectronics. Up the Nano Creek, S. Luryi, J. Xu, and A. Zaslavsky (eds), Wiley , pp.: 4–18. arXiv:quant-ph/0610117.

(23)^ Freedman, Michael H.; Kitaev, Alexei; Larsen, Michael J.; Wang, Zhenghan (2003). "Topological quantum computation". Bulletin of the American Mathematical Society 40 (1): 31–38. arXiv:quant-ph/0101025. doi:10.1090/S0273-0979-02-00964-3. MR 1943131.

(24)^ Monroe, Don, Anyons: The breakthrough quantum computing needs?, New Scientist, 1 October 2008

(25)^ Das, A.; Chakrabarti, B. K. (2008), "Quantum Annealing and Analog Quantum Computation", Rev. Mod. Phys. 80 (3): 1061–1081, doi:10.1103/RevModPhys.80.1061